NOAA Technical Memorandum NMFS-SEFC-117

SYNOPSIS OF DATA ON THE IMPACT OF HABITAT ALTERATION ON SEA TURTLES AROUND THE SOUTHEASTERN UNITED STATES

Linda Coston-Clements and Donald E. Hoss

June, 1983

U. S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
John V. Byrne, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator
for Fisheries

This TM series is used for documentation and timely communication of preliminary results, interim reports, or similar special purpose information. Although the memos are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.

TABLE OF CONTENTS

ABSTRACT..... 1

INTRODUCTION..... 1

LIFE HISTORY..... 4

Nesting..... 4

Developmental Stages..... 6

Foraging and Resting..... 8

Migration and Movements..... 9

IMPACT OF HUMAN ACTIVITIES..... 10

Nesting Success..... 10

Coastal zone development..... 12

Alteration of beach topograpy..... 12
Artificial illumination..... 13
Human activity and noise..... 13
Litter..... 14
Temperature alterations..... 14
Exotic vegetation..... 14

Pollution..... 14

Oil spills..... 14
Pesticides..... 15
Heavy metals..... 15
PCB's..... 15
Thermal effluents..... 15

Military activities..... 15

Ordance..... 15
Lights..... 15
Troops and equipment..... 15
Noise..... 16

Oceanic Survival..... 16

Pollutant discharges..... 18

Oil spills..... 18
Pesticides..... 19
Heavy metals..... 19
Radionuclides..... 19
PCB's..... 19
Domestic discharges..... 19

Energy development..... 19

Power plant intake canals..... 19
Thermal effluent..... 19
Lighted offshore structures..... 19

Dredging and mining..... 20

Alteration of topography..... 20
Channel dredging..... 20
Disposal of spoil..... 20

Fishing activities..... 20

General..... 20
Trawls..... 20
Longlines..... 20
Pound nets..... 20
Gill nets..... 20
"Bleach" and "dynamite fishing"..... 20
Recreational..... 21

Miscellaneous..... 21

Litter and garbage..... 21
Increased runoff..... 21
Military activities..... 21

RECOMMENDATIONS..... 21

ACKNOWLEDGMENTS..... 22

BIBLIOGRAPHY..... 24

Synopsis of Data on the Impact of Habitat Alteration on Sea Turtles Around the Southeastern United States

Linda Coston-Clements and Donald E. Hoss

ABSTRACT

A brief description is given of the life histories of five species of marine turtles and the habitats they utilize. The impact of man's activities on these habitats is discussed with respect to their potential for harmful effects on turtles. Recommendations are made for additional research.

INTRODUCTION

Increasing industrial development, population growth, and recreational activities in marine and estuarine waters are encroaching on the natural habitats of sea turtles. Many areas once used by these animals have been lost or severely altered by man's activities. Construction and maintenance of navigation channels, dredge and fill for land development, sea bed mining, and siting and operations of industrial plants are just a few of the activities that contribute to the loss or degradation of habitat. Man's activities may impact on all stages of sea turtle life history (Fig. 1). Although it is obvious that these activities are changing and destroying habitat, there is at present no comprehensive listing of the types of activities that are most damaging to the various species. There is, however, a need for this information in order to develop plans for quantifying and assessing the impacts of specific alterations (e.g., oil spills) and for developing recovery plans for these animals.

The Endangered Species Act of 1973 requires the National Oceanic and Atmospheric Administration (NOAA) to conserve and protect endangered species. The Southeast Fisheries Center (SEFC) is responsible for conducting research to provide information to NOAA that is necessary for management of endangered and threatened species in the Western Atlantic and Gulf of Mexico.

Our objectives were to develop a comprehensive list of human activities that are known or suspected to impact on sea turtles and to prepare a bibliography of pertinent literature. This report deals specifically with human activities and habitat alterations that may affect five species of sea turtles in the southeastern United States (Table 1). Literature on related species and from other areas is cited.

Migration and Movements

"Migrations of sea turtles begin at the hatchling stage and continue throughout life. Their travels range in scope from daily commuting between feeding and sleeping places to periodic reproductive journeys that take them across a thousand miles of ocean or more" (Carr 1982). Migrations range from the relatively short movements of hawksbills (Pritchard and Stubbs 1982) to the over 2000 km journeys of leatherbacks (Pritchard 1971, 1976b; Baker 1981). Environmental cues that may influence the ability of turtles to find the way to some nesting beaches year after year are poorly understood (Owens et al. 1982), but they may include specific beach odors and water current direction (Bennett and Kleerekoper 1978). Mortimer (1981b) suggests the green turtles in Costa Rica are guided by river effluents.

Movements of females during the nesting season are similar for most species. Green turtles at Ascension Island that had successfully nested followed different travel patterns than those that had been prevented from nesting (Mortimer 1981a). The latter stayed close to the beach and traveled back and forth in nearshore waters, while successful nesters moved to a depth of 12-16 m then travelled parallel to the shoreline. Meylan (1982a), who found similar patterns for green turtles at the Tortuguero colony, also found that travel offshore was restricted to about 4.8 km from land. Hopkins and Murphy (1981) found that loggerhead turtles nesting in South Carolina tended to move parallel to the coast and to use shoals and areas of high relief intensively. They were inactive at night and tended to move in either long straight lines or randomly during daylight. Williams-Walls et al. (1983) thought that offshore reefs at Hutchinson Island, Florida might be attractive to females during intervals between nestings and thereby influence selection of nesting sites. There also appear to be areas where females take refuge from courting males (Booth and Peters 1972). Dizon and Balazs (1982), using radio telemetry, tracked Hawaiian green turtles and found that males and females remained close to the basking and nesting islands where they were tagged. They believed that the nesting beach and its environs are imprinted on both males and females and that the green turtle, once imprinted, is unlikely to switch its breeding habitat.

Movements of turtles at times other than during nesting seem to be influenced by a variety of factors, such as age, availability of food, and water temperature. The green turtle appears to be the most regular long distant migrant. Nearly always their nesting beaches and forage grounds are spatially separated (Meylan 1982b). Although tagged loggerheads have traveled long distances (Carr et al. 1979; Timko and Kolz 1982), they seem to be wanderers influenced more by prey distribution than by migratory instincts (Carr et al. 1978). Bell and Richardson (1978) speculate that after breeding, loggerheads follow a

path from Georgia to the Cape Hatteras area and ultimately to warm mid-Atlantic water, perhaps the Sargasso Sea. Young ridleys have been observed as far north as New England (Lazell 1980). In Virginia, peak abundance of ridleys in June and October (Lutcavage 1981) implies a movement north in spring and a movement south in fall. Movements may be influenced by water temperature or prey distribution (Lutcavage 1981). Adults appear restricted to the Gulf of Mexico (Woodard 1980c). Leatherbacks, the most pelagic species, move between northerly feeding ranges and tropical nesting sites and overwintering areas (Lazell 1979). Records from the central Atlantic states indicate that many remain close to shore during migrations and are seasonally common (Lee and Palmer 1981; Hoffman and Fritts 1982). They appear to travel northward in the western Atlantic with the spring season, perhaps influenced by the Gulf Stream, and return south through the bays and sounds of New England (Lazell 1980). Rhodin and Schoelkopf (1982), who support Lazell's (1980) theory that postnesting leatherbacks can make at least a short migration northward, suggest that they are also capable of making very long, swift migrations. They do not know if females reach boreal waters only in non-nesting years. Little is known about movements of hawksbills. Once an adequate feeding area is located, they appear to be relatively sedentary, since feeding grounds such as coral reefs are typically close to nesting beaches. Tag returns have shown that some may be long-distance migrants (Pritchard and Stubbs 1982).

IMPACT OF HUMAN ACTIVITIES

Human-related activities such as urban and industrial development, petroleum exploitation, mineral sands mining, dredging, and commercial fishing appear to pose the greatest threats to turtle habitat (Shabica 1982). These activities may affect the well-being and survival of sea turtles in a variety of ways, from the direct physical destruction of nests to the more subtle effects of chemical pollutants on longevity and reproductive capacity. Because turtles come in close contact with man during their nesting cycle, they have been studied more thoroughly during this phase than they have during other periods of their life in the ocean. For convenience, we may divide the known information of man's activities on turtle populations into two major categories: the impact on nesting success, and the impact on oceanic survival.

Nesting Success

For turtles to nest successfully, they must breed successfully. Yet during breeding, turtles are quite susceptible to capture or injury

by man. Although some turtles from the Tortuguero colony have been observed to copulate 100 miles from known nesting sites (A. Meylan pers. comm.), most turtles probably mate in water just off the nesting beaches (Hirth 1971; Bustard 1973b; Ernst and Barbour 1972; Rebel 1974; Dizon and Balazs 1982), where they are most likely to be disturbed by man's activities.

Copulating pairs, which have been observed floating on the surface for hours, are vulnerable to capture by trawls, drop nets, and harpoons. Some Culebra Island fishermen use decoys to attract male turtles, which become entangled in nets when they attempt to copulate with the decoy (Carr 1977). Nicaraguan fishermen set traps to catch copulating pairs (A. Meylan pers. comm.). A substantial group of loggerheads that may moblize and breed in the Florida Keys before moving to the mainland to nest causes problems for spiny lobster fishermen, who claim they damage their traps (Higman and Davis 1978).

Destruction or modification of beaches where turtles nest probably has the greatest impact on the ability of turtle populations to sustain their numbers. Increased light from industrial or domestic development may cause a reduction in the number of females coming ashore to nest (Worth and Smith 1976), and it may cause newly-hatched turtles to become disoriented and move away from the water (McFarlane 1963; Philibosian 1976; Baker 1981) to die of desiccation or predation. Increased recreational use may destroy nests and eggs; deep tire tracks may prevent hatchlings from reaching the surf (Hosier et al. 1981). Construction may deter or prevent females from coming ashore, as it did on Hutchinson Island, where a cofferdam offshore was a barrier to turtles swimming along the shoreline (Williams-Walls et al. 1983). Clearing vegetation from the beach may reduce shade and increase nest temperatures, whereas large buildings may lower nest temperatures by increasing the time an area is shaded. Since temperature is an important factor in hatching success and sex determination, even small changes could cause increased mortality, delays in hatching, or sex ratio imbalance.

In the intervals between depositing clutches of eggs, females remain close to shore, where they are particularly vulnerable to being killed or injured by accidents or being caught by fishermen seeking them for food. Shrimp trawlers, which usually operate close to shore, may catch large numbers incidentally. Recreational fishermen or boaters may accidentally injure turtles with propellers. The potential impact from these and ecological disasters, such as oil spills, on the survival of a colony is greatly increased since it is the reproductive contingent that is affected (Meylan 1982a).

Oil spills have received considerable attention in recent years because of their high visibility, yet their effects remain largely

unknown. If they occur during the nesting season, the resulting cleanup operations could have harmful effects by 1) frightening gravid females from an area, 2) destroying nests, and 3) creating physical obstructions that would prevent hatchlings from reaching the ocean. Eggs, embryos, and hatchlings are more vulnerable than adults since volatile and water-soluble contaminants can be absorbed into the egg (see Ackerman and Prange 1972; Prange and Ackerman 1974; Bustard and Greenham 1968). There is considerable time for eggs to be affected, since incubation time is about two months (Frazier 1971, 1980). Fritts and McGehee (1982) found that fresh crude oil will cause significant mortality in incubating embryos, but that weathered oil might not. Any factors which would impair gas exchange in natural nests, and hence increase incubation time, would prolong the exposure of eggs to predators and the uncertanities of weather, disturb the synchrony of hatching, and increase egg mortality (Ackerman 1980).

Beach habitat is critical to turtle survival, for without it turtles could not reproduce. Yet it is extremely sensitive, directly or indirectly, to a wide range of human activities, including military exercises carried out on remote beaches. The following is an annotated list of human activities that may be harmful to nesting turtles:

Coastal Zone Development

Alteration of beach topography by

Artificial barriers - seawalls, riprap and jetties

- Impede turtle access to the nesting beach (Mann 1977; Witham 1982)
- Alter suitability of nesting beach by changing sand supply (Hopkins and Richardson 1981)

Traffic - vehicular, beach cleaning, pedestrian, horse, cattle

- Compression damage to nests (Mann 1977; Rainey 1978; Witham 1982; A. Carr pers. comm; P. Pritchard pers. comm.)
- Beach cleaning, including oil spill cleanup that may destroy nests (Mann 1977; Fletemeyer 1979a, Bureau of Land Management 1981) or cause compaction lowering hatching success (Ackerman 1980, 1981; Fletemeyer 1979a, 1982)

- Physical obstacles, such as tire tracks and sand piles that slow the rate of sea-approach for hatchling turtles and increase their susceptibility to stress and predation (Mann 1977; Hosier et al. 1981; Witham 1982)

Beach nourishment

- Dredge spoil may affect hatching success if deposited over nests (Mann 1977; Ackerman 1980; Witham 1982; Hopkins and Richardson 1981) or provide unsuitable nesting medium (Mann 1977; Fletemeyer 1979b, 1980a, 1981; Mortimer 1981a; Ehrhart and Raymond 1982; Witham 1982)
- Mechanical earth moving may damage nests by compression or excavation (see Traffic), increase chance of storm washover, or expose buried sediments unsuitable for nesting (Rainey 1978)

Sand mining

- May leave unsuitable nesting beach (Baker 1981; Mortimer 1981a; Sella 1982; A. Carr pers. comm.), or increase chance of storm washover (Rainey 1978)

Dredging

- Reduces sand supply that replenishes beaches (P. Pritchard pers. comm.)

Artificial illumination

- May discourage adults from nesting (Worth and Smith 1976; Mann 1978; Rainey 1978; Mortimer 1982a; Witham 1982) or disorient hatchlings (McFarlane 1963; Ehrenfeld 1968; Philibosian 1976; Mann 1977, 1978; Mrosovsky 1978; Rainey 1978; Towle 1978; Fletemeyer 1979a; Van Rhijn 1979; Frazier 1980)

Human activities and noise

- May deter nesting (Williams-Walls et al. 1983) or affect hatchling emergence (Balazs and Ross 1974 as cited in Rainey 1978)

Litter

- Solid debris may obstruct, entrap, or injure nesting females (Rainey 1978)
- May block progress of hatchlings from nest to water (K. Bjorndal, pers. comm.)
- Food may increase predation by attracting scavengers (Rebel 1974; Stancyk 1982)

Temperature alterations, such as shading of nesting beach by buildings or tar on beach

- May influence incubation time (Bustard and Greenham 1968; Fowler 1979; Mrosovsky 1980; Yntema and Mrosovsky 1980), sex determination in embryos (Yntema and Mrosovsky 1979; Limpus and Miller 1980; Mrosovsky 1980; Mrosovsky and Yntema 1980; Miller and Limpus 1981; Morreale et al. 1982; Mrosovsky 1982; Yntema and Mrosovsky 1982), or may influence hatchling emergence from nest (Mrosovsky 1968)

Exotic vegetation

- May increase shading of nesting beach, or form dense root mats that prevent nest excavation (Hopkins and Richardson 1981)

Pollution - including energy development, industrial discharges, and runoff from agribusiness (see Frazier 1980 for review on pollutants)

Oil spills and subsequent tar balls (Keller and Adams 1983)

- Act as deterrent to nesting (Bureau of Land Management 1981), affect hatching success (Fritts and McGehee 1982), irritate eyes and respiratory system of hatchlings (Bureau of Land Managment 1981), or interfere with olfactory imprinting (Manton et al. 1972 as cited in Rainey 1978)

Pesticides

- Have been detected, but minimum levels that will have adverse effects are unknown (Hillestad et al. 1974; Clark and Krynitsky 1980; Fletemeyer 1980a, 1980b; Hall 1980)

Heavy metals

- Have been detected, but effects unknown (Hillestad et al. 1974; Hall 1980; Stoneburner et al. 1980)

PCB's

- Have been detected, but effects unknown (Thompson et al. 1974; Hall 1980)

Thermal effluents

- May affect cues important in nest site selection (Lenarz and Stoneburner 1980; Stoneburner and Richardson 1981)

Military Activities

Ordnance impact and detonation

- May destroy or damage nests (Rainey 1978; Pritchard and Stubbs 1982) and pressure waves might reduce hatching success or trigger nest cave-ins (Rainey 1978)

Lights, either fixed or flares

- May discourage adults from nesting or disorient hatchlings (Rainey 1978; Pritchard and Stubbs 1982; see artificial illumination)

Troops and equipment

- May damage nests (Pritchard and Stubbs 1982)

Noise

- Possibly affect emergence of hatchlings (Balazs and Ross 1974 as cited in Rainey 1978)

Oceanic Survival

Throughout their life in the sea, turtles are subject to a host of human activities that threaten their survival. At some period in their life, however, they may be more susceptible than others. Young turtles may be more sensitive to lower levels of toxic pollutants than older turtles. Females may be more vulnerable to capture during the nesting season, when more of their time is spent close to shore. Human activities that are detrimental to turtle survival may be classified broadly into three categories: 1) those that destroy or damage habitat, 2) those that debilitate or weaken turtles physically, and 3) those that injure or kill turtles directly.

Any activity that reduces or contaminates the food supply or destroys resting places will reduce the ability of turtles to survive. For example, damage to sea grasses by dredging, thermal effluents, and chemical pollutants (Thorhaug 1981) will affect green turtles, which depend on grasses for food. Reef habitat destroyed by pollution (Johannes 1975), siltation (Rogers 1977), or mechanical damage (marine construction, dredging, ordnance) (Rainey 1978) constitutes not only a loss of foraging areas, but also a loss of resting places for adult and immature hawksbills and loggerheads. Many filter feeding detrital organisms that concentrate pollutants are important prey items for turtles in neritic habitats. The effect of these pollutants on the turtles is currently unknown. Large areas of hypoxia in the northwest Gulf of Mexico (Boesch 1983) associated with areas of "dead" bottom might influence the feeding of ridley and loggerhead turtles (B. Gallaway pers. comm.). At present, it is not known if these areas are natural and just previously undocumented, or are a result of the increased flow of nutrients from the Mississippi River discharge.

Possible effects of exploratory gas and oil drilling on sea turtles are largely speculative at present. Spoil disposal and oil development on live bottom areas may disrupt turtle feeding by smothering benthic organisms with sediments and drilling muds. Oil contamination may cause irritation or permanent damage to a turtle's eyes and respiratory system, resulting in abnormal behavior (Bureau of Land Management 1981). Loggerheads in the Gulf of Mexico are attracted to reef-like structures formed by drilling platforms for resting and

feeding (Hastings et al. 1976; Rabalais and Rabalais 1980; B. Gallaway pers. comm.). This could be a positive facet of habitat alteration unless turtles remaining in these areas were exposed to pollutants or increased predation. Dredging, otter trawls, ordnance impact, and boat propellers can damage coral reefs and seagrass beds directly by altering bottom topography and indirectly by increasing sediment and changing patterns of water flow. Rogers (1977) documents responses of corals to sediments, and Thorhaug (1981) and Zieman (1982) consider the responses of seagrasses. Altering physical features close to nesting beaches could alter nesting distribution of turtles.

A wide number of activities contribute directly to the death or injury of turtles. Garbage discharge from ships may attract turtles and cause them to ingest plastic and other foreign material accidentally, or to become entangled with refuse. Green turtles near Ascension Island have been found with turnip tops in their stomachs (A. Carr pers. comm.), and young turtles have been found with styrofoam pellets in their guts (A. Meylan pers. comm.). Turtles have been observed trying to swim with sheet plastic wrapped around their shells (Morris 1980a). Activities associated with fishing cause much damage. All species are susceptible to injury by power boats (Fletemeyer 1979b), and those attracted to refuse from seafood packing plants are particularly vulnerable (Shoop and Ruckdeschel 1982). Incidental catches of turtles in various fishing gear, particularly in shrimp trawls, have received much attention (see Shrimp Management Plan, Fuller 1981; Recovery Plan for Marine Turtles, Hopkins and Richardson 1981; Roithmayr 1981; and Crouse 1982). Immature loggerheads and ridleys in bays and sounds are particularly susceptible to shrimp trawls, pound nets or other nets in nearshore areas. Loggerheads are attracted to fish culled from shrimp trawlers and to refuse from seafood packing plants (Shoop and Ruckdeschel 1982), making them more vulnerable to trawls. Leatherbacks, the most pelagic species, may become entangled in long lines and drift gill nets (Balazs 1982a, 1982b). Increased strandings in South Carolina during the nesting season correspond to the gill net fishery for sturgeon (S. Hopkins pers. comm.). Power plants ranging from Florida to New Jersey have reported turtles, mostly immature loggerheads, that have been collected on cooling water intake screens. The average annual incidental catch for one power plant in Florida was 134 from 1977-1979 (Wilcox 1980 as cited in Roithmayr 1981). The offshore intake structure of Florida Power and Light Company's St. Lucie Plant may appear as a reef or a suitable resting area, from which some attracted turtles may be subsequently drawn into the cooling system. Also, turtles may follow prey into the canals (Witham 1982). Over a 12-month period a power plant on the Cape Fear River, N.C., collected three loggerheads and a green turtle from an upstream intake (W. Hogarth pers. comm.). Lighted permanent structures at sea (OTEC, oil rigs), which may attract young sea turtles (Witham 1982), may also attract pelagic fish that will prey on the turtles. Hatchling turtles have been observed aggregating near a lighted vessel anchored off a major nesting beach (Rainey 1978).

Pollutants from industrial and residental development are perhaps the most pervasive and subtle threats to the oceanic survival of all species. The effects are difficult to detect and evaluate and may not show up until the turtles have been exposed for many years. Chemicals, including oil, may mask olfactory cues or interfere with their perception (Kleerekoper and Bennett 1976; Frazier 1980), and may cause chronic and insidious problems in the ability of turtles to reproduce and maintain their numbers. In the Sargasso Sea, young turtles may be exposed to a high level of pollutants (A. Carr pers. comm.) as a result of the entire community being exposed to long-term heavy contamination (Morris et al. 1976). It has been estimated that tankers introduce 86,000 metric tons of tar into the western north Atlantic and that 75% of that tonnage is in the Sargasso Sea (Lineaweaver 1979). It is difficult to interpret the significance of measured levels of trace metals since little is known about their baseline levels and physiological effects. Lead found at elevated levels in some green turtles (Witkowski and Frazier 1982) is considered to be the only important trace metal to be at elevated levels in the Sargasso Sea (H. Windom pers. comm.). Frazier (1980) questions whether the decline of Kemp's ridley, a species characteristically found in waters where organic content and turbidity are high and prawns are abundant, is related to high levels of pollutants in discharges from the Mississippi River. Thermal pollution, as heated effluents from power plants or desalination plants, may affect turtles indirectly by being detrimental to their food supply or more directly by causing hatchlings to become disoriented and reduce their swimming speed (O'Hara 1980). In addition to high temperatures, effluents from desalination plants may have higher salinities and may contain heavy metals, particularly copper. Some biological effects are given by Chesher (1975). The following is an annotated list of the important activities that may impact oceanic habitat:

Pollutant Discharges

Oil spills - including major spills (e.g., IXTOC-I), "deliberate" spills e.g., bilging operations (Travers and Luney 1976), refinery spills, and spills from exploratory drilling (Keller and Adams 1983)

- Interference with olfactory perception (Kleerekoper and Bennett 1976; Bennett and Kleerekoper 1978; Witham 1978)
- Ingestion (Witham 1978; A. Meylan pers. comm.)
- Fouling of body with possibility of interference with respiratory functions (Fletemeyer 1980a; Frazier 1980; Bureau of Land Management 1981)
- Direct mortality (Hall 1980; Hooper 1981 Shabica 1982)
- Effects on seagrass habitat (Diaz-Piferrer 1962; Nadeau and Berquist 1977; Lopez 1978; Zieman et al. 1981; Zieman 1982;), reef habitat (Johannes 1975) Sargasso Sea habitat (Morris et al. 1976; Wade et al. 1976)

Pesticides from agricultural and urban runoff and spills

- o Levels have been measured in tissues but effects have not been evaluated (Thompson et al. 1974; Hall 1980)

Heavy metals from power plants, desalination plants, other industrial effluents and urban runoff

- o Levels have been measured in tissues but effects have not been evaluated (Hillestad et al. 1974; Hall 1980; Stoneburner et al. 1980; Witkowski and Frazier 1982)

Radionuclides from power plant or industrial effluents

- o Levels have been measured in tissues but effects have not been evaluated (Hillestad et al. 1974)
- o Laboratory experiments on uptake of radionuclides by Thalassia, which is eaten by green turtles (Parker 1962, 1966)

PCB's

- o Levels have been measured in tissues but effects have not been evaluated (Thompson et al. 1974; Hall 1980)

Domestic discharges, including sewage

- o Effects on seagrass habitat (Zieman 1982)
- o Effects on reef habitat (Johannes 1975)

Energy Development, including OTEC

Power plant intake canals

- o Entrainment and/or impingement (Fletemeyer 1979b, 1980a, 1981; Roithmayr 1981; Witham 1982; W. Hogarth, pers. comm.)

Thermal effluent

- o Influence on hatchling swimming speed (O'Hara 1980)
- o Possible attraction to thermal plume in cooler water (reference to work done by M. Stinson from M. Lutcavage pers. comm.)
- o Degradation of seagrass habitat (Zieman 1970; Thayer et al. 1975; Zieman and Wood 1975; Thorhaug 1981; Zieman 1982)
- o Degradation of reef habitat (Johannes 1975)

Lighted offshore structures

- o By attracting young turtles may make them more susceptible to predation (Rainey 1978)

Dredging and mining, including oil drilling

Alteration of nearshore bottom topography

- o May affect internesting behavior (Hopkins and Murphy 1981; Williams-Walls et al. 1983)

Channel dredging

- o Resting and/or hibernating turtles may be vulnerable (Richardson and Hillestad 1979; Carr et al. 1980; Raymond 1980; Rudloe 1981)

Disposal of spoil or drilling muds on live bottom areas

- o May smother benthic organisms (Bureau of Land Management 1981)
- o May destroy reef communities utilized by turtles (Johannes 1975; Rogers 1977)
- o May affect seagrass ecosystems used by turtles (Odum 1963; Taylor and Saloman 1968; Thayer et al. 1975; Thorhaug 1981; Zieman 1982)

Fishing Activities (see incidental catch overview in Recovery Plan for Marine Turtles (Hopkins and Richardson 1981)

General (Hildebrand 1980; Roithmayr 1981; Crouse 1982)

Trawls

- o Incidental catch (Cox and Mauermann 1976; Hillestad et al. 1977, 1978, 1982; Ogren et al. 1977; Carr et al. 1978; Ulrich 1978; Rabalais and Rabalais 1980; Coleman 1981; Fuller 1981)
- o Attraction to culling by trawlers and disposal from seafood processing plants makes turtles more vulnerable to being caught (Shoop and Ruckdeschel 1982)
- o Bottom and reef destruction (Thayer et al. 1975)

Incidential catch by set nets (Talbert et al. 1980) longlines (Balazs 1982a, 1982b), pound nets (Lutcavage 1981; R. Byles pers. comm.), and gill nets (L. Ehrhart pers. comm.)

"Bleach fishing" and "dynamite fishing" in Caribbean (A. Carr pers. comm.)

Recreational fisheries

- o Rod and reel fishermen (Hildebrand 1980; Roithmayr 1981)
- o Spearfishing on reefs (Carr and Stanyck 1975)
- o Reef habitat destruction from anchoring, collection of shells and corals, and leaving litter (Johannes 1975; Hopkins and Richardson 1981)

Miscellaneous

Litter and garbage dumping by ships

- o Ingestion of styrofoam pellets and fouling by plastic etc. (Carpenter and Smith 1972; Morris 1980; Mrosovsky 1981; Van Dolah et al. 1980; Horsman 1982; A. Carr pers. comm.)

Increased runoff from overgrazing by cattle or from poor land management

- o May cause subsequent degenerative changes in reef system and hawksbill habitat (Carr 1977)

Military activities

- o Disturbance and destruction of reef habitat (Carr 1978)

RECOMMENDATIONS

Major gaps in the knowledge of life histories and habitat requirements of all species of sea turtles are major impediments to the development of programs for turtle protection and management. In our literature survey and in our conversations with investigators we found continual reference to a lack of knowledge on habitat utilization (except for beach habitat) and a need for additional research in this area. We agree with Carr et al. (1978) that additional information on habitat requirements is urgently needed, especially for developmental and internesting habitats.

Much of the results from sea turtle research and surveys is not published in journals, but is published in not readily available reports or resides in file cabinets of investigators. We suggest a central listing, preferably computerized, of: 1) available unpublished information and where it may be obtained, and 2) names and addresses of people involved in research and their area of interest. Access to this type of information would aid people involved in making recommendations, such as those in environmental impact statements, that could affect the future of all turtle populations.

Despite increased interest in recent years in the well being of sea turtles, the amount of "new" information being generated is low. Many of the reports are repetitious and tend to paraphrase the same source material. We feel that research, particularly long-term research with adequate funding, is needed in the following areas:

1. Development of fishing gear, such as the Turtle Excluder Device (TED), that will mitigate or prevent turtle capture, and funding for education in the use of the new equipment.

2. The impact of fisheries other than the shrimp fishery (e.g., the roller trawl fishery) on turtles and turtle habitat.

3. Research on practical techniques, such as the use of audible or ultrasonic noise, to divert turtles from danger areas.

4. The long-term effects of chemicals, such as synthetic organics, on reproduction, survival, and other phases of turtle life history.

5. Continuing studies on the dispersal mechanisms of hatchlings, effects of currents on their transport and the developmental habitat subsequently occupied.

6. Methods of externally sexing young turtles

7. The size and sex structure of turtles in specific inshore areas.

8. Tolerance levels during embryonic development to petrochemicals, bilge effluents, and chemicals used in cleanup of oil spills.

9. The effects of an oil spill on breeding and nesting success.

ACKNOWLEDGMENTS

The preparation of this report was only possible due to the cooperation of a large number of investigators in the field. We are especially indebted to the following people for providing us with unpublished data and their personal comments:

Karen Bjorndal - Department of Zoology, University of Florida, Gainesville, FL 32611

Richard Byles - Virginia Institute of Marine Science, Gloucester Point, VA 23062

Archie Carr - Department of Zoology, University of Florida, Gainesville, FL 32611

Lew Ehrhart - Department of Biological Science, University of Central Florida, Box 25000, Orlando, FL 32861

Benny Gallaway
Greg Boland - LGL Ecological Research Associates, 1410 Cavett St., Bryan, TX 77801

W. T. Hogarth - Carolina Power and Light Company, Harris E E Center, Route 1, Box 327, New Hill, NC 27562

Sally Hopkins - South Carolina Marine Resources Center, P.O. Box 12599, Charleston, SC 29412

Molly Lutcavage - RSMAS/BLR, 4600 Rickenbacker Causeway, Miami, FL 33149

Erik Martin - Applied Biology, Inc., P.O. Box 974, Jensen Beach, FL 33457

Anne Meylan - Department of Zoology, University of Florida, Gainesville, FL 32611

Walter Nelson - NMFS, SEFC, Pascagoula, MS 39567

Larry Ogren - NMFS, SEFC, Panama City, FL 32407

James O'Hara - Environmental and Chemical Sciences Inc., Atlanta, GA

Peter Pritchard - 1101 Audubon Way, Maitland, FL 32751

Charlie Roithmayr - NMFS, SEFC, Pascagoula, MS 39567

Jim Richardson - Institute of Ecology, University of Georgia, Athens, GA 30602

Frank Schwartz - Institute of Marine Sciences, UNC-CH, Morehead City, NC 30602

Herbert Windom - Skidaway Institute of Oceanography, Savannah, GA 31406

Ross Witham - Florida Dept. of Natural Resources, Marine Research Laboratory, PO Box 941, Jensen Beach, FL 33457

In addition, a special thanks to Ann Hall, Beaufort Laboratory Librarian for extensive literature searches and help with the bibliography and to Drs. Karen Bjorndal and Frank Schwartz for reviewing this manuscript.

BIBLIOGRAPHY

Ackerman, R.A.

1980. Physiological and ecological aspects of gas exchange by sea turtle eggs. American Zoologist, vol. 20, p. 575-583.

Ackerman, R.A.

1981. Growth and gas exchange of embryonic sea turtles (Chelonia, Caretta). Copeia, 1981, p. 757-765.

Ackerman, R.A., and H.D. Prange.

1972. Oxygen diffusion across a sea turtle (Chelonia mydas) egg shell. Comparative Biochemistry and Physiology, vol. 43A, p. 905-909.

Baker, G.S.

1981. Recovery plan for the St. Croix population of the leatherback turtle. Prepared by U.S. Fish and Wildlife Service, Region 4. Available from Fish and Wildlife Reference Service, Unit I, 3840 York Street, Denver, Colo. 80205, 20 p.

Baker, R.R.

1978. The evolutionary ecology of animal migration. Holmes Meier, New York. 1012 p.

Balazs, G.H.

1982a. Annotated bibliography of sea turtles taken by longline fishing gear. Unpublished manuscript, U.S. National Marine Fisheries Service Laboratory, P.O. Box 3830, Honolulu, Hawaii 96812. 4 p.

Balazs, G.H.

1982b. Driftnets catch leatherback turtles. Oryx, vol. 16, p. 428-430.

Balazs, G.H., and E. Ross.

1974. Observations on the preemergence behavior of the green turtle. Copeia, 1974, p. 986-988.

Bell, R., and J. I. Richardson.

1978. An analysis of tag recoveries from loggerhead sea turtles (Caretta caretta) nesting in Little Cumberland Island, Georgia. Florida Marine Research Publications, no. 33, p. 20-24.

Bennett, J.A., and H. Kleerekoper.

1978. A preliminary investigation of the effects of chemical stimulation on the locomotor behavior of hatchling green turtles (Chelonia mydas). Florida Marine Research Publications, no. 33, p. 3-7.

Bjorndal, K. A.

1980. Nutrition and grazing behavior of the green turtle Chelonia mydas. Marine Biology, vol. 56, p. 147-154.

Bjorndal, K. A.

1982. The consequences of herbivory for the life history of the Caribbean green turtle, Chelonia mydas, p.111-116. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Bjorndal, K. A., A.B. Meylan, and B. J. Turner.

1983. Sea turtles nesting at Melbourne Beach, Florida, 1. Size, growth and reproductive biology. Biological Conservation, vol. 26, p. 65-77.

Blanck, C. E., and R. H. Sawyer.

1981. Hatchery practices in relation to early embryology of the loggerhead sea turtle, Caretta caretta (Linne). Journal of Experimental Marine Biology and Ecology, vol. 49, 163-177.

Boesch, D. F.

1983. Implications of oxygen depletion on the continental shelf of the northern Gulf of Mexico. Coastal Ocean Pollution Assessment News, vol. 2, no. 3, p. 25-28.

Booth, J., and J. A. Peters.

1972. Behavioural studies on the green turtle (Chelonia mydas) in the sea. Animal Behaviour, vol. 20, no. 4, p. 808-812.

Brongersma, L. D.

1969. Miscellaneous notes on turtles. II. A and B. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series C, Biological and Medical Sciences, vol. 72, p. 76-102.

Brongersma, L. D.

1972. European Atlantic turtles. Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie te Leiden, no. 121. 318 p.

Bullis, H. R., Jr., and S. B. Drummond.

1978. Sea turtle captures off the southeastern United States by exploratory fishing vessels, 1950-1976. Florida Marine Research Publications, no. 3, p. 45-50.

Bureau of Land Management.

1981. Final environmental impact statement, proposed 1981 Outer Continental Shelf Oil and Gas Lease sale 56. U. S. Bureau of Land Management, Outer Continental Shelf Office, New Orleans, La. 576 p.

Burnett-Herkes, J.

1974. Returns of green turtles (Chelonia mydas) tagged at Bermuda. Biological Conservation, vol. 6, p. 307-308.

Bustard, H. R.

1967. Mechanism of nocturnal emergence from the nest of green turtle hatchlings (Chelonia mydas). Nature, vol. 214, p. 317.

Bustard, H. R.

1973a. Saving the hawksbill turtle. Oryx, vol. 12, no. 1, p. 93-98.

Bustard, H. R.

1973b. Sea turtles, natural history and conservation. Taplinger Publishers, New York. 220 p.

Bustard, H. R.

1976. Turtles of coral reefs and coral islands, p. 343-368. In O. A. Jones and R. Endean (eds.), Biology and geology of coral reefs, vol. 3, biology 2. Academic Press, New York.

Bustard, H. R., and P. M. Greenham.

1968. Physical and chemical factors affecting hatching in the green sea turtle, Chelonia mydas (L.). Ecology, vol. 49, p. 269-276.

Caldwell, D. K.

1959. The loggerhead turtles of Cape Romain, South Carolina. Bulletin of the Florida State Museum, vol. 4, p. 319-348.

Caldwell, D. K.

1960. Sea turtles of the United States. U.S. Fish and Wildlife Service, Fishery Leaflet 492. 20 p.

Caldwell, D. K.

1962. Comments on the nesting behavior of Atlantic loggerhead sea turtles, based primarily on tagging returns. Quarterly Journal of the Florida Academy of Sciences, vol. 25, p. 287-302.

Caldwell, D. K.

1968. Baby loggerhead turtles associated with sargassum weed. Quarterly Journal of the Florida Academy of Sciences, vol. 31, p. 271-272.

Carlson, C. E.

1978. An abridged history of the threatened and endangered species program in the U.S. Georgia Department of Natural Resources, Game and Fish Division, Technical Bulletin WL 4, p. 3-6.

Carpenter, E. J., and K. L. Smith, Jr.

1972. Plastics on the Sargasso Sea surface. Science, vol. 175, p. 1240-1241.

Carr, A. F.

1952. Handbook of turtles: The turtles of the United States, Canada, and Baja California. Comstock Publishing Associates, Cornell University Press, Ithaca, N. Y. 542 p.

Carr, A. F.

1965. The navigation of the green turtle. Scientific American, vol. 212, p. 79-86.

Carr, A. F.

1967. So excellent a fishe. Natural History Press, Garden City, N.Y. 248 p.

Carr, A. F.

1972a. Great reptiles, great enigmas. Audubon, vol. 74, no. 2, p. 24-34.

Carr, A.

1972b. The case for long-range chemoreceptive piloting in Chelonia, p. 469-483. In S. R. Galler, K. Schmidt-Koening, G. J. Jacobs and R. E. Belleville (eds.), Animal orientation and navigation. National Aeronautics and Space Administration, Washington, D.C.

Carr, A.

1980. Some problems of sea turtle ecology. American Zoologist, vol. 20, p. 489-498.

Carr, A.

1982. Notes on the behavioral ecology of sea turtles, p. 19-26. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Carr, A., and D. K. Caldwell.

1956. The ecology and migration of sea turtles. 1. Results of field work in Florida, 1955. American Museum Novitates, no. 1793. 23 p.

Carr, A., and D. K. Caldwell.

1958. The problem of the Atlantic ridley turtle (Lepidochelys kempi) in 1958. Revista Biologica Tropical, vol. 6, no. 2, p. 245-262.

Carr, A., and M. H. Carr.

1972. Site fixity in the Caribbean green turtle. Ecology, vol. 53, p. 425-429.

Carr, A., M. H. Carr, and A. B. Meylan.

1978. The ecology and migrations of sea turtles. The west Caribbean green turtle colony. Bulletin of the American Museum of Natural History, vol. 162, no. 1, p. 1-46.

Carr, A. F., and L. Giovannoli.

1957. The ecology and migrations of sea turtles. 2. Results from field work in Costa Rica, 1955. American Museum Novitates, no. 1835. 32 p.

Carr, A. F., H. F. Hirth, and L. Ogren.

1966. The ecology and migration of sea turtles. 6. The hawksbill turtle in the Caribbean Sea. American Museum Novitates, no. 2248. 29 p.

Carr, A. F., D. R. Jackson, and J. B. Iverson.

1979. Marine turtles, p. 1-45. In A summary and analysis of environmental information on the Continental Shelf and Blake Plateau from Cape Hatteras to Cape Canaveral (1977), vol. 1, bk. 3. Center for Natural Areas, Washington, D. C. (NTIS PB80-184112).

Carr, A., and A. B. Meylan.

1980a. Evidence of passive migration of green turtle hatchlings in sargassum. Copeia, 1980, p. 366-368.

Carr, A., and A. B. Meylan.

1980b. Extinction or rescue for the hawksbill? Oryx, vol. 25, no. 5, p. 449-450.

Carr, A., A. Meylan, J. Mortimer, K. Bjorndal, and T. Carr.

1982. Surveys of sea turtle populations and habitats in the western Atlantic. U.S. National Marine Fisheries Service, Southeast Fisheries Center, NOAA Technical Memorandum NMFS SEFC-91. 82, 9 p.

Carr, A. F., and L. Ogren.

1960. The ecology and migrations of sea turtles. 4. The green turtle in the Caribbean Sea. Bulletin of the American Museum of Natural History, vol. 121, no. 1, p. 1-48.

Carr, A., L. Ogren, and C. McVea.

1980. Apparent hibernation by the Atlantic loggerhead turtle Caretta caretta off Cape Canaveral, Florida. Biological Conservation, vol. 19, p. 7-14.

Carr, A., P. Ross, and S. Carr.

1974. Internesting behavior of the green turtle, Chelonia mydas, at a mid-ocean island breeding ground. Copeia, 1974, p. 703-706.

Carr, A. F., and S. Stancyk.

1975. Observations on the ecology and survival outlook of the hawksbill turtle. Biological Conservation, vol. 8, no. 3, p. 161-172.

Carr, D., and P. H. Carr.

1977. Survey and reconnaissance of nesting shores and coastal habitats of marine turtles in Florida, Puerto Rico, and the U.S. Virgin Islands. Report to U.S. National Marine Fisheries Service, St. Petersburg, Fla. Contract no. 03-6-042-35129. 34 p.

Carr, T.

1977. The marine turtles and terrestrial reptiles Culebra Island. Report to the U.S. Fish and Wildlife Service. 43 p.

Carr, T.

1978. A survey of marine turtles at Vieques Island. Report to the Department of Natural Resources, Commonwealth of Puerto Rico. 16 p.

Chesher, R. H.

1975. Biological impact of a large-scale desalination plant, p. 99-153. In E. J. Ferguson Wood and R. E. Johannes (eds.), Tropical marine pollution. Elsevier, New York.

Clark, D. R., and A. J. Krynitsky.

1980. Organochlorine residues in eggs of loggerhead and green turtles nesting at Merritt Island Florida - July and August 1976. Pesticides Monitoring Journal, vol. 14, no. 1, p. 7-10.

Coleman, F. C.

1981. Shrimpers and turtles. Oceans, vol. 14, no. 2, p. 67-68.

Cornelius, S. E.

1975. Marine turtle mortalities along the Pacific coast of Costa Rica. Copeia, 1975, p. 186-187.

Cox, B. A., and R. G. Mauermann.

1976. Incidental catch and disposition by Brownsville - Port Isabel Gulf shrimp fleet. Cameron County Extension Service, San Benito, Tex. and Texas Shrimp Association, Brownsville, Tex. 5 p.

Crouse, D. T.

1982. Incidental capture of sea turtles by U.S. commercial fisheries. Report to the Center for Environmental Education, Washington, D.C. 19 p.

Dean, J. M., and O. R. Talbert.

1975. The loggerhead turtles on Kiawah Island, S.C., p. T1-T19. In W. M. Campbell and J. M Dean (eds.), An environmental inventory of Kiawah Island, S.C. Environmental Research Center, Columbia, S.C.

Delikat, D. S.

1980. IXTOC I oil spill and Atlantic ridley survival, p. 312-319. In B. L. Edge (ed.), Coastal Zone '80: Proceedings of the Second Symposium on Coastal and Ocean Management, vol. 1. American Society of Civil Engineers, New York.

Diaz-Piferrer, M.

1962. The effects of an oil spill on the shore of Guanica, Puerto Rico. (Abstract). Association of Island Marine Laboratories, 4th Meeting, Curacao, p. 12-13.

Dizon, A. E., and G. H. Balazs.

1982. Radio telemetry of Hawaiian green turtles at their breeding colony. U.S. National Marine Fisheries Service, Marine Fisheries Review, vol. 44, no.5, p. 13-20.

Dobie, J. L., L. H. Ogren, and J. F. Fitzpatrick, Jr.

1961. Food notes and records of the Atlantic ridley turtle (Lepidochelys kempi) from Louisiana. Copeia, 1961, p. 109-110.

Dodd, C. K., Jr.

1978. Terrestrial critical habitat and marine turtles. Bulletin of the Maryland Herpetological Society, vol. 14, no. 4, p. 233-240.

Dodd, C. K., Jr.

1982. Nesting of the green turtle, Chelonia mydas (L.), in Florida: Historic review and present trends. Brimleyana, no. 7, p. 39-54.

Ehrenfeld, D. W.

1968. The role of vision in the sea-finding orientation of the green turtle (Chelonia mydas). 2. Orientation mechanism and range of spectral sensitivity. Animal Behaviour, vol. 16, p. 281-287.

Ehrenfeld, D. W.

1970. Conserving the edible sea turtle: Can mariculture help? American Scientist, vol. 62, no. 1, p. 23-31.

Ehrenfeld, D. W. and A. Carr.

1967. The role of vision in the sea finding orientation of the green turtle (Chelonia mydas). Animal Behaviour, Vol. 15, p. 25-36.

Ehrhart, L. M.

1977a. Cold water stunning of marine turtles in Florida east coast lagoons: Rescue measures, population characteristics and evidence of winter dormancy. Paper read at 1977 American Society of Ichthyology and Herpetology Meeting, Gainesville, Fla.

Ehrhart, L. M.

1977b. Marine turtle studies, p. 1-91. In A continuation of baseline studies for environmentally monitoring space transportation systems (STS) at John F. Kennedy Space Center. Annual report to National Aeronautics and Space Administration, Kennedy Space Center, Contract no. NAS 10-8986.

Ehrhart, L. M.

1979. Reproductive characteristics and management potential of the sea turtle rookery at Canaveral National Seashore, Florida, p. 397-399. In R. M. Linn (ed.), Proceedings of the first conference on scientific research in the national parks, vol. 1. U.S. National Park Service, Transactions and Proceedings Series, no. 5.

Ehrhart, L. M.

1980. Threatened and endangered species of the Kennedy Space Center, vol. 4, pt. 1, Marine turtle studies. National Aeronautics and Space Administration, Contract Report 163122, KSC TR 51-2. 106, 301 p.

Ehrhart, L. M.

1982. A review of sea turtle reproduction, p. 29-38. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Ehrhart, L. M., and P. W. Raymond.

1982. 1981 Marine turtle reconnaissance interim progress report to the Department of the Army, Jacksonville District, Corps of Engineers. Professional engineering services for monitoring and preparation of a reconnaissance report of turtle nesting success, Melbourne and Indialantic Beaches, Brevard County, Florida. Contract no. DACW17-81-C-0014. University of Central Florida, Orlando. 27, 19 p.

Ehrhart, L. M., and R. G. Yoder.

1978. Marine turtles of Merritt Island National Wildlife Refuge, Kennedy Space Center, Florida. Florida Marine Research Publications, no. 33, p. 25-30.

Ernst, C. H., and R. W. Barbour.

1972. Turtles of the United States. University Press of Kentucky, Lexington. 347 p.

Felger, R. S., K. Cliffton, and P. J. Regal.

1976. Winter dormancy in sea turtles: Independent discovery and exploitation in the Gulf of California by two local cultures. Science, vol. 191, p. 283-285.

Fletemeyer, J. R.

1978a. The lost year. Sea Frontiers, vol. 24, p. 23-26.

Fletemeyer, J. R.

1978b. Underwater tracking evidence of neonate loggerhead sea turtles seeking shelter in drifting sargassum. Copeia, 1978, p. 148-149.

Fletemeyer, J. R.

1979a. Ascension Island development: Lessons from Florida. Marine Turtle Newsletter, no. 11, p. 2-3.

Fletemeyer, J. R.

1979b. Sea turtle monitoring project, 1979 report. Broward County Environmental Quality Control Board, Fort Lauderdale, Fla. 64 p.

Fletemeyer, J. R.

1980a. Sea turtle monitoring project, 1980 report. Cooperative Sea Turtle Monitoring Program, Nova University and Broward County Environmental Quality Control Board, Fort Lauderdale, Fla. 88 p.

Fletemeyer, J. R.

1980b. A preliminary analysis of sea turtle eggs for DDE. Marine Turtle Newsletter, no. 15, p. 6-7.

Fletemeyer, J. R.

1980c. The leatherback, turtle without a shell. Sea Frontiers, vol. 26, p. 302-305.

Fletemeyer, J. R.

1981. Sea turtle monitoring project, 1981 report. Cooperative Sea Turtle Monitoring Program, Nova University and Broward County Environmental Quality Control Board, Fort Lauderdale, Fla. 82 p.

Fletemeyer, J. R.

1982. Sea turtle monitoring project, 1982 report. Cooperative Sea Turtle Monitoring Program, Nova University and Broward County Environmental Quality Control Board, Fort Lauderdale, Fla. 95 p.

Fowler, L. E.

1979. Hatching success and nest predation in the green sea turtle, Chelonia mydas, at Tortuguero, Costa Rica. Ecology, vol. 60, p. 946-955.

Frazier, J. G.

1971. Observations on sea turtles at Aldabra Atoll. Philosophical Transactions of the Royal Society of London, B, Biological Sciences, vol. 260, p. 373-410.

Frazier, J. G.

1980. Marine turtles and problems in coastal management, p. 2395-2411. In B. L. Edge (ed.), Coastal Zone '80: Proceedings of the Second Symposium on Coastal and Ocean Management, vol. 3. American Society of Civil Engineers, New York.

Frick, J.

1976. Orientation and behaviour of hatchling green turtles (Chelonia mydas) in the sea. Animal Behaviour, vol. 24, p. 849-857.

Fritts, T. H.

1982. Plastic bags in the intestinal tracts of leatherback marine turtles. Herpetological Review, vol. 13, no. 3, p. 72-73.

Fritts, T. H. and W. Hoffman.

1982. Diurnal nesting of marine turtles in southern Brevard County, Florida. Journal of Herpetology, vol. 16, no. 1, p. 84-86.

Fritts, T. H., and M. A. McGehee

1982. Effects of petroleum on the development and survival of marine turtle embryos. U.S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-82/37. 41 p.

Fritts, T. H., and R. P. Reynolds.

1981. Pilot study of the marine mammals, birds and turtles in OCS areas of the Gulf of Mexico. U. S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-81/36. 139 p.

Fuller, D. A.

1981. The habits, distribution and incidental capture of sea turtles in the Gulf of Mexico, p. A1-A36. In Final environmental impact plan for the shrimp fishery of the Gulf of Mexico, United States waters. Gulf of Mexico Fishery Management Council, Tampa, Fla.

Gallaway, B. J.

1981. An ecosystem analysis of oil and gas development on the Texas - Louisiana continental shelf. U.S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-81/27. 89 p.

Hall, R. J.

1980. Effects of environmental contaminants on reptiles, a review. U.S. Fish and Wildlife Service, Special Scientific Report - Wildlife, no. 228. 12 p.

Hastings, R. W., L. H. Ogren, and M. T. Mabry.

1976. Observations on the fish fauna associated with offshore platforms in the northeastern Gulf of Mexico. U. S. National Marine Fisheries Service, Fishery Bulletin, vol. 74, p. 387-402.

Hendrickson, J.

1980. The ecological strategies of sea turtles. American Zoologist, vol. 20, p. 597-608.

Hendrickson, J. R.

1982. Nesting behavior of sea turtles with emphasis on physical and behavioral determinants of nesting success or failure, p. 53-57. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Higman, J. B., and C. Davis.

1978. Damage to spiny lobster gear by sea turtles and other predators. Final report 29 August 1977 - 31 January 1978 to Southeast Fisheries Center, National Marine Fisheries Service, Contract no. 03-7-042-35136. 37 p.

Hildebrand, H. H.

1980. Report on the incidental capture, harassment and mortality of sea turtles in Texas. Report to U.S. National Marine Fisheries Service. Contract no. 03 6 042 3519. 34 p.

Hildebrand, H. H.

1982. A historical review of the status of the sea turtle populations in the western Gulf of Mexico, p. 447-453. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Hill, G. D., Jr.

1979. The troubled turtle. NOAA, vol. 9, no. 1, p. 21-23.

Hillestad, H. O., R. J. Reimold, R. R. Stickney, H. L. Windom, and J. H. Jenkins.

1974. Pesticides, heavy metals and radionuclide uptake in loggerhead sea turtles from South Carolina and Georgia. Herpetological Review, vol. 5, no. 3, p. 75.

Hillestad, H. O., J. I. Richardson, C. McVea, Jr., and J. M. Watson, Jr.

1982. Worldwide incidental capture of sea turtles, p. 489-495. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Hillestad, H. O., J. I. Richardson, and G. K. Williamson.

1977. Incidental capture of sea turtles by shrimp trawlermen in Georgia. Report to U.S. National Marine Fisheries Service, Miami, Fla. Contract no. 03-7-042-35129. 106 p.

Hillestad, H. O., J. I. Richardson, and G. K. Williamson.

1978. Incidental capture of sea turtles by shrimp trawlermen in Georgia. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies, no. 32, p. 167-178.

Hirth, H. F.

1971. Symposis of biological data on the green turtle Chelonia mydas (Linnaeus) 1758. FAO Fisheries Synopsis, no. 85. [77] p.

Hoffman, W. and T. H. Fritts.

1982. Sea turtle distribution along the boundary of the Gulf Stream current off eastern Florida. Herpetologica, vol. 38, no. 3, p. 405-409.

Hooper, C. H., editor.

1981. The IXTOC I oil spill: The federal scientific response. U.S. National Oceanic and Atmospheric Administration, Hazardous Materials Response Project, Boulder, Colo. 202 p.

Hopkins, S. R., and T. M. Murphy.

1981. Reproductive ecology of Caretta caretta in South Carolina. S. C. Wildlife and Marine Resources Department, Study Completion Report E-1, study no. VI-A-1, October 1, 1976 through September 30, 1980, Charleston. 96 p.

Hopkins, S. R., T. M. Murphy, Jr., K. B. Stansell, and P. M. Wilkinson.

1978. Biotic and abiotic factors affecting nest mortality in the Atlantic loggerhead turtle. Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies, no. 32, p. 213-223.

Hopkins, S. R., and J. I. Richardson, editors.

1981. Recovery plan for marine turtles. Prepared by the Marine Turtle Recovery Team, April 1981. Technical draft. 266 p.

Horsman, P. V.

1982. The amount of garbage pollution from merchant ships. Marine Pollution Bulletin, vol. 13, p. 167-169.

Hosier, P. E., M. Kochhar, and V. Thayer.

1981. Off-road vehicle and pedestrian track effects on the sea-approach of hatchling loggerhead turtles. Environmental Conservation, vol. 8, p. 158-161.

Hughes, G. R.

1974. The sea turtles of south-east Africa. II. The biology of the Tongaland loggerhead turtle Caretta caretta L. with comments on the leatherback turtle Dermochelys coriacea L. and the green turtle Chelonia mydas L. in the study region. Oceangraphic Research Institute (Durban), Investigational Report, no. 36. 96 p.

Ireland, L. C.

1979. Optokinetic behavior of the hatchling green turtle (Chelonia mydas) soon after leaving the nest. Herpetologica, vol. 35, p. 365-370.

Ireland, L. C., J. A. Frick, and D. B. Wingate.

1978. Nighttime orientation of hatchling green turtles (Chelonia mydas) in the open ocean, p. 420-429. In K. Schmidt-Koenig and W. T. Keeton (eds.), Animal migration, navigation and homing. Springer-Verlag, New York.

Johannes, R. E.

1975. Pollution and degradation of coral reef communities, p. 13-51. In E. J. Ferguson Wood and R. E. Johannes (eds.), Tropical marine pollution. Elsevier, New York.

Keller, C. E., and J. K. Adams, editors.

1983. Proceedings of a workshop on cetaceans and sea turtles in the Gulf of Mexico: Study planning for effects of Outer Continental Shelf Development. Prepared by the U.S. Fish and Wildlife Service for the Minerals Management Service, Metairie, La. FWS/OBS/03. 42 p.

Kingsmill, S.F., and N. Mrosovsky.

1982. Sea-finding behaviour of loggerhead hatchlings: The time course of transient circling following unilateral and asynchronous bilateral blindfolding. Brain, Behaviour and Evolution, vol. 20, p. 29-42.

Kleerekoper, H., and J. A. Bennett.

1976. Some effects of the water soluble fraction of Louisiana crude on the locomotor behavior of juvenile green turtle (Chelonia mydas) and sea catfish (Arius felis), preliminary results. Report on Contract 206-76 to American Petroleum Institute, Washington D. C. 7 p.

Klukas, R.

1967. Factors affecting nesting success of loggerhead turtles at Cape Sable Everglades National Park. File N1415, National Park Service, P.O. Box 279, Homestead, Fla. 33030.

Kraemer, J. E., and R. Bell.

1980. Rain induced mortality of eggs and hatchlings of loggerhead sea turtles (Caretta caretta) on the Georgia coast. Herpetologica, vol. 36, p. 72-77.

Kraemer, J. E., and S. H. Bennett.

1981. Utilization of posthatching yolk in loggerhead sea turtles, Caretta caretta. Copeia, 1981, p. 406-411.

Lazell, J. D., Jr.

1979. Boreal and temperate migratory regimes of Atlantic marine turtles. American Zoologist, vol. 19, p. 954.

Lazell, J. D., Jr.

1980. New England waters: Critical habitat for marine turtles. Copeia, 1980, p. 290-295.

LeBuff, C. R., Jr.

1976. Tourist turtle. Florida Wildlife, vol. 30, no. 2, p. 16-17.

Lee, D. S., and W. M. Palmer.

1981. Records of leatherback turtles, Dermochelys coriacea (Linnaeus), and other marine turtles in North Carolina waters. Brimleyana, no. 5, p. 95-106.

Lenarz, M. S., and D. L. Stoneburner.

1980. Temperature as an environmental cue in nesting loggerhead sea turtles (Chelonidae). (Abstract). ASB (Association of Southeastern Biologists) Bulletin, vol. 27, p. 45.

Limpus, C. J., V. Baker, and J. D. Miller.

1979. Movement induced mortality of loggerhead eggs. Herpetologica, vol. 35, p. 335-338.

Limpus, C. J., and J. D. Miller.

1980. Potential problems in artificial incubation of turtle eggs. Herpetofauna, vol. 12, no. 1, p. 24.

Lineaweaver, T. H., III.

1979. The Sargasso - Sea of fascinations. Sea Frontiers, vol. 26, p. 154-164.

Meylan, A.

1982a. The behavioral ecology of the west Caribbean green turtle (Chelonia mydas) in the internesting habitat, p. 67-80. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Meylan, A.

1982b. Sea turtle migrations - Evidence from tag returns, p. 91-100. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Miller, J. D., and C. J. Limpus.

1981. Incubation period and sexual differentiation in the green turtle Chelonia mydas L., p. 66-73. In C.B. Banks and A.A. Martin (eds.), Proceedings of the Melbourne Herpetological Symposium. Zoological Board of Victoria, Parkville.

Morreale, S. J., G. J. Ruiz, J. R. Spotila, and E. A. Standora.

1982. Temperature-dependent sex determination: Current practices threaten conservation of sea turtles. Science, vol. 216, p. 1245-1247.

Morris, B. F., J. Cadwallader, J. Geiselman, and J. N. Butler.

1976. Transfer of petroleum and biogenic hydrocarbons in the sargassum community, p. 235-239. In H. L. Windom and R. A. Duce (eds.), Marine pollutant transfer. Lexington Books, Lexington, Mass.

Morris, R. J.

1980a. Floating plastic debris in the Mediterranean. Marine Pollution Bulletin, vol. 11, p. 125.

Morris, R. J.

1980b. Plastic debris in the surface waters of the south Atlantic. Marine Pollution Bulletin, vol. 11, p. 164-166.

Mortimer, J. A.

1976. Observations on the feeding ecology of the green turtle, Chelonia mydas, in the western Caribbean. M.S. thesis, University of Florida, Gainesville. 100 p.

Mortimer, J. A.

1979. Influence of beach characteristics on nesting density, site fixity and hatching success of green turtles at Ascension Island. (Abstract). American Zoologist, vol. 19, p. 954.

Mortimer, J. A.

1981a. Reproductive ecology of the green turtle, Chelonia mydas, at Ascension Island. Ph.D. thesis, University of Florida, Gainesville. 163 p.

Mortimer, J. A.

1981b. The feeding ecology of the west Caribbean green turtle (Chelonia mydas) in Nicaragua. Biotropica, vol. 13, p. 49-58.

Mortimer, J. A.

1982a. Factors influencing beach selection by nesting sea turtles, p. 45-51. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Mortimer, J. A.

1982b. The feeding ecology of the sea turtles, p. 103-109. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Moulding, J. D.

1981. Sea turtles vs. dredging: A conflict with the endangered species act. (Abstract). Florida Scientist, vol. 44 (suppl. 1), p. 35.

Mrosovsky, N.

1968. Nocturnal emergence of hatchling turtles: Control by thermal inhibition of activity. Nature, vol. 220, p. 1338-1339.

Mrosovsky, N.

1977. Individual differences in the sea-finding mechanisms of hatchling leatherback turtles. Brain, Behavior and Evolution, vol. 14, no. 4, p. 261-274.

Mrosovsky, N.

1978. Orientation mechanisms of marine turtles, p. 413-419. In K. Schmidt-Koening and W. T. Keeton (eds.), Animal migration, navigation and homing. Springer-Verlag, New York.

Mrosovsky, N.

1980. Thermal biology of sea turtles. American Zoologist, vol. 20, p. 531-547.

Mrosovsky, N.

1981. Plastic jellyfish. Marine Turtle Newsletter, no. 17, p. 5-7.

Mrosvosky, N.

1982. Sex ratio bias in hatchling sea turtles from artificially incubated eggs. Biological Conservation, vol. 23, p. 309-314.

Mrosovsky, N., and S. J. Shettleworth.

1974. Further studies on the sea-finding mechanism in green turtle hatchlings. Behaviour, vol. 51, p. 195-208.

Mrosovsky, N., and C. L. Yntema.

1980. Temperature dependence of sexual differentiation in sea turtles: Implications for conservation practices. Biological Conservation, vol. 18, p. 271-280.

Nadeau, R. J., and E. T. Berquist.

1977. Effects of the March 18, 1973 oil spill near Cabo Rojo, Puerto Rico on tropical marine communities, p. 535-538. In Proceedings of the 1977 Oil Spill Conference, (prevention), behavior, control, cleanup). Sponsored by American Petroleum Institute, Environmental Protection Agency and United States Coast Guard, New Orleans, La.

National Marine Fisheries Service.

1978. Final environmental impact statement listing and protecting the green sea turtle (Chelonia mydas), loggerhead sea turtle (Caretta caretta), and Pacific ridley sea turtle (Lepidochelys olivacea) under the Endangered Species Act of 1973. U.S. National Marine Fisheries Service, Washington, D.C. 144 p.

Odum, H. T.

1963. Productivity measurements in Texas turtle grass and the effects of dredging an intracoastal channel. Publications of the Institute of Marine Science, University of Texas, vol. 9, p. 48-58.

Odum, W. E.

1970. Insidious alterations of the estuarine environment. Transactions of the American Fisheries Society, vol. 99, p. 836-847.

Ogden, J. C., L. Robinson, K. Whitlock, H. Daganhardt, and R. Cebula.

1983. Diel foraging patterns in juvenile green turtles (Chelonia mydas L.) in St. Croix United States Virgin Islands. Journal of Experimental Biology and Ecology, vol. 66, p. 199-205.

Ogren, L., and C. McVea.

1982. Apparent hibernation by sea turtles in north American waters, p.127-132. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Ogren, L. H., J. W. Watson, Jr., and D. A. Wickham.

1977. Loggerhead sea turtles, Caretta caretta, encountering shrimp trawls. U.S. National Marine Fisheries Service, Marine Fisheries Review, vol. 39, no. 11, p. 15-17.

O'Hara, J.

1980. Thermal influences on the swimming speed of loggerhead turtle hatchlings. Copeia, 1980, p. 773-780.

O'Hara, J., and H. J. Kania.

1981. Avoidance responses by sea turtles exposed to electrical fields. Prepared for Florida Power Light Company. Environmental Chemical Sciences, Atlanta, Ga. 32 p.

Owens, D. W.

1980. Introduction to the symposium: Behavioral and reproductive biology of sea turtles. American Zoologist, vol. 20, p. 485-486.

Owens, D. W., M. A. Grassman, and J. R. Hendrickson.

1982. The imprinting hypothesis and sea turtle reproduction. Herpetologica, vol. 38, p. 124-135.

Parker, P. L.

1962. Zinc in a Texas bay. Publications of the Institute of Marine Science, University of Texas, vol. 8, p. 75-79.

Parker, P. L.

1966. Movement of radioisotopes in a marine bay: Cobalt-60, iron-59 manganese-54, zinc-65, sodium-22. Publications of the Institute of Marine Science, University of Texas, vol. 11, p. 102-107.

Philibosian, R.

1976. Disorientation of hawksbill turtle hatchlings, Eretmochelys imbricata, by stadium lights. Copeia, 1976, p. 824.

Prange, H. D., and R. A. Ackerman.

1974. Oxygen consumption and mechanisms of gas exchange of green turtle (Chelonia mydas) eggs and hatchlings. Copeia, 1974, p. 758-763.

Pritchard, P. C. H.

1971. The leatherback or leathery turtle. International Union for Conservation of Nature and Natural Resources, Monograph no. 1. 39 p.

Pritchard, P. C. H.

1976a. Endangered species: Kemp's ridley turtle. Florida Naturalist, vol. 49, no. 3, p. 15-19.

Pritchard, P. C. H.

1976b. Post-nesting movements of marine turtles (Cheloniidae and Dermochelyidae) tagged in the Guianas. Copeia, 1976, p. 749-754.

Pritchard, P. C. H.

1980. The conservation of sea turtles: Practices and problems. American Zoologist, vol. 20, p. 609-617.

Pritchard, P. C. H., and R. Marquez.

1973. Kemp's ridley turtles or Atlantic ridley, Lepidochelys kempi. International Union for Conservation of Nature and Natural Resources, Monograph No. 2: Marine Turtle Series. 30 p.

Pritchard, P. C. H., and T. H. Stubbs.

1982. An evaluation of sea turtle populations and survival status on Vieques Island. Final Report to Department of the Navy, Naval Ocean Systems Center, San Diego, Calif. 92152, on contract no. N66001-80-C-0560 (including modification no. P00001). 102 p.

Rabalais, S. C., and N. N. Rabalais.

1980. The occurrence of sea turtles on the south Texas coast. Contributions in Marine Science, vol. 23, p. 123-129.

Rainey, W. E.

1978. Status of sea turtles at Vieques Island. Report to U.S. Navy Legal Department. 64 p.

Raymond, P. W.

1980. Marine turtle observations aboard dredge Long Island, Port Canaveral, Florida, 19 July-1 August 1980. Report to U.S. National Marine Fisheries Service. Unpublished report. [39 p.]

Rebel, T. P., editor.

1974. Sea turtles and the turtle industry of the West Indies, Florida, and the Gulf of Mexico. University of Miami Press, Coral Gables, Fla. 250 p.

Rhodin, A. G. H., and R. C. Schoelkopf.

1982. Reproductive data on a female leatherback turtle, Dermochelys coriacea, stranded in New Jersey. Copeia, 1982, p. 181-183.

Richardson, J. I., and H. O. Hillestad.

1978. Ecology of a loggerhead sea turtle population in Georgia. Georgia Department of Natural Resources, Game and Fish Division, Technical Bulletin WL 4, p. 22-37.

Richardson, J. I., and H. O. Hillestad.

1979. Wintering marine turtles in South Carolina and Georgia. Report to U.S. National Marine Service, Miami, Fla. 22 p.

Richardson, T. H., J. I. Richardson, C. Ruckdeschel, and M. W. Dix.

1978. Remigration patterns of loggerhead sea turtles (Caretta caretta) nesting on Little Cumberland and Cumberland Islands, Georgia. Florida Marine Research Publications, p. 53-55.

Riedman, S. R., and R. Witham.

1974. Turtles, extinction or survival? Abelard-Schuman, New York. 156 p.

Robles, A.

1976. The endangered hawksbill. Bulletin of the Chicago Herpetological Society, vol. 10, no. 3/4, p. 6-10.

Rogers, C.

1977. Response of a coral reef to sedimentation. Ph.D. thesis, University of Florida, Gainesville. 195 p.

Roithmayr, C.

1981. Sea turtle incidental catch and mortality project. Milestone report to Endangered Species Program, U.S. National Marine Fisheries Service, Southeast Fisheries Center, Miami, Fla. 33149.

Rose, W.

1950. The reptiles and amphibians of southern Africa. Maskew Miller, Cape Town, South Africa.

Royce, C., and R. Robertson.

1979. IXTOC I blowout draws massive control effort. Offshore, vol. 39, no. 9, p. 43-48.

Rudloe, J.

1981. From the jaws of death. Sports Illustrated, vol. 54, p. 60-70.

Schmidt, J.

1916. Marking experiments with turtles in the Danish West Indies. Meddelelser fra Kommissionen for Havundersogelser, Serie Fiskeri, vol. 5, no. 1, p. 1-26.

Schwartz, F. J.

1976. Status of sea turtles, Chelonidae and Dermochelidae, in North Carolina. Journal of the Elisha Mitchell Scientific Society, vol. 92, p. 76-77.

Schwartz, F. J.

1977. Species accounts, marine turtles, p. 303-308. In J. E. Cooper, S. S. Robinson and J. B. Funderburg (eds.), Endangered and threatened plants and animals of North Carolina. North Carolina State Museum of Natural History, Raleigh.

Schwartz, F. J.

1978. Behavioral and tolerance responses to cold water temperatures by three species of sea turtles (Reptilia, Cheloniidae) in North Carolina. Florida Marine Research Publications, no. 33, p. 16-18.

Schwartz, F. J.

1982. Correlation of nest sand asymmetry and percent loggerhead sea turtle hatch in North Carolina by geological sorting analyses. (Abstract). ASB (Association of Southeastern Biologists) Bulletin, vol. 29, p. 83.

Schwartz, F. J., C. Peterson, and H. Passingham.

1980. Consequences of natural and artificial incubation of sea turtle eggs laid in North Carolina. (Abstract). ASB (Association of Southeastern Biologists) Bulletin, vol. 27, p. 61.

Sella, I.

1982. The sea turtles in the eastern Mediterranean in danger of extinction, p. 417-423. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D.C.

Seyfert, F.

1978. The plight of the loggerhead. Sea Frontiers, vol. 24, p. 19-22.

Shabica, S. V.

1978. Sea turtle nesting at Cape Hatteras and Cape Lookout National Seashores, p. 16-23. In North Carolina Workshop on Sea Turtles, Marine Resources Center, Atlantic Beach, N.C., November 17, 1978.

Shabica, S. V.

1982. Planning for protection of sea turtle habitat, p. 513-518. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Shoop, C. R., and C. Ruckdeschel.

1982. Increasing turtle strandings in the southeast United States: A complicating factor. Biological Conservation, p. 213-215.

Smith, M. H., H. O. Hillestad, M. N. Manlove, D. O. Straney, and J. M. Dean.

1977. Management implications of genetic variability in loggerhead and green sea turtles. Transactions of the International Congress of Game Biologists, no. 13, p. 302-312.

Stancyk, S.

1982. Non-human predators on sea turtles and their control, p. 139-152. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Stancyk, S. E., O. R. Talbert, Jr., and J. M. Dean.

1980. Nesting activity of the loggerhead turtle Caretta caretta in S.C. II. Protection of nests from raccoon predation by transplantation. Biological Conservation, vol. 18, p. 289-298.

Stancyk, S. E., O. R. Talbert, Jr., and A. B. Miller.

1979. Estimation of loggerhead turtle nesting activity in South Carolina by aerial surveys. (Abstract). American Zoologist, vol. 19, p. 954.

Stoneburner, D. L., N. H. Micora, and E. R. Blood.

1980. Heavy metals in loggerhead sea turtle eggs Caretta caretta. Evidence to support the hypothesis that demes exist in the western Atlantic population. Journal of Herpetology, vol. 14, no. 2, p. 171-176.

Stoneburner, D. L., and J. I. Richardson.

1981. Observations on the role of temperature in loggerhead turtle nest site selection. Copeia, 1981, p. 238-241.

Talbert, O. R., Jr., S. E. Stancyk, J. M. Dean, and J. M. Will.

1980. Nesting activity of the loggerhead turtle (Caretta caretta) in South Carolina. I. A rookery in transition. Copeia, 1980, p. 709-718.

Taylor, J. L., and C. H. Saloman.

1968. Some of the effects of hydraulic dredging and coastal development in Boca Ciega Bay, Florida. U.S. Fish and Wildlife Service, Fishery Bulletin, vol. 67, p. 213-241.

Thayer, G. W., D. W. Engel and K. A. Bjorndal.

1982. Evidence for short-circuiting of the detritus cycle of seagrass beds by the green turtle, Chelonia mydas. L. Journal of Experimental Marine Biology and Ecology, vol. 62, p. 173-183.

Thayer, G. W., D. A. Wolfe, and R. B. Williams.

1975. The impact of man on seagrass systems. American Scientist, vol. 63, p. 288-296.

Thompson, N. P., P. W. Rankin, and D. W. Johnston.

1974. Polychlorinated biphenyls and p,p' DDE in green turtle eggs from Ascension Island, South Atlantic Ocean. Bulletin of Environmental Contamination and Toxicology, vol. 11, p. 399-406.

Thorhaug, A.

1981. Seagrasses in the Gulf of Mexico, Florida and Caribbean including restoration and mitigation of seagrasses, p. 161-178. In R. C. Carey and J. B. Kirkwood (eds.), Proceedings of U.S. Fish and Wildlife Service Workshop on Coastal Ecosystems of the Southeastern United States. U.S. Fish and Wildlife Service, Office of Biological Services, FWS/OBS-80/59.

Timko, R. E., and D. DeBlanc.

1981. Radio tracking juvenile marine turtles. U.S. National Marine Fisheries Service, Marine Fisheries Review, vol. 43, no. 3, p. 20-24.

Timko, R. E., and A. L. Kolz.

1982. Satellite sea turtle tracking. U.S. National Marine Fisheries Service, Marine Fisheries Review, vol. 44, no. 4, p. 19-24.

Torrey, L.

1980. Black tide from the Bay of Campeche. New Scientist, vol. 85, no. 1191, p. 243-245.

Towle, E. L.

1978. Report on sea turtle nesting, sighting, eggs and hatchlings for 1978 in the U.S. Virgin Islands and as recommended research methodology for dealing with hatchling disorientation on the beach (with specific reference to leatherback nests at Sandy Point St. Croix). Pursuant to NOAA/NMFS Purchase Order no. 01-8-D08-00187, submitted for Island Resources Foundation, St. Thomas, U.S. Virgin Islands. 39 p.

Travers, W. B., and P. R. Luney.

1976. Drilling, tankers, and oil spills on the Atlantic outer continental shelf. Science, vol. 194, p. 791-796.

Travis, W.

1967. The voice of the turtle. George Allen and Unwin, London. 203 p.

Ulrich, G. F.

1978. Incidental catch of loggerhead turtles by South Carolina commercial fisheries. Report to U.S. National Marine Fisheries Service, Contract nos. 03-7-042-35151 and 03-7-042-35121. 36 p.

Van Dolah, R. F., V. G. Burrell, Jr., and S. B. West.

1980. The distribution of pelagic tars and plastics in the South Atlantic Bight. Marine Pollution Bulletin, vol. 11, p. 352-356.

Van Rhijn, F. A.

1979. Optic orientation in hatchlings of the sea turtle, Chelonia mydas. II. Apparatus for prolonged recording of orientation in turtles walking or swimming on the spot. Marine Behavior and Physiology, vol. 6, p. 243-255.

Wade, T. L., J. G. Quinn, W. T. Lee, and C. W. Brown.

1976. Source and distribution of hydrocarbons in surface of the Sargasso Sea, p. 270-281. In Sources, effects and sinks of hydrocarbons in the aquatic environment. American Institute of Biological Sciences, Washington, D. C.

Wilcox, J. R.

1980. Endangered species: Compatability with a utility, p. 179-185. In B.L. Edge (ed.), Coastal Zone '80: Proceedings of the Second Symposium on Coastal and Ocean Management, vol. 1. American Society of Civil Engineers, New York.

Williams-Walls, N., J. O'Hara, R. M. Gallagher, D. F. Worth, B. D. Peery, and J. R. Wilcox.

1983. Spatial and temporal trends of sea turtle nesting on Hutchinson Island, Florida, 1971-1979. Bulletin of Marine Science, vol. 33, p. 55-66.

Witham, P. R.

1976. Evidence of ocean-current mediated dispersal in young green turtles, Chelonia mydas (Linnaeus). M.S. thesis, University of Oklahoma, Norman.

Witham, R.

1978. Does a problem exist relative to small sea turtles and oil spills, p. 629-632. In Proceedings of the Conference on Assessment of Ecological Impacts of Oil Spills. American Institute of Biological Sciences, Keystone, Colo.

Witham, R.

1980. The "lost year" question in young sea turtles. American Zoologist, vol. 20, p. 525-530.

Witham, R.

1982. Disruption of sea turtle habitat with emphasis on human influence, p. 519-522. In K. A. Bjorndal (ed.), Biology and conservation of sea turtles. Proceedings of World Conference on Sea Turtle Conservation. Smithsonian Institution Press, Washington, D. C.

Witkowski, S. A., and J. G. Frazier.

1982. Heavy metals in sea turtles. Marine Pollution Bulletin, vol. 13, 254-255.

Woodard, D. W.

1980a. Selected vertebrate endangered species of the seacoast of the United States - green sea turtle. U.S. Fish and Wildlife Service, Biological Services Program, FWS-OBS 80/01.13. 9 p.

Woodard, D. W.

1980b. Selected vertebrate endangered species of the seacoast of the United States - hawksbill turtle. U.S. Fish and Wildlife Service, Biological Services Program, FWS-OBS 80/01.22. 6 p.

Woodard, D. W.

1980c. Selected vertebrate endangered species of the seacoast of the United States - Kemp's (Atlantic) ridley sea turtle. U. S. Fish and Wildlife Service, Biological Services Program, FWS-OBS 80/01.30. 7 p.

Woodard, D. W.

1980d. Selected vertebrate endangered species of the seacoast of the United States - leatherback sea turtle. U.S. Fish and Wildlife Service, Biological Services Program, FWS-OBS 80/01.12. 7 p.

Worth, D. F., and J. B. Smith.

1976. Marine turtles nesting on Hutchinson Island, Florida, in 1973. Florida Marine Research Publications, no. 18. 17 p.

Yntema, C. L., and N. Mrosovsky.

1979. Incubation temperature and sex ratio in hatchling loggerhead turtles: A preliminary report. Marine Turtle Newsletter, no. 11, p. 9-10.

Yntema, C. L., and N. Mrosovsky.

1980. Sexual differentiation in hatchling loggerheads (Caretta caretta) incubated at different controlled temperatures. Herpetologica, vol. 36, p. 33-36.

Yntema, C. L., and N. Mrosovsky.

1982. Critical periods and pivotal temperatures for sexual differentiation in loggerhead sea turtles. Canadian Journal of Zoology, vol. 60, p. 1012-1016.

Zieman, J. C., Jr.

1970. The effects of a thermal effluent stress on the seagrasses and macroalgae in the vicinity of Turkey Point, Biscayne Bay, Florida. Ph.D. thesis, University of Miami, Coral Gables, Fla. 129 p.

Zieman, J. C., Jr.

1982. The ecology of the sea grasses of south Florida: A community profile. U.S. Fish and Wildlife Service, Biological Services Program, FWS/OBS-82/25 123. 26 p.

Zieman, J. C. and E. J. Ferguson Wood.

1975. Effects of thermal pollution on tropical-type estuaries, with emphasis on Biscayne Bay, Florida, p. 75-98. In E. J. Ferguson Wood and R. E. Johannes (eds.), Tropical marine pollution. Elsevier, New York.

Zieman, J.C., R. Orth, R. C. Phillips, G. Thayer, and A. Thorhaug.

[1981]. The effects of oil on seagrass ecosystems. Draft manuscript. Recovery and Restoration of Habitat Impacted by Oil Spills and/or Oil Spill Cleanup Workshop, Nov. 8-11, 1981, Virginia Polytechnic Institute and State University, Blacksburg.

Zwinenberg, A. J.

1974. The leatherback (Dermochelys coriacea), one of the largest living reptiles. Bulletin of the Maryland Herpetological Society, vol. 10, no. 2, p. 42-48.

Zwinenberg, A. J.

1975. The green turtle (Chelonia mydas), one of the reptiles most consumed by man, needs immediate protection. Bulletin of the Maryland Herpetological Society, vol. 11, no. 2, p. 45-63.

Zwinenberg, A. J.

1977. Kemp's ridley, Lepidochelys kempii (Garman 1880), undoubtedly the most endangered marine turtle today (with notes on the current status of Lepidochelys olivacea. Bulletin of the Maryland Herpetological Society, vol. 13, 170-192.