

file copy

NOAA Technical Memorandum NMFS

DECEMBER 1990

PROCEEDINGS OF THE SECOND INTERNATIONAL
CONFERENCE ON MARINE DEBRIS
2-7 APRIL 1989, HONOLULU, HAWAII

VOLUME I

Richard S. Shomura
Mary Lynne Godfrey
(Editors)



NOAA-TM-NMFS-SWFSC-154

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southwest Fisheries Science Center
University of Hawaii Sea Grant College Program

NOAA Technical Memorandum NMFS

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.

NOAA Technical Memorandum NMFS

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information; and have not received complete formal review, editorial control, or detailed editing.

DECEMBER 1990

PROCEEDINGS OF THE SECOND INTERNATIONAL CONFERENCE ON MARINE DEBRIS 2-7 APRIL 1989, HONOLULU, HAWAII

Richard S. Shomura and Mary Lynne Godfrey (Editors)
University of Hawaii
School of Ocean and Earth Science and Technology
Hawaii Institute of Marine Biology
1000 Pope Road
Honolulu, Hawaii 96822

Canada Department of Fisheries and Oceans
Council for Solid Waste Solutions
Intergovernmental Oceanographic Commission (UNESCO)
National Coastal Resources Research and Development Institute (U.S.A.)
Pacific Rim Fishing Industries
Sea Grant College Program, University of Hawaii
SUNY, Stony Brook, Marine Sciences Research Center's
Waste Management Institute
School of Ocean and Earth Science and Technology, University of Hawaii
U.S. Environmental Protection Agency
U.S. Fish and Wildlife Service
U.S. Marine Mammal Commission
U.S. Minerals Management Service
U.S. National Oceanic and Atmospheric Administration
U.S. Navy

NOAA-TM-NMFS-SWFSC-154

U.S. DEPARTMENT OF COMMERCE

Robert A. Mosbacher, Secretary

National Oceanic and Atmospheric Administration

John A. Knauss, Under Secretary for Oceans and Atmosphere

National Marine Fisheries Service

William W. Fox, Jr., Assistant Administrator for Fisheries

1

PREFACE

The Second International Conference on Marine Debris is a sequel to the Workshop on the Fate and Impact of Marine Debris which was held in Honolulu, Hawaii, 26-29 November 1984. The workshop was the first meeting of its kind to address the issue of marine debris as an international problem, and focused primarily on the impact of debris on the living marine resources. This second meeting provides a status review of wider dimensions, including the source, impact, economics, technology, law and policy, and educational aspects of marine debris and its impact on society as well as on the resources of the sea. The details leading to this second meeting and the general conclusions drawn from this meeting are provided in the Executive Summary.

The size and scope of the Second International Conference on Marine Debris make it necessary to publish the proceedings in two volumes. The first volume includes the Executive Summary, the full text of 5 overview papers, and the full text of 40 papers presented in the technical sessions on sources (Session I), entanglement and ghost fishing (Session II), and ingestion (Session III). The second volume includes the text of 31 papers presented in the technical sessions on economics (Session IV), technology (Session V), law and policy (Session VI), and education (Session VII). Additionally, volume two includes the reports of the eight working groups (assessment, entanglement, ghost fishing, ingestion, economics, technology, law and policy, and education) and abstracts of the poster and video presentations. All technical papers were reviewed by one or more referees. Reference to several of the papers is by abstract only, the full text of these presentations not being submitted in time for publication or scheduled for publication elsewhere.

The Appendixes comprise the membership of the steering group, the agenda of the conference, and a list of conference participants.

The senior editor, who had the task of organizing this Second International Conference on Marine Debris, is indebted to many organizations and individuals for helping to make this conference a success. Special thanks are due the members of the steering group for their guidance and assistance in the technical aspects of the conference, and Ms. Christine Woolaway of the University of Hawaii Sea Grant College Program for her outstanding effort in working out the logistics and conference arrangements. Sponsors who provided financial and personnel support included (1) Canada Department of Fisheries and Oceans, (2) Council for Solid Waste Solutions, (3) Intergovernmental Oceanographic Commission (UNESCO), (4) National Coastal Resources Research and Development Institute (U.S.A.), (5) Pacific Rim Fishing Industries, (6) Sea Grant College Programs (Washington, D.C., and the University of Hawaii), (7) State University of New York (SUNY), Stony Brook, Marine Sciences Research Center's Waste Management Institute, (8) School of Ocean and Earth Science and Technology (University of Hawaii), (9) U.S. Environmental Protection Agency, (10) U.S. Fish and Wildlife Service, (11) U.S. Marine Mammal Commission, (12) U.S. Minerals Management Service, (13) U.S. National Oceanic and Atmospheric Administration (NOAA) (National Marine Fisheries Service Honolulu Laboratory), and (14) U.S. Navy.

Ms. Suzanne Montgomery of Washington Communications Service, 150 N. Muhlenberg Street, Woodstock, Virginia, prepared the Executive Summary.

Reference to trade names in the Proceedings of the Second International Conference on Marine Debris does not imply endorsement by the National Marine Fisheries Service, NOAA.

This proceedings is also a University of Hawaii Sea Grant College Program cooperative report, UNIHI-SEAGRANT-CR-91-02.

Finally, the editors owe the successful completion of these proceedings to two former employees of the Honolulu Laboratory, Ms. Louise Brewer and Ms. Elizabeth Young, who assumed the tremendous task of typing and checking the conference manuscripts.

TABLE OF CONTENTS

	Page
EXECUTIVE SUMMARY	3
OVERVIEW PAPERS	17
J. M. Coe	19
A review of marine debris research, education, and mitigation in the North Pacific	
B. Heneman	50
Overview: Marine debris in the northwest Atlantic Ocean	
M. R. Gregory	55
Plastics: Accumulation, distribution, and environmental effects of meso-, macro-, and megalitter in surface waters and on shores of the southwest Pacific	
P. G. Ryan	85
The marine plastic debris problem off southern Africa: Types of debris, their environmental effects, and control measures	
M. F. Tillman	103
International efforts to control marine debris in the Antarctic	
J. E. Corredor and J. M. Morell	115
Cooperative research on petroleum pollution in the Caribbean: The CARIPOL program [Abstract]	
SESSION I. AMOUNTS, TYPES, DISTRIBUTION, AND SOURCES OF MARINE DEBRIS	117
J. Cantin, J. Eyraud, and C. Fenton	119
Quantitative estimates of garbage generation and disposal in the U.S. maritime sectors before and after MARPOL Annex V	
R. H. Day, D. G. Shaw, and S. E. Ignell	182
The quantitative distribution and characteristics of marine debris in the North Pacific Ocean, 1984-88	
S. Mio, S. Takehama, and S. Matsumura	212
Distribution and density of floating objects in the North Pacific based on 1987 sighting survey	
R. H. Day, D. G. Shaw, and S. E. Ignell	247
The quantitative distribution and characteristics of neuston plastic in the North Pacific Ocean, 1984-88	

S. Matsumura, Y. Wakata, and Y. Sugimori	267
Movements of floating debris in the North Pacific	
J. A. June	279
Type, source, and abundance of trawl-caught marine debris off Oregon, in the eastern Bering Sea, and in Norton Sound in 1988	
C. A. Ribic and L. J. Bledsoe	302
Estimating the density of floating marine debris: Design considerations	
W. R. Trulli, H. K. Trulli, and D. P. Redford	309
Characterization of marine debris in selected harbors of the United States	
M. Yukinawa and S. Mio	325
Preliminary report on the distribution of small-sized marine debris in Suruga Bay	
S. W. Johnson	331
Distribution, abundance, and source of entanglement debris and other plastics on Alaskan beaches, 1982-88	
A. M. Manville II	349
A survey of plastics on western Aleutian Island beaches and related wildlife entanglement	
R. H. Podolsky	364
Entrapment of sea-deposited plastic on the shore of a Gulf of Maine island [Abstract]	
R. L. Swanson and R. Zimmer	365
Washups of floatable waste materials and their impact on New York Bight beaches [Abstract]	
K. V. Koski	366
Plastic debris and derelict fishing gear on Shackleford Banks, North Carolina [Abstract]	
A. F. Amos and P. T. Plotkin	367
Anthropogenic and natural debris on a south Texas barrier island beach [Abstract]	
A. Golik and Y. Gertner	369
Solid waste on the Israeli coast--composition, sources, and management	
K. J. O'Hara	379
National marine debris data base: Findings on beach debris reported by citizens	

C. A. Ribic and S. W. Johnson	392
Guidelines for the design of beach debris surveys	
D. J. Slip and H. R. Burton	403
The composition and origin of marine debris stranded on the shores of subantarctic Macquarie Island	
SESSION II. ENTANGLEMENT OF MARINE LIFE AND GHOST FISHING	417
N. Baba, M. Kiyota, and K. Yoshida	419
Distribution of marine debris and northern fur seals in the eastern Bering Sea	
D. P. French and M. Reed	431
Potential impact of entanglement in marine debris on the population dynamics of the northern fur seal, <i>Callorhinus ursinus</i>	
C. W. Fowler, R. Merrick, and J. D. Baker	453
Studies of the population level effects of entanglement on northern fur seals	
A. E. Kuzin	475
A study of the effects of commercial fishing debris on <i>Callorhinus ursinus</i> from breeding islands in the western Pacific	
C. A. Ribic and G. L. Swartzman	483
An index of fur seal entanglement in floating net fragments	
R. L. DeLong, P. J. Gearin, J. L. Bengtson, P. Dawson, and S. D. Feldkamp	492
Studies of the effects of entanglement on individual northern fur seals [Abstract]	
K. Yoshida, N. Baba, M. Kiyota, M. Nakajima, Y. Fujimaki, and A. Furuta	494
Studies of the effects of net fragment entanglement on northern fur seals. Part 1: Daily activity patterns of entangled and nonentangled fur seals	
K. Yoshida, N. Baba, M. Kiyota, M. Nakajima, Y. Fujimaki, and A. Furuta	503
Studies of the effects of net fragment entanglement on northern fur seals. Part 2: Swimming behavior of entangled and nonentangled fur seals	
G. L. Swartzman, C. A. Ribic, and C. P. Huang	513
Simulating the role of entanglement in northern fur seal, <i>Callorhinus ursinus</i> , population dynamics	

M. Nakajima, H. Yasuda, and K. Yoshida	531
Histological observation of damage to dermal tissue of fur seal caused by net entanglement	
J. R. Henderson	540
Recent entanglements of Hawaiian monk seals in marine debris	
B. S. Stewart and P. K. Yochem	554
Pinniped entanglement in synthetic materials in the Southern California Bight	
S. S. Sadove and S. J. Morreale	562
Marine mammal and sea turtle encounters with marine debris in the New York Bight and the northeast Atlantic	
P. A. Breen	571
A review of ghost fishing by traps and gillnets	
T. Gerrodette, B. K. Choy, and L. M. Hiruki	600
An experimental study of derelict gillnet fragments in the central Pacific Ocean	
S. Mio, T. Domon, K. Yoshida, and S. Matsumura	615
Preliminary study on change in shape of drifting nets experimentally placed in the sea	
SESSION III. INGESTION BY MARINE LIFE	621
P. G. Ryan	623
The effects of ingested plastic and other marine debris on seabirds	
H. Ogi	635
Ingestion of plastic particles by sooty and short-tailed shearwaters in the North Pacific	
D. G. Ainley, L. B. Spear, and C. A. Ribic	653
The incidence of plastic in the diets of pelagic seabirds in the eastern equatorial Pacific region	
L. Sileo, P. R. Sievert, M. D. Samuel, and S. I. Fefer	665
Prevalence and characteristics of plastic ingested by Hawaiian seabirds	
D. G. Ainley, W. R. Fraser, and L. B. Spear	682
The incidence of plastic in the diets of Antarctic seabirds	
R. H. Podolsky and S. W. Kress	692
Plastic debris incorporated into double-crested cormorant nests in the Gulf of Maine [Abstract]	


D. E. Hoss and L. R. Settle	693
Ingestion of plastics by teleost fishes	
T. Kubota	710
Synthetic materials found in the stomachs of longnose lancetfish collected from Suruga Bay, central Japan	
G. H. Balazs and B. K. Choy	718
Ecological aspects of marine turtles impacted by ocean debris: A 1989 perspective [Abstract]	
P. L. Lutz	719
Studies on the ingestion of plastic and latex by sea turtles	
P. Plotkin and A. F. Amos	736
Effects of anthropogenic debris on sea turtles in the northwestern Gulf of Mexico	
I. Uchida	744
On the synthetic materials found in the digestive systems of, and discharged by, sea turtles collected in waters adjacent to Japan [Abstract]	
R. J. Tarpley	745
Plastic ingestion in a pygmy sperm whale, <i>Kogia</i> <i>breviceps</i> [Abstract]	
N. B. Barros, D. K. Odell, and G. W. Patton	746
Ingestion of plastic debris by stranded marine mammals from Florida [Abstract]	
W. A. Walker and J. M. Coe	747
Survey of marine debris ingestion by odontocete cetaceans	
 SESSION IV. ECONOMIC IMPACTS ON VESSELS AND SHORELINES	 775
N. F. Meade, K. M. Drazek, and V. R. Leeworthy	777
An economic perspective on the problem of marine debris	
S. Takehama	792
Estimation of damages to fishing vessels caused by marine debris, based on insurance statistics	
R. E. Weisbrod	810
New York State marine debris program [Abstract]	
K. D. Wagner	811
Medical wastes and the beach washups of 1988: Issues and impacts	

SESSION V. SOLUTIONS THROUGH TECHNOLOGY	825
G. Scott	827
The philosophy and practice of degradable plastics	
A. L. Andrady	848
Environmental degradation of plastics under land and marine exposure conditions	
M. Xanthos, P. G. Kelleher, A. Patel, and W. G. Gordon	870
Recycling of marine plastics debris through melt reprocessing: A case for lost or abandoned fishing gear	
C. S. Alig, L. Koss, T. Scarano, and F. Chitty	879
Control of plastic wastes aboard naval ships at sea	
L. A. Martinez	895
Shipboard waste disposal: Taking out the trash under the new rules	
T. J. Chang	915
Low technology (burn barrel) disposal of shipboard-generated (MARPOL V) wastes	
F. Recht and S. Lasseigne	921
Providing refuse reception facilities and more: The port's role in the marine debris solution	
M. Aizawa and A. Satou	935
Disposition and recycling of plastic products including used nets	
 SESSION VI. SOLUTIONS THROUGH LAW AND POLICY	 945
D. P. Redford	947
Status of the U.S. Environmental Protection Agency marine debris activities and programs	
D. T. Edwards and E. Rymarz	956
International regulations for the prevention and control of pollution by debris from ships	
M. J. Bean	989
Redressing the problem of persistent marine debris through law and public policy: Opportunities and pitfalls	
R. E. Seebald	998
Implementation and enforcement of Annex V of MARPOL 73/78 in the United States [Abstract]	
C. A. Crampton	999
The Coast Guard's Annex V compliance report: A case study	


J. Sawaki and I. Yamaguchi	1008
Control regulations for discharge of onboard wastes from ships	
R. Rose	1020
Marine plastic debris: What Washington State has done	
G. Peet	1034
The protection of specific sea areas against marine debris	
L. M. Paul	1045
Using the protective principle to unilaterally enforce transnational marine pollution standards	
 SESSION VII. SOLUTIONS THROUGH EDUCATION	 1075
R. G. Bruner	1077
The plastics industry and marine debris: Solutions through education	
T. R. Dixon	1090
MARPOL 73/78 information, education, and training: Meeting the challenge	
P. Debenham	1100
Education and awareness: Keys to solving the marine debris problem	
B. Wallace	1115
Shipping industry marine debris education plan	
N. Yagi and Y. Otsuka	1123
Cleanup program in Japan	
L. Koss, F. Chitty, and W. A. Bailey	1132
U.S. Navy's plastics waste educational efforts	
B. Wallace	1140
How much do commercial and recreational fishermen know about marine debris and entanglement? Phase 1	
J. Neilson	1149
The Oregon experience--four years later	
A. Farias	1154
The Texas Adopt-A-Beach Program: A public/private approach to clean beaches	
J. Humphreys and P. Mullin	1165
Marine debris demonstration and education project at Squalicum Harbor, Bellingham, Washington, U.S.A.	

S. Friday	1174
Marine debris: North Carolina's solutions through education	
S. Laska	1179
Designing effective educational programs: The attitudinal basis of marine littering	
 POSTER PRESENTATION	 1191
G. J. Vauk and E. Schrey	1193
Animals as litter victims at the German North Sea coast	
J. N. Kushima and R. P. Clarke	1194
Marine debris and epipelagic fishes	
 VIDEO PRESENTATION	 1195
A. F. Amos	1197
Portrait of a barrier island beach	
Offshore Operators Committee	1198
U.S. oil industry efforts in addressing beach debris problem	
Government of Japan	
Plastic pollution: The sea is crying	
Government of Japan	
Cleanup operation of the sea floor	
Government of Japan	
Japanese research activities on marine debris	
 WORKING GROUP REPORTS	 1199
Report of the working group on methods to	1201
assess the amount and types of marine debris (C. A. Ribic, Chair)	
Report of the working group on entanglement	1207
of marine life (W. R. P. Bourne, Chair)	
Report of the working group on ghost fishing	1216
(P. A. Breen, Chair)	
Report of the working group on ingestion	1226
(L. Sileo, Chair)	

Report of the working group on economic aspects of marine debris (K. E. McConnell, Chair)	1235
Report of the working group on technology (W. G. Gordon, Chair)	1240
Report of the working group on law and policy (D. Cottingham, Chair)	1249
Report of the working group on marine debris education (K. J. O'Hara, Chair)	1256
APPENDIXES	1261
Appendix A. Steering group	1263
Appendix B. Agenda of the workshop	1264
Appendix C. List of participants	1265



EXECUTIVE SUMMARY



EXECUTIVE SUMMARY

I. INTRODUCTION

Until 10 or 15 years ago, the presence of debris in ocean and coastal areas was not recognized as a significant marine pollution issue. By the early 1980's, however, it became apparent that the amount of debris accumulating at sea and on beaches was increasing dramatically. There was also a corresponding increase in the incidence of marine species being adversely affected by ocean debris. These included marine mammals, seabirds and commercially valuable species of fish killed and injured in lost or discarded fishing gear and other debris, as well as beach-cast turtles with their digestive tracts blocked by plastic items. Clearly, marine debris was becoming a widespread marine pollution problem that could no longer be ignored.

In November 1984, the National Marine Fisheries Service, at the recommendation and with the assistance of the Marine Mammal Commission, convened a Workshop on the Fate and Impact of Marine Debris. The workshop was the first meeting ever undertaken to comprehensively assess information on the amounts, distribution, sources, effects, and management needs pertaining to problems of trash and other human-related debris lost or discarded into the ocean.

Participants in that conference concluded that many marine organisms throughout the world were being affected adversely by persistent debris and that there was an urgent need to: (1) educate vessel operators, fishermen and the public on the problem; (2) stop or reduce the deliberate disposal of persistent materials; and (3) obtain better quantitative data to assess the impact of marine debris on living organisms.

As a result of the workshop findings, a number of programs were undertaken in the United States and elsewhere to address the problem of marine debris. In view of the progress being made, in December 1986, the Marine Mammal Commission recommended to the National Marine Fisheries Service that it initiate planning for a second conference to review the marine debris issue. The Service agreed with this recommendation and, in fiscal year 1988, provided funds to begin planning and organizing a conference. In March 1988 the Marine Entanglement Research Program established a steering group to organize an international workshop. The Second International Conference on Marine Debris took place 2-7 April 1989 at the Ala Moana Americana Hotel in Honolulu, Hawaii.

Objectives

The objectives of the conference, as defined by the steering group, were to:

- evaluate new information on the types, amounts, sources, fates, and distribution of marine debris in different ocean areas;

- evaluate what has been done in the North Pacific basin as a prototype for activities that might be usefully undertaken in other regions;
- identify and evaluate existing and potential methods for gathering data on and monitoring trends in the sources, types, fates, amounts and distribution of debris at sea and on beaches;
- identify and evaluate information on the nature and extent of marine debris-related impacts on species and populations of marine life, including seals, turtles, seabirds, crustaceans, and fish, in different ocean areas;
- identify and evaluate the impacts of marine debris on human health and the safety of ships at sea;
- identify and evaluate aesthetic and other impacts of marine debris on coastal environment;
- review and evaluate information on existing and potential technological and procedural ways to reduce or eliminate the problem of marine debris;
- assess the effectiveness and future role of programs to educate the public and promote awareness of the problem;
- evaluate international, intergovernmental, domestic, and informally constituted regional authorities that might be usefully drawn upon to strengthen cooperative efforts to address regional issue;
- describe programs necessary to assess the effectiveness of measures presently being taken to address various elements of the problem; and
- prepare a report summarizing the results of the conference and the steps that should be taken to address different aspects of the problem.

Workshop Organization

The conference opened Monday, 3 April, with a keynote address and a plenary session that set the stage for discussions during the remainder of the week. Overview papers presented during the plenary described the marine debris issue in six geographic areas: the North Pacific, the north-west Atlantic; the southwest Pacific; waters off southern Africa; the Antarctic; and the Caribbean. During the next 4 days, background and experience papers were presented on aspects of the marine debris problem. The subject areas of these technical sessions included: (1) amounts, types, distribution and sources of marine debris; (2) entanglement of marine life and ghost fishing; (3) ingestion by marine life; (4) economic impacts on

vessels and shorelines; (5) solutions through technology; (6) solutions through law and policy; and (7) solutions through education. In some cases, these technical sessions ran concurrently. Beginning on Wednesday, 5 April, participants separated into eight working groups to discuss the results of the technical sessions and formulate recommendations on needed actions. The subject matter of the working groups mirrored those of the technical sessions except that entanglement and ghost fishing split into two working groups. At a final plenary session on Friday, 7 April, working group chairmen summarized the results of these deliberations for workshop participants as a whole. A conference summary was presented at the closing luncheon.

Sponsors and Participants

Sponsors of the workshop included: Canada Department of Fisheries and Oceans; Council for Solid Waste Solutions; Intergovernmental Oceanographic Commission (UNESCO); National Coastal Resources Research and Development Institute (U.S.A.); Pacific Rim Fishing Industries; Sea Grant Colleges--University of Hawaii; State University of New York at Stony Brook, Marine Sciences Research Center's Waste Management Institute; University of Hawaii, School of Ocean and Earth Science and Technology; U.S. Environmental Protection Agency; U.S. Fish and Wildlife Service; U.S. Marine Mammal Commission; U.S. Minerals Management Service; U.S. National Oceanic and Atmospheric Administration; and the U.S. Navy.

As at the 1984 conference, participants included representatives from these groups along with scientists from various disciplines, administration and management personnel from Federal and State offices, and representatives of the fishing industry, the academic community, conservation groups, and other public and private interests. Although participants were primarily from the United States, representatives from the Republic of Korea, Japan, New Zealand, Canada, Israel, The Netherlands, South Africa, and the United Kingdom were also present. This level of participation is indicative of the high degree of international interest in the problems of marine debris.

II. SUMMARY OF OVERVIEW SESSION (chaired by Brian Boyle)

Several papers were presented addressing the marine debris situation in various parts of the world. Activities were reported under way to reduce the impact of marine debris on North Pacific seabirds and marine mammals. Effects of these actions are as yet unknown. Impacts of marine debris in the northwest Atlantic were also discussed, and it was concluded that aesthetic degradation of beaches and the cost of cleaning beaches appear to be the most serious effects of marine debris in the study area. Threat to marine life appears to be limited to sea turtles. Addressing accumulation, distribution, and environmental effects of plastic pollution in the southwest Pacific, it was noted that plastic debris of all kinds and in all sizes is widespread in this region, including shores of isolated and unpopulated islands. Ingestion of plastics has been recorded for at least 7 species of mammals, 26 seabird species, and 2 marine turtles species in

the waters off southern Africa and the adjacent Southern Ocean. In addition, 5 marine mammal species and 13 seabird species have been found entangled in plastic debris. Steps are being taken by the Commission for the Conservation of Antarctic Marine Living Resources to monitor marine debris in the Antarctic. Assessment of petroleum hydrocarbons in the marine environment of the Caribbean was also discussed.

III. SUMMARY OF TECHNICAL SESSIONS

Session I: Amounts, Types, Distribution, and Sources of Marine Debris (chaired by Murray Gregory and Satsuki Matsumura)

Debris is found in all oceans in all parts of the world. Regional evaluations of the amounts, types, distributions, and sources of persistent debris are required to develop efficient strategies for its control. Such evaluations are also critical to the discovery of current and potential problems caused by marine debris. This session included 22 papers, including several on concentrations of marine debris in the North Pacific/Bering Sea areas and others addressing the debris problems in the Gulf of Maine, New York Bight, the Outer Banks of North Carolina, a Texas barrier island, the east Mediterranean, the Israeli coast, and elsewhere. Papers also discussed the National Marine Debris Data Base and suggested guidelines for the design of beach debris surveys.

Session II: Entanglement of Marine Life (chaired by Charles Fowler)

The destruction of marine organisms through encounters with synthetic debris has been widely reported, and entanglement is the most common mechanism for this destruction. While the consequences of entanglement are obvious for individual animals, the implications for the status of the populations involved have been difficult to ascertain. Sixteen papers were presented on these topics, including ten on the northern fur seal and one each on Hawaiian monk seal, pinnipeds in the Southern California Bight, and marine mammals and sea turtles in the New York Bight. Presentations also addressed ghost fishing and the preliminary results of a study on the impact of the changing shape of derelict driftnets.

Session III: Ingestion by Marine Life (chaired by Peter Lutz)

Ingestion of plastic bags, synthetic rope, plastic pellets, and other marine debris has been reported as the cause of death of individual sea turtles, seabirds, marine mammals, and fish. The extent to which such incidents occur is uncertain, and thus it is also uncertain whether such occurrences have negative impact on population levels. This session included 15 papers presenting the latest scientific findings on research on debris ingestion. Six papers addressed incidence and effects of ingestion by seabirds, four focused on ingestion of plastics by sea turtles, and others addressed ingestion by fishes and marine mammals.

**Session IV: Economic Impacts on Vessels and Shorelines
(chaired by John Sutinen)**

Netting, rope and sheeting discarded at sea can disable vessels, thus threatening human safety on the open water. Floating debris has a negative aesthetic impact on beaches and inshore waters and can pose a human health hazard. Coastal communities that depend on tourism can incur significant costs as a result of decreased tourist traffic and cleanup costs. An economic perspective on the problem of persistent marine debris addressed problems in enforcing regulations to prevent marine debris. A research agenda was proposed. One report described how the Japanese commercial fishing fleet suffered damages as a result of marine debris. Other papers addressed recent incidents of medical wastes washing up on U.S. Atlantic beaches and the New York State Marine Debris Program.

Session V: Solutions Through Technology (chaired by Bruce Perlson)

A vast range of useful applications of new and modified technology is available for reducing the marine debris problem. Areas ripe for advancement include: simplification of shipboard waste handling, control of land-based sources, port waste handling systems, plastics and other recycling systems, waste heat recovery; fishing gear loss and recovery, and controlled-lifetime plastics. There is apparently a changing attitude in the packaging industry toward the development of biodegradable plastics. A paper described results of studies of the weathering behavior of five types of plastic materials commonly found as marine debris, and another reported on a recently initiated project to encourage recycling of marine plastic debris. Other papers in this session addressed control and disposal of wastes aboard ships and the port's role in reducing marine debris by providing refuse reception facilities.

Session VI: Solutions Through Law and Policy (chaired by Timothy Keeney)

Solutions to the marine debris problem involve many disciplines, industries, institutions, and agencies of government. This session focused on models of public process that have dealt with fractionated leadership and authority structures in solving the marine debris and similar multidimensional problems. The U.S. Environmental Protection Agency's Interim National Coastal and Marine Policy, which is aimed at controlling medical wastes and other marine refuse, was described. Another paper discussed international and regional regulations to prevent and control pollution by ships and noted, for example, that a recent survey of the German Bight estimated that 95% of the 8.5 million pieces of debris dumped annually, come from ships. Five papers addressed aspects of MARPOL Annex V and its potential for reducing marine pollution, and the Marine Plastic Debris Action Plan adopted in the State of Washington was described.

Session VII: Solutions Through Education (chaired by Bernard Griswold)

Because a great deal of the persistent debris reaching the ocean is the result of actions by individuals, public education on impacts of debris and disposal alternatives is seen as an important factor in solving the

problem. Papers in this session represented a wide range of educational programs and materials in place throughout the world, and one discussed how the plastics industry has responded to the problem of marine pollution, specifically to the presence of resin pellets in the marine environment. Efforts being carried out by The Tidy Britain Group to control marine litter were documented. Other authors discussed marine education and cleanup programs being implemented by the shipping industry, the U.S. Navy, various states and localities, and Japan. The final paper pointed out the need to consider and understand public attitudes and perceptions when designing an education program.

IV. SUMMARY OF WORKING GROUP MEETINGS

Working Group 1: Methods to Assess the Amounts and Types of Marine Debris (chaired by Christine Ribic)

The working group reviewed various methods currently being used to survey debris at sea, on beaches, on the sea floor, and emanating from land. Participants agreed that certain areas should be selected for concentrated study: on an international level, MARPOL special areas were suggested as appropriate; on national or regional levels, areas should be chosen to meet local areas of interest or management.

The group proposed methodologies to be used to determine amounts of debris. For nearshore, open ocean and sea bottom areas, platforms of opportunity, or dedicated surveys are appropriate. Beach surveys could be done using either designed programs or volunteer programs.

To improve accuracy and usefulness of strip transect surveys used to assess floating debris, the group recommended: using two or three observers instead of a single observer; calibration runs to estimate strip width; and experiments to investigate color and size biases. The working group further recommended that, when accurate distance measurements can be made, line transect methodology should be used. It also noted the possibility of using low-flying aircraft to survey nearshore areas.

As regards the magnitude of bottom debris, the group identified fishermen as a potential source of baseline information. It recommended that a survey form be developed to collect information from fishermen on debris tangled in their nets. One suggestion was made to classify all debris in one of four size categories ranging from "mega" (>2-3 dm) to "micro" (powdered or unseen in general).

The working group attempted to list certain debris types for purposes of recordkeeping, such as: nets (by type); other fishing gear; strapping bands (open/closed) (cut/not cut); granulated plastic (recycled plastic); particulate plastic; fragmented plastic; plastic bags; plastic containers (country of origin, age); Styrofoam; medical wastes; rope; entanglement remains (e.g., bones). The group recommended that other lists be reviewed to develop a common list that can be tailored to individual areas.

The working group agreed that the technique currently appropriate for assessment studies on a large scale was the beach survey. On a limited scale, dedicated surveys using visual observations and neuston tows in the nearshore areas (e.g., bays, harbors) or limited ocean areas (e.g., off-shore dumping areas) could also be used for assessment. Techniques using aircraft are experimental and could probably be used for baseline studies. Bottom debris studies are currently in the baseline category. Development of techniques to study bottom debris is needed.

Most members of the working group agreed that a procedures manual should be compiled for use as a starting point for people interested in initiating marine debris studies.

Working Group 2: Entanglement of Marine Life (chaired by W. R. P. Bourne)

The working group found that there is accumulating anecdotal evidence that virtually all marine animals are occasionally entangled in debris, but that quantitative data are available for only a few species. Care is needed in the interpretation of available information because it is difficult to distinguish between the effects of marine debris and other factors such as oceanic fluctuations, disturbance of animals while breeding, the impact of fishing on both animals and their food supply, disease, and other forms of pollution.

The group found that entanglement of cetaceans appears to be infrequent, but even small numbers may represent a serious impact on the reduced population of North Atlantic right whales. Phocid seals are occasionally entangled in netting but the incidence is not high. One phocid seal for which the entanglement problem appears to be most significant is the endangered Hawaiian monk seal. Otariid seals appear to be among the marine species most prone to entanglement, young animals being particularly vulnerable. As regards marine turtles and seabirds, the group concluded that entanglement does occur but that there is little if any evidence of any impact on numbers when compared to other factors such as loss of habitat or incidental take in certain fisheries.

The working group concluded that, in view of the number of problems that require investigation and the wide area to be covered, there is need to establish an international organization to coordinate and standardize systematic collection and dissemination of information about the occurrence and impact of marine debris and possible conservation measures to mitigate its impact.

The group further recommended, among other things, that efforts be continued to monitor, remove, and destroy lost or discarded nets and other debris presenting a hazard to monk seals, marine turtles, and other wildlife in the Northwestern Hawaiian Islands; and that monitoring be continued on the numbers, survival, breeding success, and incidence of entanglement of northern fur seals. It also recommended the investigation of the impact of entanglement and other possible hazards on right whales in the northwest Atlantic and Kemp's ridley turtles in the Caribbean, and a review of the long-term evidence for the incidence of entanglement provided

by bird-banding and beach surveys. Continued analysis of population level effects of entanglement through simulation modeling are to be encouraged.

Working Group 3: Ghost Fishing (chaired by Paul Breen)

The group concluded that ghost fishing is a potentially serious problem because of the very large volume of fishing gear now in use and the increasing use of nondegradable materials such as plastic, vinyl-coated wire, and fiberglass. Traps and gillnets were seen as the primary cause of ghost fishing problems, with trawl and longline gear types presenting a lesser problem. The group concluded that mitigation of ghost fishing by traps is technologically simple, but that ghost fishing by nets may be more difficult to solve. It recommended that both time-failure devices and degradable meshes be developed and tested.

The group agreed upon a series of recommendations classified by priority. High priority was placed on the following four proposals:

1. Fishery agencies responsible for trap and tanglenet fisheries should conduct lost gear simulations to determine whether ghost fishing occurs and, if so, the rate at which target and nontarget species are killed. If ghost fishing is found to be a problem, the rates of gear loss should be estimated through logbook programs or questionnaires. Useful information might be obtained from surveys of manufacturers.
2. Where ghost fishing has been demonstrated or is suspected in a trap fishery, the fishery agency should decide what timed-failure mechanisms would be most appropriate to reduce the lifespan of traps and how soon timed failure should occur. Research under actual fishing conditions should then be conducted to determine the most appropriate regulation for timed failure.
3. Further studies with simulated lost pelagic gillnets should be conducted using nets larger than those studied to date and approximating commercial nets. These studies should examine whether ghost fishing takes place; if so, at what rate; and the rate at which the nets form a mass or otherwise cease to fish.
4. Direct observations should be made of lost pelagic gillnets to determine their shape and to determine the apparent rates at which ghost fishing for fish, birds, sea turtles, and marine mammals is taking place.

Working Group 4: Debris Ingestion by Marine Life (chaired by Louis Sileo)

The working group noted that studies to determine the prevalence of ingested plastics require monitoring. It recommended that future studies have statistically adequate sampling schemes designed to test hypotheses

that the prevalence is increasing or decreasing in given areas, taxa, etc. The group found that, regardless of the species, the same three general pathophysiological effects were proposed: (1) mechanical blockages; (2) pseudosatiety or other impairment of a chick's ability to accept a full meal; and (3) absorption of toxins from the plastic.

The working group placed priority on research on turtles, i.e., experimental feeding studies to determine how to interpret the lethality or other significance of ingested plastic and the range of pathophysiology in ingested plastic in turtles, along with continued monitoring of the prevalence of ingested plastic and its association with lesions. The group also recommended controlled experiments be carried out on birds to (1) determine if pseudosatiety does occur; (2) elucidate the duration of retention and erosion rates of ingested plastics; and (3) determine the toxicity of ingested plastics. The results of such studies will determine the need for long-term population studies.

Working Group 5: Economic Impacts of Marine Debris
(chaired by Kenneth McConnell)

The working group viewed the marine debris problem as an example of a situation where markets have failed to allocate resources efficiently, leading to the creation of nonmarket, or external, costs. The presence of an external cost indicates a problem that requires some form of public policy to solve. However, policies to reduce marine debris require people to change their behavior. Incentive schemes may be especially cost-effective in controlling debris when education and moral suasion fail. The working group proposed a list of projects to investigate the use of fees and incentives as part of the solution. These include: deposits on the return of nondegradable products, fees on the use of nondegradable materials, and incentives at the production level.

As regards compliance, the working group proposed investigating alternatives to the traditional approach of seeking compliance through persuasion. Policies combining punishment and reward and which partly subsidize the adoption of techniques are used elsewhere. For example, sewage treatment has been enhanced by Federal subsidies to construct waste treatment plants linked with the requirement that all households hook up. Methods of linking compliance to rules and regulations for handling marine debris can be linked to access to other beneficial programs.

Public campaigns to reduce pollution by moral suasion have been attempted for other forms of pollution. A study of such prior public campaigns would help understand their failures, which have been many, and their successes, which have been few.

As regards on-shore disposal, the working group proposed an investigation of the economic gains that can accrue to a particular region as a consequence of consolidating waste handling facilities.

On the matter of aesthetics, the working group noted that debris makes beaches less attractive and traps fish and wildlife. Each of these entails

an aesthetic loss to some individuals. Currently, little or nothing is known about the economic cost of either. The working group recommended two types of studies to help understand the magnitude of the economic costs of marine debris. These are a study of the economic costs of debris on a specific set of beaches and a study of the economic costs incurred when some individuals of a noncommercial species (e.g., birds, mammals) are entangled in marine debris.

When vessels and their gear are impaired by contact with marine debris, there are two kinds of costs: the cost to repair or replace damaged gear and the opportunity cost of the vessel and gear when it is not in service. The working group suggested research to investigate the incidence of impairment and the magnitude of costs for one of the following industry groups: commercial fishing, shipping, or recreational boating.

The biggest impact of marine debris on fish stocks is the ghost fishing phenomenon, but there is also the possibility that consumers' perception of contamination of fish stocks by marine debris can influence the demand and price of selected fish products. Ghost fishing has an economic cost in terms of the lost resource. The working group suggested a joint project involving economists and biologists to study the impact of perceived contamination on the price of and demand for fish.

Working Group 6: Technological Solutions (chaired by William Gordon)

The working group recognized that further work is required to quantify the types and volumes of ship-generated debris, but felt strongly that technology/methodology currently exists to address management of the majority of the wastes generated at sea. Because a large measure of the ship-generated debris ultimately will be transported ashore, satisfactory resolution of much of the marine debris issue will require rational resolution to many of the terrestrial waste management issues and problems.

The group concluded that more information should be obtained about types, quantities, and distribution of the plastic materials that will be brought ashore under MARPOL regulations for disposal ashore. Such information should be disseminated throughout the plastics industry to encourage reuse of such material.

The working group recommended that development of new technologies be encouraged. For example, in the area of low technology burning, research is needed on the environmental impacts of air emissions, development of guidelines concerning materials separation and operations, and environmental implications of ash and methods of its disposal. Regarding incineration, research is needed on environmental impacts of air emissions and the environmental implications of ash and methods of disposal.

In the area of ship design, new designs should accommodate waste management strategies and new construction should include facilities and space accommodations for waste management.

Regarding policy, no single methodology or technology will ensure compliance with waste management regulations. The working group felt that, accordingly, no policy should be established which will prohibit technologies which have potential.

**Working Group 7: Marine Debris Law and Policy
(chaired by David Cottingham)**

The working group reviewed existing legal and institutional arrangements to curtail the disposal and loss of solid wastes into the marine environment. The group concluded that solutions to the problem of marine debris should be developed and implemented in concert with efforts to address broader solid waste management issues. The most pressing needs identified include: (1) expanding participation to relevant international agreements; (2) assuring that adequate port reception facilities are available at all ports and harbors to receive ship-generated garbage returned to shore; and (3) adopting national policies and programs, such as recycling and innovative packaging, to reduce quantities of generated solid waste.

The group identified two international agreements as being of greatest importance to the problem of marine debris: MARPOL Annex V and the London Dumping Convention (formally known as the International Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter). In addition, at least 10 regional conventions control various forms of pollution, including the disposal of plastics and other solid wastes into the marine environment. The working group concluded that these international conventions establish the prohibition of disposing of plastics and other solid materials into the sea as "customary international law."

The group recognized that control of land-based sources of marine debris is a difficult problem that must be addressed domestically by individual nations. For example, it recommended that governments encourage recycling programs to reduce the volume of material that becomes solid waste.

Regarding compliance and enforcement, the working group recommended that the International Maritime Organization and pertinent governments party to MARPOL Annex V develop incentives to encourage vessels owners and operators to comply with garbage disposal regulations, and that vessel owners and operators be encouraged to report ports and harbors that do not have required port reception facilities.

Working Group 8: Marine Debris Education (chaired by Kathy O'Hara)

Education has been identified as an important way to help reduce the marine debris problem and is particularly important because land-based sources of debris are primarily nonpoint. Ethics and behavior patterns of individuals both on land and at sea must be changed, and education is a known means for effecting such changes.

The working group was charged with assembling a comprehensive list of the types of educational materials currently in use. The group

identified more than 100 different types of educational materials, including 21 brochures, 19 reports, fact sheets and special documents, 11 posters, 10 videos, 9 curriculums and guides for educators, 6 newsletters, and more than 30 other types of educational materials.

Marine debris education encompasses two key elements: the implementation of educational programs and the development of educational materials. With regard to the former, the working group recommended that marine debris education should be incorporated into three primary types of programs: (1) formal education in a structured academic setting; (2) informal education outside a formal academic setting but within structured educational events such as adult education classes and organized youth groups; and (3) general public awareness. Among the groups identified as target audiences for marine debris education, the working group concluded that five major groups are priority audiences: (1) general public, (2) media, (3) teachers and educators, (4) school children, and (5) all marine user groups.

A public awareness campaign is of utmost importance at present. Specific elements that should be addressed in developing this campaign include an initial assessment of human behavior and public perception of the marine debris problem. The working group felt that a comprehensive strategy to effectively use the media to disseminate information was paramount to the success of this campaign.

After reviewing the list of marine debris educational materials, the working group concluded that there is a wealth of materials currently available but there is a need to facilitate the dissemination of these materials to appropriate groups. In 1988, the National Oceanic and Atmospheric Administration's Marine Entanglement Research Program established two Marine Debris Information Offices, which respond to requests for information on marine debris. The working group suggested that the dissemination of educational materials would be facilitated if these offices were given increased visibility as a resource center along with sufficient quantities of educational materials to meet the demand.

Existing government distribution mechanisms should also be used to increase dissemination of materials, such as licensing and registration procedures for fishing and boating, the working group concluded. The group recognized the difficulty of disseminating educational materials on an international level due to the diversity of cultures and languages. However, it was suggested that specific international agencies, such as the United Nations Environment Programme, the Food and Agriculture Organization, and the International Maritime Organization, should be encouraged to facilitate information exchange.

Efforts should be made to include the marine debris issue on the agendas of international conferences and meetings that address the issues of marine pollution and education.

Appropriate means for evaluating the success of educational programs were discussed. The working group concluded that evaluation techniques could be conducted through long-term citizen monitoring of beach debris and


monitoring the usage of shoreside refuse reception facilities. Formal surveys should be conducted, where possible, to assess changes in attitude and behavior.

It was agreed by all working group participants that the marine debris issue is part of the larger solid waste problem and therefore we should incorporate lessons learned from dealing with solid waste into marine debris educational materials and programs.




V. RECOMMENDATIONS

Conference participants set forth a number of priority recommendations. Examples of some of the primary recommendations are:

- broad international acceptance and implementation of the terms of MARPOL (73/78) Annex V, especially the provision of port reception facilities;
- recognition of the marine debris problem as a symptom of the worldwide solid waste disposal crisis;
- pursuance of technological and procedural solutions to the marine debris and solid waste problems while avoiding policies and regulations that may restrict solutions;
- expansion of marine debris and solid waste disposal education to people and institutions worldwide, recognizing regional and cultural differences in the perception of these problems;
- development of a set of standard methods for surveys of the amounts, types, and sources of marine debris;
- establishment of an international committee or organization to further collaborative research on the impacts of entanglement on living marine resources;
- design and implementation of baseline experiments to establish the lethal and sublethal impacts of persistent debris ingestion by sea turtles and seabirds;
- design and implementation of experiments to evaluate ghost fishing in gillnet and trap fisheries with high gear loss rates, developing mitigative measures as needed; and
- evaluation of the economic impacts of marine debris, both direct, as in vessel disablement and commercial fish loss, and indirect, as in aesthetic damage and solution costs.



OVERVIEW PAPERS



A REVIEW OF MARINE DEBRIS RESEARCH, EDUCATION, AND MITIGATION IN THE NORTH PACIFIC

James M. Coe
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
Seattle, Washington 98115, U.S.A.

ABSTRACT

The earliest biological investigations and reports of the marine debris problem focused on North Pacific species, principally seabirds and marine mammals. In 1984, the Workshop on the Fate and Impact of Marine Debris in Honolulu brought together scientists and managers to evaluate information on this problem. Based on the recommendations of the workshop this paper reviews research and management activities and results since 1984 in the North Pacific. The Governments of the United States and Japan have been the primary participants in these activities. Both United States and Japanese programs include research and monitoring, mitigation technology, and education, as evidenced by the variety of papers being presented at this conference. The effective implementation of the requirements of MARPOL Annex V, especially in the fishing industries of the North Pacific, is a common goal of most Pacific Rim nations. The fishing industries themselves have made significant commitments to address their contribution to the marine debris problem in the North Pacific. The effects of these actions on the known impacts of persistent debris in the North Pacific have yet to be realized.

INTRODUCTION

The early focus of attention in the marine debris issue in the United States was the North Pacific Ocean and its shores. This was due in large part to the early documentation of the interactions of wildlife with fisheries-generated marine debris. Observations of northern fur seals entangled in debris were recorded as early as 1936 (Fiscus and Kozloff 1972). Records of Laysan albatross ingesting plastic debris date from the mid-1960's (Kenyon and Kridler 1969). In 1974, field biologists began keeping records of entanglement of highly endangered Hawaiian monk seals (Henderson 1985). Widespread ingestion of plastic particles by 37 species of North Pacific seabirds was reported in a study by Day (1980).

Starting in the early 1960's, large-scale industrial fisheries proliferated in the North Pacific and in the Bering Sea. Increasing use, loss, and discard of persistent plastic nets, packing straps, packaging, and other refuse from these vessels were evident in surveys of Alaskan beaches since 1972 (Merrell 1985). The general surface circulation patterns of the North Pacific suggest that floating debris may remain at sea for several years before being deposited on shore (Reed and Schumacher 1985). Accumulating evidence of increasing amounts of debris in the ocean combined with observations of its range of impacts on wildlife led to the convening of the first international scientific meeting on marine debris in late 1984.

The Workshop on the Fate and Impact of Marine Debris (FIMD), held in Hawaii in November 1984, was an opportunity for the scientific community to evaluate the state of knowledge about marine debris and to draw conclusions where appropriate (Shomura and Yoshida 1985). Based largely on data from the North Pacific, the working groups at the 1984 workshop concluded that persistent marine debris poses a long-term threat to certain species as well as to maritime and coastal commerce. Observations from other ocean areas suggested that similar problems could exist in all the world's ocean areas. These revelations fostered the initial concern about this new and apparently widespread form of marine and coastal pollution.

In response to the charge to participants, the 1984 workshop prepared a series of findings and recommendations that were to become the guidelines for marine debris action in the North Pacific as well as the world. The executive summary of the workshop report includes the following conclusory paragraph:

"The Workshop considered the information presented during the technical sessions and concluded that there is ample evidence that debris of both terrestrial and shipborne origin are widespread in the marine environment. While such debris is known to interact with a wide variety of marine mammals, fishes, turtles, birds, and invertebrates, in most instances the consequences and quantitative impacts of this interaction do not appear to be well understood. However, substantial qualitative evidence indicates these interactions are contributing to increased mortality over that resulting from natural causes."

These findings prompted the workshop participants to make general recommendations for information collection to elucidate the sources, distribution, amounts, fates, and impacts of persistent marine debris. Studies of the biological impacts of entanglement and ingestion on North Pacific marine mammals, seabirds, and sea turtles were specifically identified. The development of sampling methodology for beach and sea surface debris--especially fishing gear--was recommended. In concluding that marine debris is a real problem for marine life and vessels, however poorly quantified, the workshop recommended education and mitigation efforts to be undertaken concurrently with the information collection activities. The mitigation efforts were to include regulation of the types of debris most hazardous to marine life, investigations of the use of biodegradable

materials, net recycling, and the promotion of beach surveys and cleanups. Education efforts were recommended to advise user and interest groups of the nature and scope of the marine debris problem. The target groups were to include fishing and plastics industries, merchant carriers, the military, appropriate international organizations, and the public.

The workshop recognized that many of its findings were based on information from the North Pacific Ocean and recommended that the severity of the marine debris problem in other ocean regions be investigated. It also recommended a start on the evaluation of economic impacts of debris by obtaining data on worldwide vessel disablement caused by interactions with marine debris. The need for international cooperation in the investigation and solution of marine debris problems was broadly recognized by the workshop.

With the 1984 FIMD workshop as a starting point, this paper attempts to summarize and review recent marine debris research and monitoring, mitigation, and education activities affecting the North Pacific Ocean. This review includes known United States and foreign activities, brief summaries of their results, and an evaluation of developments and continuing needs in each action area. It is likely that there have been foreign government or industry actions addressing marine debris in the North Pacific that are not reported here. Any such omissions are unintentional. Many of these actions may be reported elsewhere in Shomura and Godfrey (1990).

RESEARCH AND MONITORING ACTIVITIES

Research related to marine debris problems has encompassed biological investigations, measurement of the sources, amounts and distribution of debris, and research and development for technological solutions. Under this section, biological research and the monitoring of debris sources and amounts will be reviewed. Research related to technologies for solving marine debris problems will be reviewed in the Mitigation section below.

Biological Research

Northern Fur Seal Entanglement

The United States, Japan, and the Soviet Union have carried out research related to the entanglement of the northern fur seal, *Callorhinus ursinus*. Until 1985, these research activities were coordinated under the Interim Convention on Conservation of Pacific Fur Seals. Since that time, cooperative research has continued between the United States and Japan at the Pribilof Islands, with each nation also doing independent research.

The northern fur seal population breeding at St. Paul Island in the eastern Bering Sea has been the subject of continuing study of the role of entanglement in fur seal population dynamics. Against a background of a declining population, the hypothesis that entanglement is a principal contributor to that decline was evaluated (Fowler 1985). Research to

elucidate this relationship was primarily confined to the immediate vicinity of St. Paul Island during the summer breeding season.

Scientists from the United States and Japan continued cooperative studies on juvenile male fur seals in order to count and tag entangled seals. These studies (involving roundups) were necessary to simulate the juvenile male harvests which ended in 1984 but from which all previous entanglement rates had been calculated. A sample of nonentangled seals was tagged at the same time to allow for later evaluation of differential mortality based on resighting rates in future roundups. Roundups with tagging were conducted in 1985, 1986, and 1988. Resighting of these tagged seals provides data on the differential survival of entangled and nonentangled juvenile males. Roundups with removal of debris, starting in 1989, are expected to produce the tag resighting data necessary to estimate the changes in survival that may be possible through removal of debris for the period between weaning and the seals' first return to St. Paul Island.

Interpretation of the entanglement rates calculated from the roundups has been complicated by the possible differences in behavior of entangled and unentangled seals. Observations suggest that entangled seals may spend a larger proportion of their time away from the hauling grounds than their unencumbered counterparts. Studies in 1985 of entangled females with pups showed significantly longer feeding forays for entangled females and consequently less healthy pups. An experiment was conducted in 1988 using radio tags on entangled and unentangled juvenile males to measure differences in hauling behavior. Results will be useful in interpretation of prior years', roundup-based tag returns for calculating survival. These results are currently being analyzed with preliminary results presented by Fowler et al. (1990).

Without correcting for possible behavioral differences affecting the calculations, the entanglement rate for juvenile male fur seals has been near 0.4% from the late 1970's to the mid-1980's. Preliminary results for 1988 suggest that this rate may be decreasing due to less entanglement in waste trawl netting. Preliminary results and some interpretation of the tagging and differential mortality studies is presented in the Technical Session on Entanglement.

Both Japanese and United States scientists have carried out research on the behavior of fur seals leading to their entanglement. In 1986, an experiment in which fur seal pups were allowed to swim in a tank with various-sized pieces of netting showed that newly weaned pups were highly susceptible to entanglement in netting as small as 15 cm stretched mesh and that few were able to extricate themselves (Bengtson et al. 1988). Similar experiments in Japan with captive juvenile male fur seals showed that investigative behavior often led to entanglement, but many of these entanglements were temporary. In the Japanese experiments, the materials offered to the seals reflected the ranges of sizes found on naturally entangled, living seals. These experiments demonstrate the susceptibility of fur seals to entanglement in nets of various mesh sizes, and suggest that newly weaned pups may be at particular risk.

For most of the year, the northern fur seal population ranges across the entire subarctic Pacific. This makes coherent studies of at-sea entanglement arduous, expensive, and risky in terms of information return. Consequently, research on the entanglement of northern fur seals during their 9 months away from the Pribilof Islands has been minimal. At the Pribilof Islands, only the survivors are being investigated, giving a potentially biased view of the role of entanglement in population fluctuations. Research reviews and workshops in the United States to elucidate methods of inquiry that may be feasibly and economically applied to this question have been unsuccessful.

Recent data from the Pribilof Islands suggest that the fur seal population may have stabilized. Having been unable in the last 4 years to identify directly the role of entanglement in the fur seal population decline from the 1970's through the early 1980's, further entanglement research at the Pribilof Islands may be unproductive, although monitoring of entanglement rates as an index of hazardous debris changes should continue.

The question is certainly not closed. The impact of entanglement on 0- to 2-year-old fur seals while at sea remains one of the most serious marine debris issues. Resources permit only opportunistic gathering of data in the pelagic range of these animals. Reviews of remote sensing applications and other high technology approaches to this issue have shown them to be prohibitively risky or expensive. It is apparent that a complete, scientific assessment of the role of marine debris in population fluctuations of the northern fur seal will take many more years, if it can be done at all.

Northern Sea Lions

The northern sea lion, *Eumetopias jubatus*, population in the eastern Aleutian Islands has experienced a population decline of about 7% per year, similar to the northern fur seals at the Pribilof Islands. Unlike the northern fur seal, the northern sea lion population in the eastern Aleutians appears to have declined continuously since the 1960's. Concomitant decreases in other North Pacific population centers rule out emigration as an explanation. Entanglement in marine debris was hypothesized as a possible cause for this decline along with changes in prey availability, disease, direct killing by fishermen, and rookery/haul-out disturbances.

Since there were a few observations on record of entangled northern sea lions, surveys of the eastern Aleutian and Gulf of Alaska haul-out sites were conducted in June-July and November of 1985 (Loughlin et al. 1986). In the June-July survey, just over 30,000 sea lions were counted. Six were entangled and five more showed obvious signs of previous entanglement. The entanglement rate in this survey was 0.04% of the adult population. These data were judged inadequate to assess the magnitude of entanglement of sea lion pups because the survey took place before the pups had gone to sea for the first time.

The November survey was conducted to census fur seal and sea lion pups hauled out or stranded in the Aleutian Islands after weaning. This survey

covered nine known haul-out locations and possible stranding sites but found no entangled fur seal or sea lion pups.

These results did help to clarify the role of entanglement in the northern sea lion population decline by suggesting a very low entanglement rate and a possibly high level of escapement from entanglement, at least in adults. Just as with the northern fur seal, the question of what may be happening to newly weaned sea lion pups at sea remains unclear.

Hawaiian Monk Seals

Classified as an endangered species under the U.S. Endangered Species Act (ESA), the Hawaiian monk seal, *Monachus schauinslandi*, is afforded special attention to protect it from entanglement. Since 1982, field biologists have collected, catalogued, and destroyed potentially entangling debris found at known monk seal haul-out sites in the Northwestern Hawaiian Islands (NWHI). Wherever possible, seals are freed from debris. Through 1984, records showed 35 incidents of monk seal entanglement, including 8 with scars of previous entanglements (Henderson 1985). From 1985 to 1987, another 19 entanglements have been observed, 3 of which resulted in the death of the animal (Henderson 1988). These 19 incidents represent an increase in the observed rate of monk seal entanglement despite the fact that many haul-out beaches in the NWHI are cleaned at least once a year.

Further information on the effects of entanglement on the Hawaiian monk seal are presented in Henderson (1990).

Seabirds

While there are scattered reports of seabirds being entangled in a variety of materials, the more widespread problem for seabirds is the ingestion of debris, especially floating plastics. Research on marine debris/seabird interactions undertaken in the North Pacific since the 1984 workshop has focused exclusively on the ingestion problem. In 1986, three specific studies of the impacts of plastic ingestion on the seabirds of Hawaii were undertaken cooperatively by the U.S. National Marine Fisheries Service (NMFS) and the U.S. Fish and Wildlife Service.

An evaluation of the incidence of ingested plastic in seabirds of the Hawaiian Islands was conducted between May 1986 and January 1987 (Sileo, Sievert, Samuel, and Fefer 1990). Prior to this study, only 2 of Hawaii's 22 species of seabirds had been thoroughly examined for ingested plastic. The study was able to examine 18 of Hawaii's 22 species, finding only 2 species with 0 plastic. The presence of plastic ranged from 0% in gray-backed and white terns and 1% in brown noddies to 94% in black-footed albatrosses. The data suggested that incidence of plastic in birds was related to the level of plastic in their immediate environment.

The other two studies of seabird ingestion of plastics involved the Laysan albatross population at Midway. Sampling of Laysan albatross chicks to determine diet, growth, and general health was initiated in 1987 to measure the relationship between plastic ingestion and growth, and plastic

ingestion and mortality. This study continued in 1989, as the variability in the amounts of plastic fed by parent albatrosses to their chicks has been higher than expected. Chicks in 1987 had on average nine times more plastic in their diet than did 1988 chicks. Preliminary indications are that plastic ingestion may not contribute in any obvious way to chick mortality; its impacts on growth, however, may be detectable (Sileo, Sievert, and Samuel 1990; Sileo, Sievert, Samuel, and Fefer 1990).

Papers presented at this conference indicate that several species of North Pacific shearwaters have been studied for plastic ingestion (Ogi 1990). Further, investigations of seabird ingestion are underway in the South Atlantic by Ryan, in the South Pacific by Gregory, and in the eastern tropical Pacific and the Antarctic by Ainley and others. As in the North Pacific studies, there is, as yet, little evidence of direct damage to most seabird species caused by the ingestion of plastic debris. This indication is by no means proven, although broad acceptance of such a finding may be forthcoming. The working group on ingestion of marine debris is expected to address this generalization and recommend definitive research actions.

Cetaceans

In the period between the 1984 workshop and the current conference, little progress has been made in distinguishing between evidence of cetacean entanglement in marine debris and cetacean entanglement in active fishing gear. In most cases, the animal is encumbered by some fragment of fishing gear or rope but is found stranded on shore or adrift at sea away from any direct source. There has been no accumulation of records since the 1984 workshop to indicate that North Pacific cetaceans are threatened by marine debris through entanglement. A direct review of this phenomenon using all available information has not been undertaken. The value of such a review should be discussed in the working group on entanglement.

The subject of ingestion of marine debris by odontocete cetaceans was reviewed by Walker and Coe from 1987 through 1988 and is reported in the Technical Session on Ingestion in these Proceedings. While Walker and Coe (1990) found virtually no ingestion of debris by free-ranging pelagic odontocetes, they describe several cases of severe trauma to captive dolphins due to plastic ingestion. This review also finds that Baird's beaked whale, *Berardius bairdi*, and the sperm whale, *Physeter macrocephalus*, which feed in the benthos, commonly ingest foreign materials that have sunk to the sea floor. This work also suggests that the filter-feeding mysticetes may be at much greater risk of ingesting debris than their toothed cousins. A review of current worldwide information on the ingestion of foreign materials by the mysticete whales and the impacts of ingestion seems justified, as does further investigation of the benthic-feeding odontocetes.

Sea Turtles

Balazs (1985) summarized the body of information on entanglement in and ingestion of marine debris by all species of sea turtles. While the North Pacific is home to at least four of the seven species of sea turtles,

all of which are protected under the ESA, the majority of research on the impacts of debris is being done in the southeastern United States. Because they are clearly vulnerable to entanglement (Balazs 1985), are relatively indiscriminate feeders (Lutz 1986), and are all endangered or threatened under ESA, sea turtles wherever they are found must be considered at serious risk from marine debris. Balazs and Choy (1990) provide an update on our knowledge about this problem for North Pacific sea turtles.

Research on the impacts of entanglement and ingestion on hatchling and juvenile sea turtles are being conducted by the Archie Carr Center for Sea Turtle Research at the University of Florida. This work is concentrating on the loggerhead turtle, *Caretta caretta*, in the Atlantic and will continue for at least 2 more years. The results of these studies regarding the role of convergence zones as debris sinks and sea turtle rearing areas may be generally applicable to other species of sea turtles, including those in the North Pacific.

Debris Sources and Amounts

The 1984 Hawaii workshop expressed concern about persistent debris at sea, on beaches, and on the sea floor. The principal focus of this concern was lost or discarded fishing gear, especially netting, traps, and ropes. These materials were judged to present the greatest hazard to marine life and ships through ghost fishing and entanglement. Since that time, research has been carried out to establish methods for surveying debris on beaches and at sea, systematic surveys have been conducted on beaches in Alaska, surveys of floating debris have been made from a number of vessels, a marine debris reference collection has been established, and trawl surveys of benthic debris have been completed.

Methods

A variety of approaches have been used to measure debris on beaches. Methods vary from geographic region to geographic region and from worker to worker. Most approaches have sound statistical underpinnings and reflect the experience and preferences of the survey initiator. Ribic and Bledsoe (1986) examined Alaskan beach survey data from Merrell (1980, 1985) and data from a number of sighting survey cruises in the North Pacific. They recommended methods for carrying out surveys of lost nets at sea and on beaches. Most of these recommendations focused on improving the ability to detect changes in debris density over time. From ship survey data for floating net sightings in the North Pacific and Bering Sea (Jones and Ferrero 1985), these workers calculated that net surveys in these regions would need to include at least 2,800 sampling units (1-h watches) in order to detect a reduction of 50% with 95% confidence. These surveys should be run annually and be designed to permit identification of, and stratification for, local concentrations of debris. This work also recommended that the suitability of aerial survey techniques for marine debris be evaluated.

Specific to entanglement problems in the North Pacific, the 1984 Hawaii workshop recognized the need for identifying the fishery of origin of nets and other fisheries materials found on animals and on the beaches.

In response to this need, the NMFS established a reference collection for fishing gear debris in the Alaska Fisheries Science Center in Seattle. Researchers with unidentified fisheries debris from the North Pacific may send a sample to the curator of this collection along with all pertinent information and receive an evaluation of its composition, the likely fishery of origin, and other pertinent information that may be available.

Debris Surveys

Since the work of Ribic and Bledsoe, there have been a large number of at-sea debris surveys, much of it from Japanese research and patrol vessels (Mio and Takehama 1988; Yagi and Nomura 1988). The surveys reported by Mio and Takehama involved 17 vessels covering 80,546 nmi in the North Pacific and recording 7,458 sightings, 1,584 of which were seaweed and 1,082 were wood, or 0.06 synthetic debris items of detectable size per track mile. Yagi and Nomura reported debris sighting data from vessels repeating a north-south transect from Japan to New Guinea from 1977 to 1986. This survey averaged 39 debris sightings annually in an average 4,000 km surveyed, or just under 0.01 items per track mile. An increasing trend in the number of plastic sheets and bags was identified in this series; however, no overall increase in plastic debris was obvious.

Cooperative research cruises each year since 1986 between the United States and the Republic of Korea and Taiwan have gathered data on the at-sea distribution of marine debris. These results are contained in the NMFS cruise reports (unpubl. data) but have not been consolidated or analyzed for time series or regional comparisons.

Some of the early research (Day 1980; Day et al. 1985) on ingestion of plastic particles by seabirds led to the speculation that a large amount of disintegrating plastic debris may be afloat in the convergence areas of the North Pacific. The density and characteristics of the microdebris were investigated by Day under contract to NMFS in 1987 and 1988. Samples were taken at 27 stations using neuston nets with mesh sizes in intervals down to 0.053 mm, and at 46 stations with mesh sizes in intervals down to 0.50 mm. In general, Day found areas of floating plastic particles all over the North Pacific, with the highest concentrations near Japan and just south of the Subarctic Front. The specific findings and interpretation of this work are reported in Day et al. (1990).

Ribic and Bledsoe (1986) concluded that "The usefulness of beach survey information is almost entirely dependent on the capability to infer ocean debris conditions from the beach information." The coordination of shipboard and beach surveys is essential if the utility of beach survey data is to be confirmed. Further, the lifetimes and dynamics of debris on beaches need to be understood if one is to conduct independent surveys over time in regions of interest. It may be necessary to remove or permanently mark debris to evaluate lifetime and movement as well as to ensure independence from survey to survey.

To date, there have been no coordinated ship/beach debris surveys to evaluate the relationship between amounts and types of floating and stranded

debris in any region of the North Pacific. Johnson and Merrell (1988) report on time series of beach debris surveys from selected beaches in Alaska, where they cleaned sections of beach and also tagged large debris items. From this work they have been able to estimate the rate of deposition of entangling materials on certain Alaska beaches. At Yakutat, the deposition rate of trawl nets was estimated at seven nets/km/year. As a result of his debris tagging work, Johnson discovered that large net debris may be buried, uncovered, and moved along the beach by severe winter storms. Investigations into the dynamics of beached debris are continuing in Alaska and are further reported in Johnson (1990).

As part of the effort to protect Hawaiian monk seals from entanglement in debris, the research teams routinely survey, catalog, and remove nets, ropes, etc. from beaches in the NWHI. The net materials found in these surveys from 1982 to 1986 were reported by Henderson et al. (1987) and are updated by Henderson (1990). The collections through 1985 amounted to 632 nets or net fragments, 539 of which were poly, i.e., polypropylene or polyethylene, and 66 monofilament nylon. All of the monofilament net fragments were from gillnets and 54 of these were most likely from Asian squid and salmon driftnet fisheries. It was concluded that most of the poly nets and net fragments were from North Pacific midwater and bottom trawls. The fisheries of origin of this unexpectedly high proportion of trawl net materials on the beaches, and the ocean current systems that brought them to the NWHI, are yet to be established. Since the number of nets per kilometer of beach was quite high at several of the most important pupping beaches, and monk seals, unlike northern fur seals, entangle in a wide range of mesh sizes, the sources of poly fragments in the NWHI need to be understood and minimized.

Interest in the nature of accumulations of sinking debris on the Continental Shelf led to the enumeration of debris in bottom trawl surveys in the eastern Bering Sea in 1987 and off the U.S. west coast in 1988 (June 1990). These surveys were for groundfish abundance in areas of sand or mud bottom. The survey nets were set up to fish hard on the bottom, making it likely that debris in the upper few centimeters of the sediment would be scooped into the nets and be recorded. In general, the concentrations of sunken debris reflect the level of vessel activities in the areas. As one would expect, high traffic areas have greater debris densities than low traffic areas. Also, the types of debris found on the bottom generally reflect the surface activities.

In an attempt to elucidate the sources of net fragments in the North Pacific and the Bering Sea, a study of NMFS Foreign Fishery Observer Program data was conducted (Berger and Armistead 1987). The records from 1,068 observer cruises in 1982-84 in the U.S. exclusive economic zone (EEZ) provided data from every month of the year. The amounts of net discarded, lost, retrieved, or seen floating were recorded, as were net-mending activities and fishing operations. In 1982, 1983, and 1984, respectively, 14, 31, and 17% of the net pieces discarded were in the mesh size range to entangle fur seals. During this period, a total of 1,551 pieces of net were brought up in trawls and most were discarded back into the ocean. In 1983, foreign joint venture operations lost 70 trawl nets or large portions

of nets. In 1984, this number increased to 90 nets. Foreign fishing in the U.S. EEZ has been almost completely displaced since 1984, but it is not known if the loss and discard rate of nets and net fragments has changed. Under the U.S. domestic law implementing MARPOL Annex V, it is illegal to discard net fragments, and one would therefore expect the amount of net input into the U.S. EEZ in the North Pacific to decrease in the coming years.

Another applied study of the disposition of derelict fishing gear in the North Pacific was reported by Gerrodette et al. (1987). In this study, a series of monofilament drift gillnets were attached to satellite transmitters and set adrift to simulate lost sections of squid or salmon drift-net. The purpose of the study was to gather information on the size, shape, location, and length of time in the ocean. Four nets, 50, 100, 350, and 1,000 m long were released on 12 August 1986 in the vicinity of Hancock Seamounts, northwest of the Hawaiian Islands. The nets were tracked by satellite from 57 to 309 days. The 50- and 100-m nets collapsed within hours of being deployed. The 1,000-m net was reduced to approximately 15% of its original length after 9 days adrift. It appears that there is a positive correlation between the length of the drifting section of gillnet and its ghost fishing effectiveness. The complex tracks of the nets showed that prediction of the drift paths of derelict fishing gear requires a detailed knowledge of the local surface currents and wind conditions. Recent Japanese studies of drifting gillnet (Mio et al. 1990) confirm these general findings; however, the ghost fishing characteristics of a lost, full length, pelagic driftnet (approximately 5 km) have yet to be measured.

Lastly, voluntary public beach cleanups have been organized to the extent that a uniform method of data collection is being employed in the western United States and Hawaii (Center for Environmental Education 1988). The data from these annual cleanup programs may have some utility as indices of the long-term changes in the amounts and types of beach debris in various regions. The myriad promoters of this voluntary initiative are intent on the development of a worldwide International Beach Cleanup Day using the same data collection methods. Over a 10-year period, the success of the implementation of MARPOL Annex V may be seen in the data from these extensive but infrequent (once or twice per year) samplings. They should be broadly promoted.

MITIGATION

Under this section, the collection of activities whose objective was to lessen the input of persistent debris into the ocean, and especially the North Pacific, are summarized. These actions include technical assessments and developments of waste handling and disposal technologies, as well as legal and administrative efforts. Recent progress in both categories affects most ocean areas, including the North Pacific.

Legal and Administrative Actions

On 30 December 1988, the terms of optional Annex V of the International Convention for the Prevention of Pollution from Ships as modified by

the Protocol of 1978 (MARPOL 73/78) entitled "Regulation for the Prevention of Pollution by Garbage from Ships" entered into force for 35 nations representing slightly over 50% of the world's registered shipping tonnage. The MARPOL Annex V prohibits the discharge of plastic from ships into the ocean and sets distance-from-shore limitations for the discharge of other types of ship's garbage. Table 1 summarizes the discharge requirements of Annex V. Ships are defined under MARPOL 73/78 as all surface and subsurface craft as well as all fixed and floating platforms regardless of size. Annex V also identifies five special areas in which all discharge of garbage is prohibited. There are no special areas in the Pacific Ocean.

The principal North Pacific coastal nations that have ratified, and are implementing, MARPOL Annex V are North Korea, Japan, the U.S.S.R., and the United States. The domestic implementing legislation for Annex V differs somewhat between nations but, typically, flag vessels of signatory nations must meet the discharge requirements worldwide. All vessels within the EEZ's of signatory nations must meet the discharge requirements.

The Japanese showed concern over the trashing of the Pacific as early as 1970, when they enacted Domestic Law 136, "Law Relating to the Prevention of Marine Pollution and Maritime Disaster," which prohibits discharge of nets or net fragments and promotes onboard incineration. At the urging of the Fur Seal Commission in 1983, Japan joined the United States and the U.S.S.R. in a campaign to protect fur seals from entanglement by stopping the dumping of fishing gear and by cutting plastic strapping bands before discard. In June 1987, the Fisheries Agency, the Government of Japan (formerly the Fisheries Agency of Japan) established the Fishing Ground Preservation Division to carry out a broad range of projects related to the marine debris problem and its solutions. This program sponsors the research on the types and distribution of marine debris in the North Pacific reported above, and promotes a broad range of recycling, cleanups, and public education efforts, principally through prefectural governments and regional fishing organizations. Japanese domestic regulations implementing MARPOL Annex V were set in place in March 1988.

In response to the northern fur seal entanglement problem and the 1984 FIMD workshop, the United States set up the Marine Entanglement Research Program in the National Oceanic and Atmospheric Administration (NOAA). This program is charged with formulation and execution of research, mitigation, and education activities to address the marine debris problem in U.S. waters.

At the request of 30 U.S. Senators, and under the direction of the White House Domestic Policy Council (DPC), NOAA convened the Interagency Task Force on Persistent Marine Debris, which included the Departments of Defense, the Interior, Transportation, and Agriculture as well as the Environmental Protection Agency (EPA), the Office of Domestic Policy, the Marine Mammal Commission, the Office of Management and Budget, and the Office of the President. The Task Force reviewed the problem and produced a set of recommendations for United States actions. The DPC approved and published the Task Force Report in May 1988 (NOAA 1988). As implemented, these recommendations will have far-reaching impacts on the control of persistent marine debris in the North Pacific.

Table 1.--Summary of at-sea garbage disposal regulations. Source: Guidelines for the implementation of Annex V of MARPOL 73/78 (IMO 1988). (Note: The Baltic Sea special area disposal regulations took effect on 1 October 1989.)

Types of refuse disposed	All ships except platforms ^a			Offshore platforms ^a
	Outside special areas	Special areas ^b	Offshore platforms ^a	
Plastics, including synthetic ropes and fishing nets and plastic garbage bags	Disposal prohibited	Disposal prohibited	Disposal prohibited	Disposal prohibited
Floating dunnage, lining, and packing materials	Disposal permitted >25 nmi offshore	do	do	Do.
Paper, rags, glass, metal, bottles, crockery, and similar refuse	Disposal permitted >12 nmi offshore	do	do	Do.
All other garbage including paper, rags, and glass, comminuted or ground ^c	Disposal permitted >3 nmi offshore	do	do	Do.
Food waste not comminuted or ground	Disposal permitted >12 nmi offshore	Disposal permitted >12 nmi offshore	Disposal permitted >12 nmi offshore	Do.
Food waste comminuted or ground	Disposal permitted >3 nmi offshore	Disposal permitted >12 nmi offshore	Disposal permitted >12 nmi offshore	Disposal permitted >12 nmi offshore
Mixed refuse types	(d)	(d)	(d)	(d)

^aOffshore platforms and associated ships include all fixed or floating platforms engaged in exploration or exploitation of seabed mineral resources, and all ships alongside or within 500 m of such platforms.

^bGarbage disposal regulations for special areas shall take effect in accordance with Annex V 5(4)(b).

^cComminuted or ground refuse must be able to pass through a screen with mesh size no larger than 25 mm.

^dWhen garbage is mixed with other harmful substances having different disposal or discharge requirements, the most stringent disposal requirements shall apply.

A wide range of activities have been undertaken in international organizations and commissions that recognize and address the marine debris issue. The broadest possible recognition of persistent marine debris as a legitimate marine pollutant has been a goal of the United States. As a result of actions by the United States, Japan, and others, the marine debris problem has appeared on the agendas of the International North Pacific Fisheries Commission, the International Fur Seal Commission, the Intergovernmental Oceanographic Commission, the Commission for the Conservation of Antarctic Marine Living Resources, the Food and Agriculture Organization, the United Nations Environmental Program, and, of course, the International Maritime Organization (IMO). One of the principal products of these international actions is the publication of guidelines by IMO (1988). The main objectives of these guidelines are:

- to assist governments in developing and enacting domestic laws which give force to, and implement, Annex V;
- to assist vessel operators in complying with requirements set forth in Annex V and domestic laws; and,
- to assist port and terminal operators in assessing the need for, and providing, adequate reception facilities for garbage generated on different types of ships.

All maritime nations are encouraged to ratify and implement Annex V, using the guidelines to help standardize international practice.

The provision of adequate port reception facilities to receive ships' garbage has been a significant concern expressed by port and terminal operators in the United States. Two projects were undertaken at North Pacific ports to evaluate this issue: one in the west coast fishing and logging port of Newport, Oregon, and one involving Unalaska/Dutch Harbor and Kodiak, Alaska. The Newport Marine Refuse Disposal Project found that community and port user involvement in, and ownership of, the local marine refuse problem led to a high level of usage of port reception facilities. Further, efficient waste management practice in the port was maintained by integrating the garbage reception system with recycling and reuse programs in the community and with waste oil reception sites. The lessons from the Newport Project are reported by Recht (1988). Currently, under a grant from NMFS the Pacific States Marine Fisheries Commission is conducting a program to assist eight west coast ports in their development and provision of adequate garbage reception facilities.

The results of the Unalaska/Dutch Harbor and Kodiak evaluations of port garbage handling problems were released in October 1989. Results suggest that the waste disposal facilities of these remote, highly vessel-dependent ports may be strained by the addition of vessel garbage. This is particularly true for Dutch Harbor, where the landfill life may be shortened significantly. These problems are complicated by the need to handle and dispose of waste oil, hazardous wastes, and garbage requiring special handling for pest control. Preliminary suggestions for solutions involve recycling, incineration, and regional consolidation of certain high-capital

waste handling facilities. The experiences gained in this project and the Newport Project are generally applicable to ports all across the Pacific Rim.

The State of Washington has developed and published a Marine Plastic Debris Action Plan as a guide for state agencies in addressing the marine debris problem (Washington State Department of Natural Resources 1988). This plan is an excellent model for coastal states seeking guidance on organizing to deal with marine debris issues. California and Alaska are in the process of developing state policy and action plans.

The principal maritime nations of the North Pacific that have not ratified Annex V are Canada, Mexico, the People's Republic of China, the Republic of Korea, and Taiwan. Domestic laws of these nations addressing garbage discharge from ships have not been reviewed for this paper. It is known that the Canada Shipping Act provides the Canadian Government with the authority necessary to establish garbage regulations more stringent than Annex V. The Government of Canada is currently reviewing options for marine debris programs and controls. The Republic of Korea has developed a guide for the conservation of marine mammals and salmonids in the North Pacific which requires fishing vessels to retain, and return for shore disposal, all plastics and waste fishing gear, and to maintain a record of these actions. Legislative and policy actions on marine debris in other North Pacific countries have not been widely reported.

Mitigation Technology and Procedures

In general, efforts to develop or improve technology and procedures to reduce, control, or eliminate marine debris and solid waste have been carried out by private industry, by governments, and by independent organizations. This work covers shipboard-specific waste handling procedures and equipment, fishing gear technologies, incineration, recycling, and degradable materials. Little of the research and development in these areas has been specific to the North Pacific or to the marine debris problem. However, these developments are germane to controlling marine debris input to the North Pacific and are briefly discussed in this section.

Shipboard Waste Handling

Since the 1984 FIMD workshop no primary technology has emerged to control either ship-generated or land-source debris entering the oceans. The variety of applications and needs has operated to broaden, rather than narrow, the technical and procedural options open to all who must dispose of wastes. While regulatory systems seem to have progressively restricted disposal options on land, most regulators are allowing vessel operators to choose methods most suited to their circumstances. Absent any substantive reasons to the contrary, preserving all technical options that allow disposers to meet the requirements of the law should result in higher levels of compliance.

In 1986, NMFS contracted for a review of shipboard waste handling options. The report (Parker et al. 1987) produced a table showing the

applicability of various waste disposal methods for various types of ships. Limited data on waste generation rates, ship configurations, and procedure capacities required some assumptions to be used in developing the table. The most general findings in this study were that:

- controlled incineration could be used aboard all but the smallest vessels;
- in using compactors, all but ships with very high crew complements (military vessels) should be able to store their compacted wastes on board;
- storage of uncompacted wastes on board is limited to fishing boats, research vessels, and others where the vessels are large relative to crew size; and
- waste generation rates on most vessels are too low to make recycling an economically attractive approach.

Alig et al. (1990) and Martinez (1990) review the more recent developments in shipboard waste handling technology and procedures.

The entry into force of Annex V has resulted in increased use of, and experimentation with, burn barrels for disposing of plastics and other garbage aboard ships with small crews, especially fishing boats in the North Pacific. The NMFS commissioned a study of the design and use of burn barrels to provide information on their technical feasibility, safety, and environmental considerations (SCS Engineers, Inc. 1989a). The work concludes that burn barrels are currently legal outside 12 nmi, may be regulated by coastal states inside 12 nmi, are not yet regulated by the EPA, are capable of reducing certain types of garbage to ash, and, under certain conditions, can be operated safely (Chang 1990). Operating and safety guidelines for the use of burn barrels aboard ships were prepared (SCS Engineers, Inc. 1989b). However, neither NMFS nor the contractor for these studies advocates the use of burn barrels.

Degradable Plastics

Since the 1984 FIMD workshop, the replacement of disposable plastics and fishing gear with degradable plastics has been widely discussed. This has been characterized as a potential solution for ghost-fishing problems caused by lost and discarded fishing gear, as well as a potential solution for litter and landfill capacity problems. Substantive research and development work on these types of plastics has been renewed after some initial work in the early 1970's. Most of this work is being done within the plastics industry and is proprietary. The American Society for Testing and Materials has formed a technical committee to define and develop standards for degradable plastics. In the mean time, there have been many commercial claims of product performance and applications for degradable polymers. Whether these products or future products will play a substantial role in controlling future plastic waste impacts on the environment remains to be seen.

The single recent study of the degradation properties of certain polymer types in marine and terrestrial settings was carried out by Andradý (1990). There has been no applied research on the use of degradable polymer products in the fishing industry. The primary work has been in the applications of natural fiber connectors or linkages in traps and pots to reduce their ghost-fishing life. Some coastal states around the North Pacific require natural fiber lacings or hangings in side panels or tunnels of crab, lobster, and fish traps. Ideally, these rot out shortly after the device is lost, rendering the trap harmless. A recently realized drawback to these approaches is that most natural fiber twines on the market are now fortified with some percentage of polymer fibers and do not rot as quickly or completely as expected (W. G. Gordon, New Jersey Marine Science Consortium, Sandy Hook: Executive Office, Building 22, Fort Hancock, New Jersey 07732, pers. commun. February 1988).

Fishing Gear Marking

Ghost fishing and entanglement are a widely recognized result of the loss and discard of fishing gear and gear fragments. The MARPOL Annex V explicitly excludes the accidental loss of fishing gear from its plastics discharge prohibition. This is sensible because, as a rule, fishermen balance the cost of replacing gear and the associated lost fishing opportunities against the expected value of their catch. Under most circumstances, this equation limits the risk of gear loss to economically acceptable levels. However, as long as fishing is a legitimate activity, some gear will continue to be lost. The wildlife and vessel hazards presented by this continuing accumulation will remain after all other plastic debris is controlled.

It has been suggested that the control of loss, discard, and abandonment of fishing gear could be improved through the use of marking systems. Nonremovable marks presumably could allow derelict gear to be traced back to its owner so punitive action could be considered. Thus, marking systems might add to fishermen's incentive to avoid loss, cease discard, and put more effort into recovery. The practical application of such marking systems would require a complex administration and a near-foolproof technology to succeed.

Under a grant from NMFS, a review of potential fishing gear tagging methods was conducted. The materials used in commercial fishing gear, their manufacturing and assembly techniques, and their working parameters were reviewed. Marking techniques considered were in the following categories: external tags, implants of various types, color codes, chemical codes, and bonded sheaths. This study (Northwest Marine Technology, Inc. 1989) concluded that it is technically feasible to mark fishing gear and that the optimum system will depend on the gear type. It points out that no matter what type of system is employed, extensive record keeping would be required if vessel-of-origin information is to be retained. This study did not evaluate the socioeconomic or political suitability of application of these techniques for any specific fishery or region.

It remains to be seen whether future improvements in fishing technology and procedures will actually reduce gear losses and increase

recovery rates or merely enable greater risks to be taken. This issue will be given increased attention in the United States in coming years.

The Marine Plastic Pollution Research and Control Act of 1987 required NMFS to report on the utility of using bounty systems and incentive systems to control the loss and discard of fishing gear in the ocean. To address this question, NMFS funded a workshop on these subjects in February 1988 in Portland, Oregon. The workshop concluded that artificial mechanisms to control fishermen's compliance with the regulations implementing MARPOL Annex V would be premature (Alaska Sea Grant 1988). It was recommended that education programs be continued and that such consideration wait until the required reports of compliance are made by the U.S. Coast Guard. If compliance levels are unacceptable then regulatory mechanisms should be explored in consultation with the fishing industry.

Recycling

Efforts to recycle postconsumer plastics have met with a wide variety of successes in recent years. In general, the controlling factors in the economic viability of plastic recycling appear to be the volume, supply, and purity of feedstocks. With few exceptions, subsidies have been necessary to entice recyclers into the mixed, postconsumer plastics arena. The municipal waste streams of urban areas are rich in plastics but require labor-intensive sorting. Technology for automated, mixed-waste sorting is under development, but separating polymer types may not be feasible. In response to this realization, a number of processes and products for recycled mixed plastics have been and are being developed. Current product examples include substitutes for outdoor lumber, watering troughs for farms, and fillers for pillows, padding, and insulation.

Plastics recycling in Japan dates back to 1964. Nylon six gillnets have been actively recycled by melt reprocessing since 1974 (Matsunaga 1988). Products from the recycled nylon six gillnets include automotive and appliance parts, telephones, heels for shoes, golf tees, light structural reinforcements, and plastic reinforced glass products. In recent years, nylon 6 has been largely replaced by nylon 66 in North Pacific fisheries because it is thinner and stronger. Nylon 66 cannot be recycled because of its heterogeneous properties, and Japanese net recycling capacity exceeds the supply of nylon 6 (Matsunaga 1988). Fishing gear recycling is currently unprofitable and must be subsidized by Federal and local governments (Nakamura 1988). Research programs in the Fisheries Agency, the Government of Japan are exploring new processes for recycling fishing gear (Takehama 1988). Aizawa and Satou (1990) report on the disposition and recycling of plastic products including nets.

It is noteworthy that in both Japan and the United States there appears to be a considerable demand for used fishing nets for less demanding fishery applications as well as for nonfishery uses. These uses include shellfish culture, seaweed drying, garden uses, erosion control, sports goals and backstops, and decorations. It is encouraging that domestic demand may absorb at least some of the nonrecyclable gear while recycling and other disposal alternatives are developed. Ports accepting

waste fishing gear under Annex V requirements should explore ways to encourage this demand.

MARINE DEBRIS EDUCATION

Recognizing that littering is chronic in developed and developing nations; that dumping at sea is a time-honored disposal method; that cheap, persistent plastics have changed the nature of the litter problem; and that terrestrial and marine enforcement capabilities are limited at best, realistic progress in the minimization of input of persistent wastes into the marine environment can only be brought about through gross changes in public attitudes and behavior. Education and example are repeatedly identified as the processes for effecting such changes.

Recognition of the ocean and coastlines as valuable national resources is particularly strong among the North Pacific Rim nations. Each of the cultures around the Pacific embodies an ocean ethos, the foundations of which lie in their maritime heritage. This heritage is based on resource utilization, trade, and transportation. A growing appreciation of the relationships between ocean (and environmental) health, productivity, and human use patterns appears to be making these cultural sentiments vulnerable to change. Education programs addressing the marine debris problem are intentionally or unintentionally using the broad appeal of the ocean and coasts to take advantage of this vulnerability. By moving a society's ocean ethos towards the belief that a clean ocean has value, individuals in that society will be less inclined to act counter to that belief.

The maritime heritage, hence the ocean ethos, varies widely among the cultures and subcultures around the North Pacific. To have a lasting effect on the attitudes and behaviors of a subculture (such as regional, ethnic, or industrial), education must be either so general that it does not seriously conflict with the world view or highly specific to that subculture's interests and vulnerabilities. In some cases, an education approach may fit both criteria. Marine debris education programs around the North Pacific have been combinations of both approaches.

Concern over entanglement of northern fur seals and ghost fishing by derelict gillnets in the North Pacific dictated that the first marine debris education program be focused on the fishing industries. In 1983, the North Pacific Fur Seal Commission funded the preparation and distribution of a poster requesting fishermen of Canada, Japan, the U.S.S.R., and the United States to control their discharge of net fragments and packing bands to reduce seal entanglement. Starting in 1985 in the United States, NMFS developed information, documents, and other educational materials for distribution to the fishing fleets of the Pacific Rim nations. Seminars and printed materials were given to every fishing association and fisheries management entity on the U.S. west coast. Formal presentations were made, and printed matter was distributed in Japan, the Republic of Korea, and Taiwan in 1986. Fishing industry associations independently and in conjunction with NMFS carried out marine debris awareness activities and developed and distributed information. The fishing industry sponsored a coast-wide meeting on the marine debris issue for fishermen in Newport, Oregon in July 1986.

The Newport meeting was followed by an international, industry-sponsored North Pacific Rim Fishermen's Conference on Marine Debris in Kailua-Kona, Hawaii, in October 1987. Sponsorship and participation in this conference came from fishing industry groups and associations from Canada, the Republic of China, Japan, the Republic of Korea, and the United States. The conference recommended a set of marine debris research priorities and adopted a resolution declaring the fishing industries' commitment to controlling their part of the marine debris problem. A group of fishing industry associations on the U.S. west coast used this resolution to develop an engraved plaque entitled "Fishermen's pledge for a clean ocean." The proceedings of this conference (Alverson and June 1988) are a valuable source of information about North Pacific marine debris programs and actions.

In response to all manner of inquiry from the public, NOAA established a west coast Marine Debris Information Office in San Francisco in late 1988. This office distributes 16 separate packages of general marine debris information depending on the nature of the request received. Requests for information may be mailed to:

Center for Marine Conservation
Marine Debris Information Office, NOAA
312 Sutter Street, Suite 606
San Francisco, CA 94108 U.S.A.

The broadest possible audience has been sought through the production of a variety of posters, brochures, and videotapes. An award winning 7 1/2-m video called "Trashing the oceans" was produced in 1987 and has been shown all over the world. This production is suitable for general audiences and is available from NOAA or the Marine Debris Information Offices. The NOAA, the Society of the Plastics Industry, and the Center for Marine Conservation (CMC) worked together to develop and distribute brochures and public service advertisements through marine trade journals and magazines nationwide (Bruner 1990; Debenham 1990).

Judie Neilson first organized large-scale, public, voluntary beach cleanups focusing on the persistent waste problem in Oregon in 1984 (Neilson 1985). The idea has caught on in every coastal state in the United States as well as in Japan. In 1988 in the United States, 47,500 volunteers cleaned 5,630 km (3,500 mi) of shoreline, removing almost 1,000 tons of trash. These data were collected by the volunteers and assembled, analyzed, and reported by CMC with support from Federal and private sources (CMC 1989). The results of the cleanups have been widely reported in local and national media, used in congressional testimony, and incorporated into ever-broadening education programs.

In an attempt to ensure the widest possible understanding of the requirements of MARPOL Annex V and to build the basis for compliance, NMFS has contracted for the development and implementation of a shipping and cruise lines industry education program (Wallace 1990). This activity is directed at all vessels and vessel operators plying U.S. waters, regardless of nationality. Products from this work will include model shipboard waste

management plans, summaries of the U.S. regulations implementing MARPOL Annex V, and packets of information on the impacts of marine debris. Delivery of these materials will be through corporate offices, union offices, and associations of shipowners and officers.

Programs initiated by the Fisheries Agency, the Government of Japan in recent years have included fishing industry and public education components (Yagi and Otsuka 1990). Voluntary beach cleanups have been organized in the coastal prefectures. Seabed cleanups involving fishermen and divers are being carried out in ports and high-use coastal areas. In 1986, 3,959 km² in 25 separate areas were cleaned. Several general video presentations on the marine debris problem, on cleanups, and on the national marine debris program have been produced for wide national and international distribution.

The Republic of Korea has developed an education program and a set of regulations to control the discharge of waste fishing gear from its fishing vessels (Lim 1988). Each year, vessel captains are required to attend a training session by the National Fisheries Research and Development Agency which includes marine debris education. The admiral of the Korean Deep Sea Fisheries Association is charged with educating Korean fishermen against discharging entangling materials. The full extent of Korean and other Pacific Rim countries' marine debris education activities, apart from fishing industry actions, is unknown.

Finally, in an effort to raise the world level of understanding and appreciation of all facets of the marine debris problem, the NMFS initiated and acted as principal sponsor for the Second International Conference on Marine Debris.

CONCLUSIONS

The recommendations from the 1984 FIMD workshop have not been fully met. Research on the impacts of marine debris on wildlife has not established a clear understanding of the role of entanglement and ingestion in the population fluctuations of any marine species. Efforts to measure the sources and amounts of persistent debris have been greater in the North Pacific than in any other ocean area, but a full understanding of the dynamics of input, output, and circulation remains well in the future.

However, since the FIMD workshop, mitigation and education efforts have enjoyed the highest priorities. By international standards, legal and administrative actions to address the marine debris problem have progressed rapidly. The entry into force of MARPOL Annex V marks the primary international step in controlling ship-generated debris. Technological solutions for solid waste in the oceans and on land are now receiving major attention from government and industry sectors around the Pacific Rim. Education programs continue to expand, reaching people all over the world, even though the high level of international cooperation recommended by the FIMD workshop has not been achieved. As domestic policy and problems are addressed, the opportunity for, and suitability of, international action will increase.

Research

The research community addressing persistent waste pollution of the oceans is in a period of transition. Researchers, particularly biologists, initially noted the effects of marine debris as an oddity, not necessarily associated with their primary research. Since the FIMD workshop and the entry into force of MARPOL Annex V, the study and understanding of this type of pollution has become a legitimate, although minor, activity purposely incorporated into marine research agendas. As this evolution proceeds, definitions of terms are accepted, methods of inquiry are shared and generalized, new disciplines are involved, the literature is established, and the underpinnings of a new subdiscipline are solidified. The next 5 years will undoubtedly see the recognition of a marine faction of the solid waste research community including biologists, chemists, oceanographers, engineers, economists, lawyers, and possibly an institution or two. The work of this community is likely to be more applied than basic, as the immediate problems of solid waste management at sea and along the coasts must be solved to comply with current and future statutes. This emphasis will likely result in the diversion of limited resources from research on the biological impacts of marine debris.

While North Pacific species (northern fur seal, Hawaiian monk seal, Laysan albatross) have been the most intensively studied, inability to field pelagic research programs continues to prevent full elucidation of the role of debris in population changes. Increased knowledge of the behavioral and physical mechanisms of impact and the materials involved has strengthened the deductive evaluations of effects on populations, particularly for northern fur seals. Should international field research programs be developed for North Pacific high seas driftnet fisheries, information may become available to assess further the impacts of marine debris on fur seals. The experiments necessary to finally assess the physical impact of plastic ingestion on seabirds and sea turtles should be undertaken immediately. Studies on the possible toxic effects of plastic ingestion should also be initiated.

Specific regional studies of the direct and indirect costs to coastal communities resulting from debris are overdue. Collection of information on the incidence of vessel damage caused by persistent marine debris has been sporadic and mostly anecdotal. There have been few recent studies of the impacts of ghost fishing on target or nontarget species, on fisheries production, or on profitability. These types of information are essential for evaluating the range of economic impacts of debris and for crafting appropriate solutions. Clear, broadly applicable models for evaluating the economic impacts of accumulating marine debris would be valuable tools used worldwide.

Mitigation

Marine debris mitigation is proceeding apace. Laws are being passed, attitudes are changing, and industries are getting the message. The relationship between marine debris and the overall solid waste crisis is the real key here. The marine problem will not be completely solved until the

land-based disposal problem is solved. Broad public concern for the ocean and coastal environment has allowed a start to be made; there can be no turning back. The timetable for control depends on the rate of development and implementation of rational solid waste disposal practice. For the near future, two areas of emphasis are evident. First, the remaining North Pacific Rim nations must accede to MARPOL and ratify Annex V. Second, the focus for the next several years must be on technology, policy, processes, regulation, enforcement, and education to fully implement MARPOL Annex V. On land, appropriate combinations of source reduction, recycling, incineration, substitution, and use of landfills must be sought. Within 2 to 5 years, control of persistent waste discharge into the ocean could be fully institutionalized around the North Pacific Rim. This endeavor assumes increased international cooperation in the provision of port reception facilities, in enforcement, and in promoting responsible waste management by all maritime nations.

Education

At least in the United States, the power of ocean issues to stir public action has been increasing for several decades. The marine debris issue has become a major rallying point for advocates and educators alike, catalyzing public awareness and action on an array of environmental issues. A high level of volunteerism is being achieved in public and industry education programs. Apparently, the United States and much of the developed world are ready to accept the responsibility for solving the marine debris problem. It is an issue whose time has come, one that may open the way for increased public insistence on, and acceptance of, a more responsible environmental policy.

Each nation must develop and effectively distribute information on its laws and regulations to implement MARPOL Annex V or its national equivalent. Timely, informed assistance in this implementation phase is critical to the long-term public acceptance of these requirements. All vessel operators and ports need this assistance. It should be noted that widely publicized examples of enforcement actions can be highly educational.

For the immediate future, existing education materials should be translated and adjusted as necessary for broad international use. The beach cleanup programs should be expanded to include all coastal nations. The cleanup data should be reported as widely as possible. Outlets for marine debris education materials should be established and publicized by all national and international agencies having environmental or maritime responsibilities. These are all low-cost, highly credible activities that should appeal to most governments and organizations. After all, who is willing to say, "I support marine debris?"

Finally, the Third International Conference on Marine Debris should be held in 1994 or 1995 to document world progress on this issue.

REFERENCES

Aizawa, M., and A. Satou.

1990. Disposition and recycling of plastic products including used nets. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Alaska Sea Grant.

1988. Oceans of plastic: A workshop on fisheries generated marine debris and derelict fishing gear. Report to the National Marine Fisheries Service. Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Seattle, WA 98115, 65 p.

Alig, C. S., L. Koss, T. Scarano, and F. Chitty.

1990. Control of plastic wastes aboard naval ships at sea. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Alverson, D. L., and J. A. June (editors).

1988. Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA, 98199, 460 p.

Andrady, A. L.

1990. Environmental degradation of plastics under land and marine exposure conditions. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Balazs, G. H.

1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Balazs, G. H., and B. K. Choy.

1990. Ecological aspects of marine turtles impacted by ocean debris: A 1989 perspective. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Bengtson, J. L., C. W. Fowler, H. Kajimura, R. Merrick, K. Yoshida, and S. Nomura.

1988. Fur seal entanglement studies: Juvenile males and newly-weaned pups, St. Paul Island, Alaska. *In* P. Kozloff and H. Kajimura (editors), *Fur seal investigations*, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.

Berger, J. D., and C. E. Armistead.

1987. Discarded net material in Alaskan waters, 1982-84. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-110, 66 p.

Bruner, R. G.

1990. The plastics industry and marine debris: Solutions through education. *In* R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Center for Environmental Education.

1988. Texas coastal cleanup report. Center for Environmental Education, Wash., D.C., 105 p.

Center for Marine Conservation.

1989. Trash on America's beaches: A national assessment. Center for Marine Conservation, Wash., D.C., 30 p.

Chang, T. J.

1990. Low technology (burn barrel) disposal of shipboard-generated (MARPOL V) wastes. *In* R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Day, R. H.

1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.

Day, R. H., D. G. Shaw, and S. E. Ignell.

1990. The quantitative distribution and characteristics of neuston plastic in the North Pacific Ocean, 1984-88. *In* R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Day, R. H., D. H. S. Wehle, and F. C. Coleman.

1985. Ingestion of plastic pollutants by marine birds. *In* R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-54.

Debenham, P.

1990. Education and awareness: Keys to solving the marine debris problem. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Fiscus, C. H., and P. Kozloff.

1972. Fur seals and fish netting. Appendix E. In Fur seal investigations, 1971, p. 124-132. Marine Mammal Biological Laboratory, National Marine Fisheries Service, Seattle, WA.

Fowler, C. W.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Fowler, C. W., R. Merrick, and J. D. Baker.

1990. Studies of the population level effects of entanglement on northern fur seals. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Gerrodette, T., B. K. Choy, and L. M. Hiruki.

1987. An experimental study of derelict gill nets in the central Pacific Ocean. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-18, 12 p.

Henderson, J. R.

1985. A review of Hawaiian monk seal entanglements in marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu Hawaii, p. 326-335. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFS-54.
1988. Marine debris in Hawaii. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 189-206. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
1990. Recent entanglements of Hawaiian monk seals in marine debris. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Henderson, J. R., S. L. Austin, and M. B. Pillos.

1987. Summary of webbing and net fragments found on Northwestern Hawaiian Islands beaches, 1982-86. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-11, 15 p.

International Maritime Organization.

1988. Guidelines for the implementation of Annex V of MARPOL 73/78. International Maritime Organization, 4 Albert Embankment, London SE1 7SR, United Kingdom, 34 p.

Johnson, S. W.

1990. Distribution, abundance, and source of entanglement debris and other plastics on Alaskan beaches, 1982-88. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Johnson, S. W., and T. R. Merrell.

1988. Entanglement debris on Alaskan beaches, 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-126, 26 p.

Jones, L. L., and R. C. Ferrero.

1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

June, J. A.

1990. Type, source, and abundance of trawl-caught marine debris off Oregon, in the eastern Bering Sea, and in Norton Sound in 1988. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Kenyon, K. W., and E. Kridler.

1969. Laysan albatrosses swallow indigestible matter. Auk 86:339-343.

Lim, J.

1988. Korean activities on avoidance of incidental mortality of marine animals by debris. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 304-319. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

- Loughlin, T. R., P. Gearin, R. L. DeLong, and R. Merrick.
1986. Assessment of net entanglement on northern sea lions in the Aleutian Islands, 25 June-15 July, 1985. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 86-02, 50 p.
- Lutz, P. L.
1986. Effect of ingestion of non-biodegradable debris in sea turtles. Report to the U.S. National Marine Fisheries Service, Marine Entanglement Research Program, Contract No. FSN-5-0178, 50 p. Available from Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Seattle, WA 98115.
- Martinez, L. A.
1990. Shipboard waste disposal: Taking out the trash under the new rules. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Matsunaga, S.
1988. Recycling of used nets: The status quo and its problems. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 429-433. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Merrell, T. R.
1980. Accumulation of plastic litter of beaches on Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

1985. The fish nets and other plastic litter on Alaska beaches. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 160-182. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Mio, S., and S. Takehama.
1988. Estimation of distribution of marine debris based on the 1986 sighting survey. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 64-94. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Mio, S., T. Domon, K. Yoshida, and S. Matsumura.
1990. Preliminary study on change in shape of drifting nets experimentally placed in the sea. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Nakamura, I.

1988. Japan: Tackling the marine debris problem. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 374-378. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

National Oceanic and Atmospheric Administration.

1988. Report of the interagency task force on persistent marine debris. Natl. Oceanic Atmos. Admin., Wash., D.C., 170 p.

Neilson, J.

1985. The Oregon experience. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 154-159. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Northwest Marine Technology, Inc.

1989. Marking of fishing gear. Final report to the U.S. National Marine Fisheries Service, Saltonstall-Kennedy Grant No. NA-88-ABH-00028, 107 p. Available from Northwest Regional Office, National Marine Fisheries Service, NOAA, 7600 Sand Point Way N.E., Seattle, WA 98115.

Ogi, H.

1990. Ingestion of plastic particles by sooty and short-tailed shearwaters in the North Pacific. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Parker, N. R., S. C. Hunter, and R. J. Yang.

1987. Development of methodology to reduce the disposal of non-degradable refuse into the marine environment. Report to the U.S. National Marine Fisheries Service, Marine Entanglement Research Program, Contract No. 85-ABC-00203, 87 p. Available from Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Seattle, WA 98115.

Recht, F.

1988. Dealing with Annex V - reference guide for ports. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWR-23, 132 p.

Reed, D. K., and J. D. Schumacher.

1985. On the general circulation in the subarctic Pacific. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu Hawaii, p. 483-496. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Ribic, C. A., and L. J. Bledsoe.

1986. Design of surveys for density of surface marine debris in the North Pacific. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 86-12, 69 p.

SCS Engineers, Inc.

1989a. An investigation of using burn barrel technology to dispose of shipboard-generated (MARPOL V) wastes. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 89-15, 33 p.

1989b. Operating and safety guidelines for use of burn barrels to dispose of shipboard-generated (MARPOL V) wastes. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 89-14, 15 p.

Shomura, R. S. and M. L. Godfrey (editors).

1990. Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154, 1274 p. [See this document.]

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

Sileo, L., P. R. Sievert, and M. D. Samuel.

1990. Causes of mortality in albatross chicks at Midway Atoll. J. Wildl. Dis. 26:329-338.

Sileo, L., P. R. Sievert, M. D. Samuel, and S. I. Fefer.

1990. Prevalence and characteristics of plastic ingested by Hawaiian seabirds. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Takehama, S.

1988. Marine debris program in Japan. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 372.1-372.10. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Walker, W. A., and J. M. Coe.

1990. Survey of marine debris ingestion by odontocete cetaceans. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Wallace, B.

1990. Shipping industry marine debris education plan. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Washington State Department of Natural Resources.

1988. Marine plastic debris action plan for Washington State. Available from Washington State Department of Natural Resources, Commissioner of Public Lands, Olympia, WA 98504, 45 p.

Yagi, N., and M. Nomura.

1988. Sighting survey on marine debris in the midwestern Pacific from 1977 through 1986. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 130-142. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Yagi, N., and Y. Otsuka.

1990. Cleanup program in Japan. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

OVERVIEW: MARINE DEBRIS IN THE NORTHWEST ATLANTIC OCEAN

Burr Heneman
Marine Mammal Commission
35 Horseshoe Hill Road
Bolinus Star Route, California 94924, U.S.A.

ABSTRACT

This review emphasizes recent developments (since the author's 1988 report) in regard to marine debris sources, types, amounts, and distribution, effects, and mitigation, on the Atlantic coasts of Canada and the United States.

INTRODUCTION

A substantial body of information about sources, types, amounts, and effects of marine debris exists for the northwest Atlantic Ocean, most of which is summarized in a report (Heneman 1988) distributed to participants at this conference. This presentation includes general observations based on that report but emphasizes new developments.

For our purposes, the northwest Atlantic reaches from the Atlantic coast east to midocean and south to, and including, the North Equatorial and Antilles Currents. Its western watershed, which includes the St. Lawrence and many lesser rivers, drains the most densely populated and industrialized areas of the United States and Canada.

SOURCES, TYPES, AMOUNTS, AND DISTRIBUTION

In contrast to areas of the world where a few sources account for most marine debris, the northwest Atlantic is plagued by a great variety of major debris sources. Merchant shipping, commercial fishing vessels, cruise ships, recreational boats, and naval vessels may be the largest sources, although MARPOL Annex V should cause these to diminish in importance. At the same time, inadequate storm drain and sewage treatment systems in the United States and Canada are known to dump large amounts of floatables into the marine environment, especially in periods of high rainfall; coastal landfills commonly "leak" debris into nearby waters; the plastics industry in the northeastern United States appears to have been a major source of plastic resin pellets; and beachgoers are an important source of litter. As we have seen with medical wastes for the past two summers, relatively small amounts of illegally dumped materials can have major effects. Virtually every kind of debris source that has been

identified anywhere in the world is a contributor somewhere in the northwest Atlantic. This variety of major sources obviously complicates efforts to reduce amounts of marine debris and to mitigate its effects.

It is more difficult to generalize about where debris occurs in the northwest Atlantic than in a trade wind area such as the Caribbean. The North Atlantic gyre concentrates floating debris in the Sargasso Sea and on the beaches of Bermuda. Along the gyre's southern periphery, trade winds deposit large amounts of debris from the Antilles Current onto Atlantic-facing beaches in the Bahamas. Farther north, local sources and local wind and current conditions are more important factors influencing the distribution of debris on the United States and Canadian coasts.

There is little information on trends in amounts of marine debris. Wilber (pers. commun.) points out that his data and Carpenter and Smith's (1972) data for the northern Sargasso Sea indicate a 1,000% increase in the density of plastic pieces and a 200-400% increase in plastic pellets in a period of about 15 years.

There is little recent information to report from Canada on sources, amounts, and distribution of debris. Canada's Ocean Policy of 1987 includes commitments to deal with plastic debris and lost and abandoned fishing gear, but little has been done to implement the policy. Growing public concern may be leading to a change, however. Last summer, for example, the Nova Scotia Department of the Environment conducted one of Canada's first beach cleanups. An opinion survey at the same time found increasing indignation about litter on beaches.

EFFECTS

The best-known and most serious effects of marine debris along the northwest Atlantic coast are aesthetic and economic; the summer of 1988 provided another well-documented example of that when tourist-dependent coastal economies lost tens of millions of dollars to beach closures in the New York area. This is not a new problem, however; the first major incident of this sort was in the summer of 1976, when sewage and debris closed Long Island beaches and the Governor of New York declared a disaster.

Other effects, such as damage to vessels and harm to wildlife, are either minor or are poorly documented. At the Workshop on the Fate and Impact of Marine Debris (FIMD) in 1984, participants agreed that the effects of debris on sea turtles and of derelict nets and traps on fish and shellfish deserved greater attention (Shomura and Yoshida 1985). That is especially true for the northwest Atlantic, where these subjects may represent the most important information gaps.

ACTION AND MITIGATION

Two new programs in the United States are collecting information on types, sources, and amounts of debris. The Marine Entanglement Research Program and the U.S. National Park Service are sponsoring regular data

collection at eight national seashores, including four on the Atlantic coast: Cape Cod, Assateague Island, Cape Hatteras, and Cape Canaveral.

The U.S. Environmental Protection Agency (EPA) has funded at least 1 year of a National Marine Debris Data Base, in which the Center for Marine Conservation is computerizing data from all the 1988 statewide volunteer beach cleanups. Over time, these two programs may provide a means of monitoring the success of Annex V and other mitigation measures.

On the Atlantic coast of the United States, mitigation efforts such as education and public awareness campaigns have focused on implementation of Annex V. The Marine Entanglement Research Program has funded several projects through the Center for Marine Conservation, including:

- a Marine Debris Information Office located in Washington, D.C. to respond to information requests from the Atlantic and Gulf coasts. It provides educational materials to marine user groups, industry, educators, policy makers, and the general public;
- separate public service advertisement campaigns aimed at the commercial fishing, shipping, and plastics industries, and recreational boaters and fishermen;
- a review of marine debris information for the general public, "A Citizen's Guide to Plastics in the Ocean."

The Society of the Plastics Industry helped fund the Citizen's Guide, public service announcements for television, and other marine debris educational materials produced by the Center for Marine Conservation.

Another Center for Marine Conservation project, this one in Florida and funded by the National Marine Fisheries Service Saltonstall-Kennedy program, endeavors to show that education is a cost-effective method of persuading commercial and recreational fishermen to comply with Annex V.

There have been continuing and expanding efforts to remove debris from the marine environment. For instance, most coastal states have had annual beach cleanups in recent years. The Army Corps of Engineers, the EPA, the U.S. Coast Guard, and New York and New Jersey state agencies recently announced that they have begun a cooperative program in the New York area. They will try to locate concentrations of floating debris by helicopter and use Army Corps vessels to collect it.

Canada's Department of Fisheries and Oceans convened a workshop in Halifax, Nova Scotia, 17-18 May 1989. The workshop provided an opportunity for organizations and individuals from the private sector to advise the government on the development of an action plan on marine debris (Buxton 1989; DPA 1989).

As for mitigation efforts, Canada has placed itself in an unusual position. Although Canada is a signatory to the London Dumping Convention,

it is not a signatory to MARPOL, much less to Annex V. For some years, the Canada Shipping Act has prohibited the disposal of any garbage or trash from vessels within 200 nmi of Canada's Atlantic and Pacific coasts, a provision that is stricter than Annex V. Unlike Annex V, however, the act does not restrict ocean disposal by Canadian vessels beyond 200 nmi, and it does not require ports to provide reception facilities.

Recent amendments to the Canada Shipping Act take a half step forward by *permitting* Canadian agencies to impose stricter regulations that would bring Canada into conformity with Annex V. But the agencies have not yet decided to actually adopt any new restrictions. Furthermore, there seems to be little enforcement of existing regulations and no educational programs to encourage compliance.

CONCLUSION

Although the Atlantic coast of the United States has the same marine debris problems, more or less, as other coastal areas of the country and the world, its problems receive more attention than is warranted simply by its geography. United States policy makers are concentrated in Washington, D.C. National, and to some extent international, opinion shapers are concentrated in New York City. As a result, events in that part of the world become more important.

To mention two examples: The cover story in Time magazine for 1 August 1988 is titled "Our Filthy Seas." That same week, Newsweek's cover story was "Don't Go Near the Water--Our Polluted Oceans." An issue has truly arrived on the national agenda when it makes the covers of these two magazines the same week, when it is a regular fixture on network news, and when it is an issue in a presidential campaign, as it was in 1988. The fact is, the response to marine debris problems on the Atlantic coast will continue to have a disproportionate influence on how the rest of the United States responds to its marine debris problems.

It has become abundantly clear since the 1984 FIMD workshop that the ultimate solutions to marine debris problems on the U.S. Atlantic coast are inextricably bound to solutions to the impending crisis in solid waste disposal on land. All of the elements that can contribute to reducing amounts and effects of marine debris--source reduction, recycling, degradability, changing societal attitudes towards waste--are vital in the larger arena of land disposal. That fact should inform much of our effort in regard to the marine debris subset of the problem.

REFERENCES

- Buxton, R.
 1989. Plastic debris and lost and abandoned fishing gear in the aquatic environment: A background paper from a Canadian perspective for a workshop held in Halifax, May 1989. Department of Fisheries and Oceans, Ottawa.

Carpenter, E. J., and K. L. Smith.

1972. Plastics on the Sargasso Sea surface. *Science* 175(4027):1240-1241.

DPA.

1989. Plastic debris in the aquatic environment--Halifax Workshop report. Department of Fisheries and Oceans, Ottawa.

Heneman, B.

1988. Persistent marine debris in the North Sea, northwest Atlantic Ocean, wider Caribbean Area, and the west coast of Baja California. A report to the Marine Mammal Commission and the National Ocean Pollution Program Office, National Oceanic and Atmospheric Administration, Department of Commerce, Washington, D.C., 160 p.

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

PLASTICS: ACCUMULATION, DISTRIBUTION, AND ENVIRONMENTAL
EFFECTS OF MESO-, MACRO-, AND MEGALITTER IN SURFACE
WATERS AND ON SHORES OF THE SOUTHWEST PACIFIC

Murray R. Gregory
Department of Geology
University of Auckland
Auckland, New Zealand

ABSTRACT

Plastic debris of all kinds and in all sizes is widespread in the southwest Pacific. Densities of virgin nibs exceed $1,000 \text{ km}^{-2}$ in surface waters north of New Zealand and in nearshore waters adjacent to manufacturing centers. There is a latitudinal gradient of densities, with numbers falling to less than 20 km^{-2} south of New Zealand. On shorelines, greatest numbers ($>>100,000 \text{ m}^{-1}$ of beach length) are found near large cities, although a similar latitudinal gradient shows with very low numbers from around southern New Zealand ($1-5 \text{ m}^{-1}$) and none from the subantarctic islands. In general, numbers of nibs on shores of eastern Australia are much less than they are on New Zealand shores. Significant numbers ($>1,000 \text{ m}^{-1}$) have been found as local concentrations on some trade wind-facing beaches of all Pacific islands so far examined.

Distribution of these nibs, together with that of other plastic and persistent synthetic litter, is influenced by surface current patterns and prevailing wind regimes, with greatest concentrations being noted on windward and downdrift shores, in windrows, and (tentatively) along oceanic fronts.

Larger, fabricated plastic items have been seen on the shores of all isolated and unpopulated islands so far visited around the region. Where identifiable, sources frequently lie in distant water fishing activities. On populated islands, many of which lack adequate facilities for domestic waste and garbage disposal, there is a buildup of locally sourced litter along shores. Not only is this litter aesthetically distasteful, some materials (e.g., syringes) are hygienically unacceptable. The problem is an ever-growing one and needs addressing in appropriate forums. The environmental implications of this plastic pollution are many, with the most important involving entanglement and ingestion. The longer term significance of hazardous and persistent chemical residues, originally present in plastics as additives and released in minor amounts during degradation,

is difficult to assess. Pelagic plastics also provide an important hard substrate for an encrusting biota that includes a hermatypic coral, bryozoans, coralline and filamentous algae, hydroids, barnacles, and some foraminifers, and are a largely unrecognized vector in their wider distribution.

From surface crazing and other evidence of aging such as chalkiness and embrittlement, it is inferred that degradation rates decrease progressively from lower to higher latitudes.

INTRODUCTION

It is generally accepted that surface waters of the South Pacific Ocean (Fig. 1) are relatively free from man-made pollutants, other than in the nearshore zone of more heavily populated islands (Matos 1981). Recent reviews have tended to emphasize localized incidents involving point-sources of sewage and industrial effluents (e.g., Suva Harbor, Fiji: Brodie and Morrison 1984), and toxic chemicals and pesticides (Cook Islands: Hambuechen 1973; and Tonga: Brodie and Morrison 1984; Morrison and Brodie 1985), although wider political concern has been expressed over the prospect of seabed disposal and dumping or storage of nuclear waste in the expanses of the region (Branch 1984; Carew-Reid 1988). The area lies remote from tanker routes (Waldichuk 1977) and major shipping lanes, and pelagic tar balls so common to more frequently traversed waters are rarely encountered (Butler et al., 1973, p. 24; Bourne 1976; Gregory 1977, unpubl. data; Oostdam 1984; Lee pers. commun.). The problems of marine oil pollution become more evident passing westward into southeast Asian waters (Bilal 1985). However, the island countries of the southwest Pacific have a long and commonly expressed concern over contingency planning for pollution from oil spills (Hayes 1981; Dahl and Baumgart 1983; Hayes and Kay 1986).

Plastics and other persistent synthetic materials are today a significant contaminant of both open ocean and nearshore waters, particularly those adjacent to the industrial North. The sources and environmental problems they create are many and varied (Gregory 1978, 1983; Laist 1987; Pruter 1987). Plastic artifacts as well as casual litter and solid domestic wastes have long been an acknowledged, although seldom seriously addressed, problem on several Pacific islands (Anonymous 1976; Connor 1976; Efi 1976). On Tonga, for example, plastics and cigarette and candy wrappers have been identified as ". . .the second most common form of litter and the second largest waste item for disposal" (Chesher 1984, p. 38). In all instances known to this author, the importance of local sources has been noted, with little recognition that some material may have been adrift for a time before stranding. The observations of Sachet (1955) on the wide dispersal of exotic pumice on Pacific atolls, as well as those of Bligh (1792) on coconut husks, are evidence that, over the vastness of the Pacific, floating materials can drift far from their places of origin. Drift pumice, often with an encrusting biota, is common on beaches of eastern Australia (Table 1). Similarly, in the Southern Ocean there is evidence of floating debris such as logs, pumice, and man-made artifacts being rapidly dispersed in circumpolar fashion by the strong West Wind




Figure 1.--Map of the southwest Pacific Ocean indicating principal places mentioned in text. Major oceanic features are also illustrated.

Drift Current and general oceanic circulation patterns (Barber et al. 1959; Gregory et al. 1984; Smith 1985; Gregory 1987, 1990; Lutjeharms et al. 1988).

Waters around New Zealand and its offshore islands are by any criteria relatively unpolluted, although semienclosed estuaries and harbors in the vicinity of larger urban centers give increasing cause for concern (Ridgway and Glasby 1984). Plastics and other persistent synthetic compounds,

Table 1.--Numbers of virgin plastic granules and drift pumice on selected beaches of eastern Australia (arranged from north to south). The quantities of granules are local maximums expressed in number per linear meter of shore, following the approach of Gregory (1978); p = present in low numbers ($<1 \text{ m}^{-1}$). Drift pumice: * = abundant, + = present.

Location	Plastic granules	Drift pumice
Tasmania		
Hobart to Bicheno	nil	nil
Victoria		
Portsea, Sorrento, Rosebud	p	nil
Mordialbo	>1,000	nil
St. Kilda	>500	nil
Altona	100	nil
New South Wales		
Narooma	nil	+
Batemans Bay	nil	*
Kioloa	nil	*
Jervois Bay	p	nil
Shoalhaven	>5	*
Stanwell Park	>50	*
Botany Bay	>>2,000	+
Bondi	>10	+
Manly	p	nil
Narrabeen	>20	nil
Port Macquarrie	p	*
Coffs Harbor	nil	*
Queensland		
Gold Coast	p	*
Brisbane (Red Cliffs)	nil	+
Bargara (Bundaberg)	5	*
Keppel Sands (Rockhampton)	5	*
Sarina	nil	*
Mackay	p	*
Townsville	nil	*

particularly those arising from packaging, are a significantly visible but minor part of the local waste stream (Ministry for the Environment 1987; Plastics Institute of New Zealand 1988). The environmental hazards and threats to local wildlife are varied and have been reviewed by a number of authors (Gregory 1977, 1978, 1987, 1990; Gregory et al. 1984; Cawthorn 1985, 1987; Mattlin and Cawthorn 1986; Dawson and Slooten 1987; Murray 1988).

Gregory (1977, 1978) initially recorded small virgin plastic resin granules and pellets in surprisingly high quantities on the New Zealand coast and mapped their distribution (Fig. 2). It was noted that greatest numbers occurred near metropolitan centers, suggesting that the distribution was caused by dispersal from local sources (Fig. 2), although some evidence indicated possible drift from eastern Australia waters (Gregory 1978). Changes in the composition of litter stranding on a remote northern New Zealand beach over an 8-year period have been recorded by Hayward (1984). Ever-increasing fishing activities add further to the seaborne litter load on even the most isolated shores (e.g., Auckland and Campbell Islands, Cawthorn 1985; Gregory 1987, 1990).

This paper reviews in detail the nature, characteristics, quantities, distribution and sources of pelagic plastics around the southwest Pacific region. It is based largely on the author's published studies from New Zealand and its offshore, subantarctic islands. However, the opportunity has been taken to include a corpus of previously unpublished data gathered from eastern Australia, several Pacific islands, and adjacent waters during opportunistic surveys over a number of years. The environmental consequences of this plastics pollution are evaluated and some conclusions reached on how they could be addressed.

PLASTIC MESOLITTER

In the category of plastic mesolitter I include the small, ovoidal-to-rounded and rod-shaped virgin plastic granules or nibs of polyethylene and polystyrene resins that are the raw materials or feedstock of plastic fabricators worldwide. The granules are mostly <5 mm across, are colorless to translucent or transparent, and have been described in detail previously (Gregory 1978, 1983). Intensely colored dye-carrying granules (yellow, blue, green, red, black, white) are never as common as the colorless ones. In addition, there are occasional sharply angular and jagged plastic chips of comparable size produced through granulation of larger items for recycling. These chips are variously colored but rarely transparent or translucent. Small, often flaky, fragments coming from the degradation and disintegration of larger plastic objects also fall into this category. The fragmenting and fracturing processes appear to be mostly embrittlement through oxidative aging and photodegradation rather than physical or mechanical weathering.

Gregory (1977, 1978) described the distribution of virgin plastic granules on New Zealand shores (Fig. 2). Large numbers, often $>10,000 \text{ m}^{-1}$ of beach length and in one instance $\gg 100,000 \text{ m}^{-1}$, were recorded near some of the larger metropolitan and industrial areas where plastics fabricators are located (Fig. 2). Away from these regions numbers decreased, but they were persistently and surprisingly high at some remote localities (e.g., to $>150 \text{ m}^{-1}$ near North Cape; and to 50 m^{-1} near East Cape). Only around the southernmost part of South Island were they consistently very rare or absent. For the mid-1970's it was estimated that at least 1,000 metric tons of these granules were stranded on the shores of New Zealand (Gregory 1978).




Figure 2.--Distribution of virgin plastic granules on New Zealand shores based on a 1972-78 survey. Three values given for each distribution line indicate abundance levels at which pellets were (i) reasonably consistent (lowest value, top of list), (ii) commonly encountered (middle value), and (iii) locally concentrated (greatest value). Data are from Gregory (1978). Local New Zealand sources of virgin plastic granules are after Bullen (1968); the surface currents and prevailing winds that spread them around and along the coast are after Brodie (1960).

Virgin plastic granules have been encountered on the shores of eastern Australia from Batemans Bay in New South Wales north to Townsville in Queensland (Table 1). They are also present around Melbourne and Adelaide. None have been noted on eastern Tasmanian shores northward from Hobart. Occurrences are sporadic, and numbers seldom reach those recorded from New Zealand. On remote beaches numbers are generally $<1 \text{ m}^{-1}$, and in many instances a lengthy search is required to turn up any granules at all. It is only at a few isolated localities around Sydney ($>2,000 \text{ m}^{-1}$) and Melbourne ($>1,000 \text{ m}^{-1}$) that quantities ever approach those frequently recorded near Auckland.

No virgin plastic granules have been found so far on any of New Zealand's subantarctic islands (e.g., Campbell, Auckland, Snares, Antipodes, and Bounty) (Gregory 1987), although they are not uncommon on Chatham Island (to $>100 \text{ m}^{-1}$, Gregory 1978). The granules, however, have been found on all subtropical and tropical southwest Pacific islands that were systematically searched by this author during visits over the past few years (Table 2, Figs. 3-8). In several instances the numbers are unexpectedly high for such remote, nonindustrialized places (e.g., Tonga, $>>1,000 \text{ m}^{-1}$).

The angular granules produced for recycling are never common away from the industrial centers of Australia and New Zealand, and have not been encountered on the shores of those Pacific islands so far examined.

The numbers of plastic granules and larger plastic items afloat in surface waters of the New Zealand sector of the Southern Ocean have been determined from over 50 neuston tow stations (Gregory et al. 1984; Gregory 1987, 1990). The numerous reports of Southern Ocean feeding seabirds ingesting plastic granules and other artifacts (Bourne and Imber 1982; Furness 1983; Randall et al. 1983; Brown et al. 1986; Skira 1986; Gregory 1987, 1990; Harper and Fowler 1987; Ryan 1987a) indicate these materials have circumpolar dispersal. Brief and sporadic surveys of pelagic plastic have been undertaken from research vessels on passage between New Plymouth and Norfolk Island, Tauranga and Raoul Island, and the Hauraki Gulf to Wellington by way of East Cape as well as around Auckland Harbor and its approaches. At this time data are inadequate to draw unequivocal conclusions. The data strongly suggest, however, that densities in surface waters to the north of New Zealand probably (and often substantially) exceed $1,000 \text{ km}^{-2}$ (Fig. 9). Indeed, fresh granules stranding along the most recent swash line (by inference over one tidal cycle--February 1988) on Raoul Island at $5-10 \text{ m}^{-1}$ suggest that densities approaching $10,000 \text{ km}^{-2}$ may occur sporadically! Variation in numbers between stations is very large. There is apparently a strong latitudinal gradient in the areal density of floating granules (Fig. 9). In higher latitudes between the Subtropical Convergence and the Subantarctic Front, granules occur in numbers that may barely reach 20 km^{-2} (Gregory et al. 1984; Gregory 1987, 1990). Densities farther south and in the region of seasonal pack ice are negligible. In some nearshore waters much higher densities are commonplace (e.g., $>10,000 \text{ km}^{-2}$ in Hauraki Gulf; $>20,000 \text{ km}^{-2}$ in Auckland Harbor; $>40,000 \text{ km}^{-2}$ in Cook Strait approaches to Wellington Harbor) (Gregory 1990, unpubl. data). For comparison, densities elsewhere have been $1,500-3,600 \text{ km}^{-2}$ for the Cape Basin region of the South Atlantic lying west of southern Africa (Morris 1980), and $3,640 \text{ km}^{-2}$ from over 1,000 neuston trawl stations

Table 2.--Virgin plastic granule numbers on representative southwest Pacific island shores. Numbers given are local maximums expressed in number per linear meter of shore, following Gregory (1978); p = present in low numbers (<1 m⁻¹). For locations see Figures 3-8.

Location	Number
Norfolk Island	
Emily Bay	ca. 100
Raoul Island	
North Beach	>50
Denham Bay	nil
Fiji, Viti Levu	
Lautoka	p
Singatoka	p
Korolevu	<5
Deuba	>>100
Suva	>5(?)
Fiji, Vanua Mbalavu	
East	24
West	p
Tonga, Tongatapu	
Anahulu Beach	100
Laulea Beach	>>1,000
Oholei Beach	<50
Keleti Beach	nil
Fahina Beach	nil
Western Samoa, Upolu Island	
Apia	p
Vaiala Beach	nil
Malaeia Beach	20(?)
Cook Islands, Rarotonga	
Ngatangiaa Harbor	>500
Raringaru Stream	>>10
Akapuao Stream	<10(?)
Totokoitu Stream	p
Papua Stream	10
Rarotongan Hotel	1-<10

KEY: Figures 3 - 8





Figure 3.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Norfolk.

KEY: Figures 3 - 8





Figure 4.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Raoul.

KEY: Figures 3 - 8





Figure 5.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Fiji.

KEY: Figures 3 - 8





Figure 6.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Tonga.

KEY : Figures 3 - 8





Figure 7.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Western Samoa.

KEY : Figures 3 - 8





Figure 8.--Virgin plastic granules, oceanic and locally generated megalitter on southwest Pacific island shores: Rarotonga.




Figure 9.--Regional distribution of pelagic plastic granules across the southwest Pacific is influenced by oceanic fronts and wind and surface current patterns. Based on Gregory et al. (1984); Gregory (1987, 1990), and unpublished data.

for waters in the Agulhas Current up to 100 nmi offshore from Cape Province (Ryan 1988). On the other hand, densities in surface waters of the northern Sargasso Sea include >10,000 pieces of plastic and 1,500 pellets km⁻² (Wilber 1987). Elsewhere around the eastern North Atlantic, densities of polluting plastic are lower, with only 700 pieces km⁻² and 80 pellets km⁻² being reported from waters north of the Gulf Stream (Wilber 1987).

PLASTIC MACRO- AND MEGALITTER

In the categories of plastic macro- and megalitter I include large manufactured items and artifacts fabricated from plastics and other persistent synthetic materials and the products of their fragmentation and disintegration, following the approach of McCoy (1988). Megalitter is of a size enabling visual identification of floating items by a shipboard observer (generally decimeters or larger), while macrolitter is mostly smaller items and fragmented material, larger than the previously described granules and readily seen with the naked eye during shoreline surveys. Typical examples of the former are fishing floats, containers, crates, bottles and their tops, netting, lines, hawsers, strapping bands, plastic sheeting and bags, foamed items, and confectionery wrappings. Only some of these items are readily degradable.

Significant quantities of macro- and megalitter have been seen on all shores examined to date (Tables 2 and 3, Figs. 3-8, 10-13). The amounts are highly variable, but even on uninhabited islands and the otherwise remotest of places discarded plastic is present. In a survey of New Zealand's subantarctic islands, Gregory (1987, Appendix 1) itemized a great diversity of plastic material and noted that the quantity of macrolitter was surprisingly small considering the abundance of megalitter items (Fig. 10). A similar diversity of seaborne megalitter becomes stranded on islands of the southwest Pacific. As an example, Raoul Island in the Kermadec Group some 500 km northeast of New Zealand (Fig. 1), has <10 permanent residents at a weather station, and yet large quantities of macro- and megalitter are stranded on the beaches (Table 3).

In late 1988 New Zealand's Department of Conservation, with cooperation from the Wildtrack Programme produced by the Natural History Unit of TVNZ (Television New Zealand), initiated a nationwide survey of plastic litter on beaches. Most participants are students who complete a standard record card (Fig. 14). Preliminary reviews of some 50 returns coming from widely separated places, both remote and near population centers of the North and South Islands, confirm casual observations that considerable quantities of plastic macro- and megalitter accumulate on these shores. It is surprising to note that few returns identified the small resin granules, even at places where they are reasonably common. Those items most frequently recorded were fragments of foamed and hard plastic, plastic bags and sheeting, strapping bands, bottles, and bottle tops. The following selected examples illustrate the magnitude of contamination:

74 bottles on 860 m of beach--Ohope, Bay of Plenty

426 bottles and 82 bags on 2 km of beach--Mohaka, Hawkes Bay

32 bags on 500 m of beach--Petone, Wellington Harbor

2,817 bottle tops (from repeated surveys: 4 August, 14 and 19 September 1988)--Oreti, Southland

200 packing straps on <200 m of beach--Mokomoko Inlet, Southland

Table 3.--Simplified catalogue of plastic megalitter and other artifacts found on a 3-km stretch of beach on the northern coast of Raoul Island, southwest Pacific.

Type of litter	Number
Fish boxes and crates	10
Fishery floats	26
Bottles and containers (detergent, cosmetics, etc).	40
Hawser, rope	
Long (ca. ± 10 m)	10
Short (<10 m)	5
Netting (trawl) and rigid mesh	5
Foamed material (Styrofoam)	
Small (<2 cm)	>30
Moderate (>2 to <15 cm)	>10
Large (>15 cm)	>20
Sheeting	10
Strapping bands	>20
Footware (jandals/thongs)	20
Miscellaneous	>10

Repeated surveys (1974, 1978, 1981, and 1982) at Kawerua, a remote beach on the exposed west coast of Northland, showed a gradual decrease in numbers of plastic bags and an increase in bottles and total plastic items, probably reflecting changes in types of packaging over that period (Hayward 1984). Comparable trends in plastic megalitter accompanying changing patterns in offshore fishing activities have been noted for the subantarctic islands and mainland New Zealand shores (Cawthorn 1985).

SOURCES AND DISTRIBUTION

From the approach of Ryan (1987b), it is appropriate to identify three categories of plastic debris on shores throughout the region.

1. Material having a local onshore source.
2. Material originating from nearby fishing and shipping activities.
3. Material that has drifted from afar and that can be considered oceanic.



Figure 10.--Representative plastic items washed up on New Zealand's subantarctic islands: North West Bay, Campbell Island (A) and Derry Castle Reef, Auckland Islands (B and C). The large crushed container in (A) is of French origin and the two smaller items (arrow) are of United Kingdom manufacture. Note the polypropylene strapping (1), incipient crazing on the inside of broken high-density plastic fishing floats (2), cordage (3), and parts of wooden packing crates (4) in B and C.



Figure 11.--Representative locally generated and oceanic plastic litter assembled from combing 100 m of beach at Makara, west coast near Wellington, New Zealand. Some of this collection has clearly come from fishing activities. (Photograph taken by M. Cochrane.)




Figure 12.--Plastic sandal heavily encrusted with bryozoans, coralline algae, and clumps of the pink foraminiferan, *Homotrema rubra* (arrow). (Collected by K. A. Rodgers on Tuvalu.)




Figure 13.--Encrusted oceanic plastic items from Rarotonga (A) and Raoul Island (B). Note the bryozoans (1), coralline algae (2), and calcareous annelid tubes (3).

For the New Zealand coast and inshore waters, Gregory (1978) identified industrial centers as the principal sources (Fig. 2) of plastic meso-litter (mostly granules). This litter was material that was accidentally spilled in wharf and other cargo handling areas and at processing plants, and reached the sea through sewage and storm drainage systems as well as natural waterways. Subsequent dispersal was effected by coastal currents (Fig. 2). On populated islands (e.g., Tonga, Rarotonga), as on New Zealand shores, it is possible to separate plastic megalitter into two populations. One is probably of local (or domestic) origin, and the other comes from offshore sources and may have been adrift for some time. Casual visitors as well as indiscriminate and uncontrolled garbage dumping are responsible

BEACH CLEAN UP & SURVEY RECORD CARD

Date: _____ General Location: Big Beach, New Zealand
 Exact Site on beach _____
 Approx. size of area: _____ metres long by _____ metres wide
 Character of survey area (leave out: rock, sand, mud, gravel, other (specify)) _____
 Name of surveyor or organisation: _____
 Contact address and tel. no.: _____

An easy way to record the material you pick up is by counting in groups of five like this:
 Bottles 11111 Total 7

HARD PLASTICS	Total	GLASS	Total
Bottles, containers		Bottles	
Tops		Light bulbs and tubes	
6-pack yokes		Broken fragments	
Buckets		ALUMINIUM	
Toys/combs		Cans, cartons	
Fragments		Foil/trays/wrapping	
Other		Other	
FOAM PLASTICS		TIN/STEEL	
Trays/cups/packages		Tins/cans	
Floating floats		Drums	
Foam fragments		Wire	
Other		Bottle tops	
PLASTIC SHEET AND FIBRES		Other	
Plastic sheets		PAPER	
Plastic & cellophane bags		CARDBOARD	
Fishing nets		TIMBER (leave driftwood on beach)	
Fishing lines (approx. length in metres)		CLOTH ITEMS	
Other rope/cord (approx. length in metres)		RUBBER ITEMS	
Packing tape/strapping - cut		FOREIGN DEBRIS (describe overleaf)	
Packing tape/strapping - uncut		ENTANGLED/STRANDED ANIMALS (describe overleaf)	
ESTIMATE OF TOTAL VOLUME OF RUBBISH COLLECTED FROM YOUR AREA			

THANK YOU FOR COMPLETING THIS SURVEY CARD. PLEASE RETURN IT TO YOUR BEACH CO-ORDINATOR, OR YOUR LOCAL DEPARTMENT OF CONSERVATION OFFICE, OR POST TO ANNIE WHEELER, MARINE DEBRIS NETWORK, C/O DEPARTMENT OF CONSERVATION, P O BOX 8 NEWTON, AUCKLAND.

DETAILS OF ENTANGLED/STRANDED ANIMALS

Identify bird or animal: _____
 Cause of injury or death: _____ If the creature is still alive seek help from your beach co-ordinator, or local RSPCA, DOC office or Forest and Bird. Take photographs.

DETAILS OF FOREIGN DEBRIS

Describe item and the country you think it is from: _____

WHERE DO YOU THINK MOST OF THE RUBBISH YOU HAVE COLLECTED HAS COME FROM? (Is fishing boats, picnickers, household rubbish, foreign boats etc)

The NZ Marine Debris Network is a coalition of conservation groups who are concerned about the issue of marine debris and coastal pollution and its effect on wildlife. The Network includes representatives from the Department of Conservation, Greenpeace, Royal Forest and Bird Protection Society, University of Auckland, RSPCA, and the World Society for the Protection of Animals.

One of the aims of the Network is to co-ordinate the efforts of the many groups and individuals throughout the country who are involved in beach clean ups and to carry out a survey of the debris collected off New Zealand beaches. This information can then be used to determine the main types and sources of the debris that litter our beaches. It will contribute to a national marine debris data base, and will help develop solutions to marine debris pollution.

Your participation in this beach clean up is an important contribution to an on-going nationwide effort.

GUIDELINES FOR COLLECTORS

1. Carry out your beach clean-up and survey at low tide.
2. Make sure you have warm clothing, and sun protection if necessary.
3. Washing rubber or gardening gloves can be a good idea.
4. Before you start, make sure you've got your record card and board, pen and rubbish collection bags.
5. Take something to drink and perhaps a snack.
6. Work in small groups, so some of you can be picking up and sorting the rubbish into categories for recycling (see below) while one person is recording the details on the survey record card.
7. Know who your beach co-ordinator is and where to find them. Be clear about what area you are meant to be surveying.
8. Know where the rubbish collection point is.
9. Report any entangled animals to your co-ordinator and take photos if possible.
10. Give your record card back to your co-ordinator or post to the address on the other side of this form.
11. MAKE SURE YOU SEPARATE YOUR RUBBISH INTO DIFFERENT BAGS WITH 1. PLASTICS, 2. GLASS, 3. PAPER, 4. ALUMINIUM CANS, 5. TINS; AS ALL THIS MATERIAL IS GOING TO BE RECYCLED.

Figure 14.--Standard record card used in New Zealand coastal plastic pollution surveys.

for much of the former; some must also come from vessels operating in local waters. The most unsavory items found during the course of these surveys were soiled (disposable) baby diapers, syringes, and discarded pesticide or agricultural chemical containers. Litter coming from distant offshore sources is considered oceanic. It is characteristically embrittled and sports an encrusting biota (see below).

Around the subantarctic islands, plastic macro- and megalitter concentrate on west-facing (i.e., windward) shores, whereas little reaches their eastern (leeward) coasts (Gregory 1987). The dominating influence here is the strong West Wind Drift Current of the Southern Ocean (Lutjeharms et al. 1988). The same pattern is repeated on southwest Pacific island shores. Here, however, it is the eastern shores--those facing into the southeasterly trade winds--onto which plastic meso-, macro-, and megalitter are herded (Figs. 3-8). Further, in several instances, it is possible to identify crude decreases in quantities of plastics in the downdrift direction (e.g., Viti Levu, Tongatapu, and Rarotonga; Figs. 5, 6, and 8). The encrusting biota (see below) of many megalitter items suggest they have been afloat for some time. These are part of the global oceanic population of pelagic litter. Plastic granules on Australian, New Zealand, and Fiji shores can have their major origins in local suppliers and manufacturers.

Norfolk, Raoul, Vanua Mbalavu, Tongatapu, and Rarotonga have no local sources for virgin plastic granules, and lie upwind from regional ones. Nibs on these shores must have come from the same global oceanic population of pelagic plastics and are dispersed by the southeast trade winds. A possible source exists in French Polynesia, which lies upwind, but I have no data for this region.

Although plastic macro- and megalitter on eastern Australian shores have not been surveyed, quantities of this litter and the resin granules appear to be much lower than at equivalent sites in New Zealand. This difference probably reflects coastal current and broad oceanic circulation patterns as well as persistent winds that blow offshore or parallel to the coast over this region. On the other hand, drift pumice is quite common on these shores, as it is on the shores of many Pacific islands (Sachet 1955). Much of the plastic litter on popular recreational beaches of Australia, New Zealand, and larger southwest Pacific islands comes from casual visitors and day trippers; it is dominated by food and confectionery wrappings and drink bottles. This material is seldom conspicuous on isolated shores. From these remote places, there is evidence that much plastic debris comes from fishing-related or other shipping activities (Cawthorn 1985, 1987; Mattlin and Cawthorn 1986; Gregory 1987, 1990).

Attention has already been drawn to the accumulation of plastic debris on the windward shores of several southwest Pacific islands. The materials involved are mostly of oceanic origin and also from fisheries-related and shipping activities, and their quantities on west- and north-facing (leeward) shores are minimal. The principal urban population centers of Tongatapu, Rarotonga, and Upolu (Western Samoa) are all situated on north-facing coasts along which much locally generated plastic has spread.

Plastic items, categorized by country of origin (when possible), are summarized in Table 4 for New Zealand's subantarctic islands and for subtropical Raoul Island. Some items are truly oceanic (e.g., an Argentinian fishing float reaching the Snares), but most appear related to regional fishing activities. South Korean, Taiwanese, and Japanese vessels are common in these waters, so the dominance of Asian-sourced artifacts is not unexpected. The Russians also have a considerable presence, but one that is not reflected in the seaborne litter. Personal experience on a Russian research vessel reveals that they generate very little plastic, and discarded paper and cardboard packaging are incinerated.

The regional distribution of dispersed pelagic or oceanic plastics is schematically summarized in Figure 9. It has been inferred (Gregory et al. 1984; Gregory 1990) that major oceanic fronts such as the Polar, Subantarctic, Subtropical, and Tropical Fronts, and eddies from the East Australian Current have important influences on the distribution and abundance of litter. They act as barriers arresting the spread of material, and along these barriers the material is also concentrated and carried. For example, Bourne and Clarke (1984) noted an accumulation of garbage in the Humbolt Front off Valparaiso, Chile. Observations in the Hauraki Gulf, northern New Zealand, show that densities of plastic granules taken in tows made along windrows may exceed $10,000 \text{ km}^{-2}$, whereas densities in tows transverse to the windrows may be as few as $1,000 \text{ km}^{-2}$ (Gregory, unpubl. data).

Table 4.--Summary of numbers of plastic items having identifiable countries of origin.

Country of origin	Subantarctic islands	Raoul Island
Asia	11	5
United Kingdom	5	1
New Zealand	4	6
Australia	3	2
Spain	2	--
Bulgaria	1	--
France	1	--
Norway	1	--
U.S.S.R.	1	1
Argentina	1	--

ENCrustING BIOTA

Plastics and other synthetic litter afloat on surface waters of the ocean are an important and expanding, although little studied, ecological niche for a pseudoplanktic biota of the kind commonly present on *Sargassum* (Winston 1982; Butler et al. 1983). Gregory (1978) noted that granules from beaches of northernmost New Zealand were sometimes encrusted by the bryozoan *Membranipora tuberculata*. This is a tropical species and has also been found on drift plastics from Australia, Norfolk and Raoul Islands, and Fiji, Rarotonga, and Tongatapu. It was inferred that there had been eastward dispersal across the north Tasman Sea by way of eddies in the East Australian Current (Gregory 1978). Other encrusting taxa identified during past and present studies include further bryozoan species awaiting identification, coralline algae, calcareous annelids, barnacles, a hermatypic coral, and the pink foraminiferan *Homotrema rubra* (Figs. 12, 13). Encrusters are less common on artifacts from the subantarctic, where only goose barnacles (*Lepas* spp.) and the annelid *Spirorbis* have been recognized.

It is evident that pelagic plastic litter may be an important vector in the transoceanic and regional dispersal of a varied biota and may increase the chances of migration to distant shores, including isolated islands, as contemplated by Ryan (1987b).

DISCUSSION

The general environmental problems of the southwest Pacific region, with its limited financial and natural resources, have received wide attention (e.g., Chan 1973; Salvat 1979; Izrael et al. 1981; Dahl and Carew-Reid 1985; Carew-Reid 1988). Plastics are an unnecessary additional contaminant to the region, and the environmental implications to be drawn are those that have been identified elsewhere (Laist 1987) and need no further elaboration. For animals these implications include death or

debilitation through entanglement; blockage of the intestinal tract through ingestion, leading to starvation and death; ulceration of delicate tissues by jagged plastic fragments; and reduction in quality of life and reproductive performance. In addition, large items can be hazards to shipping. The aesthetic concerns expressed about plastic pollution also must be acknowledged. Unsightly accumulations of locally generated or oceanic plastics on beaches could be to the detriment of tourism (Prasad 1987). Soiled diapers, used syringes, and medicinal and pesticide containers stranded or abandoned on beaches will discourage even the most hardy of tourists.

The oceanic problem can be addressed through MARPOL and the London Dumping Convention. The local problem needs to be approached with cultural delicacy, for traditional practice and attitudes towards refuse disposal are in many ways rather casual (Anonymous 1976). Educational efforts, directed primarily at the young (Bryant 1988), will need to draw on and develop from traditional Pacific ways.

The very attributes that mankind finds desirable in plastics--lightness, strength, manufacturing adaptability, flexibility, inertness, resistance to degradational processes, transparency, and prolonged shelf life in packaging--are also the reasons they are today a globally important marine pollutant (Andrady 1988; Johnson 1988).

It is difficult to estimate the rate at which plastics disappear or are adsorbed into the environment (Gerrodette 1985). And while the breakdown of plastic compounds in itself may create few problems, the effects of released additives such as antioxidants, retardants, and biocides have never been assessed, only speculated about (Gregory 1978). Locally generated litter is likely to be fresh in appearance, while much of the oceanic and offshore-generated plastic litter stranding on these tropical and subtropical Pacific shores is chalky, crazed, and embrittled, all evidence of oxidative aging and photodegradation. Whether this occurs while it is afloat or after it is stranded on the shore has not been established. Circumstantial evidence suggests that aging is more rapid once artifacts are stranded high and dry on a beach (Gregory 1983). On the New Zealand coast, the extent of degradation apparently decreases southwards, although a detailed survey to confirm this claim has not been undertaken. Similarly, the proportion of degraded virgin granules is much greater on the tropical shores than it is on temperate ones (Table 5) (Gregory 1983, table 1). On high-latitude subantarctic shores, crazing is less evident and much breakdown occurs through mechanical abrasion and battering (Gregory 1987).

The extent of crazing and embrittlement of plastic granules (Table 5) and megalitter items observed on Raoul, Rarotonga, and Tonga suggest that a survival time of 5 years (Gregory 1983) may be overly generous. Evidence indicates that plastics degrade more rapidly in the Australian and New Zealand region than they do in equivalent Northern Hemisphere latitudes, although contrary to popular belief, the reason is not necessarily related to higher ultraviolet values (Sharman 1987). Controlled experiments and observations on rates of plastic degradation around the world are needed if we are to understand adequately the population dynamics of pelagic plastics

Table 5.--Relative numbers (in percentages) of fresh, slightly degraded, and highly degraded plastic granules from selected localities.

Locality	Number	Increasing degradation ———>		
		Fresh	Slightly degraded	Highly degraded
Fiji	163	18	45	37
Raoul	25	24	52	24
Rarotonga	70	19	41	40
Tonga	60	20	30	50
Auckland, New Zealand	216	79	13	8
Botany Bay, Australia	73	53	40	7

and to establish whether an equilibrium state between accumulation (strandings) and losses in environmental sinks (disappearance from view) has been already reached.

CONCLUSIONS

Although pollution by plastics of the southwest Pacific marine environment has not yet reached the magnitude evident in waters adjacent to more heavily populated, industrialized, and fished regions of the Northern Hemisphere, it is a developing problem and cause for concern.

Increased fishing activities across the region, and in particular drift gillnetting, are likely to escalate presently identified problems.

Regional distribution and dispersal are influenced by proximity to sources, oceanic current and circulation patterns, and prevailing winds. Oceanic fronts may have a key role in defining boundaries to zones with broadly similar areal densities of pelagic plastics.

Population dynamics of pelagic plastics across the southwest Pacific as well as globally are not well understood, and more information is needed on the "sinks" of this material.

There is need to educate the public about the environmental problems arising from the indiscriminate disposal of plastics and other persistent synthetic compounds.

ACKNOWLEDGMENTS

My research into plastics in the marine milieu has been funded by University Grants Committee and Auckland University Research Committee. I am indebted to the captains, masters, and crews of several vessels over a number of years for their logistic support: RV *Ikatere*, RV *Tangaroa*, RV *Vulkanolog*, the U.S. Coast Guard cutters *Glacier*, *Polar Sea*, and *Polar*

Star, Her Majesty's New Zealand ship *Monowai* and *RL Proteus*. I thank all those who have assisted with this program for their varied commentary. It is also appropriate to express gratitude to the organizers of the Second International Conference on Marine Debris for facilitating my participation--I learned much.

REFERENCES

Andrady, A.

1988. The use of enhanced degradable plastics for control of plastic debris in the marine environment. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, 13-16 October, 1987, Kailua-Kona, Hawaii, p. 384-403. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Anonymous.

1976. American Samoa: Pacific paradise confronts solid wastes problems. *Solid Wastes Manage.* 19(11):16-18, 56-58.

Barber, H. N., H. E. Badswell, and H. D. Ingle.

1959. Transport of driftwood from South America to Tasmania and Macquarie Island. *Nature* 184:203-204.

Bilal, J.

1985. The state of hydrocarbon pollution in the east Asian seas based on studies in the south-east Asian Seas regions. In *Environment and resources in the Pacific*, p. 217-234. UNEP Regional Seas Reports and Studies No. 69.

Bligh, W.

1792. *A voyage to the South Sea*. George Nicol, Lond., 264 p.

Bourne, W. R. P.

1976. Seabirds and pollution. In R. Johnston (editor), *Marine pollution*, p. 403-502. Acad. Press, Lond.

Bourne, W. R. P., and G. C. Clarke.

1984. The occurrence of birds and garbage at the Humbolt Front off Valparaiso, Chile. *Mar. Pollut. Bull.* 15:343-344.

Bourne, W. R. P., and M. J. Imber.

1982. Plastic pellets collected by a prion on Gough Island, central South Atlantic Ocean. *Mar. Pollut. Bull.* 13:20-21.

Branch, J. B.

1984. The waste bin: Nuclear waste dumping and storage in the Pacific. *Ambio* 13:327-330.

Brodie, J. W.

1960. Coastal surface currents around New Zealand. *N.Z. J. Geol. Geophys.* 3:235-252.

- Brodie, J., and J. Morrison.
1984. The management and disposal of hazardous wastes in the Pacific islands. *Ambio* 13:331-333.
- Brown, R. S., F. I. Norman, and D. W. Eades.
1986. Notes on blue and Kerguelen petrels found beach-washed in Victoria, 1984. *Emu* 86:228-238.
- Bryant, J. J.
1988. Educating the South Pacific about the environment. *Environmentalist* 8:127-131.
- Bullen, W. J.
1968. The location of plastics industry in New Zealand. M.A. Thesis, Univ. Canterbury, New Zealand, 149 p.
- Butler, J. N., B. F. Morris, J. Cadwallader, and A. W. Stoner.
1983. Studies of *Sargassum* and the *Sargassum* community. Bermuda Biol. Stn. Spec. Publ. 22, 307 p.
- Butler, J. N., B. F. Morris, and J. Sass (editors).
1973. Pelagic tar from Bermuda and the Sargasso Sea. Bermuda Biol. Stn. Res. Spec. Publ. 10.
- Carew-Reid, J.
1988. Marine pollution: The institutional response in Oceania. *Mar. Pollut. Bull.* 19:356-365.
- Cawthorn, M. W.
1985. Entanglement in, and ingestion of, plastic litter by marine mammals, sharks, and turtles in New Zealand waters. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 336-343. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
1987. Rubbishing the ocean. *Forest Bird* 18(4):25-26.
- Chan, G. L.
1973. Planning for pollution control in Atolls. In *Regional Symposium on Conservation of Nature--Reefs and Lagoons*, p. 133-143. South Pac. Comm., Noumea, New Caledonia.
- Chesher, R.
1984. Pollution sources survey of the Kingdom of Tonga. South Pacific Regional Environment Programme, Topic Review 19. South Pac. Comm., Noumea, New Caledonia, 110 p.
- Connor, J. J.
1976. Pollution in paradise. *Oceans* 9(3):36-37.

- Dahl, A. L., and I. L. Baumgart.
1983. The state of the environment in the South Pacific. UNEP Regional Seas Reports and Studies No. 31, 25 p.
- Dahl, A. L., and J. Carew-Reid (editors).
1985. Environment and resources in the Pacific. UNEP Regional Seas Reports and Studies No. 69, 289 p.
- Dawson, S., and E. Slooten.
1987. The down under dolphin. *Forest Bird* 18(4):32-34.
- Efi, T.
1976. After the shoals the tin cans. *Ecologist* 6:293.
- Furness, B. L.
1983. Plastic particles in three procellariiform seabirds from the Benguela Current, South Africa. *Mar. Pollut. Bull.* 14:307-308.
- Gerrodette, T.
1985. Toward a population dynamics of marine debris. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 508-518. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Gregory, M. R.
1977. Plastic pellets on New Zealand beaches. *Mar. Pollut. Bull.* 8:82-84.

1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwater Res.* 12:399-414.

1983. Virgin plastic granules on some beaches of eastern Canada and Bermuda. *Mar. Environ. Res.* 10:73-92.

1987. Plastics and other seaborne litter on the shores of New Zealand's subantarctic islands. *N.Z. Antarct. Res.* 7(3):32-47.

1990. Environmental and pollution aspects. In G. P. Glasby (editor), *Antarctic sector of the Pacific*, p. 291-324. Elsevier, Amsterdam.
- Gregory, M. R., R. M. Kirk, and M. C. G. Mabin.
1984. Pelagic tar, oil, plastics and other litter in surface waters of the New Zealand sector of the Southern Ocean, and on Ross Dependency shores. *N.Z. Antarct. Res.* 6:12-28.
- Hambuechen, W. H.
1973. Pesticides in the Cook Islands. In *Regional Symposium on Conservation of Nature--Reefs and Lagoons*, p. 119-126. South Pac. Comm., Noumea, New Caledonia.

- Harper, P. C., and J. A. Fowler.
1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptila* spp.) 1958-1977. *Notornis* 34:65-70.
- Hayes, T. M.
1981. Activities of the Inter-Governmental Maritime Consultative Organisation in the South Pacific relating to marine pollution prevention control and response. South Pacific Regional Environmental Programme, Topic Review 12. South Pac. Comm., Noumea, New Caledonia.
- Hayes, T. M., and D. Kay (editors).
1986. Pacific regional workshop on contingency planning and oil spill response. Federal Department of Transportation, Canberra, Australia, 60 p.
- Hayward, B. W.
1984. Rubbish trends--beach litter surveys at Kawerua, 1972-1984. *Tane* 30:209-217.
- Izrael, Y. A., A. I. Simonov, and A. V. Tsyban.
1981. Scientific aspects of interdisciplinary investigations in the Pacific for the prevention of its pollution. In N. A. Shilo and A. V. Lozhkin (compilers), *Ecology and environmental protection in the Pacific region*, p. 229-234. UNEP XIV Pacific Science Congress. Publishing Office, Nauka, Moscow.
- Johnson, D.
1988. Plastics good. . .bad, ugly. *Chem. Ind. Lab. Manage.* 22(5):10-11.
- Laist, D. W.
1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. *Mar. Pollut. Bull.* 18:319-326.
- Lutjeharms, J. R. E., L. V. Shannon, and L. J. Beekman.
1988. On the surface drift of the Southern Ocean. *J. Mar. Res.* 46:267-279.
- Matos, C.
1981. Marine pollution in the South Pacific. South Pacific Regional Environment Programme, Topic Review, 11. South Pac. Comm., Noumea, New Caledonia.
- Mattlin, R. H., and M. W. Cawthorn.
1986. Marine debris--an international problem. *N.Z. Environ.* 51:3-6.
- McCoy, F. W.
1988. Floating megalitter in the eastern Mediterranean. *Mar. Pollut. Bull.* 19:25-28.
-

Ministry for the Environment.

1987. Packaging in the New Zealand environment: Issues and options. Discussion paper, Ministry for the Environment, Wellington, New Zealand, 110 p.

Morris, R. J.

1980. Plastic debris in the surface waters of the South Atlantic. Mar. Pollut. Bull. 11: 164-166.

Morrison, R. J., and J. E. Brodie.

1985. Pollution problems in the South Pacific: Fertilizers, biocides, water supplies and urban wastes. In Environment and Resources in the Pacific, p. 69-74. UNEP Regional Seas Reports and Studies No. 69.

Murray, T.

1988. Drift netters threaten South Pacific fisheries. Catch 15(11):13-15.

Oostdam, B. L.

1984. Tar pollution of beaches in the Indian Ocean, the South China Sea and the South Pacific Ocean. Mar. Pollut. Bull. 15:267-270.

Plastics Institute of New Zealand.

1988. Packaging in the New Zealand environment issues and options. Submission to the Ministry for the Environment, Plastics Institute of New Zealand, Wellington, New Zealand, 28 p.

Prasad, P. C.

1987. The impact of tourism on small developing countries. In S. Britton and W. C. Clarke (editors), Ambiguous alternatives; tourism in small developing countries, p. 9-15. Univ. South Pacific, Suva, Fiji.

Pruter, A. T.

1987. Sources, quantities and distribution of persistent plastics in the marine environment. Mar. Pollut. Bull. 18:305-310.

Randall, B. M., R. M. Randall, and G. J. Roussow.

1983. Plastic particle pollution in great shearwaters (*Puffinus gravis*) from Gough Island. S. Afr. J. Antarct. Res. 13:49-50.

Ridgway, N. M., and G. P. Glasby.

1984. Sources of marine pollution around New Zealand. N.Z. O. I., Oceanographic Summary 23, 21 p.

Ryan, P. G.

- 1987a. The incidence and characteristics of plastic particles ingested by seabirds. Mar. Environ. Res. 23:175-206.
- 1987b. The origin and fate of artifacts stranded on islands in the African sector of the Southern Ocean. Environ. Conserv. 14:341-346.

1988. The characteristics and distribution of plastic particles at the sea-surface off southwestern Cape Province, South Africa. *Mar. Environ. Res.* 25:249-273.
- Sachet, M.-H.
1955. Pumice and other extraneous volcanic materials on coral atolls. *Atoll Res. Bull.* 37, 27 p.
- Salvat, B.
1979. Trouble in paradise, Part 1: Assault on coral reefs and lagoons. *Parks* 3(2):1-4.
- Sharman, W. R.
1987. Problems with plastic durability in New Zealand buildings. *N.Z. J. Tech.* 3:145-152.
- Skira, I. J.
1986. Food of the short-tailed shearwater, *Puffinus tenuirostris*, in Tasmania. *Aust. Wildl. Res.* 13:481-488.
- Smith, R. I. L.
1985. Nothofagus and other trees stranded on islands in the Atlantic sector of the Southern Ocean. *Brit. Antarct. Surv. Bull.* 66:47-55.
- Waldichuk, M.
1977. Global marine pollution: An overview. Intergovernmental Oceanographic Commission Technical Series 18. UNESCO, Paris, p. 96.
- Wilber, R. J.
1987. Plastics in the North Atlantic. *Oceanus* 30(3):61-68.
- Winston, J. E.
1982. Drift plastic--an expanding niche for a marine invertebrate? *Mar. Pollut. Bull.* 13:348-351.

THE MARINE PLASTIC DEBRIS PROBLEM OFF SOUTHERN AFRICA: TYPES
OF DEBRIS, THEIR ENVIRONMENTAL EFFECTS, AND CONTROL MEASURES

Peter G. Ryan
Percy FitzPatrick Institute of African Ornithology
University of Cape Town
Rondebosch 7700, South Africa

ABSTRACT

Plastic debris is a global marine pollutant which is inflicting ever-increasing environmental and financial costs. In the seas off southern Africa and the adjacent Southern Ocean, entanglement has been recorded for at least 5 species of marine mammals, 13 seabird species, 2 turtles, and 6 shark species. Plastic ingestion has been recorded from 7 marine mammal species, 36 seabird species, 2 marine turtle species, and 7 shark species. The incidence in invertebrate taxa is not known. At present, entanglement does not pose a threat to the survival of any populations off southern Africa, but the recent introduction of a driftnet fishery to the South Atlantic and Indian Oceans and the suffering frequently associated with entanglement are causes for concern. By contrast, ingestion of plastic particles may adversely affect almost the entire population of species that do not regurgitate indigestible objects, with large, accumulated plastic loads reducing feeding efficiency or obstructing the digestive tract. Off southern Africa, generalist, surface-feeding, pelagic taxa such as certain procellariiform seabirds (petrels and albatrosses) and juvenile marine turtles are at risk from plastic ingestion. The incidence of ingested plastic in some species exceeds 90% of the population. The major financial cost of marine plastic debris is the reduced aesthetic appeal of coastal areas, which adversely affects the tourist industry. In South Africa alone, approximately R10 million is spent annually on cleaning beaches, where plastic makes up more than 90% of all stranded debris.

To address the problem of marine debris requires knowledge of the sources of various pollutants. Beach surveys readily assess the most abundant types of plastic debris, and from these data their sources can be inferred. Disposable packaging accounts for more than half the large plastic objects on southern African beaches, with most of the remainder composed of fishing gear. Sheet plastic (bags and wrappings) is the most abundant single type of plastic. Among small particles, virgin industrial pellets and fragments of other products predominate.

Using these findings to assign culpability and to elicit assistance, four approaches are being used to tackle the problem of marine plastic debris off southern Africa: education, product substitution, recycling, and legislation. As a short-term measure, specific types of artifacts responsible for most entanglements (e.g., hi-cone six-pack yokes, packing straps) and ingestion (e.g., virgin pellets, plastic bags) have been targeted for action. These approaches to control marine plastic pollution are discussed in relation to the highly diverse socioeconomic conditions prevailing in southern Africa.

INTRODUCTION

Much concern recently has been focused on the problems associated with anthropogenic marine debris, particularly as regards plastic (Shomura and Yoshida 1985; Laist 1987; Wolfe 1987). A variety of approaches has been adopted to tackle these problems, but have concentrated on maritime legislation (e.g., Bean 1987; Lentz 1987) and on awareness campaigns in developed, first world communities (e.g., Neilson 1985). Other than international maritime legislation, there have been few attempts to tackle the growth of marine debris arising from third-world communities. The southern African region comprises virtually the entire socioeconomic spectrum, and is to a large extent isolated from the world's major manufacturing centers. It is thus a useful area for examining the efficacy of various measures taken to limit persistent debris production. This paper reviews the occurrence of anthropogenic marine debris off southern Africa and in adjacent oceanic areas, and summarizes the known environmental effects of debris. The approaches used to identify the sources of marine debris and to control the amount of litter entering the sea are discussed.

THE SEAS OFF SOUTHERN AFRICA

Southern Africa has an unindented coastline, with few large bays or inlets (Fig. 1). Strong wave action is characteristic of much of the coast, with sandy beaches comprising almost 70% of the coastline. The continental shelf is narrow (<50 km wide) off the east coast, moderately broad (up to 150 km wide) off the west coast, and is most extensive off the south coast, where the Agulhas Bank extends more than 200 km offshore.

There are two main current systems. The cool (10°-16°C) Benguela Current flows north along the west coast, and is characterized by localized upwelling of cold, nutrient-rich bottom water when surface waters are advected offshore (Shannon 1985). The warm (22°-28°C) Agulhas Current flows south, close inshore along the east coast until it reaches the Agulhas Bank, where it moves offshore. South of the subcontinent, the Agulhas Current retroflects to flow eastward in oceanic waters to the north of the Subtropical Convergence (Lutjeharms 1981). However, large (500-km diameter) eddies formed at the retroflexion zone frequently transport Agulhas Current water into the South Atlantic (Lutjeharms 1988; Lutjeharms and Valentine 1988). Elsewhere to the south of the subcontinent, the predominant surface flows are eastward, associated with the West Wind Drift (Lutjeharms et al. 1988).




Figure 1.--The southern African region and adjacent oceans. The stippled line indicates the approximate edge of the continental shelf.

Most merchant ships in the area travel around southern Africa following the Cape sea route between Europe-North America and the Persian Gulf-Southeast Asia. This route runs close inshore along the south and east coasts of southern Africa, only moving offshore off the west coast. There is relatively little transoceanic merchant trade to either South America or Australia.

Commercial fisheries off southern Africa are concentrated on the broad continental shelves off the south and west coasts, where there are extensive demersal (bottom trawl) and pelagic (purse seine) fisheries (Crawford et al. 1987). There is a limited prawn fishery off the east coast, and a longline tuna fishery in oceanic waters. Gillnets were little used in the region until 1989, when oriental vessels started using driftnets more extensively in both the South Atlantic and South Indian Oceans (Ryan and Cooper in press).

THE DISTRIBUTION OF PLASTICS AND OTHER DEBRIS AT SEA

Little has been recorded of the distribution and abundance of plastic and other debris at sea off southern Africa. The abundance, distribution, and movements of tar balls at sea have been documented for two regions off the southern African coast, with a view towards identifying coastal areas

vulnerable to oil pollution (Shannon et al. 1983). The abundance of small plastic particles (<10 mm diameter) also has been estimated from surface neuston trawls. Morris (1980) reported densities ranging between 1,300 and 3,600 virgin industrial pellets km^{-2} in oceanic waters of the Cape Basin west of Cape Town.

The density of plastic debris in coastal waters off the southwestern Cape averaged 3,640 particles km^{-2} , derived from over 1,200 neuston trawls conducted at monthly intervals during 1977-78 (Ryan 1988a). Seasonal patterns of distribution and abundance were related to probable source areas and transport at sea. However, the highly clustered dispersion of particles, presumably due to fine-scale convergence zones, resulted in great variances in debris abundance estimates (range 0-445,000 particles km^{-2}), largely as a consequence of the relatively small sampling area (190 m^2 per trawl). Foamed plastics and fragments of manufactured articles were the most abundant types of particles, but virgin industrial pellets accounted for most of the mass (mean 42.4 g km^{-2} ; Ryan 1988a). It appears that at least a significant proportion of the debris arises from local sources, with concentrations inshore and close to harbors (Ryan 1988a, cf. Morris 1980). However, there is also evidence that the Agulhas Current is an important debris vector (Ryan 1988a).

A small number of neuston trawls (39) in oceanic waters south of southern Africa in the Agulhas retroflexion area collected only two plastic particles, both fragments of manufactured articles (P. G. Ryan unpubl. data). This gives a density estimate for the area of only 50 particles km^{-2} , similar to the density in the Southern Ocean south of New Zealand (Gregory et al. 1984).

Ryan (1988a) found that the density of large (>100 mm diameter) objects counted from a low-flying plane was an order of magnitude greater 10 km from the shore (19.6 objects km^{-2}) than 50 km offshore (1.6 km^{-2}) in the area between Cape Town and Saldanha Bay, where the merchant shipping lane runs close inshore. This offshore gradient is likely to be less marked farther north off the west coast where the continental shelf is broader (hence fishing grounds more extensive) and the shipping lane runs farther offshore. In oceanic waters south of the subcontinent, in the region of the Agulhas retroflexion, ship-based counts provided density estimates of between 0.04 and 0.09 large objects km^{-2} (P. G. Ryan unpubl. data).

These data refer only to floating debris. Virtually nothing is known about debris on the seabed off southern Africa. Debris comprised of materials denser than seawater presumably does not disperse far from source areas. Such items occasionally are caught in bottom trawls off the west coast (B. Rose pers. commun.; pers. observ.). Floating debris can also sink if it supports sufficient sessile organisms or entangles enough animals to increase the density above that of seawater. Such objects have a much greater dispersal capability than do plastics that are denser than seawater.

IMPACTS OF MARINE DEBRIS

Marine debris has both environmental impacts and financial costs. The major financial burden results from the reduced aesthetic appeal of polluted marine systems. Beaches are important for the >R2,000 million per annum tourist industry in southern Africa, and their appeal is reduced when they are littered with stranded debris. In South Africa alone some R10 million is spent annually on cleaning litter off beaches.

Apart from the accumulation of unsightly debris, the main environmental impact associated with marine debris is animal mortality through entanglement in and ingestion of debris. In addition, it has been suggested that anthropogenic debris is having some ecological effect by increasing the amount of available substratum onto which sessile organisms can settle (Carpenter and Smith 1972; Winston 1982), and it is possible that debris has increased the rate of propagule dispersal to islands (Ryan 1987b). However, the significance of the latter two impacts has not been determined.

Entanglement off Southern Africa

Entanglement involves animals becoming enmeshed in objects that impede movement, causing drowning, starvation, or reduced fitness, or restrict growth, cutting deep wounds into growing animals. This typically involves fairly large pieces of debris, and the apparent suffering associated with entanglement engenders considerable public concern.

Representatives of five marine vertebrate classes are known to have become entangled in debris off southern Africa (Table 1); there are no data for invertebrate groups. Most records are from coastal waters (where there are most observers), but a few entangled seals and birds have been found at subantarctic islands. Overall, the incidence of entanglement is fairly low, with only one species, the great white shark, *Carcharodon carcharias*, having more than 1% of individuals examined entangled in debris. There may be some cause for concern along the south coast, where 14% of stranded birds are entangled in debris (n = 97), with 28% of the vulnerable jackass penguin, *Spheniscus demersus*, entangled (n = 32, P. G. Ryan unpubl. data).

However, interpretation of the incidence of entanglement is complicated by different sampling techniques. For example, recoveries of banded crested terns, *Sterna bergii*, in southern Africa indicate that 14.2% (n = 267) of birds are captured after being entangled in debris, whereas only 2.2% (n = 46) of stranded birds were found entangled ($\chi^2 = 5.23$, $P < 0.05$; FitzPatrick Institute unpubl. data). And yet it is to be expected that the proportion of entanglement among stranded animals is higher than that among the general population, although the exact relationship is unclear. Also, it is not possible to infer the consequences of a given level of entanglement on population trends. Northern fur seal, *Callorhinus ursinus*, numbers have been decreasing apparently at least partly as a result of a 0.4% frequency of entanglement (Fowler 1987). A similar entanglement frequency has been recorded at some Cape fur seal,

Table 1.--A summary of the known incidence of entanglement of marine animals in plastic objects and other debris off southern Africa, excluding the by-catch of nontarget species during fishing operations (including shark exclusion nets). Based on Shaughnessy (1980), Balazs (1985), and (unpubl. data) from G. Avery, P. B. Best, N. Rice, G. J. B. Ross, and the Natal Sharks Board.

Taxon	Type of debris	Frequency of occurrence
Cetaceans	Ropes, nets	Three plus species, apparently infrequent.
Seals	Ropes, nets, line 92% Packing straps 6% O-rings 2% Wire <1%	Cape fur seals 0.12%, but 0.6% in one colony. Two records at subantarctic islands.
Birds	Nets, rope, line 89% Plastic bags 11%, Six-pack yokes	Thirteen species, 0.6% of stranded animals but up to 14% locally.
Turtles	Rope	Two species.
Fish	All packing straps	Six species of sharks, 0.2% of shark-net catch, incidence ranges 0-1.4%.

Arctocephalus pusillus, colonies (Shaughnessy 1980), and yet this species' population is increasing by 3.7% per year (David 1987).

The types of objects causing entanglement off southern Africa vary among taxa (Table 1). However, most items are either fishing gear (rope, netting, and fishing line) or disposable packaging (primarily plastic packing straps and plastic bags). Only one item was not made of plastic; a single seal was found with a piece of wire caught around its neck (Shaughnessy 1980).

These data on entanglement ignore the incidental catch of animals during commercial fishing operations. Some birds and mammals are caught in demersal trawls (e.g., Ryan and Moloney 1987) and by the longline fishery (e.g., Ryan and Rose 1988), but for at least these taxa the fishery by-catch is relatively small (cf. Tull et al. 1972; Piatt and Nettleship 1987), due largely to the limited use of gillnets. The impact of the recent expansion of oriental driftnet fisheries in oceanic waters of the South Atlantic and South Indian Oceans needs urgent investigation. The killing of seabirds for food by fishermen is an ongoing problem (Cooper 1977; Ryan and Rose 1988).

Table 2.--A summary of the known incidence of marine animals ingesting plastic objects and other debris off southern Africa. Based on Hughes (1973), Ryan (1987a), and unpubl. data from P. B. Best, J. H. M. David, G. J. B. Ross, and the Natal Sharks Board.

Taxon	Type of debris	Frequency of occurrence
Cetaceans	Plastic bags 8 Plastic bottles 1 Packing strap 1	Seven species, 3.0% of stranded animals.
Seals	--	No records.
Birds	Virgin plastic pellets 56% Plastic user fragments 43% Wood, tar balls, paint, glass, and aluminium foil 1%	Thirty-six species, incidence ranges 0-92% with 10 species >50% and 4 species >80%.
Turtles	Plastic bags 75% Virgin plastic pellets 25% Glass 1 piece	Two species, 11.1% of stranded animals.
Fish	Plastic bags 82% Plastic bottles 12% Nets and line 6%	Seven species of shark, 0.3% of shark-net catch, incidence ranges 0-6%.

Debris Ingestion off Southern Africa

The effects of debris ingestion are seldom as dramatic as those of entanglement, but ingested debris can cause death or debilitation by obstructing the digestive tract (e.g., Balazs 1985; Fry et al. 1987) or reducing meal size and the urge to eat (e.g., Ryan 1988b). Ingested plastic may also be a source of toxic chemicals (e.g., Ryan et al. 1988).

Ingestion of marine debris has been recorded for four vertebrate classes off southern Africa (Table 2); there are no data for invertebrate groups. Debris ingestion is much more prevalent than is entanglement, affecting over 90% of individuals of blue petrels, *Halobaena caerulea*, and great shearwaters, *Puffinus gravis*, breeding at oceanic islands (Ryan 1987a). The incidence of debris ingestion among southern African seabirds is among the highest in the world, largely due to the predominance of generalist, surface-feeding procellariiform seabirds (petrels, storm-petrels, shearwaters, and albatrosses) that do not frequently regurgitate indigestible objects and thus accumulate ingested plastic (Ryan 1987a, 1988c). The present incidence of debris ingestion by turtles may be greater than the 11% indicated in Table 2, because there are no observations subsequent to 1973. Debris ingestion by birds has increased since the late 1970's off southern Africa (Ryan 1988c).

Ingestion of debris off southern Africa by large proportions of populations of birds and turtles in particular is cause for concern. Almost all debris ingested is plastic that floats in seawater (Table 2). The few nonfloating debris items found in animals apparently are eaten ashore (e.g., gulls at refuse dumps, giant petrels at their breeding islands). Although the types of objects ingested are influenced by an animal's size (e.g., Ryan 1987a), two types of plastic objects make up the majority of ingested debris: virgin industrial pellets and plastic bags (Table 2). Reducing the abundance of these items at sea is the only long-term solution to the problem of debris ingestion.

TACKLING THE MARINE DEBRIS PROBLEM

Marine debris is extremely heterogeneous in terms of both the size and composition of artifacts and the wide range of their sources. This diversity makes the control of marine debris problematic. Examining the various impacts of marine debris highlights the types of debris responsible for most environmental problems. Off southern Africa these are discarded fishing gear, various types of plastic packaging (notably bags and packing straps), and small plastic particles (chiefly virgin industrial pellets). These types of debris warrant most attention, but the implementation of effective measures to reduce the amount of debris entering the sea requires knowledge of the sources of marine debris. It is evident that the general source of discarded fishing gear is the various fishing industries, providing a ready target for action. However, the sources of packaging and, to a lesser extent, industrial pellets are highly diffuse, complicating the assessment of culpability.

Using Beach Surveys to Identify Debris Sources

Beach surveys offer the simplest and most practical way to assess the relative abundance of various types of marine debris and to identify their probable sources (e.g., Merrell 1980; Vauk and Schrey 1987). However, one problem with stranded debris surveys is controlling for the selective removal of debris by beachcombers (see Ryan 1987b). Surveys at uninhabited islands avoid this problem. Figure 2 shows the numerical dominance of plastic articles and the much faster growth in amount of plastic debris compared with other debris types at Inaccessible Island in the Tristan group, central South Atlantic Ocean. Most of the debris identifiable as to country of origin derives from South America, and the proportion has been increasing: 32% in 1984, 36% in 1987, and 48% in 1988. Given the limited merchant trade across the South Atlantic, it is likely that much of the plastic debris reaching Inaccessible Island has drifted more than 3,000 km from South America (Ryan and Watkins 1988).

This contrasts with the situation on southern African beaches, where most identifiable debris derives from local sources. A survey of stranded debris at 50 sandy beaches between Cape Town and the Transkei was undertaken during June 1984. All large (>20 mm) articles within representative 50-m stretches of beach were collected (P. G. Ryan unpubl. data). Plastic made up more than 90% of stranded debris, and was recorded at all beaches sampled. Disposable packaging (e.g., bags, bottles; Fig. 3) comprised more




Figure 2.--The densities of various types of stranded debris at uninhabited Inaccessible Island during 1984 (Ryan 1987a), 1987 (Ryan and Watkins 1988), and 1988 (P. G. Ryan unpubl. data). Dashed lines between 1984 and 1987 indicate the lack of samples during this period.

than half of all plastic articles (57%), with fishing gear (netting, ropes, monofilament line, floats, traps, and fish boxes) making up most of the remainder (31%; Fig. 3). Almost half of the packaging was sheet plastic (bags and wrappings constituting 47% of packaging; Fig. 3), whereas polypropylene rope made up most of the fishing gear (85%).

The relative proportions of packaging and fishing gear among stranded plastic debris varied with distance from human settlements. Beaches in urban areas had a much greater proportion of packaging than either rural or island beaches (Fig. 4). This indicates that dumping of garbage from ships is not the only source of debris; urban areas in coastal South Africa also contribute significantly to marine debris loads (although selective removal of fishery-related products by beachcombers may contribute to the

TYPES OF PLASTIC (N = 2 661)



TYPES OF PACKAGING (N = 1 507)




Figure 3.--The proportions (by number) of various functional groups of plastic articles found stranded on 50 South African beaches during June 1984.

differences). This is evident to anyone examining storm-water outlets draining urban areas, and concurs with current thinking that land-based sources may be more important contributors of debris to the marine environment than are vessels (Bean 1987, but see Pruter 1987; Wirka 1988). The mean density of packaging at urban beaches in South Africa (0.66 articles m^{-1} of beach) was greater than that at rural beaches (0.53 m^{-1}), although variances were great.




Figure 4.--The relative proportions (by number) of packaging and fishing gear stranded at urban ($n = 26$) and rural ($n = 24$) beaches in South Africa, compared with the situation at oceanic islands.

Virgin industrial pellets and fragments of plastic articles also are widespread and abundant on southern African beaches, with exceptional densities of up to 43,350 particles per meter of beach (88% industrial pellets, P. G. Ryan unpubl. data). Determining the origin of these items is more difficult than determining the origin of larger articles. However, at least some industrial pellets derive from local sources, where poor handling practices result in spillages, with transport to the sea in wastewater (pers. observ.).

Measures to Control Marine Debris in Southern Africa

The diverse nature of marine debris requires a multilevel approach to mitigating the problem. Four basic control tools are available: education, product substitution, recycling, and legislation. However, not all of these approaches are appropriate to tackle the different facets of the marine debris problem.

Ship-Based Sources

Ships are responsible for fishing gear (with the exception of mono-filament line and other wastes from shore-based anglers) and a proportion of general refuse (packaging and other operational wastes). This source of debris has received more attention than have land-based sources (e.g., Dixon and Dixon 1981; Horsman 1982; Low et al. 1985; Pruter 1987), and is the subject of several international conventions (e.g., Lentz 1987). South Africa has agreed in principal to sign Annex V of the International

Convention for the Prevention of Pollution from Ships (MARPOL), which came into force at the end of 1988 and prohibits the dumping of all plastic products at sea. This will be a major advance, and South Africa's ratification of Annex V warrants expediting. Priority should also be given to Namibia, which became independent in 1989, acceding to MARPOL.

However, there are problems associated with enforcing Annex V of MARPOL (e.g., Bean 1987) which necessitate that its implementation in South Africa be coupled with an intensive education campaign aimed at all mariners. A representation to this effect has been made to the South African committee working on incorporating Annex V into national legislation (Dolphin Action and Protection Group 1989). In the interim, favorable responses have been received from several merchant lines and the South African Navy in response to requests to reduce the amount of debris dumped at sea (Dolphin Action and Protection Group 1988a, 1988b, 1989).

One problem area not covered by Annex V of MARPOL is the accidental loss of fishing gear at sea. There is no simple solution to this problem. The dumping at sea of damaged nets and other persistent debris by fishing vessels has officially been outlawed in South Africa since 1986 (Dolphin Action and Protection Group 1988a). However, captains of commercial fishing vessels currently are paid bonuses based on the cleanliness of vessels returning to port. This is perceived by the industry as being responsible for considerable dumping at sea, an action that could be avoided by linking bonuses to the amount of persistent debris returned to shore.

Land-Based Sources

There are two main types of marine debris derived from land-based sources: virgin industrial pellets and the diverse array of manufactured articles, principally disposable objects such as packaging and convenience items (Bean 1987). The loss of industrial pellets into the environment is limited to the plastics industry, which in southern Africa is a fairly small target for control measures. There is only one polymer producer in southern Africa (linked to the oil-from-coal plant at Secunda in the Transvaal), and almost all converters (manufacturers that convert industrial plastics into user products) are based in South Africa (Fig. 5). The industry has been apprised of the problem and is sympathetic. The recent large increases in the price of virgin pellets apparently have resulted in improved handling practices leading to reduced losses, but this needs verifying, and, if necessary, supporting with punitive legislation against accidental spillages.

A more intractable problem is that of general refuse being washed or blown into the sea. This type of debris derives from such a variety of sources that there is no simple target for control measures (Pruter 1987). Ultimately, the only solution is to educate the public to dispose of refuse correctly. There are ongoing antilittering campaigns in most southern African states, but these are proving insufficient to the task. The problem is complicated by the difficulty of communicating to a broad cultural and economic spectrum simultaneously. South Africa has the potentially disastrous combination of a burgeoning third-world population shopping in first-world supermarkets for products wrapped in first-world packaging,

SOUTH AFRICAN PLASTICS INDUSTRY




Figure 5.--The status of the South African plastics industry during 1988, showing the magnitude (metric tons) of flows between producers, users, and the environment (derived from data from the Plastics Federation of South Africa).

mostly plastic. Some 20% of South Africans live adjacent to urban areas in informal settlements where adequate waste disposal facilities are lacking. The same problem occurs in neighboring states such as Botswana and Namibia as a result of the large number of products imported from South Africa.

To counter these problems, attempts are being made to reduce the amount of plastic used in disposable applications (260,000 tonnes in South Africa in 1988, Fig. 5). This "source-reduction" approach (Wirka 1988) can be successful, judging by the small amount of litter found in Zimbabwe, where strict currency exchange regulations limit the use of plastics and almost all containers are returnable on a deposit basis. However, there is considerable industry resistance to such changes, despite support for a reduction in superfluous and environmentally damaging packaging by consumer bodies. Concerted public pressure is needed to stem the growth of plastics in disposable applications (Wirka 1988). At present, product substitution is preferred to the use of degradable plastics, which have attendant problems (e.g., Taylor 1979; Wirka 1988).

Almost 10% of South Africa's annual plastic production was recycled during 1988 (including factory scrap; Fig. 5), a greater proportion than that recycled in the United States (1%; Wirka 1988). One mixed-plastics recycling plant producing a wood substitute has recently been established in Cape Town, and there are several primary recycling operations throughout South Africa. However, there is much scope for further recycling, and incentives to return used plastics for recycling are likely to prove successful in limiting littering. There are problems associated with recycling plastics in southern Africa. The relatively small volume of material and the widely scattered markets render many recycling operations uneconomic. Also, in most areas of southern Africa, solid waste disposal using landfill sites remains by far the cheapest disposal technique, although groundwater contamination by leachates from landfills is a potential problem.

Legislation in South Africa is starting to address the problem of inadequate waste disposal. The recently promulgated Environmental Conservation Bill provides for heavy fines and, in some cases, jail sentences for littering and other disposal contraventions. However, it is hoped that voluntary measures taken by the business sector will obviate the need for further legislation. Awareness campaigns focusing public concern have had considerable success in promoting the use of more environmentally friendly products and practices (e.g., the phasing out of six-pack yokes and shrink-wrapped packaging for bricks, and the printing of warning labels on a variety of disposable plastic products; Dolphin Action and Protection Group 1989), but many problems remain to be solved. It is only through the whole-hearted support of the entire community that the marine debris problem can be diminished.

ACKNOWLEDGMENTS

Nan Rice generated much lively discussion and commented on an earlier draft. I am grateful to the following for supplying unpublished information: Graham Avery, Peter Best, Sheldon Dudley, Nan Rice, Barrie Rose, Graham Ross, the Natal Sharks Board, the Packaging Council of South Africa,

and the Plastics Federation of South Africa. I am grateful to the organizers of the Second International Conference on Marine Debris for the opportunity to attend the conference. Financial and logistical support for my work was received from the Packaging Council of South Africa, the South African Council for Scientific and Industrial Research, the South African Department of Environment Affairs, and the South African Scientific Committee for Antarctic Research.

REFERENCES

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Bean, M. J.
1987. Legal strategies for reducing persistent plastics in the marine environment. Mar. Pollut. Bull. 18:357-360.
- Carpenter, E. J., and K. L. Smith, Jr.
1972. Plastics on the Sargasso Sea surface. Science 175:1240-1241.
- Cooper, J.
1977. Editorial. Cormorant 3:3.
- Crawford, R. J. M., L. V. Shannon, and D. E. Pollock.
1987. The Benguela ecosystem. 4. The major fish and invertebrate resources. Oceanogr. Mar. Biol. Annu. Rev. 25:353-505.
- David, J. H. M.
1987. South African fur seal, *Arctocephalus pusillus pusillus*. In J. P. Croxall and R. L. Gentry (editors), Status, biology, and ecology of fur seals. Proceedings of an International Symposium and Workshop, 23-27 April 1984, Cambridge, England, p. 65-71. U.S. Dep. Commer., NOAA Tech. Rep. 51.
- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. Mar. Pollut. Bull. 12:289-295.
- Dolphin Action and Protection Group.
1988a. Save Our Sealife - prevent plastic pollution campaign. Report 1:1-6.
1988b. Save Our Sealife - prevent plastic pollution campaign. Report 3:1-9.
1989. Save Our Sealife - prevent plastic pollution campaign. Report 5:1-11.
- Fowler, C. W.
1987. Marine debris and northern fur seals: A case study. Mar. Pollut. Bull. 18:326-335.

- Fry, D. M., S. I. Fefer, and L. Sileo.
1987. Ingestion of plastic debris by Laysan albatrosses and wedge-tailed shearwaters in the Hawaiian Islands. *Mar. Pollut. Bull.* 18:339-343.
- Gregory, M. R., R. M. Kirk, and M. C. G. Mabin.
1984. Pelagic tar, oil, plastics and other litter in the surface waters of the New Zealand sector of the Southern Ocean, and on Ross Dependency shores. *N.Z. Antarct. Rec.* 6:12-28.
- Horsman, P. V.
1982. The amount of garbage pollution from merchant ships. *Mar. Pollut. Bull.* 13:167-169.
- Hughes, G. R.
1973. The sea turtles of south east Africa. Ph.D. Thesis, Univ. Natal, Pietermaritzburg.
- Laist, D. W.
1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. *Mar. Pollut. Bull.* 18:319-326.
- Lentz, S. A.
1987. Plastics in the marine environment: Legal approaches for international action. *Mar. Pollut. Bull.* 18:361-365.
- Low, L.-L., R. E. Nelson, Jr., and R. E. Narita.
1985. Net loss from trawl fisheries off Alaska. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 130-153. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Lutjeharms, J. R. E.
1981. Features of the southern Agulhas circulation from satellite remote sensing. *S. Afr. J. Sci.* 77:231-236.

1988. Meridional heat transport across the Subtropical Convergence by a warm eddy. *Nature* 331:251-254.
- Lutjeharms, J. R. E., and H. R. Valentine.
1988. Eddies at the Subtropical Convergence south of Africa. *J. Phys. Oceanogr.* 18:761-774.
- Lutjeharms, J. R. E., L. V. Shannon, and L. J. Beekman.
1988. On the surface drift of the Southern Ocean. *J. Mar. Res.* 46:267-279.
- Merrell, T. R.
1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

- Morris, R. J.
1980. Plastic debris in the surface waters of the South Atlantic. *Mar. Pollut. Bull.* 11:164-166.
- Neilson, J.
1985. The Oregon experience. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii. p. 154-159. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Piatt, J. F., and D. N. Nettleship.
1987. Incidental catch of marine birds and mammals in fishing nets off Newfoundland, Canada. *Mar. Pollut. Bull.* 18:344-349.
- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Ryan, P. G.
1987a. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.
1987b. The origin and fate of artefacts stranded on islands in the African sector of the Southern Ocean. *Environ. Conserv.* 14:341-346.
1988a. The characteristics and distribution of plastic particles at the sea-surface off the southwestern Cape Province, South Africa. *Mar. Environ. Res.* 25:249-273.
1988b. Effects of ingested plastic on seabird feeding: Evidence from chickens. *Mar. Pollut. Bull.* 19:125-128.
1988c. Intraspecific variation in plastic ingestion by seabirds and the flux of plastic through seabird populations. *Condor* 90:446-452.
- Ryan, P. G., A. D. Connell, and B. D. Gardner.
1988. Plastic ingestion and PCBs in seabirds: Is there a relationship? *Mar. Pollut. Bull.* 19:174-176.
- Ryan, P. G., and J. Cooper.
In press. Rockhopper penguins and other marine life threatened by driftnet fisheries at Tristan da Cunha. *Oryx*.
- Ryan, P. G., and C. L. Moloney.
1987. Effect of trawling on bird and seal distributions in the southern Benguela region. *Mar. Ecol. Prog. Ser.* 45:1-11.
- Ryan, P. G., and B. Rose.
1988. Migrant seabirds. In A. I. L. Payne and R. J. M. Crawford (editors), *Oceans of life off southern Africa*, p. 274-287. Vlaeberg, Cape Town.

- Shaughnessy, P. D.
1980. Entanglement of Cape fur seals with man-made objects. Mar. Pollut. Bull. 11:332-336.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Taylor, L.
1979. Degradable plastics: Solution or illusion? Chemtech 9:542-548.
- Tull, C. E., P. Germain, and A. W. May.
1972. Mortality of thick-billed murres in the west Greenland salmon fishery. Nature 237:42-44.
- Vauk, G. J. M., and E. Schrey.
1987. Litter pollution from ships in the German Bight. Mar. Pollut. Bull. 18:316-318.
- Winston, J. E.
1982. Drift plastic--an expanding niche for a marine invertebrate? Mar. Pollut. Bull. 13:348-351.
- Wirka, J.
1988. Wrapped in plastics: The environmental case for reducing plastics packaging. Environmental Action Foundation, Wash. D.C., 159 p.
- Wolfe, D. A.
1987. Persistent plastics and debris in the ocean: An international problem of ocean disposal. Mar. Pollut. Bull. 18:303-305.
-

INTERNATIONAL EFFORTS TO CONTROL MARINE DEBRIS IN THE ANTARCTIC

Michael F. Tillman
National Marine Fisheries Service, NOAA
1335 East-West Highway
Silver Spring, Maryland 20910, U.S.A.

ABSTRACT

Since much of the Antarctic, including the surrounding seas, remains in a relatively pristine state, the monitoring of environmental changes in this area often provides early warning of hazardous global phenomena, e.g., the stratospheric depletion of ozone. Reacting to a U.S. initiative, members of the Commission for the Conservation of Antarctic Marine Living Resources have taken steps to monitor the potential problem of marine debris, particularly from fishing operations. The Commission is joining with the Scientific Committee for Antarctic Research in establishing a program to monitor the effect of plastic pollution and entanglement on marine animals. The initiatives undertaken to establish monitoring programs for marine debris, the results to date, the reasons for their success and future needs in the Antarctic are discussed in this review.

INTRODUCTION

The 1984 Workshop on the Fate and Impact of Marine Debris provided ample warning that marine debris of terrestrial and shipborne origin was widespread in the marine environment and was apparently capable of contributing substantially to increased mortality of marine life (Shomura and Yoshida 1985). Of particular concern was the implication of debris arising from fishing operations (including lost or discarded net fragments, plastic packing bands, lines, and rope) in the harmful entanglement of substantial numbers of animals from many North Pacific populations of pinnipeds: northern fur seal, *Callorhinus ursinus* (Scordino 1985); Steller sea lion, *Eumetopias jubatus* (Calkins 1985); northern elephant seal, *Mirounga augustirostris*, California sea lion, *Zalophus californianus*, and harbor seal, *Phoca vitulina richardsi* (Stewart and Yochem 1985); and Hawaiian monk seal, *Monachus schuainslandi* (Henderson 1985). Fowler's (1985 1987) analyses of the substantial database for northern fur seals even suggested that the mortality of fur seals due to entanglement may be contributing significantly to declining trends (4-8% per year since the mid-to-late 1970's) of the population on the Pribilof Islands.

To begin addressing the uncertainties surrounding the marine debris problem while mitigating the known impacts, the 1984 workshop recommended, among other things, that educational efforts be undertaken to advise user and interest groups of the nature and scope of the issue. It was thought appropriate to include relevant international groups in this educational approach. The 1984 workshop also agreed that additional efforts should be undertaken to establish the severity of the debris problem in areas other than the North Pacific. Consequently, the stage was set for aggressive initiatives at several international forums to determine if the marine debris problem was occurring in other ocean basins.

Given the apparent adverse impact of marine debris, especially from fishing operations, upon North Pacific pinniped populations, it seemed reasonable to focus attention upon the Antarctic, where large populations of pinnipeds also occurred. In response to the establishment of a substantial international trawl fishery in the Antarctic during the 1970's, the Convention and Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) had come into force in 1982. The United States was a founding member of CCAMLR and brought the marine debris issue to the Commission's attention at its third annual meeting, in September 1984, 4 months after the convening of the marine debris workshop.

U.S. ANTARCTIC INITIATIVES

Organization and Mandate of CCAMLR

The CCAMLR is a unique international agreement which implements an ecosystem approach to the conservation and management of marine living resources found in the Antarctic. The CCAMLR convention area includes the marine area south of the Antarctic Convergence, the boundary between lat. 48° and 60°S which separates cold Antarctic waters from warmer subantarctic waters (Fig. 1). The area south of this boundary is considered the Antarctic marine ecosystem. The convention applies to "the populations of finfish, mollusks, crustaceans, and all other species of living organisms, including birds, found south of the Antarctic Convergence" (Anonymous 1988a).

The CCAMLR currently comprises 20 member nations, and an additional 4 nations have acceded to the convention but have not yet been accorded membership (Anonymous 1988a). The major operational units which undertake the convention's responsibilities (Fig. 2) are the Commission for the Conservation of Antarctic Marine Living Resources (the "Commission") and the Scientific Committee for the Conservation of Antarctic Marine Living Resources (the "Scientific Committee"). The work of these bodies is facilitated by a permanent secretariat which resides at CCAMLR headquarters in Hobart, Tasmania, Australia.

The convention mandates a management regime which ensures that harvesting of Antarctic species, such as finfish and krill, is conducted in a manner that considers ecological relationships among dependent and related species. Article II of the convention specifically requires the Commission to follow four basic principles of conservation (Sherman and Ryan 1988):




Figure 1.--Boundary of the area under the jurisdiction of the Convention for the Conservation of Antarctic Marine Living Resources (Anonymous 1988a).




Figure 2.--Organizational structure of the Convention for the Conservation of Antarctic Marine Living Resources (after Sherman and Ryan 1988).

1. To prevent any harvested population from falling below the level that ensures the greatest net annual increment to stable recruitment;
2. to maintain the ecological relationships between harvested, dependent, and related populations of Antarctic marine living resources;
3. to restore depleted populations; and
4. to prevent or minimize the risk of changes in the Antarctic marine ecosystem that are not potentially reversible over two or three decades.

It was within this ecosystem context that the United States was able to raise the marine debris issue. In particular, the fourth principle gave rise to a powerful argument that the Commission must act to prevent irreversible changes in the Antarctic marine ecosystem which might arise from harvesting activities, including the loss or disposal of debris resulting from those activities. At least the Commission found itself compelled to give the issue due consideration when the United States introduced it at the 1984 annual meeting.

U.S. Proposals and CCAMLR Response

1984 Initiative

In 1984, the U.S. delegation submitted and the Commission considered a paper entitled "Assessment and avoidance of incidental mortality of Antarctic marine living resources." This document indicated that, while there did not seem to be any problem with entanglement of animals in lost or discarded fishing gear and other marine debris in the convention area, there was growing evidence in other areas, e.g., the North Pacific, that significant numbers of nontarget marine organisms were being caught and killed in such debris, as well as being caught and killed incidentally during certain fishing operations. The Commission agreed with these conclusions, and asked its members to undertake steps to study and assess the possible sources, fates, and effects of marine debris in the convention area, including (Anonymous 1984):

- reviewing and reporting on past encounters with marine debris at sea or at coastal research stations;
- reporting on the nature of problems arising from debris such as fouled propellers or entangled animals, and
- periodically surveying beaches at research stations or other areas to ascertain the types, quantities, and sources of debris accumulating there.

The Commission also agreed that members should report on the number of birds, marine mammals, and other nontarget species taken incidentally during fishing operations. Moreover, members were asked to inform their nationals of international and national laws prohibiting or restricting the disposal of netting and other potentially hazardous materials at sea and to report on measures taken to assess, avoid, and mitigate incidental mortality of Antarctic marine life. Finally, it was agreed to include this item on the agenda for the 1985 meeting and to consider the desirability of marking fishing gear for identification purposes, as well as restricting the use of gillnets in the convention area.

In 1985, the Commission received formal reports from four members, including the United States, on steps taken in response to the basic monitoring program established in 1984. A number of oral reports were received as well, and the United States submitted a preliminary report of the proceedings of the 1984 Workshop on the Fate and Impact of Marine Debris. Based upon this information, the Commission again concluded that there was no evidence that significant quantities of fishing gear, binding material, or other hazardous debris had been or were being lost or discarded in the convention area (Anonymous 1985). However, given the compelling evidence for such debris in other ocean areas, including areas adjacent to the convention area, and of the extent of its harmful effects to marine life and of its hazards to navigation, the Commission agreed to continue its monitoring program.

The Commission further agreed that members should continue studying the feasibility and desirability of marking fishing gear and of maintaining inventories of such material brought into the convention area. However, given that there were no substantial gillnet operations in the area at the time, the Commission concluded that prohibiting the use of gillnets as a preventative measure could interfere unnecessarily with the Commission objective of assuring the rational use of resources. The Commission did agree to keep the matter under review.

1986 Initiative

At the 1986 meeting, the Commission received reports from members on monitoring results and the United States submitted a paper proposing additional steps for ensuring that accidental and incidental mortality of marine life did not become a problem in the convention area. While the information provided continued to indicate that incidental and accidental mortality of living marine resources did not appear to be a problem, the Commission recognized that such mortalities, including those resulting from entanglement in or ingestion of marine debris, could interfere with efforts to achieve the objectives of the convention (Anonymous 1986). As a consequence, the Commission agreed to new measures to reduce or prevent the at-sea discarding of fishing and other hazardous debris:

- Members would take steps to ratify and implement both optional Annex V of the 1978 Protocol to the International Convention for the Prevention of Pollution from Ships (MARPOL) and the International Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (London Dumping Convention); and
- the secretariat would prepare drafts of an information brochure to advise fishermen, researchers, and others working in the convention area of the hazards of marine debris; and of a placard for displaying on ships which listed the "do's and don'ts" for storing, handling, and discarding refuse.

The Commission agreed to continue its monitoring provisions and the collection of incidental catch data. Moreover, it agreed to undertake three new monitoring steps (Anonymous 1986):

1. recording and reporting fishing gear lost in the convention area;
2. if feasible, collecting and safely disposing of marine debris encountered; and
3. collecting samples of marine debris along with pertinent data, including species and numbers of entangled marine animals, for archival by the secretariat.

At the 1987 meeting, progress on all agreed monitoring measures was reviewed, and the Commission closely examined the information on lost or

discarded fishing gear obtained from national reviews of such data and from beach surveys in the convention area. Although several members observed no marine debris or entanglement problems, others reported sightings of debris consisting of fishing buoys, gas bottles, plastic containers, trawl net fragments, and plastic packing bands (Anonymous 1987). Moreover, two fur seals, *Arctocephalus gazella*, were seen entangled in derelict fishing nets and a third in longline gear. The Commission agreed not only to continue all elements of the monitoring program, including new steps agreed upon in 1986, but also to establish the issue of incidental/accidental mortality of Antarctic marine living resources as a standing item on the agenda for subsequent annual meetings.

The Commission also reviewed in 1987 the secretariat's drafts of an information brochure and a placard for display on vessels operating in the convention area. The secretariat was authorized to publish the agreed texts and members were urged to give these the widest possible circulation. Moreover, given that Annex V to the MARPOL Convention would prohibit or control the disposal of debris arising from fishing operations in the convention area, members were again specifically urged to ratify and implement this international measure.

In 1988, the Commission received further reports from members regarding loss of trawl cod ends and sightings of other derelict debris, including net fragments and packing bands. Moreover, five fur seals, *A. gazella*, were seen entangled in derelict fishing gear and two adult male fur seals died after becoming entangled in trammel nets (Anonymous 1988b). The Commission agreed to continue all elements of its monitoring program but noted that the reporting of incidental mortality as recommended in 1986 had been inadequate so far.

Also in 1988, the secretariat published and distributed the information brochure and placard for display on the ships of all member nations. As requested by the Commission, the U.S. has made these available to scientists and others working in the Antarctic and to the operators of vessels entering the convention area, including the National Science Foundation, the U.S. Coast Guard, and the National Oceanic and Atmospheric Administration.

FUTURE NEEDS AND ACTIVITIES

Improving Monitoring Efforts

The assumption is often made that much of the Antarctic, including the surrounding seas, remains unsullied by human activities. Consequently, if significant environmental changes are observed there, it is often presumed that these may be resulting from significant environmental perturbations occurring elsewhere on the globe, e.g., the stratospheric depletion of ozone resulting from the production and use of chlorofluorocarbon compounds in the Northern Hemisphere (Anonymous 1988e). The evidence reviewed so far by the Commission would tend to indicate that the marine debris problem in the Antarctic is minimal. That is, it would appear that the levels of debris discarded by vessels in the convention area or the amount brought in

by circulation or by other means from other ocean basins have not yet been sufficient to generate major problems for Antarctic marine life.

However, recent information suggests that the level of CCAMLR's monitoring efforts to date may not have been sufficient to ascertain the levels and consequences of marine debris effectively. The Bird Biology Subcommittee of the International Council of Scientific Unions, Scientific Committee for Antarctic Research (SCAR) concluded that a high proportion of Antarctic seabirds had ingested plastic particles, that the incidence was increasing in at least some species in the Southern Ocean and that the problem was particularly acute for procellariiform species which accumulate rather than excrete plastics (Anonymous 1988d). Van Franeker and Bell (1988) and Ainley et al. (1990) suggested that the source of the ingested plastic is from wintering areas outside the Antarctic. The SCAR Group of Specialists on Seals also noted that entanglements of Antarctic fur seals in discarded fishing gear had been reported from several areas around the Antarctic, including South Georgia, the South Shetland, Crozet, Marion, Heard, and Bouvet Islands (Anonymous 1988c). Consequently, one might conclude that CCAMLR has so far been seeing only the tip of the marine debris iceberg.

Taking note of CCAMLR's early monitoring initiatives in this area, both SCAR groups requested the Commission's assistance in examining the problem further. The SCAR's Bird Biology Subcommittee requested that CCAMLR consider initiating programs to monitor the level and effects of plastic pollution in subantarctic and Antarctic seabirds, considering both ingestion of plastic particles and entanglement. The SCAR Group of Specialists on Seals also requested that CCAMLR seek detailed information on the frequency of occurrence and nature of entanglement events involving seals in order to identify the causes of entanglement and trends in the frequency and extent of such entanglement over time (Anonymous 1988b).

At its 1988 meeting, however, the Commission noted that its monitoring program had three shortcomings relevant to SCAR's requests (Anonymous 1988b):

1. It did not address the problem of ingestion of plastics.
2. It did not specifically provide for quantitative and detailed reports of entanglement when fishing operations were not directly involved.
3. It may not provide adequately detailed information on incidental mortality during fishing operations to enable assessment of the problem or to monitor changes quantitatively.

To see if these shortcomings could be rectified so that assistance might be given to SCAR, the Commission authorized the chairman of the Scientific Committee to open a dialogue with the relevant SCAR groups (Anonymous 1988b). In particular SCAR's advice was sought (and provided at the 1989 meeting (Anonymous 1989)) on how the levels and effects of ingestion of plastics by Antarctic seabirds could be monitored, how quantitative

surveys could be conducted to determine the incidence, causes, and effects of marine mammal entanglements, and how the CCAMLR system of reporting incidental mortality might be improved in order to precisely determine the incidence, causes, and effects of such mortality. This new interaction between the Commission and SCAR should pave the way for greatly improving CCAMLR's pioneering efforts to monitor the marine debris problem.

Improving the Coordination of Efforts

The CCAMLR's exhortations on behalf of MARPOL apparently paid off, since Annex V came into force in December 1988 (Anonymous 1988e). It is now illegal for ships registered in the 35 ratifying nations, including the United States, to dump plastic debris such as that arising from fishing operations into the sea.

To become even more effective in controlling the marine debris problem in the Antarctic, it would seem desirable for the Commission to begin coordinating its actions with the International Maritime Organization (IMO). The IMO is the specialized agency of the United Nations which oversees implementation of MARPOL and the London Dumping Convention. This possible coordination, along with the pending cooperation between the Commission and SCAR, points out a growing need for an effective coordinating mechanism on this and other Antarctic issues.

In fact there has been a continuing debate among the Antarctic Treaty consultative parties (ATCP's) regarding the need for an Antarctic Treaty secretariat (Kimball 1987). The ATCP's favoring such a secretariat point to the increasing variety and complexity of issues being dealt with which require more numerous and more frequent communications within and between instruments of the Antarctic Treaty system, including CCAMLR, as well as with other relevant international organizations and elements of the outside world. The growing number of players becoming involved in dealing effectively with the issue of marine debris in the Antarctic (CCAMLR, SCAR, and IMO) may well provide another argument in favor of a secretariat.

DISCUSSION

Despite possible shortcomings and problems, it would appear that substantial progress has been made in trying to deal with the issue of marine debris in the Antarctic. The CCAMLR's monitoring program has evolved quite rapidly since the United States introduced the issue in 1984. Although the program is, perhaps, not yet as quantitative as some scientists would wish, the Commission is at least in a very good position to ascertain and evaluate trends in levels of debris and entanglements of marine life.

Under the convention, the Commission must take all of its decisions by consensus, which has led at times to a lowest-common-denominator-syndrome and resulted in somewhat ineffectual measures. So, the progress made with respect to marine debris might seem all the more remarkable unless one considered it in the light of the unique nature of the convention itself. The CCAMLR not only requires an ecosystem approach to the conservation and management of living marine resources but also sets forth the principle

that the Commission must act to prevent or minimize irreversible changes to that ecosystem. More than anything, these unique provisions probably account for the success achieved on the issue.

The philosophy behind CCAMLR provides great flexibility and a basis for dealing with many kinds of marine conservation issues, not just those dealing with the use of resources. This is a powerful tool, and the convention should be taken seriously as a model for all future resource use conventions and agreements in other ocean areas.

REFERENCES

- Ainley, D. G., W. R. Fraser, and L. B. Spear.
 1990. The incidence of plastics in the diets of Antarctic seabirds. In R. S. Shomura and M. L. Godfrey (editors) Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Anonymous.
 1984. Report of the third meeting of the Commission. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
1985. Report of the fourth meeting of the Commission. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
1986. Report of the fifth meeting of the Commission. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
1987. Report of the sixth meeting of the Commission. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
- 1988a. Basic documents (3d ed.). Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
- 1988b. Report of the seventh meeting of the Commission. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.
- 1988c. Report of the meeting of the SCAR group of specialists on seals. International Council of the Scientific Unions, Scientific Committee in Antarctic Research, 23-25 August 1988, Hobart, Tasmania, 23 p. [Unpubl. MS]
- 1988d. Scientific Committee on Antarctic Research Bird Biology. Subcommittee: Minutes of 1988 meeting. Cormorant 16:138-58.
- 1988e. World resources 1988-89. Basic Books, N.Y., 372 p.

1989. Report of the eighth meeting of the scientific committee. Commission for the Conservation of Antarctic Marine Living Resources, Hobart, Australia.

Calkins, D. G.

1985. Stellar sea lion entanglement in marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 308-314. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Fowler, C. W.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 27-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. Mar. Pollut. Bull. 18:326-335.

Henderson, J. R.

1985. A review of Hawaiian monk seal entanglements in marine debris. In R.S. Shomura and H.O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 326-335. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Kimball, L.

1987. Report on Antarctica. Int. Inst. Environ. Dev., 34 p. [Unpubl. MS]

Scordino, J.

1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Sherman, K., and A. F. Ryan.

1988. Antarctic marine living resources. Oceanus 31(2):59-63.

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

Stewart, B. S., and P. K. Yochem.

1985. Entanglement of pinnipeds in net and line fragments and other debris in the Southern California Bight. *In* R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 315-325. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Van Franeker, J. A., and P. J. Bell.

1988. Plastic ingestion by petrels breeding in Antarctica. *Mar. Pollut. Bull.* 19:672-674.

COOPERATIVE RESEARCH ON PETROLEUM POLLUTION
IN THE CARIBBEAN: THE CARIPOL PROGRAM

J. E. Corredor and J. M. Morell
Department of Marine Sciences
University of Puerto Rico
Mayaguez Puerto Rico 00709-5000


ABSTRACT

The CARIPOL program is a cooperative regional effort to assess the state of pollution of the marine environment in the Caribbean and adjacent regions. In its initial phase, CARIPOL has concentrated on the assessment of petroleum pollution through the monitoring of three easily determined variables: the occurrence of tar aggregates on beaches, of floating tar at sea, and of dissolved or dispersed petroleum hydrocarbons. A data base of greater than 7,000, 680, and 1,460 data points, respectively, for the three variables has been accumulated through submissions from 14 countries in the region.


Tar on beaches is a serious problem in the region, especially in the Windward islands, the Cayman islands, and the archipelago of Aruba, Curacao, and Bonaire where loads of up to 1 kg of tar/m of beach front have been reported. Other affected areas include the east coast of Florida, the Yucantan peninsula, and Campeche Sound. The occurrence of floating tar has been closely correlated with tanker traffic in areas such as the south coast of Puerto Rico and the Straits of Florida. Dissolved and dispersed petroleum hydrocarbons reach critical levels only in enclosed waters such as bays and harbors subject to intense maritime traffic or industrial petroleum sites.

The CARIPOL program has now embarked upon a second phase to assess the accumulation of petroleum hydrocarbons in sediments and organisms. Initial results indicate that although these compounds are rapidly degraded when released to the water, they may persist for extended periods upon reaching marine sediments.

SESSION I



AMOUNTS, TYPES, DISTRIBUTION, AND SOURCES OF MARINE DEBRIS



QUANTITATIVE ESTIMATES OF GARBAGE GENERATION AND DISPOSAL
IN THE U.S. MARITIME SECTORS BEFORE AND AFTER MARPOL ANNEX V

J. Cantin, J. Eyraud, and C. Fenton
Eastern Research Group, Inc.
Arlington, Massachusetts 02174, U.S.A.

ABSTRACT

Annex V of MARPOL 73/78 is regarded as an important instrument for reducing the amounts of plastics and other debris discarded into the ocean. Estimates of the aggregate quantities of garbage discarded are outdated, however, and represent only order of magnitude efforts. In this paper, the authors present updated estimates of the amounts of plastics and other debris generated in the U.S. maritime sectors.

The analysis covers both public and private sectors, including merchant marine vessels active in U.S. trade; commercial fishing vessels; recreational boats; research and industrial vessels; U.S. Navy, Coast Guard, and Army ships; and vessels and structures associated with offshore oil and gas operations. Current disposal practices as well as disposal practices under Annex V are analyzed and used to develop estimates of how the disposition of garbage generated at sea, i.e., the amounts dumped overboard, brought back to shore for disposal, and incinerated, will change under the new regulations.

INTRODUCTION

Two questions which underlie the debate over the U.S. ratification of MARPOL Annex V are: (1) How much garbage is being dumped overboard from vessels? and (2) What effect would the specific restrictions contained in Annex V have on the overall marine debris problem? Throughout the numerous congressional hearings which led up to U.S. ratification, only one source of aggregate data, a 1975 study by the National Academy of Sciences (NAS), was identified which addressed these questions. That study, however, examined the entire world fleet and included sources of debris which will not be regulated by Annex V (Table 1). It also made no attempt to account for actual disposal practices, reporting instead on the quantities of garbage "potentially" dumped (NAS 1975).

The present study utilizes current information for the U.S. maritime sectors to develop current and more comprehensive estimates of garbage

Table 1.--Global marine litter estimates
(National Academy of Science 1975).

Garbage types and sources	Ton/year	Percent
Sources regulated under MARPOL Annex V		
Crew-related wastes		
Merchant marine	11,000	1.8
Passenger vessels	2,800	0.4
Commercial fishing	34,000	5.4
Recreational boats	10,300	1.6
Military	7,400	1.2
Oil drilling and platforms	400	0.1
Commercial wastes		
Merchant cargo wastes or dunnage	560,000	89.5
Annex V subtotal	625,900	100.0
Sources not regulated under MARPOL Annex V		
Fishing gear loss	100	1.0
Loss due to catastrophe ^a	10,000	99.0
Non-Annex V subtotal	10,000	1000.0
Total	636,000	100.0

^aIncludes debris from shipwrecks or marine storms.

quantities. The main components of the model are: vessel populations, entrances to U.S. ports, crew sizes, garbage generation factors, and plastics as a percent of total garbage. It also fills a gap left in previous studies by addressing historical shipboard disposal practices and changes in practice expected to result from implementation of Annex V. It must be noted, however, that few direct measurements of garbage generation and disposal practices exist. The methodological improvements offered in this study are based on updated data where they exist, and on substantial anecdotal information collected throughout the course of broader regulatory studies of MARPOL Annex V (Eastern Research Group (ERG) 1988a, 1988b).

The study first reviews information related to the sources, types, and quantities of garbage generated in the various maritime sectors. Data on per capita garbage generation, crew size, voyage length, and annual ship utilization factors are used to derive estimates of per voyage, per vessel, and annual aggregate garbage quantities generated. Estimates are made for the following sectors:

- Merchant shipping.
- Commercial fishing.

- Commercial passenger vessels.
- Recreational boating.
- Offshore oil and gas operations.
- Research and other miscellaneous vessels.
- U.S. Navy, Coast Guard, Army, and other government vessels.

An analysis of historical garbage disposal practices in these sectors is used to estimate the pre-Annex V garbage quantities dumped overboard, brought back to shore for disposal, or burned in marine incinerators. Under MARPOL Annex V, ships may be forced to alter their current disposal practices. An analysis of options available for compliance with Annex V is used to derive the post-Annex V disposition of vessel garbage.

GARBAGE ESTIMATION PARAMETERS

This section reviews the types and quantities of garbage generated in the maritime sectors. This information is combined with data on vessel populations, crew sizes, and voyage lengths from the supporting statistical section to produce estimates of the per voyage, per vessel, and aggregate annual garbage quantities generated.

Types of Garbage Generated

Several types of garbage are generated by vessels operating at sea. In this study, "domestic" garbage refers to general garbage such as galley refuse (food wastes, food packaging materials) and garbage from the hotel areas of the vessel (discarded items such as smoking materials and packaging, shampoo bottles, and razors). Wastes associated with normal ship operations, such as rags and containers, are also included. Some vessels generate an additional amount of "commercial" waste. Examples include cargo dunnage, spent fishing gear, and disposable or single-use oceanographic research instruments. These are discussed separately.

Domestic Wastes

Several sources of information on the quantities of garbage generated by ships are available. A series of studies done for the Intergovernmental Maritime Consultative Organization, the predecessor to the International Maritime Organization (IMO), are judged to represent the best available estimates (IMO 1987). These rates, representing per capita daily quantities, were reported as follows:

- Harbor vessels--1.0 kg (2.2 lb).
- Inland and coastal vessels--1.5 kg (3.3 lb).
- Oceangoing cargo vessels--2.0 kg (4.4 lb).
- Oceangoing cruise vessels--2.4 kg (5.3 lb).

As shown, the rates vary depending upon the type of vessel and where the vessel operates.

These rates appear consistent with those obtained elsewhere. The U.S. Navy, for example, examined garbage generation aboard naval ships in 1971 (Naval Ship Engineering Center 1971) and again in 1988 (L. Koss and Lt. Mullenhard, U.S. Navy, pers. commun. 1988), and reported estimates of 1.39 and 1.43 kg/person/day (3.05 and 3.15 lb), respectively. On land, the U.S. Environmental Protection Agency (EPA) estimates that each person generates an average of 1.82 kg (4.0 lb) of garbage per day (National Solid Waste Management Association (NSWMA) n.d.). Thus, the rates from the IMO studies appear consistent with those obtained elsewhere.

In this study, the IMO rates for oceangoing cargo and cruise vessels are applied respectively to cargo and passenger vessels which operate over the open ocean. The IMO rate for inland and coastal vessels is applied to crafts which travel inland or along the coastline.

Commercial Wastes

Several classes of vessels generate wastes associated with their commercial activities. These wastes are distinct from any operational wastes generated through normal vessel repair and maintenance activities, which are included under the category of domestic wastes.

Commercial cargo wastes or dunnage.--The term "dunnage" covers materials such as timber, plywood, pallets, rope, and plastic sheeting used to protect, separate, and secure cargo carried in break-bulk form. In the study (NAS 1975), these types of cargo wastes dominated the aggregate waste quantity estimates (Table 1). Since the mid-1970's, however, the trend towards containerization, and changes in cargo handling methods, are believed to have greatly reduced the amount of dunnage used. Cargo carried by containerships or other types of intermodal ships (barge carriers, roll-on/roll-off) is generally unloaded for transshipment in port without being disturbed. According to officials of the American Institute of Merchant Shipping, therefore, the "vast majority" of cargo ships now produce no dunnage waste (J. Cox, American Institute of Merchant Shipping, pers. commun. 1988).

The quantities and types of dunnage still used in general cargo trade vary, and depend upon the type of cargo being carried. Break-bulk shipments of food products, for example, use mainly cardboard for separation and protection of the cargo. For a highly explosive shipment of ammunition, however, extensive wooden encasements are constructed to protect against movement of the cargo. In such cases, tens of thousands of board feet of lumber may be used. Palletized cargoes may be shrink-wrapped and secured with steel or plastic strapping, but these are not normally removed prior to final delivery at the customer's facilities. Lumber and plywood are the most common materials used.

Marine terminal operators familiar with the loading and unloading of break-bulk ships indicate that very little plastic waste is generated. One reported use for plastic is to capture leaks of moisture or hydraulic

fluids in the vessel's cargo hatches. This plastic would likely be removed when the vessel is being unloaded. As indicated, plastic shrink-wrap is used with palletized cargo, but this is generally not removed either on board the ship or in port.

Estimates of per-vessel and aggregate dunnage quantities are difficult to make based upon the limited data. The study (NAS 1975) estimated that general cargo ships generate up to 285 tons of dunnage per year. This estimate contrasts significantly with information provided by marine terminal operators and shipping interests. Due largely to the trend towards containerization of cargo, there appears to be much less dunnage used today. Relying on current reports, this study assumes that two-thirds of the general cargo vessels entering U.S. ports generate dunnage in the form of lumber, one third generate only cardboard, and that 10% generate plastic waste. The quantities used are estimated as follows:

- Lumber--48.6 m³ (approximately 20,000 board ft) per vessel entrance.
- Cardboard--23.6 m³ (30 yd³) per vessel entrance.
- Plastic--assumed to be generated by only 1 of every 10 break-bulk ships entering U.S. ports, in minimal quantities of 0.12 m³ (4.0 ft³) per entrance.

As the estimates presented later will show, under these assumptions the amount of waste generated by general cargo ships represents only a small proportion of the total garbage volume regulated by Annex V. Furthermore, overall estimates of plastics are not particularly sensitive to assumptions regarding dunnage volumes.

Fishing gear wastes.--Commercial fishing activity also contributes to the problem of marine debris, and to plastics in particular. Whether trawl gear, set nets, or lines are being used, occasional fouling of equipment, such as the tearing or twisting of nets and lines, will occur. During repair, portions of nets, excess line, floats, and other gear wastes may be generated. All such materials are nowadays made of synthetics, and are prohibited from disposal under Annex V. In addition to these items, substantial quantities of fishing gear are also lost accidentally. Annex V, however, does not cover this category of debris, hence no estimates are made here.

Limited information is available on the amounts of gear waste generated. The Foreign Fisheries Observer Program, National Marine Fisheries Service (NMFS) has monitored fishing gear repair operations on board foreign vessels active in Alaskan waters. Berger and Armistead (1987) analyzed data from this program and found that fishing gear repair operations took place about once every 4.9 days. The U.S. vessels active in the same fisheries report a similar incidence of gear repair (J. Gnagey, Alaska Trawl Fisheries, pers. commun. 1988; Z. Grader, Pacific Coast Federation of Fishermen's Associations, pers. commun. 1988). Discarded webbing was found to be typically small, with only 21.2% of pieces deemed to be "of a mesh

size or area thought most likely to entangle marine mammals" (Berger and Armistead 1987). Some 57.9% of discarded pieces consisted of "primarily loose strands of twine or pieces with a mesh size of less than 100 mm."

Working with these data, the amount of gear waste generated over a typical cruise is estimated to be relatively small. Assuming that (1) the average piece of webbing produced during net repair is 1 m², (2) webbing is composed of 15 mm diameter twine (Uchida 1985), and (3) the incidence of net repair and discard is twice per week, than the volume of waste generated over even a 30-day fishing expedition would amount to just over 0.12 m³ (4 ft³) of net material (ERG estimates).

Evidence suggests that these estimates from the Alaskan fisheries may represent an upper bound on the frequency and amount of fishing gear waste discarded. Beach surveys have found the concentration of fishing gear waste off the Alaskan islands to be among the highest noted anywhere (e.g., Merrell 1985). Discussions with fishery representatives elsewhere in the United States have indicated that net repair operations while at sea are relatively less common. Moreover, net fragments and floats in other fisheries are reported to be retained on board for use in repair operations, rather than discarded. Spare nets may also be carried in order to avoid at-sea repairs completely (ERG 1988a, 1988b). In this study, it is assumed that the average volume of fishing gear waste generated in all U.S. fisheries is half of that calculated for Alaska. An estimate of 0.03 m³ (1 ft³) of gear waste per week at sea is assumed.

Nongear fishing wastes.--Certain fisheries produce additional quantities of wastes due to their use of specialized fishing techniques. These include longline fishing, which uses packaged bait (B. Alverson, Longliner Vessel Owners Association, pers. commun. 1988), and the herring fishery, which utilizes quantities of packaged salt (J. Kaelin, Associated Fisheries of Maine, pers. commun. 1988). Longline bait is sold frozen, with packages wrapped in plastic, packed in cardboard boxes, and secured with plastic strapping. Chemical light or Cyalume sticks are also used to attract fish. These sticks, about the size of a pencil, are themselves made of plastic. Salt used in the herring fishery comes in large plastic bags.

Available estimates of marine waste disposal do not address this source of waste. The Center for Environmental Education (CEE) reports that longline gear is used in at least five different fisheries (CEE 1986), and that longline vessels can bait up to 5,000 hooks per day. In order to capture the additional wastes produced by vessels in these fisheries, it is assumed in this study that they generate twice the normal volume of fishing gear waste.

Research vessel wastes.--Research vessels may generate additional plastic wastes in the form of packing materials from research instruments brought on board, and from disposable measuring instruments used in monitoring oceanic experiments. Based on discussions with representatives at various research institutions, an additional 0.12 m³ (4 ft³) of plastic waste per voyage is assumed.

Dry Versus Wet Garbage

The nature of the MARPOL Annex V regulations makes it necessary to estimate separately the amounts of "wet" garbage (food waste) and "dry" garbage generated. Many vessels are expected to separate plastics from their dry garbage and then dispose of the remainder while in areas where no Annex V restrictions apply. Some, however, may have to retain all dry garbage for onshore disposal (and even wet garbage, in some cases) depending on whether they operate in Annex V "special areas" or in coastal waters. (Special areas currently include the Mediterranean Sea, Baltic Sea, Red Sea, Black Sea, and Persian Gulf Areas; Table 2.)

The dry garbage component of the overall solid waste stream is estimated based on the recent U.S. Navy study (Koss and Mullenhard, pers. commun. 1988). In this study, dry garbage accounted for 59.4% of domestic waste by weight, while wet garbage accounted for 40.6%. These percentages are similar to those found in the earlier, more extensive Navy studies, where the dry garbage component was estimated at 43.6%.

Plastics as a Percentage of Total Wastes

Annex V places a complete prohibition on the overboard disposal of plastics. Estimates of the percentage of the overall vessel waste stream accounted for by plastics are needed in order to develop projections of the quantity of garbage that may be brought back to shore for disposal.

The EPA estimates that plastics represent 6.5% (by weight) of all household and commercial solid waste on land (NSWMA n.d.). The relevance of this estimate to vessel operations is uncertain, however, because of likely differences in the types of waste generated at sea. In the national EPA estimate, paper and paperboard waste makes up 42% of the total, and yard waste accounts for another 16%. Garbage generated at sea is likely to contain much less paper waste and no yard waste. Under these assumptions, plastics would represent a larger share of the waste stream at sea than it would on land. At the same time, though, national estimates would include discards of durable plastic objects and industrial plastic waste, very little of which is generated at sea.

Studies done by the U.S. Navy in 1971 (Naval Ship Engineering Center 1971) and 1988 (Koss and Mullenhard, pers. commun. 1988) represent the only direct estimates of plastic wastes based on actual operating experience. In 1971, plastics were found to account for only 0.3% of total garbage by weight. This study covered numerous vessels and was used by the NAS in their estimates (NAS 1975). In 1988, however, the Navy found that the plastics share of total garbage weight had risen to 6.7%--an apparent twentyfold increase. It must be noted that the more recent study is based on an analysis of a single Navy vessel operating over a short (32-h) cruise. Thus, the figures may not be representative.

In reviewing the data from the Navy studies, the question of potential differences in plastics usage between Navy and other vessels arises. Navy vessels carry extensive electronic equipment on board which may be wrapped in plastic "bubble" wrap and other cushioning materials. This source of

Table 2.--Summary of MARPOL Annex V restrictions
(International Maritime Organization 1987).

Garbage type	All vessels		Offshore platforms and associated vessels ^b
	Outside special areas ^a	Within special areas ^a	
Plastics	Disposal prohibited	Disposal prohibited	Disposal prohibited
Floating dunnage, lining and packing materials	>25 nmi from land	Disposal prohibited	Disposal prohibited
Paper, rags, glass, etc., not ground	>12 nmi from land	Disposal prohibited	Disposal prohibited
Paper, rags, glass, etc., ground ^c	>3 nmi from land	Disposal prohibited	Disposal prohibited
Food waste, not ground	>12 nmi from land	Disposal prohibited	Disposal prohibited
Food waste, ground ^c	>3 nmi from land	>12 nmi from land	>12 nmi from land
Mixed refuse types	(d)	(d)	(d)

^aAnnex V special areas include the Mediterranean, Baltic, Red, and Black Seas, and the Persian Gulf Areas.

^bOffshore platforms and associated vessels include all fixed or floating platforms engaged in exploration or exploitation and associated offshore processing of seabed mineral resources, and all vessels alongside or within 500 m of such platforms.

^cGround waste must be able to pass through a screen with mesh size no larger than 25 mm (0.1 in).

^dWhen garbage is mixed with other harmful substances having different disposal or discharge requirements, the more stringent disposal requirements shall apply.

plastics is not generally present on board other types of vessels. Second, considerably more at-sea repair occurs on board Navy ships than aboard merchant marine or fishing vessels. Tools and replacement parts may be packaged in plastic, and parts themselves may be plastic. Wire and cable have plastic insulation. On the other hand, due to the large crew sizes, food supplies on board Navy vessels are generally purchased in bulk. The reduced packaging associated with this bulk purchasing would suggest smaller

plastic generation rate. Navy vessels may, therefore, generate more plastics from operational sources, but less from galley refuse.

In the absence of any conclusive evidence on the amount of plastics contained in the ship's waste stream, the results from the most recent Navy study (Koss and Mullenhard, pers. commun. 1988) have been used. This study indicates that plastics represent 6.7% of all wet and dry solid waste, and 11.3% of dry solid waste, by weight. Additional studies of this nature relative to shipping and other maritime sectors would certainly be welcome.

Garbage Densities

Further analysis of garbage generation patterns requires estimates of the density of garbage. These are needed in order to convert the weight of a given accumulation of garbage to volume terms. At sea, it may be the volume of garbage, rather than its weight, that figures in decisions regarding disposal options.

Table 3 shows estimates of garbage density for shipboard types of garbage. It will be noted that no sources of data specific to plastics were identified. Studies done for the State of New York by Franklin Associates (V. Sellers, Franklin Associates, pers. commun. 1988) found that 1,000 kg (2,200 lb) of uncompressed plastic soda containers had an average volume of 20.8 m³ (325 ft³), suggesting a density of 48.1 kg/m³ (3.07 lb/ft³). Navy officials, however, have suggested that a much lower density of 15.4 kg/m³ (1 lb/ft³) is appropriate (Koss and Mullenhard, pers. commun. 1988). This would imply that plastics used on board ship weigh one-third as much as empty soda bottles--an apparently generous volume estimate. In the absence of any data specific to ships, however, the Navy's estimate of 15.4 kg/m³ (1 lb/ft³) is incorporated. Again, this estimate is the more generous of those available in terms of estimating the volume of plastics generated on board.

Table 3.--Estimates of density for shipboard-generated garbage.

Source of estimate and garbage type	Density	
	kg/m ³	lb/ft ³
Society of Naval Architects and Marine Engineers (1982):		
Dry rubbish	100.0	6.3
Dry garbage	120.0	7.5
Refuse, 70% wet	640.0	40.0
Food waste	400 to 1,000	25.0 to 68.8
Gassan (1978):		
Hotel solids	277.0	17.2

Table 4 summarizes the numerous estimates and assumptions used in calculating garbage weights and volumes. The reader should note that densities are calculated for each of the different components of mixed garbage. Any conversions from garbage weight to volume made in this study must be considered, therefore, in the context of the density values used.

GARBAGE HANDLING AND DISPOSAL PRACTICES BEFORE AND AFTER MARPOL ANNEX V

This section summarizes the more substantial review of garbage handling and disposal practices contained in ERG (1988b). Estimates of the percentage of vessels using each of the various garbage handling and disposal methods, both historically and under Annex V, are used to evaluate the disposition of the aggregate garbage quantities under pre- and post-Annex V assumptions.

Pre-Annex V Garbage Handling and Disposal Practices

The historical methods employed by shipboard crews to dispose of garbage provide a basis for determining the current disposition of the garbage generated on board, i.e., how much is discarded overboard, how much brought back to shore for disposal, and how much is burned in onboard incinerators.

In most sectors, garbage handling practices vary depending on where the ship operates. Over deep-sea routes, garbage is typically collected throughout the ship and discharged daily. Closer to shore, crews are more likely to retain garbage for onshore disposal. The historical practice of ocean dumping while out at sea has been confirmed in most sectors. A representative of the American Institute of Merchant Shipping, for instance, states that: "Generally aboard merchant vessels on the high seas, waste generated as a result of vessel operations and deck maintenance is disposed of directly overboard" (Corrado 1986).

The predominance of ocean disposal is also indicated by statistics kept by the Department of Agriculture's Animal and Plant Health Inspection Service (APHIS). This agency requires ships entering the United States from foreign ports to incinerate, sterilize, or otherwise sanitize any garbage prior to disposing of it on shore. The APHIS inspection records for fiscal year 1986, for example, show that only 2.5% of vessels entering the United States from foreign ports off-loaded any garbage (A. Langston, U.S. Department of Agriculture, pers. commun. 1988).

Most commercial fishing groups also acknowledge that garbage dumping has traditionally been the most widely used means of getting rid of any trash which accumulates.

The use of garbage handling equipment such as grinders, compactors, or incinerators has not been widespread in the maritime sectors. Only some newer ships are equipped with such equipment. Until now, overboard disposal while well out at sea has been the most convenient and inexpensive method available. Based on discussions with operators in the merchant

Table 4.--Assumptions and estimates used
in garbage generation calculations.

Domestic garbage generation rates (International Maritime Organization 1987)		
Vessel category	Per capita per day	
	kg	lb
Oceangoing	2.0	4.4
Coastal	1.5	3.3
Inland/harbor	1.0	2.2
Passenger cruise	2.4	5.3

Fishing waste generations rates (Eastern Research Group estimates)				
Vessel category	Per vessel per day			
	m ³	kg	ft ³	lb
Normal vessels	0.004	0.064	0.140	0.140
Longliners, etc.	0.008	0.127	0.280	0.280

Domestic waste components, by weight (Koss and Mullenhard, pers. commun. 1988)		
Garbage type	As percent	As percent
	of all garbage by weight	of dry garbage by weight
Wet (food waste)	40.6	--
Dry (nonfood waste)	59.4	100.0
Plastic	6.7	11.3
Glass	4.1	6.9
Metal	13.0	21.9
Rubber	0.3	0.5
Paper, other	35.2	59.3

Garbage density (Society of Naval Architects and Marine Engineers 1982),
except for plastic density, which was suggested by Navy personnel

Garbage type	kg/m ³	m ³ /kg	lb/ft ³	ft ³ /lb
Total garbage	174.2	0.006	10.89	0.10
Dry garbage	100.0	0.010	6.30	0.16
Food waste	640.0	0.002	40.00	0.03
Plastics	16.0	0.063	1.00	1.00

marine, commercial fishing, and government sectors, it is assumed here that for most ships, overboard disposal is the predominant method used.

Vessels which spend more time operating close to shore are less likely to rely on overboard disposal. There are several possible reasons for this: Laws and regulations may already prohibit dumping in such areas; vessels are away from port for shorter periods and thereby generate less garbage; or operators may be conscious about dumping close to shore. Several categories of vessels have been identified as using alternative disposal means. These include segments of the coastal trade fleet, tug and towboat operators, recreational boaters, offshore oil and gas operations, some industrial and research vessels, and some Coast Guard vessels.

Post-Annex V Garbage Handling and Disposal Practices

Under MARPOL Annex V, vessel operators may have to implement changes in garbage handling procedures in order to achieve compliance with the requirements of the regulations. The actions taken by an individual operator will depend upon a number of factors, including (1) where the vessel operates and the specific restrictions of Annex V which apply in those areas, (2) the quantities and types of garbage generated by each vessel, and (3) the cost and noncost factors which influence the selection of compliance methods.

Each vessel owner or operator will evaluate his operations relative to the requirements of MARPOL Annex V. Table 2 presented a summary of the restrictions introduced by Annex V for the various types of garbage. Disposal of plastics is prohibited everywhere, and disposal of other types of wastes is restricted for vessels operating near shore. Vessels operating in special areas are prohibited from disposing of anything except food wastes, and then only beyond 12 mi from shore. A separate set of rules apply to offshore oil and gas operations.

Alternative compliance options have been analyzed in terms of their relative costs and conveniences in the regulatory analysis prepared for the Coast Guard (ERG 1988b). Among the compliance methods examined were: substitution of plastics, storage of garbage for onshore disposal, use of compactors to reduce garbage volumes, with subsequent disposal on shore, and installation of onboard garbage incinerators. The model used for comparing these alternatives took into account all of the relevant costs associated with each option, including the volumes and types of garbage generated; equipment, installation, and operating costs; the opportunity costs of current garbage handling and disposal procedures (i.e., not paying crews to dump garbage); and costs associated with off-loading and disposing of garbage in port.

The cost comparison model shows that for most vessels, onboard separation and storage of plastic garbage, with eventual disposal in port, is the least costly alternative (see ERG 1988b). As the garbage generation tables below will show, the quantities of plastics generated by most vessels would not present extreme storage difficulties. Where garbage volumes may cause inconveniences or storage problems, compactors can be used to reduce the volumes.

Several factors not captured by the cost comparison may steer vessel operators towards more costly compliance methods. If, for whatever reason, vessels anticipate the accumulation of large quantities of garbage on board, they may consider methods that reduce or eliminate this burden, even if it increases their costs. Operators may be concerned about situations where onshore garbage disposal would not be possible for extended periods of time, due to delays or the inability to obtain removal service in port. Finally, the cost comparisons do not consider issues caused by operations in special areas, where additional restrictions on the disposal of garbage will apply.

When both the cost and noncost issues are considered, most smaller vessels are still projected to choose separation and storage of garbage which they will no longer be able to dump overboard. Extensive use of onboard garbage compaction equipment is forecast, however, for larger commercial fishing vessels and for a majority of domestic trade merchant ships. Such equipment will be used to reduce the volume of garbage retained on board and to facilitate handling and disposal in port. Equipment manufacturers indicate that equipment suitable for onboard use can achieve a compaction ratio of between 500 and 1,000%, although for pure plastics the ratio is lower unless the material is first shredded. Only ships in the merchant shipping foreign trade category and some larger research and passenger ships are expected to select onboard incinerators. In the case of foreign trade vessels, the decision to invest in incinerators will not be based simply on economics, as incinerators represent the most expensive means of compliance, but rather upon the increased convenience afforded to the vessel. Time spent in port is extremely costly, thus incinerators may be viewed as "insurance" against the possibility of being delayed due to difficulties in obtaining garbage disposal services. It must be noted, however, that current or future air pollution standards for marine incinerators could greatly increase the cost of this option.

Special mention should be made of the solution expected to be adopted by U.S. Navy ships. As shown below, the Navy has particular garbage disposal problems due to the large number of crewmen on board. According to the most recent reports, Navy ships are expected to be outfitted with thermal extrusion equipment specially designed for shipboard application. This technology will enable Navy crews to melt down all plastics generated on board and extrude them into a storable form.

Pre- and Post-Annex V Garbage Disposition

Table 5 presents estimates of the pre- and post-Annex V distribution of vessels in the merchant shipping sector according to the garbage handling and disposal practices used. The distributions reflect ERG conclusions from the review of disposal practices and options described above. Similar distributions have been developed for each of the maritime sectors under study, but are not shown here.

Aggregate quantities of domestic garbage derived in the supporting statistical section are shown in the first column of Table 6 below. The table shows the pre- and post-Annex V disposition of these garbage

Table 5.--Current garbage handling and disposal practices and projected practices under MARPOL Annex V merchant shipping sector (Eastern Research Group estimates).

Merchant shipping	Current compliance rate (%)	Current compliance choices (%)			Annex V compliance choices (%)			
		Dump	Store	Compact	Incinerate	Store ^a	Compact	Incinerate
Foreign trade								
U.S. vessels								
Atlantic/Gulf/Pacific ports	5	95	0	0	5	5	70	25
Noncontiguous ports	5	95	0	0	5	5	70	25
Foreign vessels								
Atlantic/Gulf/Pacific ports	5	95	0	0	5	5	70	25
Noncontiguous ports	5	95	0	0	5	5	70	25
Noncontiguous trade	5	95	0	0	5	5	80	15
Great Lakes vessels								
1,000 GT and over	100	0	25	50	25	25	50	25
Under 1,000 GT	100	0	25	50	25	25	50	25
Military Sealift Command charter	5	95	0	0	5	5	75	20
Temporarily inactive vessels	5	95	0	0	5	5	75	20
Coastal shipping								
Ships								
1,000 GT and over	25	60	40	0	0	10	75	15
Under 1,000 GT	5	95	5	0	0	15	75	10
Tow/tugboats								
Large (inspected)	20	80	20	0	0	50	45	5
Small	20	80	20	0	0	60	40	0

^aRefers to storage of all garbage that vessels would not be permitted to dump. Assumes other garbage will be dumped where allowed under Annex V.

Table 6.--Final disposition of vessel-generated domestic waste, aggregated sector totals
(annual quantities) (GT = gross tons; MT = metric tons).

Maritime sector	Total generated annually (MT)	Pre-Annex V					
		Off-loaded in port		Incinerated at sea		Dumped overboard	
		(MT)	(m ³) ^a	(MT)	(m ³) ^a	(MT)	(m ³) ^a
Merchant shipping	30,949	3,302	39,794	1,148	14,971	26,499	349,304
Commercial passenger vessels	258,074	232,121	3,026,799	638	8,322	25,315	330,095
Commercial fishing	233,177	0	0	0	0	233,177	3,040,564
Recreational boating	636,055	424,036	5,529,325	0	0	212,018	2,764,662
Offshore oil and gas operations	16,710	10,733	139,958	0	0	5,977	18,656
Miscellaneous vessel classes	1,637	5	60	0	0	1,633	20,778
U.S. Navy	57,596	0	0	0	0	57,596	751,040
U.S. Coast Guard	4,317	1,452	28,786	0	0	2,864	8,941
U.S. Army	490	0	0	0	0	490	6,388
NOAA	317	7	165	88	1,146	222	2,463
Total	1,239,322	671,656	8,764,887	1,874	24,439	565,791	7,292,892

Table 6. --Continued.

Maritime sector	Post-Annex V ^b									
	Off-loaded in port				Incinerated at sea					
	Plastics (MT)	(m ³) ^a	(MT)	(m ³) ^a	Other (MT)	(m ³) ^a	Incinerated at sea (MT)	(m ³) ^a	Dumped overboard (MT)	(m ³) ^a
Merchant shipping	1,626	311,353	2,737	6,255	4,381	57,132	22,204	103,685		
Commercial passenger vessels	22,490	2,304,400	233,340	1,060,557	1,117	14,564	1,128	5,265		
Commercial fishing	15,373	1,352,768	0	0	3,723	48,542	214,081	999,660		
Recreational boating	39,848	4,975,109	554,892	2,771,045	0	0	41,315	128,964		
Offshore oil and gas operations	398	49,740	5,547	72,799	0	0	0	0		
Miscellaneous vessel classes	109	5,372	0	0	306	3,986	1,223	5,709		
U.S. Navy	3,859	2,409,124	0	0	0	0	53,737	250,929		
U.S. Coast Guard	289	126,913	765	2,604	0	0	3,262	10,183		
U.S. Army	33	9,143	0	0	0	0	199	621		
NOAA	11	331	0	0	148	1,926	158	737		
Total	84,037	11,544,253	797,282	3,913,261	9,674	126,150	337,306	1,505,752		

^aWeight-to-volume conversions reflect (1) the densities of the various types of garbage (see Table 4), (2) the composition of the vessel waste stream, and (3) the degree to which compaction equipment is used in each sector.

^bAssumes full compliance with Annex V requirements.

quantities. Both the weight and volume of garbage are indicated. Weight-to-volume conversions reflect assumptions about the types of garbage generated and the use of compaction to reduce garbage volume. This table shows aggregated sector totals only. A set of more detailed disposition tables is found in the Appendix.

The first columns of Table 6 indicate the current disposition of vessel-generated domestic garbage. The relative quantities of garbage currently brought back to shore, incinerated, or dumped overboard vary from sector to sector. A small amount of at-sea incineration occurs in portions of the merchant shipping and cruise ship sectors as well as on some National Oceanic and Atmospheric Administration (NOAA) research vessels. The percentage of domestic garbage brought back to shore for disposal is relatively high in the commercial passenger and recreational boating sector. Of the 1.2 million metric tons (MT) generated in all of the sectors, however, 566,000 MT or 45% by weight is still dumped overboard.

Under Annex V, some increased use of marine incinerators will occur, but the percentage of domestic garbage disposed of via incineration at sea will remain below 1%. All plastics, with the exception of that destroyed in incinerators, will be returned to shore for disposal. The current methodology predicts that 84,037 MT of plastics will be brought ashore for disposal. This will account for only 9.5% of all garbage brought ashore on a tonnage basis. Because of its low density, however, in volume terms plastics will represent close to 75% of the waste. Restrictions on the disposal of other types of garbage for vessels operating close to shore or in special areas will also increase the quantity of nonplastics brought ashore. Overall, the net increase in plastics and nonplastics brought ashore under Annex V will be 209,663 MT.

SUPPORTING STATISTICAL ANALYSIS OF MARITIME SECTORS SUBJECT TO COAST GUARD ENFORCEMENT OF MARPOL ANNEX V REGULATIONS

This section provides supporting data on the populations of ships covered in this study and used to generate the estimates of aggregate garbage generation shown in Table 6. Seven separate maritime sectors are identified as falling under the jurisdiction of the Coast Guard under MARPOL Annex V. These sectors are: merchant shipping, commercial fishing, commercial passenger vessels, recreational boating, offshore oil and gas operations, research and other miscellaneous vessels, and vessels operated by the U.S. Government. Each of these is profiled below in terms of the number and types of ships, onboard employment, and the frequency and duration of voyages. This information is then combined with data from earlier sections to derive per voyage, per vessel, and aggregate annual garbage quantity estimates for each of the sectors.

Merchant Shipping

Merchant vessels are those ships involved in the waterborne transport of cargo and passengers over established transoceanic, coastwise, inter-coastal, and inland water routes. Under the provisions of the Jones Act, domestic waterborne commerce (cargoes moving between U.S. ports) is

reserved exclusively for U.S. vessels. The U.S. import and export trade, however, is dominated by foreign vessels. The foreign and domestic trade sectors are discussed separately below.

Foreign Trade Vessels

According to the U.S. Maritime Administration (MARAD), the U.S. oceangoing merchant fleet numbers approximately 823 vessels of 1,000 gross tons (GT) and over. Of these, however, some 391 are inactive. Of the 432 active U.S. vessels, 122 or 28% are active in foreign trade. Another 54 vessels are active in Marine Sealift Command (MSC) operations, and will have voyage patterns comparable to foreign trade ships (see Table 7).

In addition to the vessels covered by MARAD, the Coast Guard's Marine Safety Information System (MSIS) data base shows there to be 43 vessels under 1,000 GT that are certificated for operation over open ocean routes. Thus, a total of 219 U.S.-flagged vessels operate over foreign trade routes.

The MARAD reports show that foreign-flagged vessels dominate the foreign trade sector, accounting for 95.6% of all U.S. import and export trade by tonnage (MARAD 1987b). The number of foreign vessels involved is commensurate. Data from the Coast Guard indicate that in 1987 a total of 6,751 foreign vessels, representing 110 different shipping nations, were inspected at U.S. ports (Coast Guard 1987b).

We assumed that vessels without incinerators will off-load all garbage in their final foreign port of call prior to setting sail for the United States. Under this assumption, vessels will retain on board all garbage they are prevented from dumping, and seek to off-load it upon return to the United States.

In order to estimate how much garbage is generated by these ships while en route to the United States, we examined data from the U.S. Customs Bureau's AE-975 file, Vessel Entrances and Clearances (U.S. Bureau of the Census 1987). This data base includes information on the final foreign port of call of vessels arriving at U.S. ports.

Four months of data (January, April, July, and October) were examined and used to derive annual estimates. In 1987, U.S. vessels made a total of 3,969 entrances to U.S. ports, while an estimated 33,087 entrances were made by foreign vessels. Table 8 shows a breakdown of these entrances by U.S. coastal area and foreign region of origin. Along the Atlantic and Gulf coasts, the largest number of entrances, 55 and 40%, respectively, were recorded by vessels clearing Customs from Caribbean ports. Entrances at Pacific coast ports were dominated by vessels arriving from Pacific Rim countries (46.6%) and from Pacific Canadian ports (28.8%).

A weighted average voyage length for foreign trade vessels arriving at U.S. ports was developed by calculating typical voyage lengths for each of the U.S.-foreign region pairings from Table 8. The estimated voyage lengths are based upon representative voyages from each foreign region to the U.S. coast and an assumed vessel speed of approximately 500 nmi per

Table 7.--Deployment status of U.S. flag merchant fleet, vessels 1,000 gross tons (GT) and over (U.S. Maritime Administration (MARAD), Merchant Marine Data Sheet, 1 March 1987).

Deployment status	Passenger/ combination		General cargo		Intermodel vessels		Bulk carriers		Tankers		Total	
	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt
Active vessels												
Privately owned	2	15	36	531	112	2,694	22	934	192	11,772	364	15,946
Oceangoing	--	--	26	380	69	1,703	13	684	14	806	122	3,573
U.S. foreign trade	--	--	--	--	4	72	--	--	8	571	12	643
Foreign-to-foreign	2	15	1	18	20	358	7	188	146	9,671	176	10,250
Domestic trade	--	--	--	--	1	17	5	133	97	4,062	103	4,212
Coastal	2	15	1	18	19	341	2	55	49	5,609	73	6,038
Noncontiguous												
Military Sealift												
Command charter	--	--	9	133	19	561	2	62	24	724	54	1,480
Great Lakes ^a	--	--	--	--	--	--	55	1,819	3	20	58	1,840
Government-owned	4	32	3	26	--	--	--	--	1	17	8	75
BB charter and other custody	4	32	3	26	--	--	--	--	1	17	8	75
Subtotal--active fleet	6	47	39	557	112	2,694	22	934	193	11,789	372	16,021
Inactive vessels												
Privately owned	6	59	13	164	35	1,124	39	937	33	3,040	126	5,324
Oceangoing	6	59	13	164	35	1,124	4	336	33	3,040	91	4,723
Temporarily inactive	--	--	3	43	1	33	1	63	2	131	7	270
Laid up	6	59	7	82	32	1,085	3	273	29	2,841	77	4,340
Laid up (MARAD custody)	--	--	3	39	2	6	--	--	2	68	7	113
Great Lakes	--	--	--	--	--	--	35	601	--	--	--	601

Table 7.--Continued.

Deployment status	Passenger/ combination		General cargo		Intermodel vessels		Bulk carriers		Tankers		Total	
	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt	No.	Dwt
Government-owned (MARAD)	24	183	184	2,102	28	761	--	--	21	623	257	3,669
National defense												
Reserve fleet	21	166	184	2,102	23	593	--	--	18	401	246	3,262
Ready research force (RRF)	1	9	53	661	17	438	--	--	8	141	79	1,249
Other reserve	6	57	122	1,356	6	155	--	--	9	244	143	1,812
Special programs	1	5	3	28	--	--	--	--	--	--	4	33
Nonretention	13	95	6	57	--	--	--	--	1	16	20	168
In processing for RRF	--	--	--	--	4	152	--	--	--	--	4	152
Other government-owned	3	17	--	--	1	16	--	--	3	222	7	255
Subtotal--Inactive fleet	30	242	197	2,266	63	1,885	39	937	54	3,663	383	8,993
Total--Active and inactive	36	289	236	2,823	175	4,579	61	1,871	247	15,452	755	25,014

*Includes ships normally active but laid up due to the winter freeze.

Table 8.--Entrances to U.S. ports by U.S. and foreign vessels, and estimated days at sea by U.S. coastal area, 1987 (U.S. Bureau of the Census 1987; Eastern Research Group estimates).

U.S. coastal area	Vessel origin	Estimated number of entrances			Estimated voyage length ^a (days)
		Foreign vessels	U.S. vessels	Total vessels	
Atlantic	Caribbean	5,895	954	6,849	3
	Scandinavia and N. Europe	1,497	204	1,701	9
	Canada--Atlantic	963	3	966	3
	Mediterranean	765	45	810	9
	W. coast S. America	612	27	639	9
	E. coast S. America	465	36	501	8
	Pacific Rim	282	3	285	15
	W. coast Africa	267	9	276	11
	Australasia	126	--	126	17
	Indonesia and India	123	--	123	16
	Middle East	78	6	84	12
	E. coast Africa	39	9	48	16
	Canada--Great Lakes	39	--	39	4
		Total	11,151	1,296	12,447
	Weighted average voyage length	5.7	4.6		
Gulf	Caribbean	3,264	474	3,738	3
	W. coast S. America	1,794	249	2,043	6
	Scandinavia and N. Europe	1,212	51	1,263	10
	Mediterranean	645	42	687	10
	Pacific Rim	435	9	444	13
	W. coast Africa	327	27	354	15
	E. coast S. America	267	15	282	8
	Canada--Atlantic	282	--	282	6
	Australasia	99	--	99	14
	Indonesia and India	72	3	75	13
	Middle East	63	3	66	12
	E. coast Africa	27	9	36	10
	Canada--Great Lakes	33	--	33	7
	Canada--Pacific	3	--	3	7
	Total	8,523	882	9,405	
	Weighted average voyage length	6.7	5.3		
Pacific	Pacific Rim	3,285	306	3,591	11
	Canada--Pacific	1,650	567	2,217	3
	Caribbean	708	195	903	10
	W. coast S. America	321	138	459	15

Table 8.--Continued.

U.S. coastal area	Vessel origin	Estimated number of entrances			Estimated voyage length ^a (days)
		Foreign vessels	U.S. vessels	Total vessels	
	Australasia	174	3	177	11
	Indonesia and India	150	15	165	15
	Scandinavia and N. Europe	120	3	123	15
	E. coast Africa	51	--	51	11
	Mediterranean	12	--	12	18
	Total	6,471	1,227	7,698	
	Weighted average voyage length	9.2	7.7		
Great Lakes	Canada--Great Lakes	1,716	156	1,872	1
	Scandinavia and N. Europe	96	9	105	10
	Mediterranean	27	--	27	10
	W. coast Africa	12	--	12	11
	Pacific Rim	6	--	6	16
	Middle East	3	--	3	13
	Total	1,860	165	2,025	
	Weighted average voyage length	1.7	1.5		
Noncon- tiguous areas (includes Alaska, Hawaii Puerto Rico, and Virgin Islands	Caribbean	3,891	354	4,245	(b)
	Pacific Rim	426	6	432	(b)
	W. coast S. America	195	27	222	(b)
	E. coast S. America	135	3	138	(b)
	Scandinavia and N. Europe	93	--	93	(b)
	Australasia	84	--	84	(b)
	Indonesia and India	75	6	81	(b)
	W. coast Africa	60	3	63	(b)
	Canada--Atlantic	54	--	54	(b)
	E. coast Africa	42	--	42	(b)
	Mediterranean	18	--	18	(b)
	Canada--Pacific	6	--	6	(b)
	Middle East	3	--	3	(b)
	Total	5,082	399	5,481	(b)

^aThe percentage of entrances from each vessel origin is used to derive the weighted average voyage lengths for each coastal area.

^bVoyage lengths for entrances to noncontiguous ports are estimated as follows: Hawaii--6 days (60% of entrances are from Japan), Puerto Rico and the Virgin Islands--1 day (majority of entrances are from Caribbean countries).

As indicated above, the Jones Act excludes foreign vessels from competing for U.S. domestic trade. Consequently, all domestic trade moves aboard U.S. vessels. In 1987, the fleet of U.S.-flagged domestic trade vessels included the following:

- 176 vessels of 1,000 GT and over (MARAD 1987a). Of these, 103 or 59% are active in "coastal" trade or trade between ports in the contiguous United States (see Table 7). All but six of these are tankers. The remaining 73 vessels operate in "noncontiguous" trade or trade between the contiguous U.S. states and the noncontiguous states and properties. Included in this total are 49 tankers and 19 intermodal vessels, as well as 2 U.S. cruise ships (described in the next section);
- 12 freighters and 14 tankers under 1,000 GT designated for coastwise travel (Coast Guard 1987b);
- 14 freighters and 43 tankers designated for lakes, bays, and sounds operation (Coast Guard 1987b);
- 9 freighters and 6 tankers designated for river operation (Coast Guard 1987b); and
- ca. 5,000 tug and towboats (U.S. Army Corps of Engineers 1987), which operate predominantly over the inland waterways.

Great Lakes ships may operate in either domestic or foreign (United States-Canada) trade. No breakdown is reported for these ships based on trade status, hence they are analyzed in terms of the number of ships and annual operating ratios, rather than number of entrances. (Operating ratios or utilization rates refer to the percentage of days annually on which the ship is engaged in trading activities.) In 1987, there were estimated to be 58 active Great Lakes ships of 1,000 GT or over (MARAD 1987a) and 7 of under 1,000 GT (Coast Guard 1987b).

Domestic trade vessels operate exclusively within U.S. waters, and will hence be under Coast Guard jurisdiction whenever they are operating. One exception is noncontiguous vessels which may exit U.S. waters en route from the continental United States. The approach to estimating garbage quantities in the domestic sector is, therefore, somewhat different. Whereas the annual garbage quantities generated by foreign trade vessels are estimated based upon the number of voyages, in this case it is the number of ships, the crew size, and the annual ship utilization rate which are the determinants.

Crews aboard domestic trade ships also average 20-25 men. Large oceangoing tugs carry up to 10 men, while smaller tugs and motor barges carry 6-man crews.

Domestic ships over 1,000 GT are estimated to have average voyage lengths of 5 days, while those under 1,000 GT average 4 days. Trips of large tugboats are also assumed to average 4 days, while small tugs are estimated to average 2 days at sea.

All vessels in the merchant marine sector, with the exception of Great Lakes ships, are assumed to operate with 90% utilization rates. Due to the winter freeze-up, Great Lakes ships are limited to approximately 50% utilization.

Garbage Generation Estimates

Domestic garbage.--In Table 9 below, estimates of the amount and types of garbage generated over typical voyages are shown for each of the merchant shipping categories. The table shows both weight and volume estimates, and indicates that the greatest accumulation would occur on foreign trade and large domestic trade ships. Over a 7-day voyage these ships are estimated to generate 330 kg or 2.2 m³ of garbage, of which only 22 kg is plastics. One cubic meter represents approximately 8-9 large 113.5 liter (30-gal) garbage bags.

Cargo wastes.--Table 10 presents estimates of the number of entrances to U.S. ports by U.S. and foreign general cargo vessels, and of the quantities and types of dunnage generated by such ships. In the Customs data base, dry cargo ships account for 56.1% of entrances by U.S. ships and 74.4% of entrances by foreign ships (Bureau of the Census 1987). The MARAD data indicate that 28 of the 101 dry cargo ships in the U.S. foreign trade fleet (27.7%) are general cargo-type ships (MARAD 1987a). Applying this percentage to the number of dry cargo entrances, it is estimated that U.S. and foreign break-bulk ships enter U.S. ports 617 ($0.277 \times 0.561 \times 3,969$ entrances) and 6,819 ($0.277 \times 0.744 \times 33,087$ entrances) times annually. These entrances are seen in Table 10 to generate potentially close to 20,000 m³ of waste lumber, 3,815 m³ of cardboard, and 2,981 m³ of plastic.

Commercial Passenger Vessels

The category of commercial passenger vessels encompasses all for-hire passenger-carrying vessels, including cruise ships, ferries and excursion vessels, and charter boats.

Cruise Ships

The cruise ship category includes domestic ships which operate exclusively within U.S. waters and foreign ships which sail from U.S. ports on international voyages. The Customs data base identifies approximately 80 foreign cruise ships which operate regularly out of U.S. ports. In 1986, these vessels recorded an estimated 3,324 entrances to U.S. ports (see Table 11). A high proportion of these entrances (45%) was recorded by vessels entering the Miami and Tampa port districts from the Bahama Islands. Other origin and destination combinations which account for large numbers of entrances include Canada/Alaska, Mexico/Los Angeles, Mexico/Miami, and Bermuda/New York. Puerto Rico and the Virgin Islands also receive numerous cruise ships, which arrive primarily from other Caribbean or South American ports.

Based upon the predominance of short-haul trips represented by these data, an average voyage duration of 1 day (24 h) is assumed for cruise

Table 9. -- Derivation of per voyage, per vessel, and annual domestic garbage quantities generated merchant shipping (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Merchant shipping	Domestic garbage generation per voyage ^a										Annual ship utilization rate ^b (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (m ³)
	Voyage length (days)	Crew size	Person days per voyage	Per capita generation (kg/day)	Total garbage (m ³)			Plastic garbage (kg)	Dry garbage (m ³)	Annual ship utilization rate (%)						
					(kg)	(m ³)	(m ³)									
Foreign trade																
U.S. vessels																
Atlantic/Gulf/Pacific																
7	25	165	2.0	330	2	196	2	22	1	90	50	16,425	NA	3,405	1,124	7,326
Noncontiguous/foreign																
2	25	53	2.0	105	1	62	1	7	0	90	156	16,425	NA	399	42	273
Foreign vessels																
Atlantic/Gulf/Pacific																
7	25	173	2.0	345	2	205	2	23	1	90	48	16,425	NA	26,145	9,020	58,809
Noncontiguous/Great Lakes																
2	25	60	2.0	120	1	71	1	8	1	90	137	16,425	NA	6,942	833	5,431
Noncontiguous trade (U.S.--domestic)																
7	25	175	2.0	350	2	208	2	23	1	90	47	16,425	71	NA	1,166	7,603
Great Lakes (domestic and foreign trade)																
1,000 GT and over																
2	25	53	1.5	79	1	47	0	5	0	50	87	6,844	58	NA	397	2,588
Under 1,000 GT																
2	25	53	1.5	79	1	47	0	5	0	50	87	6,844	7	NA	48	312
United States																
Military Sealift charter																
7	25	175	2.0	350	2	208	2	23	1	90	47	16,425	54	NA	887	5,783
Temporarily inactive vessels																
7	25	175	2.0	350	2	208	2	23	1	90	47	16,425	7	NA	115	750
Coastal shipping																
Ships																
1,000 GT and over																
5	25	125	1.5	188	1	111	1	13	1	90	66	12,319	103	NA	1,269	8,273
Under 1,000 GT																
4	25	100	1.5	150	1	89	1	10	1	90	82	12,319	98	NA	1,207	7,871
Tow/tugboats																
Large (inspected)																
4	10	40	1.5	60	0	36	0	4	0	90	82	4,928	12	NA	59	386
Small																
2	6	12	1.5	18	0	11	0	1	0	90	164	2,957	5,000	NA	14,783	96,380
Total garbage per year															30,949	201,785

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage^c and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Table 10.--Estimates of annual quantities of cargo waste (dunnage) currently dumped and quantities dumped under MARPOL Annex V (Eastern Research Group estimates).

Basis of estimate	U.S. vessels	Foreign vessels
Number of entrances to U.S. ports per year	3,969	33,087
Dry cargo as percentage of all entrances	56.1%	74.4%
Dry cargo entrances	2,227	24,617
General cargo as percentage of all dry cargo ships	27.7%	27.7%
General cargo entrances	617	6,819
Dunnage generated per clearance from U.S. port		
Lumber		
Quantity (m ³)	48.6	48.6
Percent of entrances	66.7%	66.7%
Cardboard		
Quantity (m ³)	23.6	23.6
Percent of entrances	33.3%	33.3%
Plastic		
Quantity (m ³)	0.12	0.12
Percent of entrances	10.0%	10.0%
Total dunnage quantities generated per year		
Lumber (m ³)	19,983.4	220,930.2
Cardboard (m ³)	130,979	130,979
Plastic (m ³)	7.20	7.20
Incidence of dumping	50.0%	50.0%
Total dunnage quantities dumped in U.S. waters per year		
Current practice		
Lumber (m ³)	9,991.7	110,465.1
Cardboard (m ³)	1,768,214	1,768,214
Plastic (m ³)	3.602	3.602
Under MARPOL Annex V		
Lumber (m ³)	9,991.7	110,465.1
Cardboard (m ³)	1,768,214	1,768,214
Plastic (m ³)	0.00	0.00

Table 11.--Cruise ships entering U.S. ports (Bureau of the Census 1987; Eastern Research Group estimates).

Vessel origin	U.S. port of entrance	Estimated number of entrances (1987)	Percent of total
Bahamas	Miami	1,232	37.1
Bahamas	Tampa	276	8.3
Canada (Pacific coast)	Anchorage	244	7.3
Mexico (Pacific coast)	Los Angeles	228	6.9
Mexico (Gulf coast)	Miami	168	5.1
Bermuda	New York	152	4.6
French West Indies	Virgin Islands	116	3.5
Leeward/Windward Islands	Virgin Islands	84	2.5
Netherlands Antilles	Virgin Islands	80	2.4
French West Indies	San Juan, Puerto Rico	72	2.2
Netherland Antilles	Miami	72	2.2
Haiti	Miami	56	1.7
Netherland Antilles	San Juan, Puerto Rico	56	1.7
Bahamas	San Juan, Puerto Rico	52	1.6
Haiti	San Juan, Puerto Rico	44	1.3
Jamaica	Miami	44	1.3
Dominican Republic	Virgin Islands	40	1.2
Dominican Republic	San Juan, Puerto Rico	32	1.0
Venezuela	San Juan, Puerto Rico	32	1.0
All other origins	All other destinations	244	7.3
Total		3,324	100.0

ships arriving in the United States. While examples of much longer voyages may be found within the data, short voyages are much more typical.

Foreign cruise ships entered U.S. ports with an average passenger complement of 786. Crew-to-passenger ratios are approximately 1:2 (J. Ruers, International Committee of Passenger Liners, pers. commun. 1988), hence an average of approximately 1,000 persons are assumed to be on board such ships.

Coast Guard data indicate that approximately two dozen U.S.-flagged vessels are used in domestic cruise operations (L. Stanton, Coast Guard, pers. commun. 1988). These include two large vessels of over 1,000 GT which operate in the Hawaiian interisland trade as well as several smaller vessels active on coastal routes along both the east and west coasts. Average time between ports is estimated at 1 day, as the vessels are usually in port each night. Such vessels are estimated to carry an average of 200 passengers and crew members (E. Scharfe, Director, Small Passenger Vessel Association, pers. commun. 1988) during typical cruises.

Other Passenger-Carrying Vessels

Additional categories of passenger-carrying vessels include ferries and charter fishing and pleasure vessels, of which there are a large number. In 1987, the Coast Guard's MSIS data base contained some 49 U.S.-flagged passenger vessels of 1,000 GT and over, and 4,774 vessels under 1,000 GT.

Among the larger passenger vessels, four are ocean-designated and include the two Hawaiian cruise ships discussed above as well as two converted hospital ships that are part of the MSC. These are covered in the merchant vessel data. Ten larger passenger vessels operate with river designations (e.g., Mississippi River cruises), while the remaining 34 are designated for operation in lakes, bays, and sounds. These vessels offer ferrying services and excursion or sightseeing cruises of short duration. Thus, a total of 44 additional large ferries and riverboats operate domestically. They are assumed to carry up to 1,000 passengers on voyages averaging 1 day in duration.

Approximately 75% of the 4,774 passenger-carrying vessels under 1,000 GT are charter fishing boats, with ferries, yachts, and other small boats accounting for the remaining 25% (Stanton pers. commun. 1988). Charter fishing boats are assumed to carry an average of 20 persons, while ferries and other commercial passenger vessels are assumed to carry 200 people. Voyage lengths of 1 day or less are assumed for all vessels in this category.

Large cruise ships generate substantial quantities of garbage even on overnight voyages. Table 12 indicates that 1,000 passengers on a luxury cruise will generate over 2 MT of garbage each day. Smaller ships carrying 200 passengers may generate close to 500 kg per day.

Commercial Fishing

United States Vessels

Fishing vessels may be classified according to whether they operate in onshore, offshore, or inland fisheries. Onshore fishing, defined as fishing which takes place within 12 nmi from shore, is conducted by smaller boats making primarily day-long trips. Data sources distinguish between fishing boats, which are under 5 net tons in size, and fishing vessels, which include all craft of 5 net tons or more (see Table 13).

Boats under 5 net tons generally do not exceed 7.6 m (25 ft) in length (T. Willis, Coast Guard Documentation Branch, pers. commun. 1988), and are not eligible for Coast Guard documentation. Normally, therefore, they do not operate at significant distances from shore. For convenience, all fishing boats (i.e., <5 net tons) are assumed to operate in the onshore fisheries. The NMFS estimates there to be approximately 105,500 boats active in the U.S. fisheries (NMFS 1987). These are assumed to carry an average of three crew members, and to return to port each night.

Table 12.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated commercial passenger vessels (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Commercial passenger ships	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a			Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels	No. of entrances	Total Garbage per year ^b (MT)				
					Total garbage (kg)	Dry garbage (kg)	Plastic garbage (kg)										
					(m ³)	(m ³)	(m ³)										
Cruise ships																	
U.S. vessels																	
>1,000 GT	1	1,000	1,000	2.4	2,400	16	1,426	14	161	10	90	329	788,400	2	NA	1,577	10,281
Under 1,000 GT	1	200	200	2.4	480	3	285	3	32	2	90	329	157,680	24	NA	3,784	24,673
Foreign vessels	1	1,000	1,000	2.4	2,400	16	1,426	14	161	10	90	NA	NA	NA	3,324	7,978	52,013
Excursion vessels	1	200	200	2.4	480	2	285	3	32	2	90	329	157,680	1,194	NA	188,270	1,227,495
Charter boats	1	20	20	2.4	48	0	29	0	3	0	90	329	15,768	3,581	NA	56,465	368,148
Total garbage per year												258,074	1,682,608				

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Table 13.--Employment and craft
in the U.S. commercial fisheries
([U.S.] National Marine Fisheries
Service 1987).

Size	Number
Vessel >5 tons	24,300
Motor boats	104,000
Other boats	1,500
Total craft	129,800
Fishermen	238,800

While some larger craft also operate close to shore, fishing vessels (5 net tons and over) are assumed to operate beyond 12 nmi from shore. These vessels are capable of longer voyages, and are frequently equipped with sophisticated navigational and fish locating equipment. They also have greater onboard storage and processing capacity.

The NMFS estimates that in 1986 there were 24,300 fishing vessels of 5 net tons or more in the United States. While these may range up to 1,000 GT and over in size, relatively few are this large. Table 14 indicates that over 60% of fishing vessels are both smaller than 25 net tons in size, and <15.2 m (50 ft) in length.

Inland fishing covers commercial activity taking place on the inland waterways. At present, small commercial fisheries operate on the Great Lakes and along the Mississippi River (S. Koplín, NMFS, Statistics Branch, pers. commun. 1988), and account for only a small percentage of the national catch. States bordering the Great Lakes, for example, accounted for only 1.7% of the 1987 U.S. commercial catch (NMFS 1987). As boats active in the inland fisheries will be contained within the data presented above, the craft involved will be assumed to operate in a fashion similar to those in the saltwater fisheries. Assumptions regarding crew sizes and voyage lengths of fishing vessels are shown in Table 15, which derives the per voyage, per vessel, and aggregate annual garbage quantities.

Foreign Fishing Vessels

Foreign fishing vessels granted access to fishing stocks within the U.S. exclusive economic zone (EEZ) will also be expected to comply with MARPOL Annex V. While some restrictions on vessel discharges already apply, the requirements do not address specifically the problem of garbage dumping.

In the recent past, foreign fishing activity in U.S. waters has centered around the eastern Bering Sea and Aleutian Islands areas, where

Table 14.--Documented U.S. fishing vessels,^a by length and gross tonnage (U.S. Coast Guard, Marine Safety Information System 20 April 1986; [U.S.] National Marine Fisheries Service 1987).

Gross tonnage	Vessel length				Total
	<15.2 m (<50 ft)	15.2-19.8 m (50-65 ft)	19.8-24.1 m (65-79 ft)	>24.1 m (>79 ft)	
Less than 25	14,703	112	2	2	14,815
25-49	2,774	1,152	33	--	3,959
50-99	340	1,511	1,107	45	3,003
100-199	18	117	1,418	674	2,227
200-299	--	--	--	69	69
300-399	--	--	--	32	32
400-499	--	--	--	49	49
500-599	--	--	--	45	45
600-699	--	--	--	15	15
700-799	--	--	--	10	10
800-899	--	--	--	10	10
900-999	--	--	--	23	23
1,000-1,999	--	--	--	34	34
2,000-2,999	--	--	--	2	2
3,000-3,999	--	--	--	2	2
4,000-4,999	--	--	--	2	2
More than 5,000	--	--	--	--	--
Total	17,835	2,891	2,560	1,015	24,300

^aVessels are defined as craft of 5 net tons or over.

the most significant target species has been Alaskan pollock. The country most active in this fishery is Japan. Other fisheries with considerable foreign participation include the Pacific whiting and Atlantic mackerel fisheries.

Direct access to U.S. fishing stocks by foreign vessels has been cut back considerably in recent years. At present, foreign access is obtained primarily through joint venture permits (J. Kelley, NMFS, Office of Fishery Conservation and Management, pers. commun. 1989). Under joint venture agreements, U.S. vessels deliver their catch to large foreign motherships or other factory trawlers, which process the fish at sea.

Data on the number of foreign fishing vessel permits issued in 1987, by type of vessel, flag of vessel, and fishery, were requested from the NMFS, but were not available in time for this report. In general, though, activity by foreign fishing vessels within U.S. waters has been decreasing in recent years with the "Americanization" of the U.S. EEZ. Direct fishing

Table 15.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: commercial fishing (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Commercial fishing length (days)	Crew size	Voyage length (days)	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a					Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (m ³)	
					Total garbage (kg)	Dry garbage (kg)	Plastic garbage (kg)	Total garbage (MT)								
					(m ³)	(m ³)	(m ³)									
Undocumented	1	3	3	1.5	5	0	3	0	0	66	241	1,084	105,500	NA	114,367	745,660
Documented	7	7	49	2.0	98	1	58	1	7	66	34	3,373	14,815	NA	49,965	325,766
5-25 GT	15	15	225	2.0	450	3	267	3	30	66	16	7,227	9,258	NA	66,908	436,229
25-300 GT	15	15	225	2.0	450	3	267	3	30	66	16	7,227	188	NA	1,359	8,858
300-1,000 GT	30	30	900	2.0	1,800	12	1,069	11	121	66	8	14,454	40	NA	578	3,770
Over 1,000 GT	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0
Foreign vessels ^b																
Total garbage per year															233,177	1,520,282

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

by foreign vessels has been almost completely phased out, while joint ventures between U.S. catcher vessels and foreign processing vessels are declining. More and more, foreign access to U.S. fishery products will be in the form of exported products processed on U.S. soil. According to a report to the Alaska Department of Environmental Conservation, "it is generally assumed that there will be little, if any, joint venture activity in the North Pacific EEZ by 1991" (Pacific Associates 1988).

Garbage Generation Estimates

Domestic garbage.--Table 15 shows the derivation of the per voyage and annual domestic waste estimates. The largest ships may generate up to 1,800 kg of garbage overall per voyage. Of this amount, however, they would likely have to retain only the plastics. Small fishing boats are estimated to generate only 4.5 kg of total garbage per day at sea.

Commercial wastes.--As indicated, it is assumed that most fishing craft will generate an additional 0.028 m³ (1.0 ft³) of plastic gear waste per week (0.004 m³ (0.14 ft³) per day). Longliners and boats in the herring fisheries are assumed to generate twice this amount. Such vessels are assumed to represent 5% of all vessels in the 5-25 and 25-300 GT categories. Table 16 shows the estimated quantities of fishing wastes generated annually.

Recreational Boating

All recreational boats operating over the navigable waters of the United States are also required to comply with Annex V. Potentially, therefore, most of the approximately 14 million recreational boats in the United States might be included in an analysis of Annex V. For this study, we limit the analysis to numbered boats in coastal states or in states bordering the Great Lakes. Still, some 7.3 million recreational boats fit this criterion (see Table 17).

The majority of recreational boats are used on inland waters or, when used in the ocean, within 3 nmi from shore. When operating in these waters, boaters are prohibited from disposing of any garbage overboard. Beyond 3 nmi from shore, limited dumping may occur.

In order to identify those boats prohibited from any overboard disposal, several assumptions were made. First, only boats registered in coastal states are assumed to operate in the ocean. Secondly, only larger boats are assumed to operate beyond 3 nmi from shore. Within coastal and Great Lakes states, the size breakdown of the registered boating fleet is as follows:

- 56.3% are under 4.9 m (16 ft) long,
- 39.6% are between 4.9 and 7.9 m (16 and 26 ft) in length, and
- 3.7% are greater than 7.9 m (26 ft) in length (see Table 17).

Table 16.--Estimates of annual quantities of plastic fishing gear wastes generated in the U.S. fisheries (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Vessel category	Number of vessels	Annual quantities of fishing waste generated			
		Vessels generating normal quantities ^a		Vessels generating additional quantities ^b	
		(MT)	(m ³)	(MT)	(m ³)
Undocumented	105,500	1,779	131,781	0	0
Documented					
5-25 GT	14,815	216	12,570	23	1,323
25-300 GT	9,258	135	7,855	14	827
300-1,000 GT	188	3	160	0	0
Over 1,000 GT	40	1	34	0	0
Foreign vessels	NA	NA	NA	NA	NA
Total	129,801	2,133	152,400	37	2,150

^aVessels using trawls, set nets, or pots. Plastic waste in these fisheries is essentially gear-related.

^bVessels active in bait fisheries (i.e., longlining) or herring fisheries which generate additional quantities of plastic waste in the form of bait wrappings or salt bags.

According to the Boat Owner's Association of the U.S. (BOATUS), recreational boats under 4.9 m (16 ft) in length "are most likely confined to inland lakes, rivers, and bays," and of those over 4.9 m (16 ft), only 10% are estimated to venture beyond 3 nmi from shore (Schwartz 1987). Based on this, approximately 219,000 boats are estimated to operate in areas where some overboard disposal of garbage is permitted. The remaining 13.1 million operate in areas where no garbage disposal may occur.

Garbage Generation

Voyage lengths and onboard complements for recreational boats of various sizes are shown in Table 18, which derives the per voyage and annual garbage quantities generated.

Offshore Oil and Gas Operations

Offshore oil and gas operations such as exploratory drilling, development drilling, and oil and gas production from offshore platforms are also covered by MARPOL Annex V. The restrictions which apply to such operations are different from those applicable to commercial and recreational vessels. Under Annex V, ocean disposal of all types of garbage, with the exception of ground food wastes, is prohibited. For operations located within 12 nmi from shore, even the disposal of ground food wastes is prohibited.

Table 17.--Recreational boats in coastal and Great Lakes states
(U.S. Coast Guard 1987a).

Region	Class and size					Total
	Class A <4.9 m (<16 ft)	Class 1 4.9-7.9 m (16-26 ft)	Class 2 7.9-12.2 m (26-40 ft)	Class 3 12.2-19.8 m (40-65 ft)	Class 4 >19.8 m (>65 ft)	
Coastal states						
Number	2,548,709	1,955,105	212,458	22,754	1,806	4,740,832
Percent of total	53.8	41.2	4.5	0.5	0.0	100.0
Great Lakes states						
Number	1,540,340	916,301	55,846	4,540	232	2,517,259
Percent of total	60.7	37.2	1.9	0.3	0.0	100.0
Coastal and Great Lakes						
Number	4,089,049	2,871,406	268,304	27,294	2,038	7,258,091
Percent of total	56.3	39.6	3.7	0.4	0.0	100.0

Table 18.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: recreational boats (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Recreational boats	Domestic garbage generation per voyage ^a											Total garbage per year ^b (m ³)					
	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Total garbage (kg)			Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels		No. of entrances				
					Dry garbage (kg)	Dry garbage (m ³)	Plastic garbage (kg)										
Coastal states																	
Under 4.9 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	2,548,709	NA	223,267	1,455,671
4.9-7.9 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	1,955,105	NA	171,267	1,116,640
7.9-12.2 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	212,458	NA	18,611	121,343
12.2-19.8 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	22,754	NA	1,993	12,996
Over 19.8 m	2	6	12	1.5	18	0	11	0	1	0	0	11	197	1,806	NA	356	2,321
Subtotal																415,495	2,708,971
Great Lakes states																	
Under 4.9 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	1,540,340	NA	134,934	879,751
4.9-7.9 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	916,301	NA	80,268	523,337
7.9-12.2 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	56,105	NA	4,915	32,044
12.2-19.8 m	1	4	4	1.0	4	0	2	0	0	0	0	22	88	4,540	NA	398	2,593
Over 19.8 m	2	6	12	1.5	18	0	11	0	1	0	0	11	97	232	NA	46	298
Subtotal																220,560	1,438,022
Total garbage per year													636,055	4,146,994			

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Mobile Offshore Drilling Units

Data from the Department of the Interior's Minerals Management Service (MMS) for February 1988 showed there to be 124 mobile offshore drilling units (MODU's) active in U.S. Federal waters (L. M. Tracey, Department of the Interior, Minerals Management Services, pers. commun. 1988). All but one of these were reported to be operating beyond 12 nmi from shore. Approximately 78 MODU's were active in state waters (J. Dees, Ocean & Oil Weekly, Houston, TX, pers. commun. 1988). State waters extend out to 3 nmi from shore, except off Florida and Texas, where the state-federal boundary occurs at 3 leagues or approximately 10.35 nmi. All activity in state waters is subject to the complete ban on disposal within 12 nmi of shore, while MODU's in Federal waters would be able to dispose of ground food wastes. The MMS data indicate that MODU's in Federal waters have an average of 40 beds. This figure has been used as an estimate of the number of men aboard MODU's on a 24-h basis. Active MODU's are assumed to operate at 100% utilization.

Platforms

Approximately 3,500 production platform "complexes," consisting of one or more platforms in a single location, actively operate in U.S. Federal waters. Of these, however, only 779 are manned. A total of 124 manned platforms are situated within 12 nmi from shore, while the remaining 655 are located beyond 12 nmi. Dees (pers. commun. 1988) estimates that a maximum of 40 additional manned platforms are active in state waters.

The MMS data indicate that platform complexes have an average of 15 beds each.

Offshore Service Vessels

Service vessels employed in petroleum support activities are also covered by Annex V prohibitions. This category includes supply ships, tugs, anchor-handling vessels, crew ships, and research and survey vessels. Coast Guard data indicate that there are 484 offshore service vessels (OSV's) operating in the Federal Outer Continental Shelf region. Most crew and supply ships fall in the 50-200 dwt range. These are assumed to carry crews of five persons, and to make trips lasting an average of 1 day.

No data are available to indicate how many OSV's operate in state waters. Assuming the same ratio of structures (MODU's and platforms) to OSV's exists in state waters as in Federal waters, it is estimated that there are 63 additional OSV's active in state waters $(484 + 903) \times 118$.

Garbage Generation

Garbage quantities for the offshore oil and gas sector are calculated in Table 19 on a per day, rather than a per voyage basis, since the structures are stationary and relatively permanent. Currently, all garbage with the exception of food wastes is required by the MMS to be transported to shore for disposal.

Table 19.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: offshore oil and gas (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

Offshore oil and gas operations	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a				Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (MT)	Total garbage per year ^b (m ³)				
					Total garbage (kg)	Dry garbage (kg)	Plastic garbage (kg)	Plastic garbage (m ³)											
					(m ³)	(m ³)	(m ³)	(m ³)											
Mobile offshore drilling units																			
Within 12 nmi	1	40	40	2.0	80	1	48	0	5	0	0	0	365	29,200	74	NA	2,161	14,088	
Outside 12 nmi	1	40	40	2.0	80	1	48	0	5	0	0	0	365	29,200	123	NA	3,592	23,417	
Offshore oil and gas production platforms																			
Within 12 nmi	1	15	15	2.0	30	0	18	0	2	0	0	0	365	10,950	655	NA	7,172	46,762	
Outside 12 nmi	1	15	15	2.0	30	0	18	0	2	0	0	0	365	10,950	655	NA	7,172	46,762	
Offshore service vessels	1	5	5	2.0	10	0	6	0	1	0	0	0	365	3,650	545	NA	1,989	12,970	
Total garbage per year																		15,710	108,945

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Research and Other Miscellaneous Vessels

Several categories of miscellaneous vessels have also been included in this analysis. These include vessels operated by universities and other oceanographic research institutions, maritime academy training ships, and various "industrial" vessels such as dredges and cable-laying ships.

Research Vessels

Numerous universities as well as private and nonprofit groups (e.g., Greenpeace, the Cousteau Foundation), operate oceanographic research vessels. The Coast Guard's MSIS data base indicates that in 1987 there were 26 vessels actively involved in oceanographic research (Coast Guard 1987b).

A 1978 profile of the world's oceanographic research fleet indicated that a typical research cruise might involve 20-25 crew members and 10-20 scientists (Trillo 1978; cited in Parker et al. 1987). These estimates were deemed appropriate by individuals connected with two major oceanographic research institutes, the Woods Hole Oceanographic Institute (J. Colburn, Woods Hole Oceanographic Institute, pers. commun. 1988) and the Scripps Institution of Oceanography (G. Schorr, Associate Director, Scripps Institution of Oceanography, La Jolla, CA, pers. commun. 1987).

School Training Vessels

Seven maritime academies in the United States operate a total of 14 ships used for training (Coast Guard 1987b). Seven of these are ocean-designated, six are authorized for coastwise travel, and one carries a Great Lakes designation. Only five of the vessels are greater than 1,000 GT in size.

Training ships of 1,000 GT or over are estimated to carry 150 men, while those under 1,000 GT are estimated to carry a crew of 50. Voyage lengths are estimated at 15 and 7 days, respectively. These estimates are based on discussions with officials at the Massachusetts Maritime Academy, who are familiar with the sizes and operations of vessels used at their and other maritime academies (D. Kan, Massachusetts Maritime Academy, Buzzard's Bay, MA, pers. commun. 1987).

Industrial Vessels

The category of industrial vessels comprises an assortment of vessel types including dredges, cable-laying ships, and drilling ships. Their common characteristic is that they carry crews who perform functions other than operating the vessel. The Coast Guard's MSIS data base indicates that in 1987 there were a total of 85 such vessels. Of these, 57 were greater than 1,000 GT, while 22 were under 1,000 GT. Furthermore, 69 were ocean-designated, while 17 were designated for coastal operation only.

While it is difficult to generalize about these vessels as a group, voyage lengths and crew complements on board have been approximated.

Oceangoing industrial vessels of 1,000 GT or over are estimated to carry an average of 30 persons on board and have voyage lengths averaging 15 days. Coastal vessels of 1,000 GT are also assumed to carry crews of 30 men, but are at sea for an average of 7 days. Both oceangoing and coastal vessels under 1,000 GT are estimated to carry 15 persons and to operate over 7-day voyages.

Garbage Generation

Domestic garbage.--Estimates of garbage generation in these sectors are shown in Table 20. School training ships and research vessels over 1,000 GT generate substantial quantities of garbage and plastics. Large research vessels, for example, may generate over 10 m³ of plastics from domestic sources alone. This would be sufficient to fill an average commercial garbage dumpster.

Research vessel wastes.--The additional quantities of plastics associated with oceanographic research wastes are derived in Table 21.

U.S. Navy

Data from the Jane's Fighting Ships (1986) indicate that the U.S. Navy fleet currently numbers approximately 679 active vessels (see Table 22). Normal operational cycles for Navy vessels involve one 6-month tour of duty outside of U.S. waters every 18 months (D. Steigman, Jane's Publishing Co., pers. commun. 1988). Consequently, at any given time approximately one-third of the Navy fleet is operating outside of U.S. waters.

Crew complements on board Navy vessels range from 25 men up to as many as 5,000 on board the largest aircraft carriers. Where a range of crew sizes was reported, crew complements shown in Table 22 represent the average. Utilization factors while in U.S. waters range from 20 to 75%, depending on the vessel's strategic importance and its re-fit cycle (Mullenhard, pers. commun. 1988). Steigman (pers. commun. 1988) provided separate estimates of operating ratios for each class of Navy vessel, which are used to derive garbage quantity estimates for these ships while in U.S. waters.

Garbage Generation

Because of the large crew sizes and extended periods at sea, several categories of Navy ships are seen in Table 23 to generate extremely large quantities of wastes. Aircraft carriers with 5,000 men aboard, for example, could generate as much as 200 MT of garbage over a 20-day cruise. Several other categories of ships may generate 10 to 20 MT as well. Clearly, the Navy has particular garbage handling problems.

U.S. Coast Guard

The Coast Guard operates a large fleet of vessels, ranging from small harbor patrol boats to a pair of 121.9-m (400-ft) icebreakers. Table 24 provides a summary of the Coast Guard's fleet and indicates the number of

Table 20.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: miscellaneous vessel categories (Eastern Research Group estimates)
(GT = gross tons, MT = metric tons).

Miscellaneous vessels	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a				Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel (kg)	No. of vessels	No. of entrances	Total Garbage per year (MT)	Total Garbage per year (m ³)			
					Total garbage (kg)	Dry garbage (kg)	Plastic garbage (kg)	Plastic garbage (m ³)										
					(m ³)	(m ³)	(m ³)	(m ³)										
School training	15	150	2,250	2.0	4,500	29	2,673	26	302	19	35	9	38,325	5	NA	192	1,249	
1,000 GT and over																		
Under 1,000 GT	7	50	350	2.0	700	5	416	4	47	3	35	18	12,775	2	NA	26	167	
Ocean	7	50	350	1.5	525	3	312	3	35	2	35	18	9,581	5	NA	48	312	
Coastal																		
Industrial vessels																		
1,000 GT and over																		
Ocean	15	30	450	2.0	900	6	535	5	60	4	75	18	16,425	52	NA	854	5,569	
Coastal	7	30	210	1.5	315	2	187	2	21	1	75	39	12,319	11	NA	136	883	
Under 1,000 GT																		
Ocean	7	15	105	2.0	210	1	125	1	14	1	75	39	8,213	17	NA	140	910	
Coastal	7	15	105	1.5	158	1	94	1	11	1	75	39	6,159	5	NA	31	201	
Research vessels																		
Inspected																		
1,000 GT and over	25	50	1,250	2.0	2,500	16	1,485	15	168	10	35	5	12,775	2	NA	26	167	
300-1,000 GT	15	50	750	1.5	1,125	7	668	7	75	5	35	9	9,581	15	NA	144	937	
Uninspected																		
Under 300 GT	10	25	250	1.5	375	2	223	2	25	2	35	13	4,791	9	NA	43	281	
Total garbage per year																	1,637	10,676

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.
^bTotal garbage weight per year is equal to the product of either (1) per vessel population, or (2) garbage quantity per voyage and the number of entrances.
^cRefers to the percentage of days annually operating in U.S. waters.

Table 21.--Estimates of annual quantities of additional plastic wastes generated by oceanographic research vessels (Eastern Research Group estimates) (GT = gross tons).

Vessel category	Number of vessels	Voyage length (days)	Voyages per year ^a	Additional waste per year (m ³)
Private vessels				
1,000 GT and over	2	25	5	1.17
300-1,000 GT	15	10	13	22.78
Under 300 GT	9	10	13	13.67
National Oceanic and Atmospheric Administration vessels				
Large deepwater vessels	4	25	5	2.34
Small coastal vessels	20	10	13	30.37
Total	50			70.32

^aAnnual vessel utilization of 35% is assumed.

vessels in each class, the crew complement, and typical voyage durations. This table is based upon discussions with Coast Guard operations personnel.

Coast Guard vessels are assumed to operate entirely within U.S. waters. Utilization factors for Coast Guard vessels are similar to those of Navy ships, and are assumed to average 50%.

Garbage Generation

Several categories of Coast Guard cutters as well as the large polar icebreakers are estimated to generate substantial quantities of garbage over representative voyages. The relevant quantities are shown in Table 25.

U.S. Army

The U.S. Army reports a fleet of approximately 580 crafts (G. Danish, U.S. Army, pers. commun. 1988). Of these, only a small number are "sea deployable." As shown in Table 26, these include four logistic support vessels approximately 91.4 m (300 ft) in length, 35 utility class landing craft capable of extended trips at sea, and 10 large oceangoing tugs.

The rest of the Army's fleet is made up of approximately 490 "ship-to-beach" craft of various types, used mainly for shuttling troops and supplies to and from larger vessels anchored offshore. In addition, the

Table 22.--U.S. Navy vessels by type and status (Jane's Fighting Ships 1986; Navy League of the United States 1987).

Vessel type	Active	Building/ reactivating conversion	Approximate onboard complement	^a Estimated manpower total
Strategic missile submarines	38	5	150	5,700
Attack submarines	101	15	140	14,140
Aircraft carriers	13	3	5,000	65,000
Battleships	2	2	1,500	3,000
Cruisers	31	13	500	15,500
Destroyers	68	1	350	23,800
Frigates	100	4	300	30,000
Light forces	7	0	25	175
Light amphibious warfare ships	57	7	700-2,800	99,750
Mine warfare ships	3	6	70	210
Auxiliary ships	79	3	100-1,000	35,550
Military Sealift Command	72	18	25-120	5,220
Ready reserve force	73	0	40-1,200	45,260
Naval reserve	35	0	NA	NA
Total	679	77		343,305

^aWhere crew complements vary within a class, the arithmetic mean of the range is used. Total estimated complement is derived by dividing average complement by the number of active vessels.

Army maintains 15 small harbor tugs and about 25 small outboard motor-powered J boats.

These craft are used only intermittently during peacetime in logistics exercises. A utilization rate of 35% is assumed for all vessels.

Garbage Generation

The largest Army ships, the logistic support vessels, carrying 40 persons on board for up to 30 days, may generate close to 2 MT of garbage and 10 m³ of plastics alone. Other vessel classes generate considerably smaller quantities of garbage.

National Oceanic and Atmospheric Administration Research Vessels

The NOAA operates a fleet of approximately two dozen vessels which are engaged in atmospheric and oceanographic research (B. Cunningham, Office of NOAA Corps, NOAA, pers. commun. 1988). These vessels range in size from 250 to 4,000 GT. Smaller vessels are estimated to carry approximately 10 persons on board, and to remain at sea for periods of approximately 1 week.

Table 23.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: U.S. Navy vessels (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

U.S. Navy vessels	Voyage length (days)	Crew size	Person days per voyage	Per capita generation (kg/day)	Domestic garbage generation per voyage ^a				Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel per year (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (MT)	
					Total garbage (kg)	Dry garbage (m ³)	Plastic garbage (kg)	Plastic garbage (m ³)							
															Per capita generation (kg/day)
Strategic missile submarines	3	150	450	2.0	900	6	535	5	60	4	3	38	NA	113	737
Attack submarines	7	140	980	2.0	1,960	13	1,164	12	131	8	6	101	NA	669	4,362
Aircraft carriers	20	5,000	100,000	2.0	200,000	1,304	118,800	1,177	13,400	837	22	13	NA	10,544	68,748
Battleships	20	1,500	30,000	2.0	60,000	391	35,640	353	4,020	251	33	2	NA	487	3,173
Cruisers	20	500	10,000	2.0	20,000	130	11,880	118	1,340	84	22	31	NA	2,514	16,394
Destroyers	20	350	7,000	2.0	14,000	91	8,316	82	938	59	22	68	NA	3,861	25,172
Frigates	20	300	6,000	2.0	12,000	78	7,128	71	804	50	22	100	NA	4,867	31,730
Light forces	15	25	375	2.0	750	5	446	4	50	3	22	7	NA	28	185
Light amphibious warfare ships	5	1,750	8,750	2.0	17,500	114	10,395	103	1,173	73	22	16	NA	16,182	105,502
Mine warfare ships	15	70	1,050	2.0	2,100	14	1,247	12	141	9	22	3	NA	34	222
Auxiliary ships	15	500	7,500	2.0	15,000	98	8,910	88	1,005	63	22	79	NA	6,408	41,778
Military Sealift Command	15	75	1,125	2.0	2,250	15	1,337	13	151	9	22	5	NA	876	5,711
Ready reserve force	15	620	9,300	2.0	18,600	121	11,048	109	1,246	78	33	73	NA	11,013	71,805
Naval reserve	15	620	9,300	2.0	18,600	121	11,048	109	1,246	78	0	35	NA	0	0
Total garbage per year														57,596	375,520

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.
^bTotal garbage weight per year is equal to the product of either (1) per vessel population, or (2) garbage quantity per voyage and the number of entrances.
^cRefers to the percentage of days annually operating in U.S. waters.

Table 24.--U.S. Coast Guard fleet by type and status (Jane's Fighting Ships 1986; Navy League of the United States 1987).

Vessel type	Active	Reserve	Under construction	Approximate complement
Cutters, high endurance	15	--	--	171
Cutters, medium endurance	34	--	7	82
Icebreakers	6	--	--	161
Icebreaking tugs	8	--	1	17
Surface effect craft	3	--	--	18
Large patrol craft	83	--	8	11
Training cutter	1	--	--	245
Buoy tenders, seagoing	28	--	--	53
Buoy tenders, coastal	12	--	--	20
Buoy tenders, inland	6	--	--	20
Buoy tenders, river	18	--	--	20
Construction tenders, inland	17	--	--	20
Harbor tugs, medium	4	--	--	10
Harbor tugs, small	14	--	--	10
Total	249	--	16	--

Larger vessels make voyages of up to 1 month and typically carry some 20 officers, 55 to 60 crewmen, and up to 30 scientists.

Approximately half of these vessels operate out of the NOAA base in Seattle, while the other half are stationed in Newport News. Other bases maintained by NOAA include Woods Hole, Miami, Pascagoula, and San Diego, as well as one each in Alaska and Hawaii.

Garbage Generation

The largest NOAA ships may generate as much as 4 MT of garbage over a typical 20-day voyage, and close to 20 m³ of plastics from domestic sources alone, as shown in Table 27.

Wastes associated with the research activities of these ships are derived along with those of private research vessels in Table 21.

Other Government Vessels

Other Federal Government agencies such as the Customs Bureau and the U.S. Army Corps of Engineers, as well as numerous state and local government departments and agencies, may operate modest fleets of boats. No large craft, however, are estimated to be operated by agencies other than those discussed above. Smaller boats are included in the data presented in the section on recreational boats, but are not separately analyzed here.

Table 25.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated:
U.S. Coast Guard vessels (Eastern Research Group estimates) (GT = gross tons, MT = metric tons).

U.S. Coast Guard vessels	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a				Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel per year (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (MT)	
					Total garbage (kg)	Dry garbage (kg)	Plastic garbage (kg)	(m ³)							
					(m ³)	(m ³)	(m ³)	(m ³)							
Icebreakers															
Polar class 121.9 m	90	140	12,600	2.0	25,200	164	14,969	148	1,688	105	50	2	51,100	102	666
Mackinaw class 73.2 m	6	75	450	2.0	900	6	535	5	60	4	50	30	27,375	1	178
Bay class 42.7 m	6	17	102	2.0	204	1	121	1	14	1	50	30	6,205	9	364
High endurance cutters															
115.2 m	60	156	9,360	2.0	18,720	122	11,120	110	1,254	78	50	3	56,940	12	683
82.3 m	60	109	6,540	2.0	13,080	85	7,770	77	875	55	50	3	39,785	10	398
Medium endurance cutters															
64.0 m	30	71	2,130	2.0	4,260	28	2,530	25	285	18	50	6	25,915	16	415
61.9-64.9 m	30	75	2,250	2.0	4,500	29	2,673	26	302	19	50	6	27,375	10	274
Patrol boats															
33.5 m	10	16	160	2.0	320	2	190	2	21	1	50	18	5,840	23	134
29.0 m	3	13	39	1.5	59	0	35	0	4	0	50	61	3,559	15	53
25.0 m	2	10	20	1.5	30	0	18	0	2	0	50	91	2,738	15	41
Buoy tenders															
Seagoing Coastal	5	50	250	1.5	375	2	223	2	25	2	50	37	13,688	27	370
River	4	32	128	1.5	192	1	114	1	13	1	50	46	8,760	12	105
Inland	5	18	90	1.0	90	1	53	1	6	0	50	37	3,285	18	59
Construction	3	14	42	1.0	42	0	25	0	3	0	50	61	2,555	6	15
	5	8	40	1.0	40	0	24	0	3	0	50	37	1,460	16	23
Harbor tugs															
Medium	1	4	4	1.0	4	0	2	0	0	0	50	183	730	4	3
Small	1	4	4	1.0	4	0	2	0	0	0	50	183	730	2,120	NA
Search and rescue <19.8 m	1	4	4	1.0	4	0	2	0	0	0	50	183	730	2,120	NA
Total garbage per year															4,317

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Table 26.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: U.S. Army vessels (Eastern Research Group estimates) (GT = gross tons; MT = metric tons).

U.S. Army vessels	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a			Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel per year (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (MT)	Total garbage per year ^b (m ³)	
					Total garbage (kg)	Dry garbage (m ³)	Plastic garbage (kg)								
					(m ³)	(kg)	(m ³)								
Logistic support vessels	30	40	1,200	2.0	2,400	16	1,426	14	161	10	35	4	10,220	41	267
Landing craft, utility class 2000	20	10	200	2.0	400	3	238	2	27	2	35	6	2,555	89	583
Large oceangoing tugs	20	8	160	2.0	320	2	190	2	21	1	35	6	2,044	20	133
Other small landing craft	2	5	10	1.0	10	0	6	0	1	0	35	64	639	314	2,045
Small harbor tugs	2	5	10	1.0	10	0	6	0	1	0	35	64	639	10	62
J-boats	2	5	10	1.0	10	0	6	0	1	0	35	64	639	16	104
Total garbage per year													490	3,194	

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

Table 27.--Derivation of per voyage, per vessel, and annual domestic garbage quantities generated: National Oceanic and Atmospheric Administration (NOAA) research vessels (MT = metric tons, NA = not applicable). (Source: Eastern Research Group estimates.)

NOAA research vessels	Voyage length (days)	Crew size	Person days per voyage	Per capita generation rate (kg/day)	Domestic garbage generation per voyage ^a			Annual ship utilization rate ^c (%)	Voyages per year	Garbage per vessel per year (kg)	No. of vessels	No. of entrances	Total garbage per year ^b (MT)	Total garbage per year ^b (m ³)	
					Total garbage (kg)	Dry garbage (m ³)	Plastic garbage (kg)								
					(m ³)	(kg)	(m ³)								
Large deepwater vessels	20	110	2,200	2.0	4,400	29	2,614	26	295	18	35%	6	28,105	10	281
Coastal research vessels	5	10	50	2.0	100	1	59	1	7	0	35%	26	2,555	14	36
Total garbage per year													317	2,066	

^aDry garbage is calculated as 59.4% of total garbage by weight. Plastic garbage is 6.7% of dry garbage by weight. See Table 4.

^bTotal garbage weight per year is equal to the product of either (1) per vessel annual garbage quantity and the vessel population, or (2) garbage quantity per voyage and the number of entrances.

^cRefers to the percentage of days annually operating in U.S. waters.

ACKNOWLEDGMENTS

The research contained in this paper is based upon work performed under contract for the Coast Guard and the U.S. Department of Transportation's Transportation Systems Center in Cambridge, Massachusetts. The authors wish to acknowledge the cooperation and support provided by these agencies.

REFERENCES

- Berger, J. D., and C. E. Armistead.
1987. Discarded net material in Alaskan waters, 1982-84. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-110, 66 p.
- Center for Environmental Education.
1986. Marine wildlife entanglement in North America. Center for Environmental Education, Wash., D.C., 219 p.
- Corrado, E. J.
1986. Hearings before the Subcommittee on Oversight and Investigation of the Committee on Merchant Marine and Fisheries, House of Representatives. Letter from Ernest J. Corrado, Vice President, American Institute for Merchant Shipping to the Honorable Walter B. Jones, Chairman, Committee on Merchant Marine and Fisheries, U.S. House of Representatives, October 28, 1986.
- Eastern Research Group.
1988a. Development of estimates of garbage disposal in the maritime sectors. Final report. Report prepared by the Eastern Research Group, Inc. for the Transportation Systems Center, Research and Special Programs Administration, U.S. Department of Transportation, December 1988, 139 p.

1988b. A regulatory evaluation of regulations implementing Annex V to MARPOL 73/78. Final report. Report prepared by the Eastern Research Group, Inc. for the Port Safety and Security Division, Program Development Branch, U.S. Coast Guard, December 1988, 333 p.
- Gassan, M. C.
1978. Integrated waste disposal systems. Management Digest of Marine Research, Report No. E11.15TM.78 (October 1978). Marine Research Program, Exxon International Company, 22 p.
- International Maritime Organization.
1987. Draft guidelines for the implementation of Annex V regulations for the prevention of pollution by garbage from ships. Marine Environmental Protection Committee, MEPC 26/10, 12 December 1987. International Maritime Organization: Lond.
- Jane's Fighting Ships.
1986. Jane's fighting ships, 1986-87. Jane's Publishing Co.

Merrell, T. R. Jr.

1985. The fish nets and other plastic litter on Alaskan beaches. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 160-182. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

National Academy of Science.

1975. Marine litter. In Assessing potential ocean pollutants. A report of the study panel on assessing potential ocean pollutants to the Ocean Affairs Board, Commission on Natural Resources, National Research Council, p. 405-438. Natl. Acad. Sci., Wash., D.C.

National Solid Waste Management Association.

- Undated. Solid waste disposal overview. National Solid Waste Management Association.

Naval Ship Engineering Center.

1971. Naval shipboard refuse study. Naval Ship Engineering Center, Hyattsville, MD, June 1971, 98 p.

Navy League of the United States.

1987. The almanac of seapower. Navy League of the United States, Arlington, VA.

Pacific Associates.

1988. A report to the Alaska Department of Environmental Conservation on the effects of MARPOL Annex V on the ports of Kodiak and Unalaska. Pacific Associates, September 1988, 64 p.

Parker, N. R., S. C. Hunter, and R. J. Yang.

1987. Development of methodology to reduce the disposal of non-degradable refuse into the marine environment. KVB, Inc., Engineering and Research Division, January 1987, 89 p.

Schwartz, R.

1987. Letter of 16 July 1987 from Richard Schwartz, President, Boat Owner's Association of the United States to Congressman Jerry Studds, Hearings before the Subcommittee on Coast Guard and Navigation and the Subcommittee on Fisheries and Wildlife Conservation and the Environment of the Committee on Merchant Marine and Fisheries, House of Representatives, 100th Congress, 17 June and 23 July 1987.

Society of Naval Architects and Marine Engineers.

1982. Guide for the disposal of shipboard wastes. Society of Naval Architects and Marine Engineers, Tech. Res. Bull. 3-33, January 1982.

Trillo, R. L. (editor).

1978. Jane's ocean technology. Franklin Watts, Inc., N.Y.

Uchida, R. N.

1985. The types and estimated amounts of fish net deployed in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 37-108. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

U.S. Army Corps of Engineers.

1987. Summary of United States flag passenger and cargo vessels operating or available for operation on 1 November 1986. Unpublished table. U.S. Army Corps of Engineers, Waterborne Commerce Statistics Center.

U.S. Bureau of the Census.

1987. File AE-975, vessel entrances and clearances. Data base of monthly port entrances and clearances by vessels.

U.S. Coast Guard.

1987a. Boating statistics. Office of Boating Safety, Consumer and Public Affairs, 1987.

1987b. Unpublished statistical tables from the Marine Safety Inspection System data base. July 1987.

U.S. Maritime Administration.

1987a. Merchant marine data sheet. March 1, 1987, 7 p.

1987b. United States oceanborne foreign trade routes. U.S. Department of Transportation. December 1987.

[U.S.] National Marine Fisheries Service.

1987. Fisheries of the United States, 1986. Current Fishery Statistics 8385, 116 p.

APPENDIX

**Detailed Garbage Generation Tables
for 10 Maritime Sectors**

(ERG estimates)

Appendix Table 2.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (GT - gross tons, MT - metric tons). Commercial passenger ships.

Commercial passenger ships	Total MT generated	Before MARPOL Annex V						After MARPOL Annex V*											
		Off-loaded in port			Incinerated at sea			Dumped overboard			Off-loaded in port			Incinerated at sea			Dumped overboard		
		MT			m ³			MT			m ³			MT			m ³		
		Plastics	Other	MT	Plastics	Other	MT	Plastics	Other	MT	Plastics	Other	MT	Plastics	Other	MT	Plastics	Other	MT
Cruise ships																			
U.S. vessels																			
>1,000 GT	1,577	20,561	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Under 1,000 GT	3,784	46,879	0	0	189	2,467													
Foreign vessels	7,978	5,744	74,899	638	8,322	1,596	20,805												
Excursion vessels	188,270	178,856	2,332,241	0	0	9,413	122,750												
Charter boats	56,465	42,349	552,219	0	0	14,116	184,073												
Total	58,0742	232,121	3,026,799	638	8,322	25,315	330,095												

*Assumes full compliance with Annex V requirements.

Appendix Table 3.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (GT = gross tons, MT = metric tons). Commercial fishing.

	Before MARPOL Annex V						After MARPOL Annex V ^a												
	Off-loaded in port			Incinerated at sea			Dumped overboard			Off-loaded in port			Incinerated at sea			Dumped overboard			
	Total MT Generated	MT		m ³	MT	m ³		MT	m ³	MT	m ³		MT	m ³	MT	m ³		MT	m ³
		Plastics	Other			Plastics	Other				Plastics	Other							
Commercial fishing																			
Undocumented	114,367	0	0	0	0	114,367	1,491,320	7,663	880,206	0	0	0	0	0	0	106,705	498,263		
Documented																			
5-25 GT	49,965	0	0	0	0	49,965	651,532	3,348	317,669	0	0	0	0	0	0	46,617	217,682		
25-300 GT	66,908	0	0	0	0	66,908	872,457	4,259	151,124	0	0	3,345	43,623	59,304	276,921				
300-1,000 GT	1,359	0	0	0	0	1,359	17,717	77	3,092	0	0	204	2,658	1,077	5,031				
Over 1,000 GT	578	0	0	0	0	578	7,539	27	677	0	0	173	2,262	378	1,763				
Foreign vessels ^b	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total for sector	233,177	0	0	0	0	233,177	3,040,564	15,373	1,352,768	0	0	3,723	48,542	214,081	999,660				

^aAssumes full compliance with Annex V requirements.

^bData unavailable in time for this report. See Section 2.3 for discussion.

Appendix Table 4.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). Recreational boating.

Recreational boating	Before MARPOL Annex V										After Annex V*										
	Total MT generated	Off-loaded in port			Incinerated at sea			Dumped overboard			Off-loaded in port			Incinerated at sea			Dumped overboard				
		MT	m ³	m ³	MT	m ³	m ³	MT	m ³	m ³	Plastics	Other	MT	m ³	m ³	MT	m ³	m ³	MT	m ³	m ³
Coastal states	223,267	148,845	1,940,895	0	0	74,422	970,447	14,959	1,867,751	208,308	972,704	0	0	0	0	0	17,127	53,461	0	0	0
Under 4.9 m	171,267	114,178	1,488,853	0	0	57,089	744,427	10,327	1,289,470	43,813	746,158	0	0	0	0	0	1,861	5,809	0	0	0
4.9-7.9 m	18,611	12,408	161,791	0	0	6,204	80,896	1,122	140,125	15,628	81,084	0	0	0	0	0	199	622	0	0	0
7.9-12.2 m	1,993	1,329	17,328	0	0	664	8,664	120	15,007	1,674	8,684	0	0	0	0	0	68	211	0	0	0
12.2-19.8 m	356	237	3,094	0	0	119	1,547	19	2,219	269	1,509	0	0	0	0	0	68	211	0	0	0
Over 19.8 m																					
Subtotal	415,495	276,996	3,611,961	0	0	138,498	1,905,981	26,548	3,314,573	369,692	1,810,140	0	0	0	0	0	19,255	60,103	0	0	0
Great Lakes states	134,934	89,956	1,173,001	0	0	44,978	586,500	8,137	1,015,916	113,304	587,865	0	0	0	0	0	13,493	42,119	0	0	0
Under 4.9 m	80,268	53,512	697,782	0	0	26,756	348,891	4,840	604,337	67,401	349,703	0	0	0	0	0	8,027	25,055	0	0	0
4.9-7.9 m	4,915	3,277	42,725	0	0	1,638	21,363	296	37,004	4,127	21,412	0	0	0	0	0	491	1,534	0	0	0
7.9-12.2 m	398	265	3,457	0	0	133	1,729	24	2,994	334	1,733	0	0	0	0	0	40	124	0	0	0
12.2-19.8 m	46	30	398	0	0	15	199	2	285	35	194	0	0	0	0	0	9	27	0	0	0
Over 19.8 m																					
Subtotal	220,560	147,040	1,917,363	0	0	73,520	958,682	13,299	1,660,537	185,200	960,906	0	0	0	0	0	22,060	68,860	0	0	0
Total for sector	636,055	424,036	5,529,325	0	0	212,018	2,764,662	39,848	4,975,109	554,892	2,771,045	0	0	0	0	0	41,315	128,964	0	0	0

*Assumes full compliance with Annex V requirements.

Appendix Table 6.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (GT = gross tons, MT = metric tons). Miscellaneous vessels.

Miscellaneous vessels	Total MT generated	Before MARPOL Annex V						After MARPOL Annex V*										
		Off-loaded in port			Incinerated at sea			Incinerated at sea			Off-loaded in port			Dumped overboard				
		MT	m ³	m ³	MT	m ³	m ³	MT	m ³	m ³	Plastics	Other	MT	m ³	MT	m ³	MT	m ³
School training	192	0	0	0	0	0	192	2,499	9	224	0	0	57	750	125	584		
1,000 GT and over																		
Under 1,000 GT	26	0	0	0	0	0	26	333	1	51	0	0	5	67	19	89		
Ocean																		
Coastal	48	0	0	0	0	0	48	625	3	96	0	0	10	125	36	167		
Industrial vessels																		
1,000 GT and over																		
Ocean	854	3	37	0	0	0	851	10,780	60	3,255	0	0	128	1,665	667	3,112		
Coastal	136	0	6	0	0	0	135	1,710	10	516	0	0	20	264	106	494		
Under 1,000 GT	140	0	6	0	0	0	139	1,762	10	625	0	0	14	181	116	541		
Ocean																		
Coastal	31	0	1	0	0	0	31	389	2	138	0	0	3	40	26	119		
Research vessels																		
Inspected																		
1,000 GT and over	26	0	1	0	0	0	25	322	2	45	0	0	13	166	11	51		
300-1,000 GT	144	0	6	0	0	0	143	1,814	10	324	0	0	43	560	91	423		
Uninspected																		
Under 300 GT	43	0	2	0	0	0	43	544	3	97	0	0	13	168	27	127		
Total for sector	1,637	5	60	0	0	0	1,633	20,778	109	5,372	0	0	306	3,986	1,223	5,709		

*Assumes full compliance with Annex V requirements.

Appendix Table 7.--Final disposition of vessel-generated garbage before and after MARPOL Annex V
(annual quantities) (MT - metric tons). U.S. Navy.

U.S. Navy vessels	Total MT Generated	Before MARPOL Annex V						After MARPOL Annex V*													
		Off-loaded in port			Incinerated at sea			Dumped overboard			Off-loaded in port			Incinerated at sea			Dumped overboard				
		MT	m ³	m ³	MT	m ³	m ³	MT	m ³	m ³	MT	m ³	m ³	Plastics	Other	MT	m ³	m ³	MT	m ³	m ³
Strategic missile submarines	113	0	0	0	0	0	113	1,473	0	0	0	8	4,725	0	0	0	0	0	105	492	0
Attack submarines	669	0	0	0	0	669	8,724	0	0	0	45	27,984	0	0	0	0	0	624	2,915	0	
Aircraft carriers	10,544	0	0	0	0	10,544	137,497	0	0	0	706	441,051	0	0	0	0	0	9,838	45,939	0	
Battleships	487	0	0	0	0	487	6,346	0	0	0	33	20,356	0	0	0	0	0	454	2,120	0	
Cruisers	2,514	0	0	0	0	2,514	32,788	0	0	0	168	105,174	0	0	0	0	0	2,346	10,955	0	
Destroyers	3,861	0	0	0	0	3,861	50,345	0	0	0	259	161,492	0	0	0	0	0	3,602	16,821	0	
Frigates	4,867	0	0	0	0	4,867	63,460	0	0	0	326	203,562	0	0	0	0	0	4,541	21,203	0	
Light forces	28	0	0	0	0	28	370	0	0	0	2	1,187	0	0	0	0	0	26	124	0	
Light amphibious warfare ships	16,182	0	0	0	0	16,182	211,005	0	0	0	1,084	676,843	0	0	0	0	0	15,097	70,498	0	
Mine warfare ships	34	0	0	0	0	34	444	0	0	0	2	1,425	0	0	0	0	0	32	148	0	
Auxiliary ships	6,408	0	0	0	0	6,408	83,556	0	0	0	429	268,023	0	0	0	0	0	5,978	27,917	0	
Military Sealift Command	876	0	0	0	0	876	11,423	0	0	0	59	36,641	0	0	0	0	0	817	3,816	0	
Ready reserve force	11,013	0	0	0	0	11,013	143,610	0	0	0	738	460,660	0	0	0	0	0	10,275	47,981	0	
Naval reserve	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	57,596	0	0	0	0	57,596	751,040	0	0	0	3,859	2,409,124	0	0	0	0	0	53,737	250,929	0	

*Assumes full compliance with Annex V requirements.

Appendix Table 8.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). U.S. Coast Guard.

U.S. Coast Guard vessels	Total MT generated	Before MARPOL Annex V				After MARPOL Annex V ^a															
		Off-loaded in port		Incinerated at sea	Dumped overboard	Off-loaded in port			Incinerated at sea	Dumped overboard											
		MT	m ³			Plastics	Other	MT			m ³										
				MT	m ³				MT	m ³		MT	m ³								
Icebreakers																					
Polar class 400 ft	102	0	0	0	102	319	7	4,275	0	0	0	0	0	0	0	0	0	0	0	95	298
Mackinaw class 240 ft	27	16	322	0	11	35	2	1,145	9	29	0	0	0	0	0	0	0	0	0	17	53
Bay class 140 ft	56	33	657	0	23	71	4	467	17	59	0	0	0	0	0	0	0	0	0	35	108
High endurance cutters																					
378 ft	683	0	0	0	683	2,133	46	28,580	0	0	0	0	0	0	0	0	0	0	0	638	1,990
270 ft	398	0	0	0	398	1,242	27	16,641	0	0	0	0	0	0	0	0	0	0	0	371	1,159
Medium endurance cutters																					
210 ft	415	0	0	0	415	1,294	28	17,343	0	0	0	0	0	0	0	0	0	0	0	387	1,208
203-213 ft	274	0	0	0	274	855	18	11,450	0	0	0	0	0	0	0	0	0	0	0	255	797
Patrol boats																					
110 ft	134	80	1,581	0	55	170	9	5,618	42	143	0	0	0	0	0	0	0	0	0	83	260
95 ft	53	32	628	0	22	68	4	2,233	17	57	0	0	0	0	0	0	0	0	0	33	103
82 ft	41	24	483	0	17	52	3	1,718	13	44	0	0	0	0	0	0	0	0	0	25	79
Buoy tenders																					
Seagoing	370	220	4,351	0	150	468	25	15,458	116	394	0	0	0	0	0	0	0	0	0	229	715
Coastal	105	62	1,238	0	43	133	7	4,397	33	112	0	0	0	0	0	0	0	0	0	65	203
River	59	35	696	0	24	75	4	2,473	19	63	0	0	0	0	0	0	0	0	0	37	114
Inland	15	9	180	0	6	19	1	641	5	16	0	0	0	0	0	0	0	0	0	10	30
Construction	23	14	275	0	9	30	2	977	7	25	0	0	0	0	0	0	0	0	0	14	45
Harbor tugs																					
Medium	3	2	34	0	1	4	0	122	1	3	0	0	0	0	0	0	0	0	0	2	6
Small	10	6	120	0	4	13	1	427	3	11	0	0	0	0	0	0	0	0	0	6	20
Search and rescue boats <65 ft	1,548	919	18,219	0	628	1,961	104	12,947	484	1,648	0	0	0	0	0	0	0	0	0	959	2,995
Total	4,317	1,452	28,786	0	2,864	8,941	289	126,913	765	2,604	0	0	0	0	0	0	0	0	0	3,262	10,183

^aAssumes full compliance with Annex V requirements.

Appendix Table 9.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). U.S. Army.

U.S. Army vessels	Total MT Generated	Before MARPOL Annex V				After MARPOL Annex V*									
		Off-loaded in port		Incinerated at sea		Dumped overboard		Off-loaded in port			Incinerated at sea		Dumped overboard		
		MT	m ³	MT	m ³	MT	m ³	Plastics	Other	MT	m ³	MT	m ³	MT	m ³
U.S. Army vessels															
Logistic support vessels	41	0	0	0	0	41	533	3	1,710	0	0	0	0	17	52
Landing craft, utility class 2	89	0	0	0	0	89	1,166	6	3,740	0	0	0	0	36	113
Large oceangoing tugs	20	0	0	0	0	20	267	1	855	0	0	0	0	8	26
Other landing craft	314	0	0	0	0	314	4,090	21	2,624	0	0	0	0	127	397
Small harbor tugs	10	0	0	0	0	10	125	1	80	0	0	0	0	4	12
J-boats	16	0	0	0	0	16	208	1	134	0	0	0	0	6	20
Total	490	0	0	0	0	490	6,388	33	9,143	0	0	0	0	199	621

*Assumes full compliance with Annex V requirements.

Appendix Table 10.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). National Oceanic and Atmospheric Administration (NOAA) research vessels.

NOAA research vessels	Before MARPOL Annex V						After MARPOL Annex V ^a							
	Total MT generated	Off-loaded in port		Incinerated at sea		Dumped overboard	Off-loaded in port			Incinerated at sea		Dumped overboard		
		MT	m ³	MT	m ³	MT	m ³	Plastics	Other	MT	m ³	MT	m ³	
Large deepwater vessels	281	6	141	84	1,099	191	9	235	0	0	141	1,832	131	612
Coastal research vessels	36	1	24	4	47	31	2	96	0	0	7	93	27	125
Total	317	7	165	88	1,146	222	11	331	0	0	148	1,926	158	737

^aAssumes full compliance with Annex V requirements.

Appendix Table 11.--Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). Summary table.

Sector	Before MARPOL Annex V										After MARPOL Annex V*														
	Off-loaded in port				Incinerated at sea			Dumped overboard			Off-loaded in port				Incinerated at sea			Dumped overboard							
	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	Plastics	Other	MT	m ³	MT	m ³	MT	m ³	MT	m ³	
Merchant shipping	30,949	3,302	39,794	1,148	14,971	26,499	349,304	1,626	311,353	2,737	6,255	4,381	57,132	22,204	103,685										
Commercial passenger ships	258,074	232,121	3,026,799	638	8,322	25,315	330,095	22,490	2,304,400	233,340	1,060,557	1,117	14,564	1,128	5,265										
Commercial fishing	233,177	0	0	0	0	233,177	3,040,564	15,373	1,352,768	0	0	3,723	48,542	214,081	999,660										
Recreational boating	636,055	424,036	5,529,325	0	0	212,018	2,764,662	39,848	4,975,109	554,892	2,771,045	0	0	41,315	128,964										
Offshore oil and gas operations	16,710	10,733	139,958	0	0	5,977	18,656	398	49,740	5,547	72,799	0	0	0	0										
Miscellaneous vessels	1,637	5	60	0	0	1,633	20,778	109	5,372	0	0	306	3,986	1,223	5,709										
U.S. Navy	57,596	0	0	0	0	57,596	751,040	3,859	2,409,124	0	0	0	0	53,737	250,929										
U.S. Coast Guard	4,317	1,452	28,786	0	0	2,864	8,941	289	126,913	765	2,604	0	0	3,262	10,183										
U.S. Army	490	0	0	0	0	490	6,388	33	9,143	0	0	0	0	199	621										
National Oceanic and Atmospheric Administration research vessels	317	7	165	88	1,146	222	2,463	11	331	0	0	148	1,926	158	737										
Total	1,239,322	671,656	8,764,887	1,874	24,439	565,791	7,292,892	84,037	11,544,253	797,282	3,913,261	9,674	126,150	337,306	1,505,752										

*Assumes full compliance with Annex V requirements.

THE QUANTITATIVE DISTRIBUTION AND CHARACTERISTICS OF
MARINE DEBRIS IN THE NORTH PACIFIC OCEAN, 1984-88

Robert H. Day,* David G. Shaw
Institute of Marine Sciences
University of Alaska
Fairbanks, Alaska 99775, U.S.A.

and

Steven E. Ignell
Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

*Present address Alaska Biological Research, P.O. Box 81934, College,
Alaska 99708, U.S.A.

ABSTRACT

The distribution, abundance, and characteristics of marine debris in the North Pacific, Bering Sea, and Japan Sea were studied during the 5-year period 1984-88 using standardized observations at 181 daily transect stations encompassing approximately 21,420 km of observations, for a total of 1,070 km² of sampling. The most abundant debris type was plastic, which composed 89.3% of the total 2,127 debris items seen on transect; other debris items consisted of glass (3.3%), wood (3.2%), paper/fiber (2.4%), metal (0.5%), rubber (0.2%), and unidentified debris objects (1.0%). The most abundant plastic type was fragments (34.2%); other main plastic types were Styrofoam objects (22.5%), sheets and bags (18.2%), gillnet floats (5.0%), polypropylene line (3.1%), miscellaneous floats (2.8%), and miscellaneous/unidentified plastic objects (12.3%). Gillnet fragments, trawl net fragments, unidentified net fragments, and uncut plastic strapping, which were minor components of the plastic debris, were recorded a total of 46 times, primarily between lat. 37° and 44°N, in and near the Subarctic Front. The distribution and characteristics of the 6 general debris types are presented, as well as the distribution and characteristics of the 11 main plastic types. The highest densities of marine debris generally occurred in Japan Sea and nearshore Japan Water, Transitional Water, and Subtropical Water. Densities of most types of marine debris generally were low in Subarctic Water and Bering Sea Water. Heterogeneous geographic input, currents, and winds are important in locally concentrating marine debris.

Appendix Table 11.---Final disposition of vessel-generated garbage before and after MARPOL Annex V (annual quantities) (MT = metric tons). Summary table.

Sector	Before MARPOL Annex V										After MARPOL Annex V*																				
	Off-loaded in port					Incinerated at sea					Dumped overboard					Off-loaded in port					Incinerated at sea					Dumped overboard					
	MT	m ³	MT	m ³	MT	MT	m ³	MT	m ³	MT	MT	m ³	MT	m ³	Plastics	Other	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	MT	m ³	
Merchant shipping	30,949	3,302	39,794	1,148	14,971	26,499	349,304	1,626	311,353	2,737	6,255	4,381	57,132	22,204	103,685																
Commercial passenger ships	258,074	232,121	3,026,799	638	8,322	25,315	330,095	22,490	2,304,400	233,340	1,060,557	1,117	14,564	1,128	5,265																
Commercial fishing	233,177	0	0	0	0	233,177	3,040,564	15,373	1,352,768	0	0	3,723	48,542	214,081	999,660																
Recreational boating	636,055	424,036	5,529,325	0	0	212,018	2,764,662	39,848	4,975,109	554,892	2,771,045	0	0	41,315	128,964																
Offshore oil and gas operations	16,710	10,733	139,958	0	0	5,977	18,656	398	49,740	5,547	72,799	0	0	0	0																
Miscellaneous vessels	1,637	5	60	0	0	1,633	20,778	109	5,372	0	306	3,986	1,223	5,709																	
U.S. Navy	57,596	0	0	0	0	57,596	751,040	3,859	2,409,124	0	0	0	0	53,737	250,929																
U.S. Coast Guard	4,317	1,452	28,786	0	0	2,864	8,941	289	126,913	765	2,604	0	0	3,262	10,183																
U.S. Army	490	0	0	0	0	490	6,388	33	9,143	0	0	0	0	199	621																
National Oceanic and Atmospheric Administration research vessels	317	7	165	88	1,146	222	2,463	11	331	0	0	148	1,926	158	737																
Total	1,239,322	671,656	8,764,887	1,874	24,439	565,791	7,292,892	84,037	11,544,253	797,282	3,913,261	9,674	126,150	337,306	1,505,752																

*Assumes full compliance with Annex V requirements.

THE QUANTITATIVE DISTRIBUTION AND CHARACTERISTICS OF
MARINE DEBRIS IN THE NORTH PACIFIC OCEAN, 1984-88

Robert H. Day,* David G. Shaw
Institute of Marine Sciences
University of Alaska
Fairbanks, Alaska 99775, U.S.A.

and

Steven E. Ignell
Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

*Present address Alaska Biological Research, P.O. Box 81934, College,
Alaska 99708, U.S.A.

ABSTRACT

The distribution, abundance, and characteristics of marine debris in the North Pacific, Bering Sea, and Japan Sea were studied during the 5-year period 1984-88 using standardized observations at 181 daily transect stations encompassing approximately 21,420 km of observations, for a total of 1,070 km² of sampling. The most abundant debris type was plastic, which composed 89.3% of the total 2,127 debris items seen on transect; other debris items consisted of glass (3.3%), wood (3.2%), paper/fiber (2.4%), metal (0.5%), rubber (0.2%), and unidentified debris objects (1.0%). The most abundant plastic type was fragments (34.2%); other main plastic types were Styrofoam objects (22.5%), sheets and bags (18.2%), gillnet floats (5.0%), polypropylene line (3.1%), miscellaneous floats (2.8%), and miscellaneous/unidentified plastic objects (12.3%). Gillnet fragments, trawl net fragments, unidentified net fragments, and uncut plastic strapping, which were minor components of the plastic debris, were recorded a total of 46 times, primarily between lat. 37° and 44°N, in and near the Subarctic Front. The distribution and characteristics of the 6 general debris types are presented, as well as the distribution and characteristics of the 11 main plastic types. The highest densities of marine debris generally occurred in Japan Sea and nearshore Japan Water, Transitional Water, and Subtropical Water. Densities of most types of marine debris generally were low in Subarctic Water and Bering Sea Water. Heterogeneous geographic input, currents, and winds are important in locally concentrating marine debris.

INTRODUCTION

Marine debris, especially plastic debris, increasingly is recognized as a national and international pollution problem (Shomura and Yoshida 1985; Wolfe 1987). Debris presents problems on beaches, where it is aesthetically displeasing, is expensive (and probably impossible) to remove, causes unnecessary mortality of coastal wildlife, and (in the case of some medical, military, and industrial wastes) is potentially toxic. Debris also can cause problems at sea, where it can damage vessels, entangle marine animals, and result in the deaths of some animals that mistake it for food. Although the general nature of the marine debris problem is understood, the actual magnitude of the problem is unknown, because much of the information about it is anecdotal. For instance, we know that the northern fur seal, *Callorhinus ursinus*, become entangled and die in derelict fishing nets at sea, but estimates of the abundance of derelict nets at sea are highly uncertain (Pruter 1987). Consequently, estimates of both the true mortality rate of fur seals due to entanglement and the true effects of this mortality on fur seal populations also are uncertain (but see Fowler 1982, 1985, 1987).

During the last two decades, several workers have systematically observed floating debris and lost plastic nets in the North Pacific Ocean (Venrick et al. 1973; DeGange and Newby 1980; Dahlberg and Day 1985; Jones and Ferrero 1985; Yoshida and Baba 1985; Baba et al. 1986; Day and Shaw 1987; Mio and Takehama 1987; Yagi and Nomura 1987) and stranded debris on coastal beaches (Merrell 1980, 1984). These studies have shown that marine debris is distributed widely, is of several types, and is distributed by surface currents and winds.

The objective of this study was to improve our knowledge of the quantitative distribution and characteristics of marine debris in the North Pacific Ocean. Specifically, we wanted to: (1) describe the quantitative distributions of the six main types of marine debris; (2) describe the comparative at-sea densities of the main debris types; (3) describe the mean dimensions of the main debris types; (4) describe the quantitative distributions of the 11 main types of plastic debris; (5) describe the frequencies of colors of the main plastic types; and (6) examine the effects of input, currents, and winds on the quantitative distribution of marine debris. Because of the extensive geographic coverage of the work, this study constitutes the first complete analysis and the most detailed synoptic picture of marine debris anywhere in the world ocean.

METHODS

We collected data on the density (number per square kilometer), types, sizes, and colors of marine debris at 181 debris transect stations in the Bering and Japan Seas and the North Pacific Ocean north of Hawaii. At each station, we counted, identified, and estimated the two largest dimensions (at least 2.5×2.5 cm) of marine debris within 50 m of one side of a ship moving forward at a known rate of speed for a known period of time (Dahlberg and Day 1985; Day and Shaw 1987). The only types of debris that were sampled as far as we could see from either side of a moving ship were

gillnet fragments, trawl net fragments, unidentified net fragments, and uncut pieces of plastic strapping. This paper includes some published data from 38 stations in 1984 (Dahlberg and Day 1985) and 49 stations in 1985 (Day and Shaw 1987); the data from the other 94 stations are from 1986 to 1988 and have been combined with the 1984-85 data for a broader overview of patterns in the North Pacific.

The sampling surveyed approximately 21,425 km of ocean, for a total of approximately 1,073 km² of sampling (Fig. 1). The total effort consisted of 854 h 47 min (854:47) of sampling at 152 of the stations during which observation conditions were recorded. Effort by observation condition was: poor 21:50 (2.6% of the total effort of known conditions); fair 163:30 (19.1%); moderate 253:00 (29.6%); good 320:17 (37.5%); and very good 96:10 (11.3%). We decreased sampling effort when conditions were less than moderate (21.7% of total effort during known conditions) and sampled extensively when conditions were moderate to very good (78% of total effort during known conditions). Sampling was not conducted during periods when high waves could affect sightability of debris.

General debris types were standardized and consisted of glass, metal, paper/fiber, plastic, rubber, wood, or miscellaneous/unidentified debris. Plastic debris types also were standardized: fragment, Styrofoam (which may include foamed plastics of other chemical composition), polypropylene line fragment (which may include synthetic lines of other chemical composition), gillnet float, miscellaneous float, gillnet fragment, trawl net fragment, unidentified net fragment, uncut plastic strapping, sheet/bag, and miscellaneous/unidentified plastic debris. The two largest dimensions of pieces of debris were estimated in centimeters. Pieces of plastic debris were identified to the same standardized colors that were used for neuston plastic (Day et al. 1990): black/gray, blue, brown, green, orange, red/pink, tan, transparent, white, yellow, and mixed/unidentified colors.

Data were compiled as the density (number/km²) of total marine debris, of each general type of marine debris, and of each type of plastic debris at each station. We stratified the density data into five oceanographic water mass strata: Bering Sea Water, Subarctic Water (north of the Subarctic Front or north of approximately lat. 42°N), Subtropical Water (south of the Subtropical Front or south of approximately lat. 31°N), Japan Sea/nearshore Japan Water (west of approximately long. 150°E), and Transitional Water (Subarctic Front, Transition Zone, and Subtropical Front). We then subjected the stratified density data (total density, the 6 general debris types, and the 11 plastic debris types) to Kruskal-Wallis tests (Conover 1980; Zar 1984). For each data set, we tested the hypothesis:

H_0 : The density does not vary among water masses.

When test results were significant, we conducted multiple comparison tests (Conover 1980) to determine which water masses were significantly different.

The size data were combined into 10-cm size classes for sizes up to 100 cm; larger debris items were combined into size classes 101-200 cm,




Figure 1.--Cruise tracks for marine debris sampling, 1984-88.

>200 cm, or unknown. The size data were compiled for each of the six general debris types but not for the individual plastic types. The color data were compiled as frequencies of each color of plastic; subsequently, these frequencies were divided by the total number of plastic items to determine percentages of each color type.

RESULTS

Total Debris

We recorded 2,127 debris objects on the 181 debris transects. Plastic was the most common general type of debris, being recorded 1,899 times (89.3% of the total number of debris objects). Glass was next in frequency (72 objects; 3.3%), followed by 68 wood objects (3.2%), 53 paper/fiber

objects (2.4%), 10 metal objects (0.5%), and 4 rubber objects (0.2%). Miscellaneous/unidentified marine debris was recorded 22 times (1.0%).

Marine debris was widespread in occurrence, but occurred in greatest densities in the Japan Sea and off the eastern coast of Japan; it also was common along the Subarctic Front and in southern Transitional Water (Fig. 2). Lowest densities were in the central Alaska Gyre, in the Bering Sea, and in the vicinity of the Hawaiian Islands. The highest density of total marine debris was 36.7 pieces/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. Densities of total marine debris differed significantly among water masses ($H = 66.735$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Transitional Water = Subtropical Water > Subarctic Water = Bering Sea Water.

Glass Debris

Glass objects were recorded 72 times. Glass containers of various types (miscellaneous bottle, sake bottle, jar, beer bottle, and Japanese whisky bottle, in decreasing order of frequency) were recorded 54 times (75.0% of total glass); bottles were the most abundant, being recorded 42 times (58%). The second main class of glass objects was light bulbs (11 objects; 15.3%), which were represented (in decreasing order) by incandescent bulbs, fluorescent bulbs, and floodlights. The remaining seven (9.7%) glass objects consisted of glass fishing floats (glass balls). The mean dimensions of glass debris were 17.9 × 33.7 cm ($n = 34$ objects of known dimensions).

Glass debris was widespread south of the Subarctic Water, occurring in greatest densities in southern Transitional Water, in the Japan Sea, and off eastern Japan; it was uncommon in Subarctic Water and absent in the Bering Sea (Fig. 3). The highest density was 1.3 pieces/km² at lat. 30°34'N, long. 173°10'W in Subtropical Water northwest of the Hawaiian Islands. Densities of glass differed significantly among water masses ($H = 34.744$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons tests were confusing, however, in that those water masses with the largest difference in mean ranks were not significantly different, whereas water masses with smaller differences in mean ranks were significantly different. The two water masses that were significantly different were Transitional Water > Subarctic Water, the two with the largest sample sizes (49 and 99, respectively). We suspect that other water masses were different but that sample sizes in most were too small for the multiple comparisons test to find significant differences. The pattern of mean ranks (in descending order) was: Japan Sea/nearshore Japan Water, Subtropical Water, Transitional Water, Subarctic Water, and Bering Sea Water.

Metal Debris

Metal objects were recorded 10 times. Metal cans of various sizes were the most common metal debris, being recorded eight times (80% of total metal). The remaining two metal objects were a 208.2 L (55-gal) drum and a metal trawl float (10% each). The mean dimensions of metal debris were 41.5 × 64.5 cm ($n = 5$ objects of known dimensions).




Figure 2.--Densities of total marine debris, 1984-88. Solid black circles indicate stations at which debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 36.7 pieces/km².

Metal debris was sporadic in occurrence and almost certainly originated from ships. The main areas of occurrence were the Japan Sea and off eastern Japan, with other records in the northern Gulf of Alaska and the eastern subarctic Pacific (Fig. 4). The highest density was 0.5 piece/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. Densities of metal debris appeared to differ significantly among water masses ($H = 10.106$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons, however, indicated that none of the water masses were significantly different; we suspect that densities were too low overall for the multiple comparisons to find significant differences. The pattern of mean ranks (in descending order) was: Japan Sea/nearshore Japan Water, Transitional Water, Subarctic Water, and none in Subtropical and Bering Sea Waters.

Table 1.--Densities (number/km²) of general types of marine debris in five water masses of the North Pacific, 1984-88.

Parameter	Bering Sea Water		Subarctic Water		Transitional Water		Subtropical Water		Japan Sea and nearshore Japan Water	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
n	7		99		49		18		8	
Distance sampled (km)	872.6		11,010.5		6,072.0		2,408.0		1,061.9	
Area sampled (km ²)	43.7		551.7		303.6		120.4		53.1	
Total density	0.3	0.3	0.4	0.6	3.6	3.8	2.4	3.5	11.5	12.8
Glass	0.0	0.0	<0.1	0.1	0.1	0.2	0.2	0.3	0.2	0.2
Metal	0.0	0.0	<0.1	<0.1	<0.1	<0.1	0.0	0.0	0.1	0.2
Paper/fiber	<0.1	0.1	<0.1	0.1	0.1	0.2	<0.1	0.1	0.1	0.8
Plastic	0.2	0.2	0.3	0.4	3.3	3.7	2.1	3.1	10.5	11.7
Rubber	0.0	0.0	0.0	0.0	<0.1	<0.1	0.0	0.0	0.1	0.1
Wood	<0.1	<0.1	0.1	0.3	<0.1	0.1	<0.1	0.1	0.2	0.2
Miscellaneous/unidentified	<0.1	0.1	<0.1	<0.1	0.1	0.1	<0.1	0.1	0.0	0.0




Figure 3.--Densities of glass debris, 1984-88. Solid black circles indicate stations at which glass debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 1.3 pieces/km².

Paper/Fiber Debris

Paper/fiber objects were recorded 53 times. Paperlike objects were the most common, being recorded 34 times (66.0% of total paper/fiber); of these, cardboard (fragments, boxes, sheets, and tubes) was recorded 19 times (35.8%), and paper (fragments, towels, cups, magazines, and cigarette packs) was recorded 16 times (30.2%). Hemp line was recorded 11 times (20.8%); it consisted of fragments of hemp deck lines from ships and of 1




Figure 4.--Densities of metal debris, 1984-88. Solid black circles indicate stations at which metal debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 0.5 piece/km².

piece of twine. Woven debris was the least common kind of paper/fiber, being recorded seven times (13.2%); this category included cloth fragments and bags, canvas fragments and bags, and one carpet fragment. The mean dimensions of paper/fiber objects were 23.1 × 75.6 cm (n = 42 objects of known dimensions); the mean dimensions excluding objects >200 cm long were only 23.1 × 50.6 cm (n = 42 and n = 39, respectively), however.

Because paper decomposes rapidly at sea, paper/fiber debris occurred primarily near shore (e.g., the Japan Sea, off eastern Japan, the Bering

Sea) or in areas that are fished heavily (e.g., southeastern Bering Sea, flying squid fishery near the Subarctic Front east of Japan), where numerous fishing boats provide constant input of paper debris; most of the records of this debris type in southern Transitional Water and northern Subtropical Water are of hemp deck lines (Fig. 5). The highest density was 2.3 pieces/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. Densities of paper/fiber differed significantly among water masses ($H = 38.676$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water > Transitional Water = Subtropical Water = Subarctic Water = Bering Sea Water.

Rubber Debris

Rubber objects were recorded only four times, the least of all general debris types. All four (100%) objects were rubber gloves, which frequently are used on fishing boats. The mean dimensions of rubber objects were 15.5 × 25.5 cm ($n = 3$ objects of known dimensions).

Rubber debris was recorded at only three stations: at lat. 32°15'N, long. 141°36'E in Transitional Water east of Japan; at lat. 36°55'N, long. 132°30'E in the Japan Sea; and at lat. 27°59'N, long. 157°13'W in Subtropical Water north of the Hawaiian Islands. The highest density was 0.3 piece/km² in the Japan Sea. Densities of rubber appeared to differ significantly among water masses ($H = 28.715$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons, however, found no significant differences; we suspect that densities were too low overall for the multiple comparisons to find differences. The pattern of mean ranks (in descending order) was: Japan Sea/nearshore Japan Water, Transitional Water, and none in Subtropical, Subarctic, and Bering Sea Waters.

Wood Debris

Wood objects were recorded 68 times. Sawed or milled wood objects were the most common (63 objects; 92.6% of total wood); this category consisted (in decreasing order of frequency) of boards, sawed logs (for shipping to sawmills), dock pilings, and large timbers or blocks that frequently are used as dunnage on ships. Bamboo objects (3; 4.4%) were next in abundance and consisted of flagpoles (for marking the ends of drift gillnets) and fragments. Finally, fabricated objects (2; 2.9%) were represented by one wooden pallet and what appeared to be a wooden ladder. The mean dimensions were 23.5 × 183.0 cm ($n = 62$ objects of known dimensions); the mean dimensions excluding objects >200 cm long were 23.5 × 78.8 cm ($n = 62$ and $n = 45$, respectively), however.

Wood debris occurred primarily near shore, probably because of its tendency to become waterlogged and sink with time. The highest densities were in the northern Gulf of Alaska, where harvested logs were common in the Alaska Coastal Current and farther offshore, in the Japan Sea and off the eastern shore of Japan; little wood debris was recorded far from shore, however (Fig. 6). The highest density was 2.8 pieces/km² at lat. 59°47'N, long. 148°17'W near the coast of the northern Gulf of Alaska. Densities of wood differed significantly among water masses ($H = 19.830$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities




Figure 5.--Densities of paper/fiber debris, 1984-88. Solid black circles indicate stations at which paper/fiber debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 2.3 pieces/km².

were: Japan Sea/nearshore Japan Water > Subarctic Water = Subtropical Water = Transitional Water = Bering Sea Water.

Plastic Debris

Types of Plastic Debris

Of the 1,899 plastic debris objects recorded on transect, fragments were the most common type (649 objects). Styrofoam was next in abundance




Figure 6.--Densities of wood debris, 1984-88. Solid black circles indicate stations at which wood debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 2.8 pieces/km².

(428 objects), followed by 346 sheets/bags, 95 gillnet floats, 59 polypropylene line fragments, 54 miscellaneous floats, 12 gillnet fragments (plus 8 seen off transect), 11 trawl net fragments (plus 3 seen off transect), 8 uncut plastic straps, and 3 unidentified net fragments (plus 1 seen off transect). Miscellaneous/unidentified plastic debris was recorded 234 times. The mean dimensions of plastic objects were 13.3 × 24.3 cm (n = 1,569 objects of known dimensions); the mean dimensions excluding objects >200 cm were 11.7 × 19.1 cm (n = 1,564 and n = 1,557, respectively).

As might be expected from its abundance overall, plastic debris was the most widespread of all debris types (Fig. 7). The highest densities were in the Japan Sea and off the eastern coast of Japan, with lower densities in the Subarctic Front east of Japan and in southern Transitional Water; the lowest densities were near the Hawaiian Islands, in Subarctic Water (especially in the Alaska Gyre), and in the Bering Sea. The highest density of total plastic debris was 32.6 pieces/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea; local densities here were so high that Day was unable to census all marine debris, so he stopped sampling here. The only other high densities of plastic debris were 23.8 pieces/km² at lat. 41°50'N, long. 146°12'E and 18.2 pieces/km² at lat. 40°15'N, long. 150°46'E, both off the eastern coast of Japan. Densities of plastic differed significantly among water masses ($H = 74.168$; $n = 181$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Subtropical Water = Transitional Water > Subarctic Water = Bering Sea Water.

Fragments were irregular pieces of plastic (other than the specific categories discussed here) that apparently had been broken from other, larger pieces. They were the most abundant plastic type, being recorded 649 times (34.2% of total plastic). Fragments occurred in highest densities off eastern Japan and in the Japan Sea, with lower densities in northern Subtropical Water near the Subtropical Front; in contrast, they were uncommon in Subarctic Water and the Bering Sea (Fig. 8). The highest density was 18.9 pieces/km² at lat. 41°50'N, long. 146°12'E off the eastern coast of Japan. The density of plastic fragments differed significantly among water masses ($H = 62.887$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Transitional Water > Subtropical Water = Subarctic Water = Bering Sea Water.

Styrofoam included all objects made of foamed polystyrene, including fragments, sheets, boxes or other containers, and fishing floats; based on observed colors and textures, we believe that none of this debris consisted of other types of foamed plastics (e.g., polyurethane). Styrofoam objects were recorded 428 times (22.5% of total plastic), making them second in abundance of all plastic types. As was seen for neuston plastic (Day et al. 1990), Styrofoam debris also is a "nearshore Japan/transitional species," with few records in Subarctic Water or the Bering Sea (Fig. 9). The highest density was 4.9 pieces/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. The density of Styrofoam differed significantly among water masses ($H = 58.655$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Transitional Water = Subtropical Water > Subarctic Water = Bering Sea Water.

Polypropylene line is used more commonly than are other synthetic lines and largely has replaced hemp line on ships; consequently, we categorized all lines that appeared to be synthetic as polypropylene. Debris of this type consisted of intact lines and line fragments. Polypropylene lines were recorded 59 times (3.1% of total plastic). These lines were absent in the Bering Sea, were recorded in Subarctic Water only three times, and peaked in abundance in and around the Subarctic Front, in




Figure 7.--Densities of plastic debris, 1984-88. Solid black circles indicate stations at which plastic debris was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 32.6 pieces/km².

the Japan Sea, and in and near the Subtropical Front (Fig. 10). The highest density was 1.2 pieces/km² at lat. 40°00'N, long. 175°17'E near the Subarctic Front in the central Pacific. Densities of polypropylene line differed significantly among water masses ($H = 27.068$; $n = 131$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons were confusing, however, in that those water masses with the largest difference in mean ranks were not significantly different, whereas water masses with smaller differences in mean ranks were significantly different. The two water masses that were significantly different were Transitional Water > Subarctic Water, the two




Figure 8.--Densities of plastic fragments, 1984-88. Solid black circles indicate stations at which plastic fragments was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 18.9 pieces/km².

with the largest sample sizes (49 and 99, respectively). We suspect that other water masses were different but that sample sizes in most were too small for the multiple comparisons to show significant differences. The pattern of mean ranks (in descending order) was: Transitional Water, Subtropical Water, Japan Sea/nearshore Japan Water, Subarctic Water, and Bering Sea Water.

Table 2.--Densities (number/km²) of types of plastic debris in five water masses of the North Pacific, 1985-88.

Parameter	Bering Sea Water		Subarctic Water		Transitional Water		Subtropical Water		Japan Sea and nearshore Japan Water	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
n	7		99		49		18		8	
Fragment	0.1	0.1	0.1	0.2	0.9	1.3	0.8	1.4	5.3	7.6
Styrofoam	<0.1	0.1	0.1	0.1	0.7	0.8	0.4	0.8	2.1	1.7
Polypropylene line	0.0	0.0	<0.1	<0.1	0.1	0.2	0.1	0.2	0.2	0.4
Gillnet float	0.0	0.0	<0.1	0.1	0.2	0.2	0.2	0.3	0.1	0.1
Miscellaneous float	0.0	0.0	<0.1	<0.1	0.1	0.1	0.1	0.1	0.1	0.2
Gillnet fragment	0.0	0.0	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.0	0.0
Trawl net fragment	0.0	0.0	<0.1	<0.1	<0.1	0.1	0.1	<0.1	0.0	0.0
Unidentified net fragment	0.0	0.0	0.0	0.0	<0.1	<0.1	0.0	0.0	0.0	0.0
Uncut strapping	0.0	0.0	<0.1	<0.1	<0.1	0.1	<0.1	<0.1	<0.1	0.1
Sheet/bag	<0.1	<0.1	<0.1	0.1	0.8	1.3	0.1	0.2	2.1	2.8
Miscellaneous/unidentified	0.1	0.1	<0.1	0.1	0.5	0.5	0.3	0.4	0.6	0.5




Figure 9.--Densities of Styrofoam, 1984-88. Solid black circles indicate stations at which Styrofoam was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 4.9 pieces/km².

Floats include gillnet floats, trawl net floats, longline floats, crab pot buoys, and large boat bumpers made out of plastic other than Styrofoam. They primarily represent various types of fishing floats.

Gillnet floats were widely distributed and were common, being recorded 95 times (5.0% of total plastic). They were especially common in and around the Subarctic Front (center of the major gillnet fishery for squid--see below), in southern Transitional Water, and in and near the Subtropical




Figure 10.--Densities of polypropylene line, 1984-88. Solid black circles indicate stations at which polypropylene line was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 1.2 pieces/km².

Front; the only place they were absent was in the Bering Sea, probably because of the limited sampling there (Fig. 11). The highest density was 0.8 piece/km² at lat. 40°15'N, long. 150°46'E near the Subarctic Front east of Japan and at lat. 27°59'N, long. 157°13'W in Subtropical Water north of Hawaii. Densities of gillnet floats differed significantly among water masses ($H = 28.690$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons again were confusing, however, in that those water masses with the largest difference in mean ranks were not significantly different, whereas




Figure 11.--Densities of plastic gillnet floats, 1984-88. Solid black circles indicate stations at which gillnet floats were not recorded. Sizes of hollow circles indicate relative densities. The highest density was 0.8 piece/km².

water masses with smaller differences in mean ranks were significantly different. The two water masses that were significantly different were Transitional Water > Subarctic Water, the two with the largest sample sizes (49 and 99, respectively). Again, we suspect that other water masses were different, but that sample sizes in most were too small for the multiple comparisons to show significant differences. The pattern of mean ranks (in descending order) was: Transitional Water, Subtropical Water, Japan Sea/nearshore Japan Water, Subarctic Water, and Bering Sea Water.

Miscellaneous floats also were widespread at sea. These floats were concentrated in southern Transitional Water and Subtropical Water, with records scattered everywhere but the Bering Sea, again probably because of the limited sampling there (Fig. 12). These floats were rare in Subarctic Water as a whole, however. The highest density was 0.6 piece/km² at lat. 33°01'N, long. 158°31'W in Transitional Water north of Hawaii. Densities of miscellaneous floats differed significantly among water masses ($H = 29.842$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons again were confusing, in that those water masses with the largest difference in mean ranks were not significantly different, whereas water masses with smaller differences in mean ranks were significantly different. The two water masses that were significantly different were Subtropical Water > Subarctic Water; one of these water masses had a moderate sample size and the other had a large sample size (18 and 99, respectively). We again suspect that other water masses were different, but that sample sizes in most were too small for the multiple comparisons to show significant differences. The pattern of mean ranks (in descending order) was: Japan Sea/nearshore Japan Water, Subtropical Water, Transitional Water, Subarctic Water, and Bering Sea Water.

Although they were recorded only 34 times on transect and another 12 times off transect, net fragments and uncut packing straps are important components of marine debris, for they are thought to cause excessive mortality of some marine animals such as northern fur seals (Fowler 1982, 1985, 1987). These four plastic types were not distributed evenly in the North Pacific, but instead were concentrated between lat. 37° and 44°N (Fig. 13).

Gillnet fragments were recorded on transect 12 times and off transect 8 times; they were seen between lat. 25°37' and 45°15'N, with the most (3) seen at lat. 38°-39° and 42°-43°N (Fig. 13). The highest density was 0.7 piece/km² in Subtropical Water northwest of the Hawaiian Islands. Densities of gillnet fragments differed significantly among water masses ($H = 14.732$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons, however, did not find significant differences. We suspect that densities were too low overall for the multiple comparisons to find significant differences. The pattern of mean ranks (in descending order) was: Subtropical Water, Transitional Water, Subarctic Water, and none in Japan Sea/nearshore Japan Water and Bering Sea Water.

Trawl net fragments were recorded on transect 11 times and off transect 3 times; they were seen between lat. 30°21' and 44°07'N, with the most (3) seen at lat. 40°-41° and 41-42°N (Fig. 13). The highest density was 0.2 piece/km², recorded at four stations near the Subarctic Front in the central and western North Pacific. Densities of trawl net fragments differed significantly among water masses ($H = 10.629$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons did not find significant differences, however, and we suspect that densities were too low overall for the multiple comparisons to find significant differences. The pattern of mean ranks (in descending order) was: Transitional Water, Subtropical Water, Subarctic Water, and none in Japan Sea/nearshore Japan Water and Bering Sea Water.




Figure 12.--Densities of miscellaneous plastic floats, 1984-88. Solid black circles indicate stations at which miscellaneous plastic floats were not recorded. Sizes of hollow circles indicate relative densities. The highest density was 0.6 piece/km².

Unidentified net fragments were recorded on transect three times and off transect once; they were seen between lat. 33°07' and 43°35'N, with the most (two) seen at lat. 33°-34°N (Fig. 13). The estimated mesh size of these nets was 4 × 4 cm. The highest density was 0.3 piece/km² at lat. 33°01'N, long. 158°31'W in Transitional Water north of Hawaii. Densities of unidentified net fragments did not differ significantly among water masses ($H = 5.418$; $n = 181$; $df = 4$; $P > 0.05$; Table 2), probably because they were so low overall.




Figure 13.--Numbers of nets and uncut plastic strapping seen, by 1° blocks of latitude, 1984-88.

Uncut straps were recorded eight times, all on transect; they were seen between lat. 29°33' and 50°03'N, with the most (four) seen at lat. 40°-41°N (Fig. 13). The highest density was 0.3 piece/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. Densities of uncut straps did not differ significantly among water masses ($H = 5.462$; $n = 181$; $df = 4$; $P > 0.05$; Table 2), probably because they were so low overall.

Sheets and bags, which are a pollution problem because they are eaten by and entangle sea turtles (Balazs 1985; Carr 1987), were recorded 346 times (18.2% of total plastic). Sheets and bags occurred in highest densities in the Japan Sea, off the eastern coast of Japan, and along the Subarctic Front east of Japan; this debris type was common in Transitional Water and was essentially absent from Subarctic Water and the Bering Sea (Fig. 14). The highest density was 7.9 pieces/km² at lat. 36°55'N, long. 132°30'E in the Japan Sea. Densities of sheets/bags differed significantly among water masses ($H = 61.202$; $n = 181$; $df = 4$; $P < 0.05$; Table 2). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Transitional Water > Subtropical Water = Subarctic Water = Bering Sea Water.

Miscellaneous/unidentified plastic consisted of both fabricated objects and truly unidentified pieces; the latter objects occurred when we encountered such high local densities that we were unable to record all details on individual plastic objects. One hundred fifty-seven containers of various kinds constituted 67.1% of this category and included bottles, jars, squeeze tubes, boxes, bowls, cups, pans, beer or soda cases, woven bags, and buckets. The remaining 77 objects were a diverse assortment of screens, sponges, lids, mats, bottle caps, sandals, trays, rings, shoe liners, shovels, pipes, toys, paddles, poles, baseball caps, handles, helmets, and unidentified plastic debris. The highest density was 1.3 pieces/km² at lat. 35°10'N, long. 176°01'E in Transitional Water in the central North Pacific, at lat. 40°15'N, long. 150°46'E near the Subarctic Front off eastern Japan, and at lat. 36°55'N, long. 132°30'E in the Japan Sea.

Colors of Plastic Debris

Plastic debris was recorded in all 10 of the standardized colors, plus miscellaneous/mixed colors (Fig. 15). White was by far the most common color, being recorded 922 times (48.6% of total plastic). The color tan was second in abundance (187; 9.9%), followed by transparent (124; 6.5%), blue (119; 6.3%), and yellow (86; 4.5%). The colors green (35; 1.8%), brown (32; 1.7%), red/pink (28; 1.5%), black/gray (25; 1.3%), and orange (17; 0.9%) were rare in occurrence. Finally, miscellaneous/mixed plastic was recorded 323 times (17.0% of total plastic), primarily in cases when local densities were too high for us to record all data on individual pieces of debris.

Frequencies of some colors of debris plastic differed strongly from those frequencies of neuston plastic (Fig. 15). The greatest difference was in transparent plastic, whose frequency in marine debris was <25% of that in neuston plastic. Similarly, the frequency of black and gray plastic in marine debris was <50% of that in neuston plastic. In




Figure 14.--Densities of plastic sheets and bags, 1984-88. Solid black circles indicate stations at which plastic sheets and bags were not recorded. Sizes of hollow circles indicate relative densities. The highest density was 7.9 pieces/km².

contrast, the frequency of white plastic was nearly 33% higher, that of tan plastic was nearly four times higher, and that of yellow plastic was nearly four times higher in marine debris than in neuston plastic. Frequencies of the other colors were relatively similar comparing the two types of plastic.

DISCUSSION

We believe that the present distribution of marine debris is controlled largely by four main phenomena: (1) the heterogeneous geographic




Figure 15.--Frequencies of colors of plastic debris, 1984-88, compared with frequencies of colors of neuston plastic, 1985-88 (the latter data from Day et al. 1990).

input of debris, (2) the subsequent redistribution of that debris by currents and winds, (3) the decomposition of the debris at sea, and (4) the beaching of the debris. The data that we report on here strongly suggest that these factors interact to yield the distributions that we observed. In the absence of precise data on rates of input, transport, and degradation, however, our conclusions about factors controlling marine debris in the North Pacific must be considered tentative.

It is clear that there is heterogeneous geographic input of marine debris, with much originating in the far western Pacific. This conclusion

is supported strongly by the high densities in and around the Japan Sea and nearshore Japan, where the highest densities of both marine debris and neuston plastic (Day et al. 1990) in the North Pacific were recorded. Debris was most abundant in Tokyo Bay (which had far more debris than Day has ever seen elsewhere in the North Pacific--it was too abundant for him to sample) and in localized areas in the Japan Sea. It is unclear to us how much of this debris comes from ships and how much comes from the land. At the other extreme were the Bering Sea and Gulf of Alaska, where low human populations probably provide little input of marine debris.

In Transitional Water to the east of Japan, the importance of transport compared to direct input from ships is difficult to evaluate. The area between lat. 35° and 45°N and from the eastern coast of Japan to long. 145°W is the site of a large pelagic fishery for neon flying squid, *Ommastrephes bartrami*. At the height of the fishery (May-December), approximately 700 gillnetting ships from Japan, Korea, and Taiwan participate (Fredin 1985), as well as an unknown number of small jigging ships. This fishery undoubtedly contributes to marine debris in the area, although its contribution relative to transport is unknown. In contrast, debris entering the ocean around Japan and Korea is moved eastward by the Subarctic Current (in Subarctic Water) and in the Kuroshio (Kawai 1972; Favorite et al. 1976; Nagata et al. 1986) into the same area. In addition to this general eastward movement, Ekman (wind) stress tends to move surface waters from the Subarctic and the Subtropic into the Transitional Water mass as a whole (Roden 1970). As a result, densities of debris in Transitional Water generally are high, but the relative importance of the two sources (i.e., local input and transport into the area) is unclear. Further, the generally convergent nature of surface water in the North Pacific Central Gyre (Masuzawa 1972) should result in high densities there also.

Surprisingly, there are differences among the distributions of types of debris. For example, Styrofoam debris clearly is a "nearshore Japan/transitional/subtropical species" (Fig. 9); neuston Styrofoam also is most abundant around Japan (Day et al. 1990). This localized distribution of Styrofoam may be a consequence of its weak, crumbly texture, which can lead to rapid disintegration; hence, it probably cannot survive long enough to be transported offshore in large quantities. Further, Styrofoam sinks when crushed and waterlogged. Thus, it may be observed only in places where input rates are high. In addition, plastic sheets and bags also seem to be "transitional" (Fig. 14), placing them directly in the range of most of the world's sea turtle species, which readily ingest this type of plastic debris (Balazs 1985). The reason for this distribution of sheets and bags is not known.

The comparison of frequencies of colors of neuston plastic and debris plastic (Fig. 15) suggests a bias in our sampling. Colors that do not contrast strongly with seawater (black/gray, transparent) are underrepresented in debris in comparison with neuston plastic. Although some bias in the color frequency data for neuston plastic probably results from color-selective ingestion of neuston plastic by seabirds (Day et al. 1985), we believe that the difficulty in observing low-contrast debris is the major cause of the differences in Figure 15. Although there is bias in

the debris sighting data, however, densities of debris plastic and neuston plastic are strongly correlated (Day and Shaw 1987). Hence, although absolute estimates of at-sea densities of marine debris plastic are affected by these sighting biases, the debris data presented here provide important information about relative abundances in various parts of the ocean.

Although Fowler (1982, 1985, 1987) claimed that entanglement in lost netting and other marine debris is the major source of mortality of northern fur seals, we find that the data on at-sea densities of lost net fragments are inadequate to determine quantitatively its true importance. We have seen fur seals entangled in net fragments only twice, both in the flying squid fishery and both during fall 1987. The first record was of a fur seal with a trawl net fragment caught over its head at lat. 44°07'N, long. 156°23'W; there were raw, open cuts on the face and gums, although this animal did not appear to be hurt in any way and swam playfully with another unentangled fur seal. The second record was of an immature female fur seal completely entangled in a gillnet fragment at lat. 43°15'N, long. 145°11'W, along with the partially eaten remains of what appeared to be a salmon shark, *Lamna ditropis*, and a yellowtail, *Seriola lalandi*. Thus, sightings of entangled fur seals in derelict net fragments at sea are quite rare, making it difficult to assess the frequency of entanglement. On the other hand, our extensive experience at sea in the North Pacific suggests to us that the probability of entanglement and subsequent mortality of fur seals is higher in nets that actively are fishing for flying squid than in net fragments. The flying squid fishery deploys approximately 3,000,000 km of drift gillnets annually and is concentrated approximately in the zone lat. 39°-46°N (Day unpubl. data). Further, many of the deployments of research nets observed by Day in this area resulted in fur seals' feeding from the nets, climbing on and swimming around the nets, and occasionally becoming caught in the nets. (Most escaped unharmed, however.) Given the high number of entanglements of fur seals in actively fishing gillnets that we have observed, the nearly 60,000 vessel-nights of net deployments in a year, the large amounts of those nets that are fished, and the low number of entanglements in lost net fragments that we have observed in over 21,000 km of observations at sea, we suggest here that the mortality of fur seals from actively fishing nets should be assessed quantitatively and compared to estimates of mortality from derelict nets.

ACKNOWLEDGMENTS

Most of this research was done while Day was an Angus Gavin Memorial Fellow, a Sea Grant Fellow, or a Resources Fellow at the University of Alaska. Additional funding came from the National Marine Fisheries Service (NMFS), NOAA (contracts 40ABNF63228 and 43ABNF702983) and two contracts from the Joint Institute for Marine and Atmospheric Research at the University of Hawaii. Data were collected aboard the TV *Oshoro Maru* of Hokkaido University, Hakodate, Japan; RV *Pusan 851* of the National Fisheries Research and Development Agency, Government of the Republic of Korea, Pusan, Korea; RV *Miller Freeman* of the Northwest and Alaska Fisheries Center, NMFS, Seattle, Washington; RV *Akademik Korolev* of the Far Eastern Regional Institute of Goskomgidromet, Government of the Soviet Union, Vladivostok, U.S.S.R.; and RV *Alpha Helix* of the Institute of Marine Sciences, University of Alaska,

Fairbanks, Alaska. We thank the captains, officers, and crews of these ships for assistance. This manuscript was improved by comments of three anonymous reviewers. This is Contribution No. 828 of the Institute of Marine Sciences, University of Alaska, Fairbanks, Alaska.

REFERENCES

- Baba, N., K. Yoshida, M. Onoda, N. Nagai, and S. Toishi.
1986. Research on ocean current and drifting stray fishing gear and net fragments in the area to the southwest of the Pribilof Islands and the southern coast of the Aleutian Islands July-August, 1985. Far Seas Fisheries Research Laboratory, Shimizu, Japan, 15 p. Document submitted to the Standing Scientific Committee of the North Pacific Fur Seal Commission.
- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Carr, A.
1987. Impact of non-degradable marine debris on the ecology and survival outlook of sea turtles. Mar. Pollut. Bull. 18:352-356.
- Conover, W. J.
1980. Practical nonparametric statistics, 2d ed. Wiley, N.Y., 493 p.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Day, R. H., and D. G. Shaw.
1987. Patterns in the abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-1985. Mar. Pollut. Bull. 18:311-316.
- Day, R. H., D. G. Shaw, and S. E. Ignell.
1990. Quantitative distribution and characteristics of neuston plastic in the North Pacific, 1985-88. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154. [See this document.]
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

DeGange, A. R., and T. C. Newby.

1980. Mortality of seabirds and fish in a lost salmon driftnet. *Mar. Pollut. Bull.* 11:322-323.

Favorite, F., A. J. Dodimead, and K. Nasu.

1976. Oceanography of the subarctic Pacific region, 1960-71. *Int. North Pac. Fish. Comm., Bull.* 33:1-187.

Fowler, C. W.

1982. Interactions of northern fur seals and commercial fisheries. *Trans. N. Am. Nat. Resour. Conf.* 47:278-292.

1985. An evaluation of the role of entanglement of population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.

Fredin, R. A.

1985. Fishing effort by net fisheries in the North Pacific Ocean and Bering Sea since the 1950's. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 218-251. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Jones, L. L., and R. C. Ferrero.

1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84, In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Kawai, H.

1972. Hydrography of the Kuroshio Extension. In H. Stommel and K. Yoshida (editors), *Kuroshio: Physical aspects of the Japan Current*, p. 235-352. Univ. Wash. Press, Seattle, WA.

Masuzawa, J.

1972. Water characteristics of the North Pacific Central Region. In H. Stommel and K. Yoshida (editors), *Kuroshio: Physical aspects of the Japan Current*, p. 95-127. Univ. Wash. Press, Seattle, WA.

Merrell, T. R., Jr.

1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.

- Mio, S., and S. Takehama.
1987. Estimation of distribution of marine debris based on the 1986 sighting survey. Fisheries Agency of Japan, Tokyo, Japan, 16 p. Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Vancouver, Canada.
- Nagata, Y., J. Yoshida, and H.-R. Shin.
1986. Detailed structure of the Kuroshio Front and the origin of the water in warm-core rings. *Deep-Sea Res.* 33:1509-1526.
- Pruter, A. T.
1987. Sources, quantities, and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Roden, G. I.
1970. Aspects of the mid-Pacific Transition Zone. *J. Geophys. Res.* 75:1097-1109.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Venrick, E. L., T. W. Backman, W. C. Bartram, C. J. Platt, M. S. Thornhill, and R. E. Yates.
1973. Man-made objects on the surface of the central North Pacific Ocean. *Nature (Lond.)* 241(5387):271.
- Wolfe, D. A. (guest editor).
1987. Special issue: Plastics in the sea. *Mar. Pollut. Bull.* 18:303-365.
- Yagi, N., and M. Nomura.
1987. Sighting survey on marine debris in the midwestern Pacific from 1977 through 1986. Fisheries Agency of Japan, Tokyo, Japan, 12 p. Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Vancouver, Canada.
- Yoshida, K., and N. Baba.
1985. Results of the survey on drifting fishing gear or fish net pieces in the Bering Sea. Far Seas Fisheries Research Laboratory, Shimizu, Japan, 12 p. Document submitted to the 28th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 4-12 April 1985, 13 p.
- Zar, J. H.
1984. *Biostatistical analysis*, 2d ed. Prentice-Hall, Englewood Cliffs, NJ, 718 p.
-

DISTRIBUTION AND DENSITY OF FLOATING OBJECTS IN THE
NORTH PACIFIC BASED ON 1987 SIGHTING SURVEY

Shin-ichi Mio*

National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

Shuichi Takehama

Division of Fishing Ground Environment Conservation
Fisheries Agency, the Government of Japan
Chiyoda-ku, Tokyo, 100 Japan

and

Satsuki Matsumura

National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

*Present address: Japan Sea National Fisheries Research Institute,
Fisheries Agency, the Government of Japan, Niigata-shi, Niigata, 951 Japan

ABSTRACT

A sighting survey has been conducted in the North Pacific since 1986 to understand the distribution of floating objects there. The survey was conducted on board various types of vessels, including a fisheries research vessel, patrol vessels, training vessels, and commercial freighters, a total of 32 vessels with a distance surveyed of 165,288 nmi.

A total of 46,706 floating objects were recorded in 1987. Of these, fishing net debris accounted for 0.7%, other fishing gears 5.9%, Styrofoam 14.0%, and other petrochemical products 18.3%. The remainder included drifting logs or lumber 7.9%, floating seaweed 42.7%, and other 10.5%.

Density of the objects was generally high in the coastal waters, but high density was also observed in areas between lat. 25° and 30°N, and long. 170° and 130°W. It is assumed that floating objects transported from various areas by ocean currents accumulated here. A belt-shaped low-density area was observed between lat. 45° and 50°N.

INTRODUCTION

A total of 32 vessels, vessels which belong to the Fisheries Agency of the Government of Japan, training ships of fishery high schools and universities, and cargo transport ships, participated in a sighting survey of floating objects in the North Pacific (Table 1). The total distance for which the sightings were conducted was 165,228 nmi, and 46,706 items of marine debris were sighted during the cruises.

This survey has been repeated continuously since 1986, with the objectives of defining patterns of marine debris, clarifying the conditions of distribution, and determining the actual volume of various types of debris floating in the sea. Although the areas surveyed extended to the Sea of Japan, Yellow Sea, South Pacific Ocean, and Gulf of Mexico, this report concentrates on the North Pacific and its adjacent waters.

METHODS

Methods of sighting and items of observation were the same as in the previous year (Mio and Takehama 1987), except for the addition of the size of debris items observed. Size is described as follows. We measured with the eye the length of the longest piece of marine debris and recorded that $S = <50$, $M = 50-200$, and $L = >200$ cm.

RESULTS

Outline of Results

The distribution pattern of the cruising distance for the surveys (henceforth referred to as effort) shows that effort was high in Japanese waters and in the western Pacific, and low in the eastern and southern Pacific (Fig. 1). By season, 57.7% of the entire effort was expended during the 4 months from June to September. In the other months, excluding December, 4 to 8% of the effort was expended.

Looking at marine debris by kind (Table 2), 310 pieces of fishing net were recovered, 0.7% of all marine debris found (gillnet 0.2%, trawl net 0.1%, and unidentified net 0.4%). The proportion of fishing gears other than nets was somewhat larger (5.9%) and accounted for 15.3% of the total petrochemicals (fishing nets, other fishing gears, Styrofoam, and other plastic debris). Styrofoam accounted for 36% of all petrochemicals and for 14% of all marine debris, being the most abundant single material. Sheets and bags made of nylon and vinyl, and other plastic debris represented by containers for detergent and drinking water, accounted for 18.3% of the total marine debris and for 47.0% of the total petrochemicals. The number of their sightings was large, and they were quantitatively the major item of marine debris. Among biodegradable marine debris, pieces of wood and drifting logs accounted for 7.9%, and floating seaweed accounted for 42.7%. Other consisted mainly of glass products and empty cans, and accounted for 10.5%.

Table 1.--Vessels engaged in marine debris sighting survey in 1987.

Name of vessel	Gross tonnage	Horsepower	Area of survey	Cruising distance (nmi)	Number of debris pieces sighted
<i>Kotaka Maru</i>	47	235	J	648.9	2,319
<i>Tankai Maru</i>	157	900	J	2,758.8	296
<i>Hokko Maru</i>	466	1,800	J P	9,435.1	971
<i>Wakataka Maru</i>	170	540	J	3,618.9	1,325
<i>Soyo Maru</i>	494	1,600	J	6,286.1	3,221
<i>Yoko Maru</i>	499	1,600	J	5,363.6	1,341
<i>Mizuho Maru</i>	150	900	J	4,979.2	3,094
<i>Shunyo Maru</i>	393	2,600	J	7,156.6	1,340
<i>Shoyo Maru</i>	1,362	2,000	J P	14,857.4	2,798
<i>Kaiyo Maru</i>	2,644	3,800	J P	9,940.2	179
<i>Wakatake Maru</i>	427	1,500	J P	4,166.7	259
<i>Shin Riasu Maru</i>	471	1,400	J P	12,194.9	910
<i>Wakasio Maru</i>	199	900	J	1,285.3	448
<i>Hoyo Maru No. 12</i>	284	1,000	J P	4,134.4	663
<i>Kanki Maru No. 58</i>	96	470	J P	4,337.7	229
<i>Hokuho Maru</i>	441	1,300	J P	7,281.7	654
<i>Shirafuji Maru</i>	138	1,000	J	386.0	876
<i>Osyoro Maru</i>	1,779	3,200	J P	4,018.5	166
<i>Hoksei Maru</i>	893	2,100	J P	4,057.2	236
<i>Tansu Maru</i>	444	1,500	J	1,561.9	779
<i>Omi Maru</i>	417	1,300	J P	4,485.1	107
<i>Shirahagi Maru</i>	366	2,600	J P	6,903.4	305
<i>Toko Maru</i>	1,513	8,000	J P	12,845.6	343
<i>Hakuryu Maru</i>	517	2,500	J P	4,086.7	230
<i>Coop</i>	2,445	3,800	J	1,237.6	52
<i>Sunbelt Dexie</i>	11,447	14,000	J P	5,366.6	15,387
<i>Nichiyo Maru</i>	995	3,000	J P	1,176.7	4,673
<i>Kumamoto Maru</i>	380	1,600	J	3,599.2	598
<i>Riasu Maru No. 1</i>	476	1,100	J P	11,732.4	251
<i>Hoyo Maru No. 78</i>	300	440	J P	5,294.1	885
<i>Taisei Maru No. 55</i>	350	3,400	J P	725.2	449
<i>Tosi Maru No. 15</i>	730	3,600	J	3,803.5	1,974

^aJ = Japanese waters north of lat. 20°N and west of long. 160°E;
P = Pacific area other than Japanese waters.

Effects of Environmental Conditions

In order to study the effects of luminous intensity and waves on the sighting survey, the numbers of sightings by wind force and by time of day were examined comparatively for eight vessels which had conducted surveys for a fairly long time in the area where the effort expended was largest (lat. 40° to 45°N and long. 140° to 150°E).




Figure 1A.--Tracks of vessels engaged in marine debris sighting survey in 1987, coastal waters of Japan.




Figure 1B.--Tracks of vessels engaged in marine debris sighting survey in 1987, neighboring waters of Japan.




Figure 1C.--Tracks of vessels engaged in marine debris sighting survey in 1987, Pacific high seas area.

Table 2.--Number of debris items sighted, by month and type of debris.

Month	Cruising distance (nmi)	Type of debris										Total
		Fishing net		Other fishing gear	Styrofoam	Other plastic debris	Wood	Floating seaweed	Others			
		Unknown	Trawl							Gill		
January	7,857.3	--	--	18	30	12	192	169	151	582		
February	7,832.1	4	--	38	128	85	34	32	31	354		
March	8,813.8	2	--	104	415	115	88	709	394	1,827		
April	6,819.6	4	2	134	360	208	207	884	62	1,875		
May	11,181.7	4	2	169	334	131	90	257	221	1,210		
June	28,435.9	13	14	313	752	1,020	1,172	2,192	833	6,617		
July	31,323.2	23	5	344	724	958	314	1,601	296	4,266		
August	20,184.3	19	8	315	873	799	295	781	309	3,408		
September	18,743.9	44	25	567	1,136	1,500	507	4,951	627	9,371		
October	13,030.9	31	3	282	1,162	2,761	612	7,987	1,784	14,641		
November	9,840.1	27	1	459	541	901	141	374	162	2,619		
December	1,225.2	2	--	31	97	59	20	14	13	236		
Total	165,287.9	173	60	2,774	6,552	8,544	3,672	19,951	4,893	46,706		

Wind Force

The evidence suggests that the number of debris items sighted per unit of effort (100 nmi) is inversely related to wind force (Beaufort scale). The maximum sightings occurred at wind force class 1 and sightings decreased as the wind force increased (Fig. 2).

Only five vessels conducted sightings surveys in wind force class 1. This effort was extremely small compared to the effort in wind force classes 2 to 5, and was only 11% of wind force class 3, which had the largest effort. Wind force which showed maximum effort varied by vessel; in the case of *Shoyo Maru*, wind force exceeding class 6 showed maximum effort. Although as a general trend the number of sightings decreased as wind force increased, the number of sightings also varied by type of vessel and kind of marine debris.

Luminous Intensity

Time of day was used as an index of luminous intensity, and data from the same time of day were compared for sighting of marine debris (Fig. 3). Using the average value of the same eight vessels, the number of sightings decreased after 1200 (time of the maximum value); the rate of decrease remained within 60%, except at 1700. Five vessels showed the maximum value between 1200 and 1400, but for the *Hokko Maru* the maximum value was obtained at 1700, and for the *Shoyo Maru* the maximum value was obtained at 0600. These findings suggest that the number of sightings by time were related to many elements, and no clear trend by time was recognized. It is considered that there was no time of day at which it was extremely difficult to find marine debris.

Sighting Rate by Distance

In this survey, we usually observed at close range from the stern the distances and angles each debris items sighted. The distance at right angles to the track of the vessel (right angle distance) was measured, and marine debris was collected by category (Fig. 4). The number of sightings decreased as the right angle distance increased, and the number of sightings per 10 m accounted for <5% when the distance exceeded 100 m. Therefore, in estimating the number of sightings, the sighting width of 200 m, 100 m on each side of the track, was also estimated.

The relationship between right angle distance and the number of sightings of marine debris indicated a distribution with the maximum value of 10 to 20 m except for pieces of wood and drifting logs. Since the position of the marine debris never changed and never showed any movement against the vessel, it was easier to spot marine debris that was closer to the observers. If an observer could stop and scan the sea completely, the number of sightings would likely be in proportion to the distance. However, when sightings are conducted from a moving vessel it is not always possible to find marine debris close to the observer. As the vessel is sailing, the closer the debris is to the observer, the shorter the time in which it remains in the observer's visual field. Also it is not possible




Figure 2.--Number of marine debris pieces sighted per 100 nmi in terms of wind force (Beaufort scale) for each vessel.




Figure 3.---Number of marine debris pieces sighted in terms of time of observation for each vessel.




Figure 4. --Number of marine debris pieces sighted in terms of right angle distance (in meters) by type of debris.

to find marine debris on the sea when it is hidden behind the vessel. It is thought that there is a preferred distance for finding marine debris when the sighting is conducted from a fixed position. This was determined by the fact that in comparison with the sightings in Japanese waters and those in the Pacific Ocean, the number of sightings at a distance of 10 to 20 m tended to be larger than the number of sightings at a distance of 0 to 10 m.

Vessels conducting surveys in the Pacific Ocean were larger in size and had higher speeds than vessels conducting surveys in Japanese waters only. For the above two reasons, it is considered to be more difficult to observe the surface of the sea that is closer to the vessel. In addition, a decrease in the number of sightings in the range of 0 to 10 m was also caused by errors which arose from rounding to the nearest whole number when the angles sighted were reported. That is, as an angle was measured with the eye, the article which was recognized in the range of 0° to 5° was mostly reported as at 5°. If a distance sighted exceeded 115 m, marine debris was located from 10 to 20 m in right angle distance. For the optimum distance sighted in the relationship between right angle distance and number of sightings, it is necessary to collect more data and to continue further studies. In this report, the effective width was calculated from the assumption that the sighting probability on the path is 1.

Distribution Density of Marine Debris

Relationship between right angle distance (Y) and sighting rate (g(Y)) is shown in the following curvilinear equation:

$$g(Y) = 1 - \text{Exp} \left(-(Y/A)^{(1-B)} \right).$$

The coefficients of each type of marine debris are shown in Table 3.

The number of individual items per unit area for each type of marine debris was calculated by blocks (5° of latitude by 10° of longitude) on the basis of the following equation (Seber 1982):

$$N = \frac{nf(0)}{2L} .$$

- N = Number of individuals per unit area.
- n = Number sighted.
- L = Steaming distance.
- f(0) = 1/effective width.

Figure 5 shows the number of individual items per unit area by block obtained in this manner, and by the type of marine debris.

Table 3.--Coefficients of each type of marine debris.

Type of marine debris	A	B	f(0)
Fishing net	46,099	4,178	0.01752
Other fishing gear	39,623	3,627	0.01923
Styrofoam	41,598	3,969	0.01891
Other petrochemical products	46,366	4,613	0.01779
Pieces of wood and drifting logs	38,330	4,165	0.02060
Floating seaweed	47,890	5,679	0.01797
Other	31,485	3,565	0.02331
Total debris	38,162	3,430	0.01951

DISCUSSION

Distribution of Effort

A glance at the distribution of effort by block tells its own story: The blocks where effort was expended abundantly were concentrated in Japanese waters (Fig. 6). There were three blocks in which the survey distance exceeded 10,000 nmi, and the blocks which exceeded 3,000 nmi were also restricted to Japanese waters and adjacent areas. Next to the Japanese waters and adjacent areas, the offshore area of California, the southern area of the Alaska Peninsula, and the Northwestern Hawaiian Islands were also areas in which a large amount of effort was expended. However, the former two areas were also completely surveyed, one by the *Kaiyo Maru* only and the other by the *Toko Maru* and the *Shin Riasu Maru*, respectively, and the survey season was biased. As the Northwestern Hawaiian Islands are in the path of vessels which come and go from Honolulu, the area of survey is restricted.

Furthermore, glancing at numbers sighted by block, even blocks of other plastic debris, which has the most abundant sightings, the blocks in which 50 or more petrochemical items were found were only 18.4% of the total number of blocks in which petrochemicals were found. In order to obtain reliable density of marine debris, ideally speaking, it is necessary to conduct surveys evenly throughout the blocks in each season. As we mentioned before, in the present surveys, effort is frequently biased by season and by block, and the number of reliable blocks are extremely few. However, we calculated tentative density using the results of sightings as they were obtained.

Distribution of Marine Debris by Type

There were only 310 individual sightings of fishing nets, and reliable results were not obtained. However, the blocks in which sighting density was high were from lat. 25° to 40°N and long. 170° to 130°W, and in that area the density increased toward the east. Sighting density was next highest in Japanese waters and the East China Sea, but was only 2% of the block in which the density was the highest. In waters of lat. 45° to 50°N,




Figure 5A. --- Estimated density distribution of fishing net debris in 1987.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 5B.--Estimated density distribution of trawl net debris in 1987.
Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 5C. --Estimated density distribution of drift net debris in 1987.
Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 5D.--Estimated density distribution of other fishing gear debris in 1987.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 rmi².




Figure 5F.--Estimated density distribution of other plastic debris in 1987.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 5H. --- Estimated density distribution of floating seaweed debris in 1987.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi^2 .




Figure 5J.--Estimated density distribution of total marine debris in 1987.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi^2 .

fishing nets were not found at all. The distribution split to both south and north with this area in between, and it was thought that fishing nets found in the south were transported by currents moving from west to east. There were many unidentified fishing nets, and characteristics by kind of fishing net were not clearly recognizable. In waters north of lat. 50°N, only trawl nets were identified.

The numbers of other fishing gear sighted were greater than the number of fishing nets, and other fishing gear was found in 70.1% of all the areas surveyed. Although a distribution pattern of other fishing gear was similar to that of fishing nets, the blocks in which density was high inclined toward the south.

A great number of Styrofoam pieces were sighted. The range of distribution was widest, Styrofoam items being found in 77.8% of the blocks in the area surveyed. The distribution pattern was different for petrochemical articles other than Styrofoam, and the areas in which density was high were found in waters off Japan, at lat. 25° to 35°N and long. 170°W, lat. 25° to 35°N and long. 160°E to 140°W, in the Gulf of Mexico, and in the coastal areas of Central America. Areas which showed a comparatively high density were scattered widely. To explain the difference in this distribution pattern, petrochemical articles except Styrofoam are transported mainly by ocean currents, while Styrofoam items are floating on the surface of the sea and are thought to be strongly influenced by wind.

Other plastic debris was sighted in the greatest numbers (8,544 items), and the number of blocks sighted was the same as for other fishing gears. The distribution pattern was also similar. Six blocks in which the density was highest were concentrated in the range of lat. 20° to 35°N and long. 160°E to 130°W, followed by blocks in Japanese waters. In addition, an area in which the density was extremely high was in the Gulf of Mexico as well as the coastal areas of Central America.

For pieces of wood and drifting logs, densities were high in the coastal areas, suggesting that pieces of wood and drifting logs come primarily from the rivers and coastal areas. Floating seaweed showed this trend remarkably, and beyond three coastal blocks it was not found at all.

Blocks of highest density of combined petrochemical articles were seen in the coastal areas of Central America, followed by blocks of high density concentrated in waters of lat. 20° to 35°N and long. 150° to 130°W. Although the number of blocks was small, there were also those that showed high density in waters of lat. 25° to 35°N and long. 170°E to 170°W. Furthermore, densities of marine debris that were <2% of the highest density block, could be found in Japanese waters and the East China Sea, but a considerably high density was shown in the wide range. As another distinctive phenomenon, density was low in any blocks in waters of lat. 45° to 50°N, and the North Pacific Ocean and the Bering Sea are separated by this area. It is believed that marine debris seldom passes from one of these areas to the other.

Figures used in the above determinations were the numbers of individual items sighted. When considering the effects of marine debris,




Figure 7.--Surveyed cruising distance in each 5° x 10° area in 1986. Unit: nautical mile.




Figure 8A. --Estimated density distribution of fishing net debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi^2 .




Figure 8C.--Estimated density distribution of Styrofoam debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 8D.--Estimated density distribution of other plastic debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi^2 .




Figure 8F. --- Estimated density distribution of floating seaweed debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi^2 .




Figure 8G.--Estimated density distribution of other marine debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 nmi².




Figure 8H.--Estimated density distribution of total marine debris in 1986.
 Unit: number of debris pieces $\times 10^{-1}$ per 1 km^2 .

not only the number of individual items, but also their volume are important elements. However, it is quite difficult to measure with the eye the volume of things having various shapes. Therefore, in the survey we set these very rough size criteria and recorded the sizes of marine debris. Judging from the results by type of marine debris, "small" showed an extremely high rate, except in fishing net, pieces of wood, and drifting logs. In particular, "small" accounted for >90% of other plastic debris and Styrofoam, which were also great in actual volume. Although the number of items of this type of marine debris was great, it is believed that there was no greater difference in quantity than in number between this marine debris and other marine debris. More than half of the "large" items were fishing nets; the number was small, but the volume of each item was large. It is necessary to obtain more information on size in future surveys. It is believed that pieces of wood, drifting logs, and floating seaweed, which occur naturally, constitute the bulk of marine debris because of their large quantity and relatively large size.

These distribution patterns were almost the same as those obtained from the experimental sighting surveys conducted in 1986 (Figs. 7 and 8). It is necessary to study relationships between movement and accumulation of marine debris and ocean currents as well as to collect more data in the future. Furthermore, in order to understand yearly changes, it is also necessary to intensify the surveys in the North Pacific Ocean and adjacent areas and to establish methods of monitoring.

Yagi and Nomura (1988) reported on yearly changes in the density of marine debris based on sighting surveys conducted by the *Ryofu Maru* of the Meteorological Agency twice in winter and summer during 1976-86 using observations lines fixed between the Equator and lat. 34°N along long. 137°E. The survey results are said to be valuable for examining the yearly changes in marine debris using the same blocks at fixed periods each year, although observation blocks were limited in number. The survey results showed that the number of marine debris pieces sighted by unit distance more than doubled from when the survey was first launched. In particular, plastic sheet fragments have shown a marked increase in recent years.

REFERENCES

- Mio, S., and S. Takehama.
1987. Distribution of marine debris based on sighting survey in 1986. Document submitted to the Thirty-fourth Annual Meeting of the International North Pacific Fisheries Commission.
- Seber, G. A. F.
1982. The estimation of annual abundance and related parameters. Chas. Griffin & Co., Ltd.
- Yagi, N., and M. Nomura.
1988. Sighting survey on marine debris in the midwestern Pacific from 1977 through 1986. In D. L. Alverson and J.A. June (editors) Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 130-142. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

THE QUANTITATIVE DISTRIBUTION AND CHARACTERISTICS OF
NEUSTON PLASTIC IN THE NORTH PACIFIC OCEAN, 1985-88

Robert H. Day
Alaska Biological Research
P.O. Box 81934
College, Alaska 99708, U.S.A.

David G. Shaw
Institute of Marine Sciences
University of Alaska
Fairbanks, Alaska 99775, U.S.A.

and

Steven E. Ignell
Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

ABSTRACT

The distribution, abundance, and characteristics of neuston plastic in the North Pacific, Bering Sea, and Japan Sea were studied during the 4-year period 1985-88 at 203 neuston stations encompassing ca. 91,000 m² of sampling. The highest total density of neuston plastic was 316,800 pieces/km² at lat. 35°59'N, long. 152°00'E in Transitional Water east of Japan. The highest total concentration of neuston plastic was 3,491.8 g/km² at lat. 40°00'N, long. 171°30'E near the Subarctic Front in the central North Pacific. Main types of neuston plastic were miscellaneous line fragments (21.7% of all stations), Styrofoam (12.8%), polypropylene line fragments (7.4%), miscellaneous or unidentified plastic (7.4%), and raw pellets (5.9%). Plastic fragments were recorded at 52.2% of all stations and at 88.3% of those stations with plastic. The highest densities (number per square kilometer) and concentrations (gram per square kilometer) of neuston plastic occurred in Japan Sea/nearshore Japan Water, in Transitional Water, and in Subtropical Water. Densities of neuston plastic in Subarctic Water and Bering Sea Water were low. Heterogeneous geographic input and currents and winds are important in distributing and concentrating neuston plastic. Microscale convergences appear to be important mechanisms that locally concentrate neuston plastic, increasing the probability of its entering food chains.

INTRODUCTION

Marine debris, especially plastic debris, increasingly is recognized as a national and international pollution problem (Shomura and Yoshida 1985; Wolfe 1987). Plastic enters the ocean in many forms and many sizes. In addition to plastic objects associated with ships (e.g., lines, nets, floats), virtually every kind of plastic packaging and plastic object used on land may be discarded or lost to the sea. Some plastics are denser than seawater and thus sink, but some are buoyant enough to float, either because of trapped gas or because of low specific gravity. At sea, plastic objects undergo mechanical breakdown or fragmentation, leading to progressively smaller pieces of floating plastic. The size fraction of plastic debris caught in nets designed to catch surface plankton (hereafter referred to as neuston plastic) is of interest for several reasons. First, small plastic objects are more abundant than are the larger ones from which they are formed. Second, collection of plastic in nets is an objective process that provides unbiased estimates of densities. Finally, objects in this size range can be mistaken for food items, with possibly important ecological consequences (Day 1980; Day et al. 1985).

Several workers have investigated the distribution of neuston plastic in the North Pacific (Wong et al. 1974; Shaw 1977; Shaw and Mapes 1979; Day et al. 1985; Day and Shaw 1987). These studies have shown that neuston plastic is widespread, is most abundant in the central and western North Pacific, and is distributed by currents and winds.

The goal of this study was to improve our knowledge of the quantitative distribution and characteristics of neuston plastic in the North Pacific Ocean. Specifically, we wanted to: (1) describe the quantitative distributions of the main types of neuston plastic, (2) compare the at-sea densities of the main neuston types, (3) describe the frequencies of colors of neuston plastic, and (4) examine the importance of currents and winds in affecting the quantitative distribution of neuston plastic. Because of the extensive geographic coverage of the work, this study provides one of the most detailed synoptic pictures of neuston plastic anywhere in the world ocean.

METHODS

We collected data on the density, concentration, and types of neuston plastic ≥ 0.500 mm in size at 203 neuston stations in the North Pacific Ocean north of lat. 21°N (i.e., Hawaii) and in the Bering and Japan Seas. At each station, a 1.3-m ring net (during 1985) or a Sameoto (Sameoto and Jaroszynski 1969) neuston sampler (1986-88) with a 0.500-mm mesh net was used to collect neuston samples. Following Day and Shaw (1987), the area of ocean's surface sampled was calculated by multiplying the width of the net opening (0.5 m for the Sameoto sampler; see Day and Shaw 1987 for information on the ring net) by the distance the ship traveled in 10 min of sampling at a known speed, corrected for the time that the net was not fishing. Samples were washed from the net and either were sorted on the ship or were preserved in formalin and sorted later in the laboratory. Although areas sampled varied among stations, we ignored these differences

among stations in the analyses. Data from 1985 that already were published (32 stations, Day and Shaw 1987) were included here because that number is small compared with the 171 stations for which the data have not been published.

During sorting, individual pieces of plastic were counted and identified as one of six standardized types: pellet, fragment, Styrofoam (which may include foamed plastics of other chemical composition), polypropylene line (which may include synthetic line of other chemical composition), miscellaneous or unidentified line, and miscellaneous or unidentified plastic. These pieces of plastic also were identified as 1 of 11 standardized colors: black/gray, blue, brown, green, orange, red/pink, tan, transparent, white, yellow, and mixed or unidentified. The samples then were placed in preweighed vials and were air-dried before being weighed to the nearest 0.001 g.

Data were compiled as the total density (number per square kilometer) and total concentration (mass per square kilometer) of neuston plastic at each station and as the density of each general type of plastic at each station. The color data were compiled as the numbers and frequencies of occurrence of each color at each station and were tabulated as total frequencies of each color. For data analysis, each station was stratified geographically into one of five water masses: Bering Sea Water, Subarctic Water (north of the Subarctic Front, or north of ca. lat. 42°N), Subtropical Water (south of the Subtropical Front, or south of ca. lat. 31°N), Japan Sea/nearshore Japan Water (the latter area consisting of water east of Japan and west of lat. 150°E), and Transitional Water (that between Subarctic Water and Subtropical Water, and including the Subarctic Frontal Zone, the Transition Zone, and the Subtropical Front).

The stratified data on total density, total concentration, and densities of each type of neuston plastic were analyzed with a Kruskal-Wallis test (Conover 1980; Zar 1984). For each data set, we tested the hypothesis:

H_0 : The density (or concentration) does not differ among water masses.

When test results were significant, we conducted multiple comparisons tests (Conover 1980) to determine which water masses were different. We also calculated means and standard deviations of each data set in each water mass. The color data were compiled as frequencies of each color of plastic. Subsequently, these frequencies were divided by the total number of plastic items to determine percentages of each color type.

RESULTS

Neuston plastic was recorded at 120 stations (59.1% of total stations); the total number of pieces recorded was 1,774. The two water masses in which plastic occurred at 100% of the stations were Subtropical Water (n = 2 stations) and Japan Sea/nearshore Japan Water (n = 11 stations). Neuston plastic also was common in Transitional Water, where it

occurred at 56 (93.3%) of 60 stations, and in Subarctic Water, where it occurred at 46 (71.9%) of 64 stations. Finally, it was uncommon in Bering Sea Water, where it occurred at only 5 (7.6%) of 66 stations.

Total Density

Total densities of neuston plastic were highest in the Japan Sea, in nearshore water east of Japan, and in Transitional Water and the Subarctic Front; total densities generally were very low in Subarctic Water (especially in the center of the Alaska Gyre) and in the Bering Sea (Fig. 1). The highest total density of neuston plastic was 316,800 pieces/km² at lat. 35°59'N, long. 152°00'E in Transitional Water east of Japan. Other stations with high total densities were 221,000 pieces/km² at lat. 38°55'N, long. 135°58'E in the Japan Sea; 217,300 pieces/km² at lat. 37°58'N, long. 52°00'E near the Subarctic Front east of Japan; and 202,700 pieces/km² at lat. 40°00'N, long. 174°30'E near the Subarctic Front in the central North Pacific. Total densities differed significantly among water masses (H = 1221.482; n = 203; df = 4; P < 0.05; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Subtropical Water = Transitional Water > Subarctic Water > Bering Sea Water.

Concentration

Total concentrations of neuston plastic generally were low, with high concentrations recorded at only four stations in Transitional Water, at two stations in nearshore water east of Japan, and at one station in Subarctic Water; total concentrations at the other stations with plastic generally were <10% of the highest concentration (Fig. 2). The highest total concentration was 3,941.8 g/km² at lat. 40°00'N, long. 171°30'E near the Subarctic Front in the central North Pacific. Other concentrations >1,000 g/km² were 3,007.9 g/km² at lat. 37°58'N, long. 152°00'E near the Subarctic Front east of Japan, 1,979.1 g/km² at lat. 35°59'N, long. 152°00'E in Transitional Water east of Japan, and 1,048.5 g/km² at lat. 28°20'N, long. 162°20'W in Subtropical Water north of the Hawaiian Islands. Total concentrations differed among water masses (H = 120.604; n = 203; df = 4; P < 0.05; Table 1). Multiple comparisons indicated that concentrations were: Subtropical Water = Japan Sea/nearshore Japan Water = Transitional Water > Subarctic Water > Bering Sea Water. The similarity in patterns between total densities and total concentrations is understandable, considering the strong correlation between these two parameters (Spearman's R = 0.905; Z = 12.861; n = 203; P < 0.05; Conover 1980; Zar 1984). The Pearson's product-moment correlation between these parameters was not as high, however (r = 0.544; n = 203; P < 0.05).

Pellets

In the plastics industry, plastic resins commonly are manufactured as cylindrical pellets a few millimeters in size. Later, these pellets are melted and molded into finished products. Pellets were uncommon, being recorded only 12 times (5.9% of total stations and 10.0% of stations with plastic). Pellets were absent in the Bering and Japan Seas, were recorded only once in Subarctic Water, and were recorded primarily in Transitional




Figure 1.--Total densities of neuston plastic, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 316,800 pieces/km².

Table 1.--Densities (number per square kilometer) and concentrations (grams per square kilometer) of neuston plastic in five water masses of the North Pacific, 1985-88.

Parameter	Bering Sea Water		Subarctic Water		Transitional Water		Subtropical Water		Japan Sea and nearshore Japan Water	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Number		66		64		60		2		11
Area sampled (m ²)	35,906		28,662		22,154		541		3,824	
Total concentration	1.0	4.2	61.4	225.5	291.6	714.4	535.1	726.1	128.2	172.2
Total density	100	600	12,800	22,300	57,900	72,800	61,000	74,000	74,700	73,800
Pellet	0	0	<100	300	300	800	3,300	4,600	500	1,200
Fragment	0	0	9,600	20,300	52,700	69,200	57,700	69,400	46,100	40,000
Styrofoam	0	0	400	1,300	1,100	3,200	0	0	26,200	37,200
Polypropylene line	100	400	400	1,500	500	1,500	0	0	0	0
Miscellaneous line/thread	100	300	2,600	6,900	2,300	4,600	0	0	1,900	3,300
Miscellaneous/unidentified	100	500	100	500	1,000	3,100	0	0	0	0




Figure 2.--Total concentrations of neuston plastic, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest concentration was 3,941.8 g/km².

Water and in nearshore water east of Japan (Fig. 3). The highest density was 6,500 pieces/km² at lat. 28°20'N, long. 162°20'W in Subtropical Water north of the Hawaiian Islands. The density of pellets differed among water masses ($H = 22.996$; $n = 203$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons were confusing, however, in that none of the individual water masses were significantly different. We suspect that the significant result was an artifact of the presence of pellets at both of the two stations in Subtropical Water. Consequently, the mean rank in this water mass was much higher than those in the other water masses, although the small sample size made it impossible to prove that significant differences actually existed.

Fragments

Fragments are small pieces of plastic broken from larger pieces (excluding Styrofoam). This category included primarily chips and pieces of sheets. Fragments were common, being recorded at 106 stations (52.2% of total stations and 88.3% of all stations with plastic). Fragments were common except in the Bering Sea and occurred in highest densities in nearshore water east of Japan and in and around the Subarctic Front; densities were lower in the Japan Sea and Subtropical Water and were much lower in Subarctic Water (Fig. 4). The highest density was 288,000 pieces/km² at lat. 35°59'N, long. 152°00'E in Transitional Water east of Japan. Other stations with high densities of fragments were 202,700 pieces/km² at lat. 40°00'N, long. 174°30'E near the Subarctic Front in the central North Pacific; and 199,000 pieces/km² at lat. 37°58'N, long. 152°00'E near the Subarctic Front east of Japan. The density of fragments differed significantly among water masses ($H = 113.587$; $n = 203$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water = Subtropical Water = Transitional Water > Subarctic Water > Bering Sea Water.

Styrofoam

This category included all pieces of pieces of foamed plastic; based on observed color and texture, we believe that all of this plastic was polystyrene. Styrofoam was uncommon, being recorded only 26 times (12.8% of total stations and 21.7% of stations with plastic). It was recorded in all locations except the Bering Sea and Subtropical Water, and occurred in highest densities in the Japan Sea and nearshore water east of Japan. It was a "transitional/nearshore Japan species," being recorded outside of this area only five times (Fig. 5). The highest density was 99,500 pieces/km² at lat. 36°37'N, long. 131°54'E in the Japan Sea. Other stations with high densities were 82,200 pieces/km² at lat. 38°55'N, long. 135°58'E in the Japan Sea; and 65,400 pieces/km² at lat. 34°49'N, long. 144°55'E off the eastern coast of Japan. Densities of Styrofoam differed significantly among water masses ($H = 52.967$; $n = 203$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons indicated that densities were: Japan Sea/nearshore Japan Water > Transitional Water = Subarctic Water = Subtropical Water.




Figure 3.--Densities of pellets, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 6,500 pieces/km².




Figure 4.--Densities of fragments, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 288,000 pieces/km².




Figure 5.--Densities of Styrofoam, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 99,500 pieces/km².

Polypropylene Line Fragments

Polypropylene line fragments are small, woven pieces of large synthetic lines that are used as deck lines on fishing boats and cargo ships. Polypropylene is the most commonly used plastic for these applications. These line fragments were uncommon, being recorded 15 times (7.4% of total stations and 12.5% of stations with plastic). Polypropylene line fragments occurred primarily in and near the Subarctic Front and in Transitional Water; they were absent in the Japan Sea and in Subtropical Water (Fig. 6). The highest density was 8,400 pieces/km² at lat. 41°09'N, long. 170°00'E near the Subarctic Front in the central North Pacific. We failed to reject the null hypothesis that the density of polypropylene line fragments did not differ significantly among water masses ($H = 3.597$; $n = 203$; $df = 4$; $P > 0.05$; Table 1), probably because densities were low everywhere.

Miscellaneous Lines/Threads

Miscellaneous lines and threads included unidentified woven line fragments and (especially) monofilament lines that were from either gillnets or monofilament fishing line. We do not know what type of plastic they were, but they probably were not nylon, as it does not float (Carpenter 1976). Miscellaneous lines/threads were somewhat common, being recorded 44 times (21.7% of total stations and 36.7% of stations with plastic). They were recorded in all but Subtropical Water, with the highest densities occurring east of Japan and near the Subarctic Front (Fig. 7); they possibly may be fragments of line used by squid jiggers, which fish in this area. The highest density was 40,500 pieces/km² at lat. 47°59'N, long. 166°41'E in western Subarctic Water. Densities of miscellaneous lines/threads differed significantly among water masses ($H = 24.607$; $n = 203$; $df = 4$; $P < 0.05$; Table 1). Multiple comparisons were confusing, however, in that those water masses with the largest difference in mean ranks were not significantly different, whereas water masses with smaller differences in mean ranks were significantly different. The two water masses that were significantly different were Transitional Water > Bering Sea Water, two with large sample sizes (60 and 66, respectively). We suspect that other water masses were different but that sample sizes in most were too small for the multiple comparisons to show significant differences. The pattern of mean ranks (in descending order) was: Japan Sea/nearshore Japan Water, Transitional Water, Subarctic Water, Bering Sea Water, and Subtropical Water.

Colors of Neuston Plastic

Most neuston plastic was transparent. This color was recorded 785 times (44.3% of the total 1,774 pieces and 44.9% of plastic of identified color). White plastic also was abundant, being recorded 610 times (34.4% of the total and 34.9% of plastic of identified color), followed by blue (128 pieces; 7.2% of the total and 7.3% of plastic of identified color), black/gray (74 pieces; 4.2% and 4.2%), green (62 pieces; 3.5% and 3.5%), and tan (45; 2.5% and 2.6%). The colors brown (17 pieces; 1.0% and 1.0%), red/pink (13 pieces; 0.7% and 0.7%), yellow (8 pieces; 0.5% and 0.5%), and orange (5 pieces; 0.3% and 0.3%) were rare in occurrence. Miscellaneous or unidentified colors occurred 27 times (1.5%).




Figure 6. --Densities of polypropylene line fragments, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 8,400 pieces/km².




Figure 7.--Densities of miscellaneous lines/threads, 1985-88. Solid black circles indicate stations at which neuston plastic was not recorded. Sizes of hollow circles indicate relative densities. The highest density was 40,500 pieces/km².

DISCUSSION

The distribution of neuston plastic results from two main phenomena, heterogeneous geographic input of plastic and subsequent redistribution by currents and winds. In addition, a phenomenon of unknown importance is the in situ decomposition of plastic in the ocean.

It appears that there is heterogeneous geographic input of neuston plastic, with much of it originating in the western Pacific. This conclusion is indicated by the high densities in and around the Japan Sea and nearshore Japan, where the highest densities of both neuston plastic and marine debris (Day et al. 1990) were recorded. The most polluted water in this area were Tokyo Bay (which had far more plastic than Day has ever seen elsewhere in the Pacific--he was unable to sample there) and localized areas in the Japan Sea. At the other extreme was the poorly populated Bering Sea area, where low rates of input probably occur. The low human population around much of the Gulf of Alaska probably contributes to the low densities there, also.

After entering the ocean, however, neuston plastic is redistributed by currents and winds. For example, plastic entering the ocean in Japan is moved eastward by the Subarctic Current (in Subarctic Water) and the Kuroshio (in Transitional Water, Kawai 1972; Favorite et al. 1976; Nagata et al. 1986). In this way, the plastic is transported from high-density areas to low-density areas. In addition to this eastward movement, Ekman stress from winds tends to move surface waters from the subarctic and the subtropics toward the Transitional Water mass as a whole (see Roden 1970: fig. 5). Because of the convergent nature of this Ekman flow, densities tend to be high in Transitional Water. In addition, the generally convergent nature of water in the North Pacific Central Gyre (Masuzawa 1972) should result in high densities there also.

One point that is not entirely clear is the cause of the low densities of neuston plastic in Subarctic Water. Part of the reason for these low densities is the apparently low input from shipping in this area: densities of both neuston plastic and marine debris in this area are low, suggesting little input from ships. The role of the divergent Alaska Gyre in helping to maintain these low densities is unclear, however. For example, neuston plastic tends to concentrate near the edges of the subarctic water mass, with little occurring in its center (Fig. 1), as would be expected for an upwelling gyre. On the other hand, the rate of vertical advection (in the low hundreds of meters/year, with downwelling occurring much of the year; T. C. Royer, Institute of Marine Sciences, University of Alaska, Fairbanks, Alaska, pers. commun.) is much lower than the rate of lateral advection (ca. 3,000 km/year at a speed of 15 cm/sec; Favorite et al. 1976), which should result in upwelling having little effect on the distribution of neuston plastic in this gyre.

A third factor, and one of unknown importance, is the in situ decomposition of larger marine debris plastic into small neuston plastic. As discussed by Day and Shaw (1987), the small percentage of raw plastic pellets and the high correlation between abundances of debris plastic and

neuston plastic suggested that in situ decomposition was occurring. Although the present study did not test this hypothesis, we believe that the in situ decomposition of plastic can be important. The large pool of debris plastic and neuston plastic (particularly fragments) in Transitional Water probably is resident for a long period of time and appears to be decomposing there. For example, our impression was that transparent neuston plastic in this area tended to be opaque on the surface, to have more surface crazing (Gregory 1978, 1983), and to be more brittle than did most from Subarctic Water, where it tended to be more transparent on the surface and more pliable. The same phenomenon was true for much of the marine debris plastic in Transitional Water, where it was heavily bleached and heavily encrusted, suggesting long residence time. In reality, however, chemical weathering (leaching of plasticizers from the plastic matrix, causing the remaining plastic to be brittle and more susceptible to mechanical weathering), thermal weathering (increasing the rate of chemical weathering), and solar weathering (from strong sunlight) probably are most important in the in situ production of fragments of neuston plastic in Transitional and Subtropical Waters, whereas mechanical weathering (from rough seas) probably is most important in stormier Subarctic Waters. Finally, thermal (i.e., freezing) and mechanical weathering probably are most important in the stormy, cool Bering Sea, which is ice-covered in winter.

Frequencies of colors of neuston plastic in the North Pacific differed from frequencies of colors of neuston plastic ingested by seabirds (Day et al. 1985). For example, white, yellow, tan, and brown neuston plastic (light colors) represented only 40.0% of total identified neuston plastic in the ocean, whereas it represented 85.0% of neuston plastic ingested by seabirds. One of the largest differences was in tan plastic, which composed only 2.6% of the identified neuston plastic in the ocean but 55.1% of the neuston plastic eaten by seabirds. The largest difference was in transparent plastic, which represented 44.9% of the identified neuston plastic in the ocean but was not found in seabirds. Transparent plastic is not eaten by birds, probably because of difficulty in seeing it at sea (Day et al. 1985).

Neuston plastic can enter food chains when it is mistaken for prey (Day et al. 1985), especially where it becomes concentrated near important, localized prey. For example, there appeared to be a relationship between high densities of neuston plastic and high densities of water-striders, *Halobates sericeus* (Insecta: Gerridae) in Transitional and Subtropical Waters. These marine insects live at the surface of the ocean and are eaten by at least nine species of tropical seabirds that breed in the Hawaiian Islands and feed in these water masses. Water-striders are especially important prey of blue-gray noddies, *Procelsterna cerulea*, Bulwer's petrels, *Bulweria bulwerii*, and Bonin petrels, *Pterodroma hypoleuca*, with the latter two species also containing significant amounts of neuston plastic (Harrison et al. 1983; Cheng et al. 1984). We suspect that these insects are moved slowly into microscale convergences at the same time that plastic and other organisms are. For example, the density of water-striders was 136,000/km² at one station where the density of neuston plastic was 113,300 pieces/km²; the highest density of water-striders was

ca. 250,000/km² (Day unpubl. data). Given the co-occurrence of water-striders and neuston plastic in some tropical seabirds, we suggest that many of these birds are feeding in these microscale convergences, where they are picking up water-striders, other plankters, and neuston plastic. Indeed, Day has seen surface-feeding planktivorous seabirds (phalaropes and storm-petrels) feeding in large numbers in microscale convergences in the Oyashio-Kuroshio Confluence. These convergences contained visible lines of kelp wrack, plastic, and other marine debris.

Another group that ingests neuston plastic as well as planktonic prey in coastal and oceanic microscale convergences is sea turtles (Carr 1987). Young turtles apparently feed in these convergences during the first year or more at sea, when they drift with the currents and hence act much like neuston plastic. (During this period they also may become entangled in marine debris plastic.) Later, as they become older, these turtles both ingest larger pieces of marine debris plastic and become entangled in marine debris plastic (Balazs 1985).

Microscale convergences may be found in many areas of the world ocean (e.g., Owen 1981; Bourne and Clark 1984), and they may occur in areas different from the general areas of concentration discussed above. From our experience, microscale convergences concentrating neuston plastic are near lat. 28°-29°N north of Hawaii; in and near the Subarctic Front as microscale ephemeral convergences; in the complex Oyashio-Kuroshio Confluence east of Japan (including the ephemeral, mobile warm-core and cold-core rings; Nagata et al. 1986); at scattered locations in the Japan Sea; and probably in and around the Subtropical Front (i.e., around lat. 30°-32°N).

Perhaps the most impressive microscale convergences are in and around the Subarctic Front. Here, dynamic instabilities in surface layers (Roden 1970) create numerous ephemeral convergences in the zone lat. 37°-42°N and in the Oyashio-Kuroshio Confluence east of Japan. This juxtaposition of high biological productivity, physical complexity, large numbers of seabirds that ingest neuston plastic, and large amounts of neuston plastic increases the possibility of ingestion of that plastic.

ACKNOWLEDGMENTS

Most of the work was done while Day was an Angus Gavin Memorial Fellow, a Sea Grant Fellow, or a Resources Fellow at the University of Alaska. Additional funding came from the National Marine Fisheries Service (NMFS), NOAA (contracts 40ABNF63228 and 43ABNF702983) and two contracts from the Joint Institute for Marine and Atmospheric Research at the University of Hawaii. Additional samples were collected by James Seger and Richard Rowlett of NMFS and by Amy Stone of the U.S. Fish and Wildlife Service (FWS). Samples were collected aboard the TV *Oshoro Maru* of Hokkaido University, Hakodate, Japan; RV *Pusan 851* of the National Fisheries Research and Development Agency, Government of the Republic of Korea, Pusan, Korea; RV *Miller Freeman* of the Northwest and Alaska Fisheries Center, NMFS, Seattle, Washington, RV *Eagle* of the FWS, Homer, Alaska; and RV *Alpha Helix* of the Institute of Marine Sciences, University of Alaska,

Fairbanks, Alaska. We thank the captains, officers, and crews of these ships for assistance in collecting the samples. This manuscript was improved by comments from three anonymous reviewers. This is Contribution No. 827 of the Institute of Marine Sciences, University of Alaska, Fairbanks, AK 99775.

REFERENCES

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Bourne, W. R. P., and G. C. Clark.
1984. The occurrence of birds and garbage at the Humboldt Front off Valparaiso, Chile. Mar. Pollut. Bull. 15:343-344.
- Carpenter, E. J.
1976. Plastics, pelagic tar, and other litter. In E. D. Goldberg (editor), Strategies for marine pollution monitoring, p. 77-89. Wiley, N.Y.
- Carr, A.
1987. Impact of nondegradable marine debris on the ecology and survival outlook of sea turtles. Mar. Pollut. Bull. 18:352-356.
- Cheng, L., M. Schultz-Baldes, and C. S. Harrison.
1984. Cadmium in ocean-skaters, *Halobates sericeus* (Insecta), and in their seabird predators. Mar. Biol. (Berl.) 79:321-324.
- Conover, W. J.
1980. Practical nonparametric statistics, 2d ed. Wiley, N.Y., 493 p.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., and D. G. Shaw.
1987. Patterns in the abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-85. Mar. Pollut. Bull. 18:311-316.
- Day, R. H., D. G. Shaw, and S. E. Ignell.
1990. Quantitative distribution and characteristics of marine debris in the North Pacific Ocean, 1984-88. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154. [See this document.]

- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54. [See this document.]
- Favorite, F., A. J. Dodimead, and K. Nasu.
1976. Oceanography of the subarctic Pacific region, 1960-71. *Int. North Pac. Fish. Comm. Bull.* 33:1-187.
- Gregory, M. R.
1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwater Res.* 12:399-414.

1983. Virgin plastic granules on some beaches of eastern Canada and Bermuda. *Mar. Environ. Res.* 10:73-92.
- Harrison, C. S., T. S. Hida, and M. P. Seki.
1983. Hawaiian seabird feeding ecology. *Wildl. Monogr.* 85:1-71.
- Kawai, H.
1972. Hydrography of the Kuroshio Extension. In H. Stommel and K. Yoshida (editors), *Kuroshio: Physical aspects of the Japan Current*, p. 235-352. Univ. Wash. Press, Seattle, WA.
- Masuzawa, J.
1972. Water characteristics of the North Pacific central region. In H. Stommel and K. Yoshida (editors), *Kuroshio: Physical aspects of the Japan Current*, p. 95-127. Univ. Wash. Press, Seattle, WA.
- Nagata, Y., J. Yoshida, and H.-R. Shin.
1986. Detailed structure of the Kuroshio Front and the origin of the water in warm-core rings. *Deep-Sea Res.* 33:1509-1526.
- Owen, R. W.
1981. Fronts and eddies in the sea: Mechanisms, interactions, and biological effects. In A. R. Longhurst (editor), *Analysis of marine ecosystems*, p. 197-233. Acad. Press, N.Y.
- Roden, G. I.
1970. Aspects of the mid-Pacific Transition Zone. *J. Geophys. Res.* 75:1097-1109.
- Sameoto, D. D., and L. O. Jaroszynski.
1969. Otter surface sampler: A new neuston net. *J. Fish. Res. Board Can.* 25:2240-2244.
- Shaw, D. G.
1977. Pelagic tar and plastic in the Gulf of Alaska and Bering Sea. *Sci. Total Environ.* 8:13-20.

- Shaw, D. G., and G. A. Mapes.
1979. Surface circulation and the distribution of pelagic tar and plastic. Mar. Pollut. Bull. 10:160-162.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Wolfe, D. A. (editor).
1987. Plastics in the sea. Mar. Pollut. Bull. 18:303-365.
- Wong, C. S., D. R. Green, and W. J. Cretney.
1974. Quantitative tar and plastic waste distributions in the Pacific Ocean. Nature (Lond.) 247(5435):30-32.
- Zar, J. H.
1984. Biostatistical analysis, 2d ed. Prentice-Hall Inc., Englewood Cliffs, NJ, 718 p.
-

MOVEMENTS OF FLOATING DEBRIS IN THE NORTH PACIFIC

Satsuki Matsumura
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

Yoshinobu Wakata and Yasuhiro Sugimori
Faculty of Marine Science and Technology
Tokai University
Shimizu-shi, Shizuoka, 424 Japan

ABSTRACT

A net fragments tracking experiment and numerical simulations using surface current data set (SCUDS) data were conducted to estimate movements of floating debris in the North Pacific.

Six driftnet sets (40 tans each) were placed in the area lat. 39°N, long. 155°E. Locations of the net sets and sea surface temperatures were collected and transmitted every day using the Argos system. Data were taken about six times a day for 4 months. At termination of the net drifting experiment, the net sets with buoys were retrieved and new Argos buoys with curtain drogues were released at the points of retrieval to continue the surface current tracking.

The buoys moved predominantly eastward, although each track line was complicated, particularly in areas near the Oyashio Front. It is considered that the movements of the nets were mainly due to surface currents and that direct influence from wind was negligible, because the underwater portion was very large (a driftnet 2,000 m long although it had formed a mass) compared with the above-water portion of the buoy. Average speed was estimated based on the buoy movements and ranged from 10 km/day to 20 km/day. Movements of floating debris in the North Pacific were simulated using a computer model based on SCUDS.

Results showed the existence of two large-scale eddies in the eastern and western parts of the mid-Pacific, and floating debris are through to accumulate in these areas.

INTRODUCTION

North Pacific currents are found in a great circle ranging from the Kuroshio through the Kuroshio Extension and the North Pacific Current to the California Current and the North Equatorial Current in the south. In the vicinity of these currents are the Oyashio, Alaska, Aleutian, and other currents.

Floating debris (excluding Styrofoam, which is mostly above the surface) moves along these currents. Movements change depending on large and small vortexes in the water and are difficult to generalize, but we can estimate average movements on the basis of surface currents.

Marine features of the North Pacific are outlined by Favorite et al. (1976). The northwestern part of the North Pacific is characterized by the Kuroshio Extension and the Oyashio. The distribution of water masses in summer is greatly affected by the southerly intrusion of cold water masses and the strength of the Aleutian Current, both of which vary from year to year, as reported by Hiramatsu (1987, 1988).

Tests using driftnets were conducted to estimate movements of floating of floating debris. Driftnets were set in the Oyashio waters in May 1988 and were recovered after about 4 months of drifting. The Argos system was used for tracking the nets and analyzing their drift routes. We have also illustrated overall currents on a computer display using the surface current data sets (SCUDS), which covers ship drift data from about 4 million ship observations over the past 40 years.

CURRENT OBSERVATIONS USING ARGOS BUOYS

Method

Six driftnets were set by the first survey ship in order to observe changes in net shapes and movements in the northwestern part of the North Pacific. Each net was equipped with an Argos buoy at one tip and a radio buoy at the other. Nets were set in waters at lat. 39°N, long. 155°E and were arranged in the shape of a star, with five nets set in a 37-km (20-mi) radius from a key net.

Each net was 2,000 m (40 tans) long. Their fluid resistance in the water was so large that the early movements of each net were presumed to indicate an average current over a distance of 2,000 m. Ten days later, the shapes of several nets had changed greatly. Some nets were either folded or "balled up." They became masses with a maximum length of roughly 50 m (Mio et al. 1990).

They were still large enough to resist winds, and buoy tracks accurately indicated the movements of nets in the sea currents. There was a thermometer inside each buoy case. The buoy was metallic and had no heat-resistant structure. Therefore, each thermometer was able to indicate surface water temperatures. Styrofoam was used as a floater at the upper part of each buoy. It reduced heat flow from sunlight. Even if the buoy

was exposed to strong sunlight, the thermometer's deviation from surface water temperature would have been very small.

Argos buoys were set on 5 and 6 May 1988. Three of the six buoys suspended transmission during drifting because of mechanical troubles, but the second survey ship recovered three of the four trouble-hit Argos buoys 4 months later by tracking radio buoys. The second survey ship recovered the nets and released three Argos buoys with curtain drogues for further observation of currents.

Results and Analysis

Tracks of buoys from May to September are illustrated in Figure 1.

Buoy 4783, which moved the greatest distance eastward in the 4 months, reach long. 180° . The slowest-moving buoy (4782) came as far as long. 162° E. An average speed of the maximum eastward movement was about 20 km/day. Any buoy movement was not straight but very complex depending on inertial forces and tides. The theoretically calculated inertial cycle at lat. 40° N is about 19 h, and it matched these observations. Figures 2-4 indicate the track of buoy 4789, which is zoomed gradually. Figure 4, where vertical and horizontal scales are almost the same, shows that the floating debris moved north to northeast at an average speed of 7 cm/sec while making a clockwise motion. The radius of the circular motion was about 1.6 km, and the flow speed was 10-50 cm/sec. Driftnets, which were set at roughly the same place, followed very different tracks, hinting at a large diffusion coefficient in the waters. One hundred days after setting, the maximum east-west distance between tracks was 1,637 km and the maximum




Figure 1.--Trajectories of drifting buoys.




Figure 2.--Western part of buoy 4780 trajectory.




Figure 3.--Part of buoy 4780 trajectory, enlarged from Figure 2.




Figure 4.--Inertia circle of buoy 4780, enlarged from Figure 3.

north-south distance was 1,021 km. The large diffusion indicates a need to incorporate a large plus or minus deviation in forecasting the movement of any object in the water on the basis of the generally known currents. The large diffusion coefficient gives evidence of great turbulence in the waters.

Data from thermometers on the buoys are shown in Figure 5. Any buoy with a large net cannot be expected to go through a water mass or across the front of a mass. Short-term changes in the water temperature, especially sudden declines, may be attributable to vertical mixing of waters.

The water temperature rose gradually during the observation period between May and September. Any sudden rise in the temperature may be attributed to a combination of strong sunlight and calm water, which can cause an increase in just the surface water temperature. It may also be ascribed to surface water which was heated by the sun and flowed into the vicinity of a buoy.

Driftnets remained at a depth of 10 m. Warm surface waters which are frequently seen in the northern part of the North Pacific in summer are usually limited to 1 or 2 m in depth.

The surface water temperature changes are seen frequently in infrared heat pictures taken by satellites and have some adverse effect on analysis of water masses, which depends on surface temperature.




Figure 5.--Daily changes in buoy temperatures (May-August 1988).

SIMULATION OF FLOATING DEBRIS MOVEMENT AND DEBRIS DENSITY USING SHIP DRIFT DATA

If the speeds of all sea currents in time and space are available, the track of floating debris like the buoy on currents such as those discussed in the previous section can be simulated. Based on a specific speed at an initial point, we can determine a point debris would reach within a given period of time. Another current speed at the arrival point can be used to estimate how fast and where the debris would move further. Repeating such estimates can lead to a possible track which floating debris would follow.

In this section, we try to solve the Lagrangean equation and simulate a buoy track on the basis of given sea current speeds. We have used the ship drift data released by Meehl (1982). The currents in any time and space can be obtained by interpolating these data. We have compared the simulated results with findings from buoy drift observations by Kirwan et al. (1978) and sightings of floating debris observed by the Fisheries Agency, the Government of Japan (Mio et al. 1990) in order to analyze the mechanism for the gathering of objects.

Our simulation results successfully matched the findings from Kirwan's observation in both time and space. The model may be available for the simulation of buoy movement.

Thus, in the following, we will discuss where buoys set in waters around Japan move and where a number of buoys set all over the Earth would gather.

Drifting in the Western North Pacific

Assume that buoys are initially dropped at lat. 30°N, long. 140°W near Torishima Island in the western part of the North Pacific. The results show how the buoy track changes depending on the season.

Spring

Figure 6(a) shows the track of a buoy set on 1 April. The buoy immediately begins to move eastward and reaches the international dateline in 6 months. However, the buoy remains around lat. 30°N, long. 150°W for nearly 2 years while its track loops. This is because it is dragged into a vortex in the eastern part of the North Pacific (Meehl 1982, fig. 2). Floating debris can remain in the water for a long time. If floating debris such as waste were dumped evenly all over the Pacific, it would tend to gather in the vortex waters.

Summer

Figure 6(b) shows the track of a buoy set on 1 July. It also moves eastward, but its speed is far faster than that of the spring buoy. It reaches the international dateline in about 4 months. Since the speed of currents at long. 140°E is fastest in autumn, the buoy set in summer is eventually moved by these fast currents. Later, it is dragged into the vortex in the eastern Pacific and remains just north of Hawaii.




Figure 6.--Simulated tracks of buoys which were set at lat. 30°N, long. 140°E in the western North Pacific in (a) spring (1 April), (b) summer (1 July), (c) autumn (1 October), and (d) winter (1 January). Each buoy took 3 months to move from one mark to another.

Autumn

The track in Figure 6(c) is for a buoy set 1 October. Unlike the above tracks, this one loops around lat. 30°N, long. 150°E before extending eastward. The Kuroshio Extension weakens, and a vortex and a southward current around lat. 30°N, long. 150°E appear in winter. A buoy dropped in autumn is dragged into this winter current. The loop indicates that floating debris stays in the water for a long time.

Winter

Figure 6(d) shows the track of a buoy put into the water on 1 January. Like the track of a buoy set in autumn, the winter buoy track includes a small loop in the western Pacific.

All of the above tracks loop in the eastern part of the North Pacific. This indicates that floating debris remains in the water for a long time and tends to gather there.

Drifting in the Eastern North Pacific

Figure 7(a,b,c) shows the tracks of buoys put into the water on 1 May. The three drop sites are lat. 50°N, long. 140°W; lat. 40°N, long. 130°W; and lat. 30°N, long. 120°W, respectively. The buoy in Figure 7(a) is set in the region of the Alaskan current system. It moves westward, traveling south of the Aleutian Islands around the Alaskan gyre and western subarctic gyre. After turning south it is picked up by the North Pacific current system and moves eastward as in Figure 6.

The buoy which is set at lat. 40°N (Fig. 7b) is carried westward by the North Equatorial Current system after remaining in the vicinity of lat. 30°N for nearly 2 years. The buoy set at lat. 30°N (Fig. 7c) immediately begins to move westward.

Debris Density

We calculated many tracks in the Pacific from starting points evenly distributed in space and time to find where buoys would gather. We tried to find not the place where a buoy dropped at a certain point would go, but the place where buoys would gather intrinsically due to currents, irrespective of setting points or time.

We simulated tracks for 7,755 buoys. Dropping points were chosen randomly so that the number of initial buoys would be the same for every unit area. The setting density was five buoys for every 3.09×10^{11} m² of ocean; buoy setting continued for 4 years. To scatter drop times evenly, they were chosen on the basis of random numbers. Presuming a lifespan of 2 years, we simulated a track for each buoy for 2 years.

Figure 8 indicates the debris density in January after 4 years of setting. In the sea, you see some points to which buoys were carried by currents. In the North Pacific, which is our area of concern, debris is




Figure 7.--Simulated tracks of buoys which were set at (a) lat. 50°N, long. 140°W; (b) lat. 40°N, long. 130°W, and (c) lat. 30°N, long. 120°W in the eastern part of the North Pacific on 1 May.




Figure 8.--Floating debris density in the North Pacific on 15 January. Contour line shows the number of buoys in unit area. Original density was give at every place.

seen to gather in the eastern part. The gathering points are to the north of Hawaii, which agrees with sighting observations of Mio et al. (1990). However, note that the gathering points shift seasonally. In the Northern Hemisphere, winds in areas of high atmospheric pressure circle to the right and sea surface currents deviate from the wind direction by 20° to 30°, as pointed out by McNally (1981). So we know that a high concentration of debris could be seen in the center of the North Pacific area of high atmospheric pressure.

CONCLUSION

The results of drift buoy observations matched fairly well those of the simulation based on ship drift data in waters like the Kuroshio Extension where the current is strong. Debris getting into the currents may cross the North Pacific in about 1 year.

There are no marked currents in waters around the Ogasawara Islands and northwest of Hawaii in any season. Figures for drift tracks and for distribution of floating debris point to those waters as locations where floating debris can gather.

REFERENCES

- Favorite, F., A. J. Dodimead, and K. Nasu.
1976. Oceanography of the subarctic Pacific region, 1960-71. *Int. North Pac. Fish. Comm. Bull.* 33:1-187.
- Hiramatsu, K.
1987. Outline of oceanographic condition of the Northwest Pacific during the summer of 1987. *Int. North Pac. Fish. Comm.*
1988. Outline of oceanographic conditions of the Northwest Pacific during the summer of 1988. *Int. North Pac. Fish. Comm.*
- Kirwan, A. D., G. J. McNally, E. Reyna, and W. J. Merrell.
1978. The near-surface circulation of the eastern North Pacific. *J. Phys. Oceanogr.* 8:937-945.
- McNally, G. J.
1981. Satellite-tracked drift buoy observations of the near-surface flow in the eastern mid-latitude North Pacific. *J. Geophys. Res.* 86:8022-8030.
- Meehl, G. A.
1982. Characteristics of surface current flow inferred from a global ocean current data set. *J. Phys. Oceanogr.* 12:538-555.
- Mio, S., T. Domon, K. Yoshida, and S. Matsumura.
1990. Preliminary study on change in shape of drifting nets experimentally placed in the sea. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii.* U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154.

TYPE, SOURCE, AND ABUNDANCE OF TRAWL-CAUGHT
MARINE DEBRIS OFF OREGON, IN THE EASTERN
BERING SEA, AND IN NORTON SOUND IN 1988

Jeffrey A. June
Natural Resources Consultants, Inc.
4055 21st Avenue West
Seattle, Washington 98199, U.S.A.

ABSTRACT

In 1988, National Marine Fisheries Service scientists collected information on type, source, and abundance of marine debris caught during annual bottom trawl surveys off Oregon, in the eastern Bering Sea, and in Norton Sound. Numbers of individual debris items caught were tallied by haul. When possible, the nationality of origin was determined. Animals entangled or associated with debris items were noted. Debris items were categorized by material (e.g., plastic, glass) and use (e.g., galley wastes, fishing equipment). Effort in square kilometers trawled was calculated for each haul from distance fished and average net width measurements. Average catch-per-unit-effort (CPUE) in numbers of items per square kilometer was calculated for individual debris items, major categories, and total debris by area and for combined areas.

Of the 696 hauls surveyed, 70 were off Oregon, 541 in the eastern Bering Sea, and 85 in Norton Sound. Marine debris was most abundant off Oregon, occurring in 70% of the hauls and averaging 149.6 items/km². In the eastern Bering Sea, 23% of the hauls caught marine debris, for an average of 7.5 items/km². Norton Sound had the least amount of debris. It occurred in 7% of the hauls and averaged 1.9 items/km². Galley wastes dominated debris in Oregon (64% of the total CPUE) and in the eastern Bering Sea (40% of the total CPUE), followed by engineering/processing wastes. Fishing equipment debris was abundant in the eastern Bering Sea (1.86 items/km²) and off Oregon (1.69 items/km²), but was not found in Norton Sound. Plastic debris was found in all three areas, but was most abundant in the eastern Bering Sea. Debris of foreign origin accounted for 70% of the total CPUE of all debris found in the eastern Bering Sea; however, domestic debris dominated off Oregon (88% of the total CPUE) and in Norton Sound (100% of the total CPUE).

INTRODUCTION

Marine debris, particularly plastic debris, has been identified as a potential threat to the marine environment world wide (Pruter 1987). To determine the magnitude of the problem, scientists must document the effects and abundance of different types of debris in the marine environment. Educators need to know the probable sources of marine debris in order to direct information campaigns at the proper audiences.

Prior to 1985, the majority of information about marine debris was anecdotal. Few studies presented scientific evidence on the abundance of marine debris or its effects on the marine environment. Recently, studies have reported on the effects of marine debris on marine mammals (Fowler 1988), marine birds (Day et al. 1985), marine turtles (Balazs 1985), and other marine wildlife (Pruter 1987).

While several studies have attempted to estimate the abundance of debris in the marine environment from at-sea disposal rates (Horseman 1982), few studies have addressed the abundance of marine debris using systematic methods. Quantitative surveys of marine debris deposited on beaches in Alaska have been conducted since 1980 (Merrell 1980; Johnson 1988). At-sea surveys have quantified floating debris in the North Pacific since 1977 (Shaw 1977; Dixon and Dixon 1983; Yagi and Nomura 1988). Berger and Armistead (1987) reported the number of pieces of net material caught in trawl nets deployed by foreign fishing vessels in the exclusive economic zone off Alaska between 1982 and 1984.

This study presents baseline information on the type, probable source, and abundance of marine debris caught on the seabed during bottom trawl surveys off Oregon, in the eastern Bering Sea, and in Norton Sound off Alaska during 1988.

METHODS

Survey Areas and Sampling Design

Marine debris was sampled by National Marine Fisheries Service (NMFS) scientists from bottom trawl hauls conducted during 1988 off the coast of Oregon in November-December, in the eastern Bering Sea from May to August, and in Norton Sound during August. A total of 696 hauls were completed covering 33.1 km² over a combined survey area of 907,851 km² (Table 1).

Seventy hauls were conducted between 45 and 110 km off the coast of Oregon between lat. 44° and 45°30'N and from 100 to 675 m deep (Fig. 1). The survey area off Oregon encompassed 7,230 km², of which 2.7 km² was actually covered by bottom trawls (Table 1).

In the eastern Bering Sea, 541 hauls were conducted from the 20 m isobath on the Alaskan coastline out to the 500 m isobath on the continental slope and north from the Alaska Peninsula to Saint Lawrence Island. Stations were sampled at the centers of 37 × 37 km (20 × 20 nmi) grids. The survey area encompassed an area of 858,941 km², of which 26.2 km² was

Table 1.--Survey area (square kilometers) and sampling density for marine debris during the NMFS bottom trawl survey off Oregon, in the eastern Bering Sea, and in Norton Sound, 1988.

Area	Effort					
	Area encom- passed by survey (km ²)	Area covered by trawls (km ²)	Percent area sampled	Total number of hauls	Number of hauls with debris	Percent hauls with debris
Oregon	7,230	2.7	0.037%	70	49	70%
Eastern Bering Sea	858,941	26.2	0.003%	541	122	23%
Norton Sound	41,680	4.2	0.010%	85	6	7%
Total	907,851	33.1	0.004%	696	177	25%

actually covered by trawl hauls. Because of differences in sampling density, the eastern Bering Sea survey area was divided into four subareas. The four subareas for analysis were the north-south shelf and slope (Fig. 2).

Eighty-five hauls were conducted in Norton Sound between the 7 and 20 m isobaths (Fig. 2). The Norton Sound survey area encompassed 41,680 km² and a total of 4.2 km² was actually surveyed.

Trawls were towed on the bottom for approximately 0.5 h at each station at a towing speed of about 5.6 km/h (3 kn). For each haul, location, depth, and distance fished were recorded. The effective path width of the trawl net on the bottom was estimated using a sonar measuring device on a subset of hauls during each survey.

Catches of 1 metric ton or less were entirely sampled. Larger catches were weighed and subsampled, and numbers of marine debris items extrapolated to the total catch. Marine debris items in the catch or subsample were sorted by type of material: plastics, glass, rubber, metal, wood, paper, cloth, and other. Debris items were also described as accurately as possible, such as "plastic strapping band" or "metal beverage can." The number of each of the items caught was recorded on a tally sheet and the vessel, cruise, and haul number indicated. When possible, the U.S. or foreign original of an item was indicated and the percent of all items from U.S., foreign, and unknown sources indicated on each haul tally sheet. The number of entangled animals was recorded by species and debris item. A complete description of NMFS sampling procedures is provided by Wakabayashi et al. (1985).




Figure 1.--Area surveyed and station locations sampled for marine debris during the National Marine Fisheries Service bottom trawl survey off Oregon, 1988.




Figure 2.--Area surveyed and station locations sampled for marine debris during the National Marine Fisheries Service bottom trawl survey in the eastern Bering Sea and Norton Sound, 1988.

Vessels and Fishing Gear

The survey off the coast of Oregon was conducted aboard the 64.6 m NOAA ship *Miller Freeman* using two nets, a modified Nor'eastern trawl and a poly-Nor'eastern trawl. The mean effective path width of the poly-Nor'eastern trawl was estimated to be 14.7 m and the modified Nor'eastern 16.4 m. The eastern Bering Sea survey was conducted using three vessels: the *Miller Freeman*, the 30.5 m RV *Alaska*, and the 37.5 m MV *Morning Star*. Two nets were used during the survey, the eastern trawl, with an estimated mean effective path width of 17.0 m, and the modified Nor'eastern trawl used on the Oregon survey. The *Miller Freeman* conducted the Norton Sound survey with the eastern trawl used in the eastern Bering Sea survey. The eastern trawl had 10.2 cm (4 in) mesh in the wings and body, 8.9 cm (3.5 in) mesh in the cod end, and a 3.2 cm (1.25 in) cod end liner. The modified and poly-Nor'eastern had construction similar to the eastern trawl except for 12.7 cm (5 in) mesh in the wings and body.

Data Analysis

It was assumed all debris 6.5 cm² (1 in²) and larger lying on the surface of the bottom and within the mean effective path width of each net was caught with equal efficiency by each net. This assumption may not necessarily be valid for all hauls, since different nets and different towing conditions can affect the ability of the net to catch objects on the bottom. However, since the NMFS has standardized fishing gear and methods used during most of its annual resource assessment surveys, results obtained from the 1988 surveys should be comparable to future surveys using the same gear and techniques. A second assumption was that scientists identified all of the marine debris caught in each haul.

Marine debris items were grouped by use and by material of composition. Use categories included galley waste, personal use waste (e.g., deodorant tubes, gloves, lighters), fishing gear, engineering and fish processing waste, and other unidentified use waste. Material categories included plastic, glass, rubber, metal, wood, paper, and other. Numbers of items caught were summed by use and material categories by haul and by combinations of the two categories, such as plastic galley waste or metal engineering and processing waste.

The effort expended in each haul was calculated in square kilometers by multiplying the distance fished in each haul by the effective path width of the net. The numbers of individual and grouped marine debris items caught in each haul were divided by the effort to give catch-per-unit-effort (CPUE) in numbers of items per square kilometer for each haul. Mean CPUE per haul was calculated for the entire survey area off Oregon and in Norton Sound and for individual subareas in the eastern Bering Sea using the following formulas:

For an individual haul, CPUE = catch in numbers per unit effort in square kilometers.

For the entire survey area,

$$\text{Mean CPUE} = \frac{\Sigma(\text{CPUE})}{N}$$

$$\text{Variance} = \frac{\Sigma((\text{CPUE} - \text{mean CPUE})^2)}{(N * (N-1))}$$

where Σ = summation for all hauls in the area,

N = the number of hauls in the area.

In the eastern Bering Sea the mean CPUE and variance for the combined subareas were weighted by the area of each subarea in square kilometers using the formulas:

$$\text{Overall mean CPUE} = \frac{\Sigma(A * \text{mean CPUE})}{\Sigma(A)}$$

$$\text{Variance} = \frac{\Sigma(A^2 * \text{variance (mean CPUE)})}{\Sigma(A)^2}$$

where Σ = summation for all subareas,

A = subarea weighting factor.

South shelf = 299,115 km²

North shelf = 520,618 km²

South slope = 17,544 km²

North slope = 21,660 km²

Estimates of CPUE for material and use categories and for total debris items were calculated independently and therefore sums of individual categories do not necessarily equal totals. A more complete description of the standard NMFS methods of calculating CPUE is given in Wakabayashi et al. (1985).

Estimates of the total number of items of debris on the bottom of each area during the 1988 surveys were calculated using an area-swept method (Wakabayashi et al. 1985). Mean CPUE and estimates of numbers of items present in each area are presented as baseline estimates for subsequent comparisons within areas and for all areas combined and were not meant to provide statistically significant comparisons between areas. The percent of debris items by use and material categories is presented for each area and for all areas combined.

RESULTS

Oregon

Of the three areas surveyed, the area off Oregon had the highest concentration of marine debris with 149.6 items/km² (Table 2, Fig. 3). A total of 399 debris items were caught in 49 out of the 70 hauls completed (Table 1). Within use categories, the mean CPUE of galley waste was 89.4 items/km², accounting for 64% of the CPUE of all debris items caught, followed by engineering and processing waste (27%), personal use waste (6%), other use waste (2%), and fishing equipment (1%). Of material categories, the mean CPUE of metal debris was 54.08 items/km² and represented 36% of the mean CPUE of all debris caught, followed by plastics (26%) (Fig. 4), glass (19%), rubber (8%), cloth (6%), wood (3%), and paper (1%) (Table 3).

Of the 399 debris items caught off Oregon, 149 or 37% were identified as of either U.S. or foreign origin. Debris of U.S. origin made up 88% of the mean CPUE of debris of identifiable national origin caught off Oregon, 100% of the CPUE for engineering and processing waste and fishing equipment (Table 4). Foreign debris was represented in the CPUE as galley waste (15%) and personal use items (11%). By material category, U.S. debris caught off Oregon dominated all categories except rubber debris, where foreign debris was 54% of the CPUE of identified items (Table 5).

No animals entangled in marine debris were found in the survey off Oregon. Anemones were attached to a glass bottle and starfish were observed on a piece of plastic rope.

Eastern Bering Sea

The mean CPUE of all debris items caught in the eastern Bering Sea was 7.52 items/km² (Table 2, Fig. 5). Out of the 541 hauls completed, 122 hauls contained a total of 255 marine debris items (Table 1). Galley waste CPUE was 3.15 items/km² or 40% of the mean total CPUE, followed by fishing equipment (24%), engineering and processing waste (24%), and personal use waste (12%). By material category, plastic dominated the total mean CPUE with 4.4 items/km² (51%) (Fig. 6), followed by metal debris (27%), rubber debris (9%), cloth debris (5%), glass debris (4%), and wood debris (1%) (Table 3).

Of the 255 debris items caught in the eastern Bering Sea, U.S. or foreign origin was identified for 60 items. Foreign debris dominated the identified items, accounting for 70% of the mean CPUE (Table 4). Foreign debris was 76% of the CPUE of identified galley waste and 93% of the personal use waste CPUE. Debris of U.S. origin was greatest in fishing equipment waste (67% of CPUE) and engineering and processing waste (64% of CPUE). Foreign debris made up most of the plastic (76% of CPUE), metal (57% of CPUE), rubber (100% of CPUE), and glass debris (84% of CPUE) (Table 5). The U.S. debris accounted for 100% of the CPUE of identified paper and other material debris.

Table 2.--Catch-per-unit-effort (CPUE) (number per square kilometer) by use category and area for marine debris caught during the National Marine Fisheries Service bottom trawl survey, 1988 (CI = confidence interval).

Use category	Oregon		Eastern Bering Sea		Norton Sound		All areas	
	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE
Galley wastes	89.40 (57.2-121.6)	64%	3.15 (2.1-7.2)	40%	0.70 (0.0-1.7)	36%	5.12 (2.7-7.6)	51%
Engineering and processing	37.87 (18.3-57.4)	27%	1.84 (1.0-2.6)	23%	0.73 (0.0-1.8)	38%	2.10 (1.3-2.8)	21%
Fishing equipment	1.69 (0.0-3.4)	1%	1.86 (1.1-2.6)	24%	0.00 --	0%	1.80 (1.1-2.5)	18%
Personal use items	8.92 (0-18.9)	6%	0.91 (0.0-1.9)	12%	0.51 (0.0-1.2)	26%	0.96 (0.0-1.9)	10%
Other debris	2.55 (0.0-6.6)	2%	0.08 (0.0-0.2)	1%	0.00 --	0%	0.05 (0.0-0.1)	<1%
Total	149.60 (97.9-201.3)	100%	7.52 (6.7-14.4)	100%	1.94 (0.3-3.6)	100%	11.26 (7.6-14.9)	100%

Note: Individual and total debris categories were calculated separately and thus are not necessarily additive.

Table 3.--Catch-per-unit-effort (CPUE) (number per square kilometer) by debris material category and area for marine debris caught during the National Marine Fisheries Service bottom trawl survey, 1988 (CI = confidence interval).

Debris category	Oregon			Eastern Bering Sea			Norton Sound			All areas		
	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE	Number/km ² (95% CI N = 70)	Percent of area total CPUE
Plastic	39.05 (20.7-57.4)	26%	4.40 (4.9-7.8)	51%	0.24 (0.0-0.7)	12%	6.37 (5.0-7.7)	57%				
Metal	54.08 (29.6-78.6)	36%	2.33 (0.0-4.8)	27%	0.96 (0.0-2.3)	49%	2.68 (0.3-5.0)	24%				
Rubber	12.10 (0.3-23.9)	8%	0.80 (0.0-1.8)	9%	0.26 (0.0-0.8)	13%	0.87 (0.0-1.8)	8%				
Glass	28.66 (16.4-40.9)	19%	0.38 (0.1-0.6)	4%	0.00 --	0%	0.59 (0.3-0.8)	5%				
Cloth	9.61 (3.6-15.6)	6%	0.41 (0.1-0.7)	5%	0.48 (0.0-1.2)	25%	0.49 (0.2-0.8)	4%				
Wood	4.24 (0.0-8.9)	3%	0.11 (0.0-0.2)	1%	0.00 --	0%	0.14 (0.0-0.3)	1%				
Paper	1.34 (0.0-2.7)	1%	0.13 (0.0-0.3)	2%	0.00 --	0%	0.13 (0.0-0.3)	1%				
Other	0.54 (0.0-6.4)	<1%	0.00 --	0%	0.00 --	0%	>0.01 (0.0-0.1)	<1%				
Total	149.60 (97.9-201.3)	100%	7.52 (6.7-14.4)	100%	1.94 (0.3-3.6)	100%	11.26 (7.6-14.9)	100%				

Note: Individual and total debris categories were calculated separately and thus are not necessarily additive.




Figure 3.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of all marine debris caught during the National Marine Fisheries Service bottom trawl survey off Oregon, 1988.




Figure 4.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of plastic marine debris caught during the National Marine Fisheries Service bottom trawl survey off Oregon, 1988.

Table 4.--Percent of catch-per-unit-effort (number per square kilometer) by foreign or domestic (U.S.) origin, use category, and area for marine debris caught during the National Marine Fisheries Service bottom trawl survey, 1988.

Use category	Oregon		Eastern Bering Sea		Norton Sound		All areas	
	U.S.	Foreign	U.S.	Foreign	U.S.	Foreign	U.S.	Foreign
Galley wastes	85%	15%	24%	76%	0%	0%	37%	63%
Engineering and processing	100%	0%	64%	36%	100%	0%	75%	25%
Fishing equipment	100%	0%	67%	33%	0%	0%	66%	34%
Personal use items	89%	11%	7%	93%	0%	0%	24%	76%
Other debris	100%	0%	100%	0%	0%	0%	100%	0%
Percent by area	88%	12%	30%	70%	100%	0%	42%	58%

Table 5.--Percent of catch-per-unit-effort (number per square kilometer) by foreign or domestic (U.S.) origin, material category, and area for marine debris caught during the National Marine Fisheries Service bottom trawl survey, 1988.

Debris material	Oregon		Eastern Bering Sea		Norton Sound		All areas	
	U.S.	Foreign	U.S.	Foreign	U.S.	Foreign	U.S.	Foreign
Plastic	100%	0%	24%	76%	0%	0%	33%	67%
Metal	85%	15%	43%	57%	100%	0%	55%	45%
Rubber	46%	54%	0%	100%	0%	0%	4%	96%
Glass	81%	19%	16%	84%	0%	0%	37%	63%
Cloth	0%	0%	0%	0%	0%	0%	0%	0%
Wood	100%	0%	0%	0%	0%	0%	100%	0%
Paper	100%	0%	100%	0%	0%	0%	100%	0%
Other	100%	0%	100%	0%	0%	0%	100%	0%
Percent by area	88%	12%	30%	70%	100%	0%	43%	57%




Figure 5.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of all marine debris caught during the National Marine Fisheries Service bottom trawl survey in the eastern Bering Sea, 1988.




Figure 6.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of plastic marine debris caught during the National Marine Fisheries Service bottom trawl survey in the eastern Bering Sea, 1988.

A Tanner crab, *Chionoecetes opilio*, and an unidentified hermit crab, Paguridae, entangled in separate pieces of plastic trawl web twine, were caught during the eastern Bering Sea survey. Numerous invertebrates including mussels, anemones, octopus, barnacles, unidentified tunicates, and starfish were associated with plastic sheeting, plastic rope, glass bottles, and a rubber shoe. Fish eggs were found attached to plastic sheeting.

Norton Sound

Of the three areas surveyed, Norton Sound had the lowest concentration of marine debris, with 1.94 items/km² (Table 2, Fig. 7). Eight items of debris were found in 6 of the 85 hauls completed. Galley waste had a mean CPUE of 0.73 items/km² or 38% of the total debris mean CPUE followed by engineering and processing waste (36%), and personal use waste (26%). No fishing equipment waste was found in Norton Sound. Metal debris accounted for 49% of the total debris mean CPUE, cloth debris 25%, rubber debris 13%, and plastic debris 12% of the total debris mean CPUE (Fig. 8).

Out of the eight debris items caught in Norton Sound, a single debris item, a metal piece of railroad track, was identified as being of U.S. origin.

No animals were found entangled or associated with marine debris in Norton Sound.

All Areas Combined

Out of a total of 696 trawl hauls examined for marine debris in the 3 areas, 177 (25%) had a total of 662 marine debris items identified in the catch. For the 3 areas combined, the mean CPUE of all debris items, weighted by surface area, was 11.3 items/km² (Table 2). Galley waste accounted for 51% of the mean CPUE of all debris items, followed by engineering and processing waste (21%), fishing equipment waste (18%), and personal use waste (10%). Over all areas surveyed, plastic was the most abundant debris material, caught with a mean CPUE of 6.37 items/km² (57% of the mean total CPUE), followed by metal debris (24%), rubber (8%), glass (5%), cloth (4%), and wood and paper (1% of the mean total CPUE) (Table 3).

Of the 210 debris items identified to national origin in the 3 areas, 58% of the mean total CPUE was foreign (Table 4). Foreign debris dominated galley waste (63%) and personal use waste (76%). The U.S. debris accounted for 75% of the mean CPUE of identified engineering and processing waste and 66% of identified fishing equipment waste mean CPUE. Foreign debris accounted for 67% of the mean CPUE of identified plastic debris, 96% of rubber debris, and 63% of the mean CPUE of identified glass debris (Table 5). The U.S. debris dominated identified debris made of metal (55% of mean CPUE) and accounted for all of the identified wood and paper debris caught in the three areas. Plastic represented the largest percentage of CPUE of galley waste (46%), engineering and processing waste (48%), and fishing equipment waste (92%) (Table 6). Rubber debris made up most of the CPUE of personal use waste (77%). A complete list of the individual marine debris items found during the survey is found in Tables 7 through 9.




Figure 7.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of plastic marine debris caught during the National Marine Fisheries Service bottom trawl survey in the Norton Sound, 1988.




Figure 8.--Relative density (catch-per-unit-effort (CPUE) in number per square kilometer) of all marine debris caught during the National Marine Fisheries Service bottom trawl survey in the Norton Sound, 1988.

Table 6.--Percent of catch-per-unit-effort (number per square kilometer) by debris material and use categories for marine debris caught during the National Marine Fisheries Service bottom trawl survey off Oregon, in the eastern Bering Sea, and in Norton Sound, 1988.

Debris	Use category					Percent by material category
	Galley wastes	Engineering and processing	Fishing equipment	Personal use	Other	
Plastic	45.8%	47.5%	91.8%	9.9%	65.4%	56.6%
Metal	42.9%	16.9%	6.1%	0.0%	30.8%	23.8%
Rubber	0.0%	6.1%	0.0%	77.4%	3.8%	7.7%
Glass	10.7%	0.0%	2.1%	0.0%	0.0%	5.2%
Cloth	0.0%	20.3%	0.0%	7.0%	0.0%	4.3%
Wood	0.0%	6.5%	0.0%	0.0%	0.0%	1.2%
Paper	0.5%	2.5%	0.0%	5.6%	0.0%	1.2%
Other	0.0%	0.2%	0.0%	0.0%	0.0%	<1%
Percent by use category	51.3%	20.8%	17.8%	9.6%	0.5%	100.0%

DISCUSSION

The three areas surveyed provide an interesting comparison of the abundance and type of marine debris found on the bottom in areas with different amounts and types of vessel use. The area off Oregon is used extensively by cargo vessels, U.S. and U.S.-foreign joint venture commercial fishing operations, and recreational boaters and fishermen. In 1985, the latest year for which data are available, approximately 1,740 commercial fishing vessels operated off the coast of Oregon (Korson and Thomson 1987) and the U.S. Coast Guard reported 143,373 commercial and recreational vessels in Oregon with Coast Guard identification numbers (Coast Guard 1986). The area surveyed off Oregon is located on one of the major north-south west coast cargo shipping lanes, with frequent vessel traffic observed during the survey (T. Dark, Alaska Fisheries Science Center, Seattle, Wash., pers. commun. 1989).

In the eastern Bering Sea, some nonfishery tug, barge, and cargo vessel operations exist, but vessel traffic is predominantly associated with the commercial fishing industry. Harvesting vessels, domestic and foreign processing vessels, and a wide variety of support vessels operate in the eastern Bering Sea each year. In 1985, the Alaska Department of Fish and Game (1986) estimated that 1,729 domestic commercial fishing vessels operated in the eastern Bering Sea, and the NMFS estimated that 254 foreign vessels fished or processed seafood in the eastern Bering

Table 7.--Description, material and use category, number caught, catch-per-unit-effort (CPUE) (number per square kilometer), and swept-area estimate of the number of debris items in the survey area for marine debris caught off Oregon during the National Marine Fisheries Service bottom trawl survey, 1988 (CI = confidence interval).

Use category	Item	Catch			Swept-area estimate	
		Number caught	Mean CPUE km ²	CPUE variance	Estimated number	95% CI
Plastic						
Galley waste	Bags	39	14.31	11.2468	103,277	54,946-151,607
	Bottles	2	0.51	0.1290	3,702	0-8,896
	Lids, caps	4	1.71	1.4426	12,325	0-29,634
	Six-pack ring	1	0.25	0.0624	1,803	0-5,405
	Vegetable sack	1	0.90	0.8185	6,529	0-19,567
	Other	2	0.72	0.2909	5,170	0-12,943
Fishing equipment	Fishing line	2	0.88	0.4228	6,347	0-15,718
	Fishing net	1	0.27	0.0732	1,952	0-5,850
	Rope	21	9.45	12.9024	68,218	0-119,983
Personal use	Lighter	1	0.27	0.0717	1,932	0-5,790
	Deodorant tube	15	4.02	16.1272	28,981	0-86,855
Engineering and processing	Sheeting	8	2.36	2.8377	17,058	0-41,339
	Strapping band	9	2.92	1.2100	21,102	0-36,955
	Duct tape	1	0.22	0.0494	1,605	0-4,809
Other	Clay pigeon	1	0.25	0.0601	1,770	0-5,303
Glass						
Galley waste	Bottle	65	25.92	32.4534	187,265	105,167-269,364
	Pieces	2	0.75	0.3088	5,447	0-13,456
	Fruit jar	4	1.95	3.0305	14,082	0-39,170
Rubber						
Personal use	Gloves	6	2.47	0.1049	17,798	3,035-32,561
	Shoe	1	0.86	0.4123	6,211	0-15,465
Engineering and processing	Tar	10	5.05	0.2546	36,411	0-109,125
	Gasket	3	0.79	0.3292	5,666	0-13,935
	Paint	4	1.80	2.4378	13,012	0-35,513
	Sheeting	2	0.87	0.4358	6,287	0-15,801
Other	Misc. pieces	1	0.27	0.0702	1,913	0-5,732

Table 7.--Continued.

Use category	Item	Catch			Swept-area estimate	
		Number caught	Mean CPUE km ²	CPUE variance	Estimated number	95% CI
Metal						
Galley waste	Beverage can	95	35.38	75.5465	255,327	130,066-380,587
	Lids, caps	4	1.46	0.8108	10,537	0-23,514
	Container	5	1.96	0.8102	14,136	1,164-27,108
	Pull tab	7	2.50	3.1064	18,075	0-43,475
	Tinfoil	2	0.50	0.1231	3,581	0-8,638
	Cook pot	2	0.54	0.1429	3,879	0-9,326
Fishing equipment	Crab trap	1	0.54	0.2867	4	0-11,581
Engineering and processing	Drum, 208.2 liter (55-gal)	1	0.50	0.2546	3,641	0-10,913
	Pieces	16	5.94	7.1185	42,849	4,398-81,299
	Instruments	3	0.97	0.3255	6,972	0-15,193
	Paint can	4	1.76	2.4200	12,715	0-35,134
Other	Bullet	4	2.04	4.1527	14,706	0-44,074
Paper						
Personal use	Newspaper	1	0.27	0.0717	1,932	0-5,709
	Pieces	2	0.53	0.1371	3,807	0-9,144
	Book	1	0.25	0.0613	1,770	0-5,303
Engineering and processing	Carton	1	0.30	0.0879	2,139	0-6,412
Wood						
Engineering and processing	Pieces	14	3.76	5.2919	26,951	0-60,103
	Broom	1	0.29	0.0813	2,058	0-6,168
	Fiberboard	1	0.22	0.0494	1,605	0-4,809
Cloth						
Engineering and processing	Pieces and rags	6	9.61	9.0228	69,330	26,041-112,619
Other						
Engineering and processing	Fire brick	26	0.54	0.2867	3,864	0-11,581

Table 8.--Description, material and use category, number caught, catch-per-unit-effort (CPUE) (number per square kilometer), and swept-area estimate of the number of debris items in the survey area in the eastern Bering Sea during the National Marine Fisheries Service bottom trawl survey, 1988 (CI = confidence interval).

Use category	Item	Catch			Swept-area estimate	
		Number caught	Mean CPUE km ²	CPUE variance	Estimated number	95% CI
Plastic						
Galley waste	Bags	49	1.10	0.0653	1,629,941	1,195,289-2,064,593
	Bottles	2	0.01	0.0017	11,209	0-33,626
	Lids, caps	3	0.11	0.0070	97,164	0-239,880
	Wrappers	4	0.21	0.0177	177,959	0-404,079
	Other	1	0.04	0.0016	34,424	0-102,581
Fishing equipment	Bait jar	2	0.09	0.0041	74,915	0-183,768
	Fishing line	17	0.78	0.0546	669,889	272,412-1,067,367
	Fishing net	7	0.35	0.0173	299,166	75,558-522,774
	Net twine	8	0.33	0.0140	285,103	83,872-486,335
	Floats	1	0.05	0.0030	46,871	0-139,671
	Light stick	2	0.11	0.0113	91,208	0-271,792
	Rope	28	1.32	0.0646	1,131,965	669,668-1,564,263
Personal use	Hard hat	1	0.01	0.0001	5,894	0-17,681
	Toothpaste tube	2	0.05	0.0017	40,270	0-109,798
	Glove liner	1	0.01	0.0001	9,100	0-27,491
Engineering and processing	Sheeting	15	0.72	0.0775	619,684	146,206-1,093,162
	Strapping band	7	0.22	0.0094	192,401	27,813-356,990
	Duct tape	1	0.05	0.0026	44,114	0-131,455
Other	Clay pigeon	1	0.04	0.0017	35,414	0-105,532
	XBT tube ^a	1	0.04	0.0013	30,766	0-91,681
Glass						
Galley waste	Bottle	8	0.31	0.0133	263,374	67,448-459,299
	Pieces	1	0.03	0.0010	28,129	0-83,823
Fishing equipment	Glass float	1	0.04	0.0016	34,424	0-102,581
Rubber						
Personal use	Gloves	14	0.67	0.2437	571,566	0-1,411,060
	Shoes	2	0.07	0.0028	64,347	0-154,201
Engineering and processing	Tar	1	0.01	0.0001	6,433	0-19,299
	Sheeting	1	0.05	0.0030	47,198	0-140,648

Table 8.--Continued.

Use category	Item	Catch			Swept-area estimate	
		Number caught	Mean CPUE km ²	CPUE variance	Estimated number	95% CI
Metal						
Galley waste	Beverage can	33	1.55	1.4860	1,328,689	0-3,401,783
	Lids, caps	1	0.05	0.0029	45,914	0-136,821
	Container	7	0.26	0.0188	223,756	0-457,004
	Tinfoil	1	0.05	0.0026	43,827	0-130,601
	Cook pot	1	0.03	0.0012	29,301	0-87,316
Fishing equipment	Crab trap	3	0.11	0.0050	94,090	0-214,290
Engineering and processing	Pieces	3	0.17	0.0280	143,604	0-427,928
	Wire	9	0.10	0.0054	89,238	0-214,176
Paper						
Galley waste	Bag	1	0.03	0.0008	24,249	0-72,261
Personal use	Piece	1	0.05	0.0024	41,663	0-124,152
Engineering and processing	Carton	1	0.05	0.0027	44,404	0-132,320
Wood						
Engineering and processing	Pieces	2	0.04	0.0014	37,988	0-101,750
	Paint brush	1	0.03	0.0010	27,047	0-80,599
	Other	1	0.03	0.0010	27,196	0-81,044
Cloth						
Personal use	Pants	1	0.06	0.0034	49,995	0-148,982
Engineering and processing	Pieces	6	0.25	0.0120	217,809	31,675-403,944
	Tarp	1	0.05	0.0026	43,544	0-129,759
	Bag	1	0.05	0.0024	41,663	0-124,152

*XBT - Expendable bathythermograph.

Table 9.--Description, material and use category, number caught, catch-per-unit-effort (CPUE) (number per square kilometer), and swept-area estimate of the number of debris items in the survey area for marine debris caught in Norton Sound during the National Marine Fisheries Service bottom trawl survey, 1988 (CI = confidence interval).

Category			Catch			Swept-area estimate	
			Number caught	Mean CPUE No./km ²	CPUE variance	Estimated number	95% CI
Material	Use	Item					
Plastic	Galley waste	Bag	1	0.24	0.0559	9,857	0-29,493
Rubber	Personal use	Shoe	1	0.26	0.0664	10,741	0-32,138
Metal	Galley waste	Beverage can	2	0.46	0.2128	19,228	0-57,530
	Engineering and processing	Railroad track	2	0.50	0.2516	20,906	0-62,557
Cloth	Engineering and processing	Pieces and rags	1	0.23	0.0539	9,674	0-28,947
	Personal use	Dress	1	0.25	0.0629	10,453	0-31,275

Sea-Aleutian Islands area (Berger et al. 1988). There are few, if any, recreational boaters operating in the eastern Bering Sea and the major cargo transit routes lie south of the Aleutian Islands.

Norton Sound has the least amount of vessel traffic of the three areas surveyed. Tug and barge traffic to Nome, Alaska, occurs during the spring and summer. A fleet of about a dozen vessels conducts a commercial red king crab fishery in the survey area for approximately 1 week each year (Alaska Department of Fish and Game 1986). During the winter, most of Norton Sound is covered by ice.

The estimated abundance of marine debris in the three areas surveyed differed by nearly two orders of magnitude, from 1.94 items/km² in Norton Sound to 149.60 items/km² off Oregon. The higher concentration of marine debris off Oregon is probably related to the extensive vessel operations in this area. Most of the marine debris off Oregon was galley waste, 89.4 items/km² (64%), and engineering and processing waste, 37.87 items/km² (27%), which are associated with the operation of most types of vessels. Fishing equipment waste abundance off Oregon, 1.69 items/km², was quite similar to that found in the eastern Bering Sea, 1.84 items/km². It is

interesting to note that the numbers of commercial fishing vessels operating off Oregon and in the eastern Bering Sea were also similar, 1,740 and 1,983, respectively. The abundance of galley waste and engineering and processing debris caught in the eastern Bering Sea may represent the average amount resulting from commercial fishing operations and minimal cargo traffic. The higher abundance of galley waste and engineering and processing waste found off Oregon may be due to the added input of cargo vessel and recreational boater debris.

RECOMMENDATIONS

- Collect marine debris data from all annual NMFS bottom trawl surveys.
- Develop a standardized data collection protocol, data base system, analysis methodology, and reporting format.
- Provide similar marine debris data forms to commercial trawl fishermen.
- Encourage foreign governments to conduct similar bottom trawl marine debris surveys.

REFERENCES

- Alaska Department of Fish and Game.
1986. Alaska 1985 catch and production. Commer. Fish. Stat. Leaflet 38, 61 p.
- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Berger, J. D., and C. E. Armistead.
1987. Discarded net material in Alaskan waters, 1982-84. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-110, 66 p.
- Berger, J. D., R. Nelson Jr., J. Wall, H. Weikart, and B. Maier.
1988. Summary of U.S. observer sampling of foreign and joint venture fisheries in the northeast Pacific Ocean and eastern Bering Sea, 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-128, 167 p.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

- Dixon, T. J., and T. R. Dixon.
1983. Marine litter distribution and composition in the North Sea. Mar. Pollut. Bull. 14:145-148.
- Fowler, C.
1988. A review of seal and sea lion entanglement in marine fishing debris. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 16-63. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Horseman, P. V.
1982. The amount of garbage pollution from merchant ships. Mar. Pollut. Bull. 13:167-169.
- Johnson, S.
1988. Deposition of entanglement debris on Alaskan beaches. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 207-231. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Korson, C. S., and C. J. Thomson.
1987. A report on the PacFIN research data base for 1985. Southwest Fish. Cent., Natl. Mar. Fish. Serv., NOAA, La Jolla, CA. Southwest Fish. Cent. Admin. Rep. LJ-87-21, 16 p.
- Merrell, T. R., Jr.
1980. Accumulation of plastic litter on beaches of Amchitka Island Alaska. Mar. Environ. Res. 3:171-184.
- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. Mar. Pollut. Bull. 18:305-310.
- Shaw, D. G.
1977. Pelagic tar and plastic in the Gulf of Alaska and Bering Sea, 1975. Sci. Total Environ. 8:13-20.
- U.S. Coast Guard.
1986. Boating statistics 1986. U.S. Dep. Trans., U.S. Coast Guard COMDTPUB P16754.1, 36 p.
- Wakabayashi, K., R. G. Bakkala, and M. S. Alton.
1985. Methods of the U.S.-Japan demersal trawl surveys. In R. G. Bakkala and K. Wakabayashi (editors), Results of cooperative U.S.-Japan groundfish investigations in the Bering Sea during May-August 1979, p. 7-29. Int. North Pac. Fish. Comm. Bull. 44.

Yagi, N., and M. Nomura.

1988. Sighting survey of marine debris in the midwestern Pacific from 1977 through 1986. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 130-142. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

**ESTIMATING THE DENSITY OF FLOATING MARINE
DEBRIS: DESIGN CONSIDERATIONS**

Christine A. Ribic* and L. J. Bledsoe
Center for Quantitative Science in Forestry, Fisheries, and Wildlife
University of Washington
Seattle, Washington 98195, U.S.A.

*Present address: Environmental Research Laboratory, U.S. Environmental Protection Agency, 200 SW 35th Street, Corvallis, OR 97333, U.S.A.

ABSTRACT

We calculated sample sizes needed to estimate the density of surface marine debris potentially injurious to marine mammal and bird populations in the Gulf of Alaska and the Bering Sea as well as sample sizes needed to specifically estimate floating nets. Using published estimates of debris density, we developed alternative sample size requirements that depended on the accuracy required based on the coefficient of variation of the density. The survey technique used was visual sighting of debris using strip transect methodology. In general, large numbers of transects are needed in order to get estimates even with large coefficients of variation. Sparsity of data and nonstandard definition of transects contribute to the problems in estimating required sample sizes.

INTRODUCTION

The problems of marine debris and its impacts on marine mammals and on human activity in the oceans have been reviewed and discussed extensively by Shomura and Yoshida (1985). There has been interest in estimating the amount of floating marine debris using visual assessment. This technique has been used by many researchers (Venrick et al. 1973; Suzuoki and Shirakawa 1979; Dahlberg and Day 1985; Jones and Ferrero 1985; Yoshida and Baba 1985a, 1985b, 1988; Baba et al. 1986; Ignell and Dahlberg 1986; Day and Shaw 1987; McCoy 1988; Mio and Takehama 1988; Yagi and Nomura 1988). The purpose of this paper is to investigate survey design to estimate the density of surface marine debris in the Gulf of Alaska and Bering Sea.

We considered the design of two surveys. The first was to estimate density for all potentially harmful floating debris that could be visually assessed (specifically nets, fragmented plastic pieces, and strapping bands). Each type of debris was assumed to be equally important. The second design was for estimating the density of floating nets only.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154, 1990.

METHODS

The strip transect was the method used for visually assessing the density of floating objects. This method was chosen because of its widespread use (references cited above). The transects have a fixed width and the assumption is that all objects within that width are seen. The method of Burnham et al. (1980) was used to estimate sample size. This method is nonparametric because it does not make an assumption about the distribution of the debris. Estimation of sample size is based on achieving a certain coefficient of variation for the density of objects.

We used the conservative estimate for total transect length:

$$L = (3 \cdot L1) / (cv(D)^2 \cdot n1)$$

where L1 (total length of transects) and n1 (total number of objects seen) come from a pilot study, and cv(D) is the coefficient of variation (Burnham et al. 1980).

We used previously published papers on the Gulf of Alaska and the Bering Sea for estimates of L1 and n1 for total floating debris and floating nets. In addition, the data for the 1984 marine mammal observer program were made available to us (L. Jones, National Marine Mammal Laboratory, Seattle, WA, pers. commun.).

RESULTS

Total Floating Debris

From Dahlberg and Day (1985), an estimate of all debris was based on a strip transect with a width of 50 m. They do not state the length of their transects but state that an average of 5.5 h/day were spent watching for debris and that 1,516 nmi were sampled from Alaska to Hawaii (Dahlberg and Day 1985). This gives an average transect length of 47 nmi covered per 5.5 h. So the sampling unit will be defined here as a transect 47 nmi long by 50 m wide. Twelve objects were seen in the Gulf of Alaska (n1) and we estimate 670 nmi (Dahlberg and Day 1985, fig. 3) was surveyed (L1). Dahlberg and Day (1985) gave a density estimate for all floating marine debris as 0.28 pieces/km², but they did not publish a variance estimate. Day and Shaw (1987) give density and variance estimates for large floating plastic for the subarctic North Pacific (Gulf of Alaska) and, separately, for the Bering Sea.

Estimates of required sample sizes (number of transects) for estimating total floating debris are presented in Table 1. In general, in order to estimate density to any degree of precision (low cv(D)), 2 months or more of daily transects (5.5 h of observation for a 47-nmi-long by 50-m-wide transect) would be needed. Dahlberg and Day (1985) carried out about 14 transects, which would put their estimate in the 0.50 cv(D) category (not a small coefficient of variation).

Table 1.--Sample size estimation for all floating marine debris using a strip transect of 47 nmi long by 50 m wide for the Gulf of Alaska and the Bering Sea for different coefficients of variation for the density. (L = total transect length and n = number of transects needed to cover that length.)

cv(D)	L (nmi)	n
0.10	16,750	994
0.25	2,680	57
0.50	670	15
0.80	262	6
1.0	167	4
1.2	117	3

Nets

From Jones and Ferrero (1985), 8,759 nmi (L1) were surveyed in 1984 with 12 pieces of net seen (n1). A density estimate of floating nets would be 0.0074 nets/km². A transect for this study was 2 nmi in length and 100 m in width. A total of 1,410 transects were made.

Estimates of total sample size (number of transects) for estimating floating nets are presented in Table 2. In all cases, a large number of transects (2 nmi length by 100 m width) would need to be made to get even an inaccurate estimate of the density of nets. There were 1,410 transects made in 1984, which would put the net density estimate in the 0.80 cv(D) category, a large coefficient of variation.

DISCUSSION

The number of transects needed to produce a reasonable estimate for floating marine debris and especially for nets is extremely large. This demonstrates that targeting for a specific type of debris that is relatively rare, like floating nets, will take a large commitment of resources. These sample size estimates, however, depend on a large number of factors.

First, the approach we used is a nonparametric approach that is extremely general and requires sighting 25 or more objects to produce estimates of means and variances with any degree of accuracy (Burnham et al. 1980). Sample sizes for estimating rare objects like floating nets will be extremely large. A parametric approach such as using a binomial distribution may lead to smaller sample sizes but then the underlying model will have to be verified (Ribic and Bledsoe 1986).

Second, there was little information on which to base preliminary estimates of density and variation. Some of this had to do with the way

Table 2.--Sample size estimation for floating nets using a strip transect of 2 nmi long by 100 m wide for the Gulf of Alaska and the Bering Sea for different coefficients of variation for the density. (L = total transect length and n = number of transects needed to cover that length.)

cv(D)	L (nmi)	n
0.10	218,975	109,488
0.25	35,036	17,518
0.50	8,760	4,380
0.80	3,422	1,711
1.0	2,190	1,095
1.2	1,521	761

the data were reported. For example, in some cases we could not determine the length of a transect so we could not use the reported data. But more importantly, there is little published information on which to base preliminary estimates. Dahlberg and Day (1985) worked along long. 155°W. Jones and Ferrero (1985) worked in the middle of the gillnet fishery. Whether these studies are representative of the rest of the unsampled area is not known.

Third, transect length and width are not standardized, so sample size estimates in this paper depend on a specifically defined transect. Density estimates depend on the dimensions of the strip transect. Therefore, generalizations are difficult, since most researchers use different transect widths and lengths for their transects (e.g., Mio and Takehama (1988) used a width of 10 m).

Fourth, due to lack of information on variation for the Gulf of Alaska and the Bering Sea, we did not consider stratification (Cochran 1977), which could be potentially very useful in determining sample allocation and the placement of transect lines. Dahlberg and Day (1985) and Ignell and Dahlberg (1986) noted the concentration of debris in downwelling areas and frontal zones. A large-scale survey such as that of Mio and Takehama (1988) for the Gulf of Alaska and the Bering Sea would greatly improve our knowledge of the distribution of marine debris and improve survey design immensely.

Further refinement of the survey objective would be helpful when we consider placement of the transect lines. If a study is a one-time occurrence, the transects can be considered temporary and location will be decided by where the ship goes. However, if the study is to be a long-term study, thought should be given to permanent transects. For example, Day and Shaw (1987) compared the density of debris along long. 155°W previously sampled by Dahlberg and Day (1985). The long. 155°W line would be an

example of a permanent transect that could be surveyed over time. Another example is the study of Yagi and Nomura (1988), where the long. 137°E line was surveyed between lat. 0° and 34°N each summer and winter for 9 years; however, they commented that their limited coverage of the area did not allow them to make conclusions about changes in marine debris distribution over time.

ACKNOWLEDGMENTS

Initial work on this problem was funded by Northwest and Alaska Fisheries Center, National Marine Fisheries Service, NOAA, Seattle, WA, Contract No. 43-ABNF-5-2498. We thank M. B. Hanson for doing the literature review and reading this manuscript. We thank L. Jones for allowing us to use unpublished data in the initial project.

REFERENCES

- Baba, N., K. Yoshida, M. Onoda, N. Nagai, and S. Toishi.
1986. Research on ocean current and drifting stray fishing gear and net fragments in the area to the southwest of the Pribilof Islands and the southern coast of the Aleutian Islands, July-August, 1985. Document submitted to the Standing Scientific Committee of the North Pacific Fur Seal Commission.
- Burnham, K. P., D. R. Anderson, and J. L. Laake.
1980. Estimation of density from line transect sampling of biological populations. *Wildl. Monogr.* 72.
- Cochran, W. G.
1977. *Sampling techniques*. Third edition. Wiley, N.Y.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Day, R. H., and D. G. Shaw.
1987. Patterns in the abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-1985. *Mar. Pollut. Bull.* 18:311-316.
- Ignell, S. E., and M. L. Dahlberg.
1986. Results of cooperative research on the distribution of marine debris in the North Pacific Ocean. Document submitted to the International North Pacific Fisheries Commission, Anchorage, AK, November 1986. Northwest and Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Auke Bay Laboratory, P.O. Box 210155, Auke Bay, AK 99821, 15 p.

Jones, L. L., and R. C. Ferrero.

1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

McCoy, F. W.

1988. Floating megalitter in the eastern Mediterranean. Mar. Pollut. Bull. 19:25-28.

Mio, S., and S. Takehama.

1988. Estimation of distribution of marine debris based on the 1986 sighting survey. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, October 13-16, 1987, p. 64-94. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Ribic, C. A., and L. J. Bledsoe.

1986. Design of surveys for density of surface marine debris in the North Pacific. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA. NWAFC Processed Rep. 267, 69 p.

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

Suzuoki, T., and K. Shirakawa.

1979. Visual observation of floating pollutants in open ocean. Oceanogr. Mag./Kisho-Cho Obun Kaiyoho 30(1-2):55-60.

Venrick, E. L., T. W. Backman, W. C. Bartram, C. J. Platt, M. S. Thornhill, and R. E. Yates.

1973. Man-made objects on the surface of the central North Pacific Ocean. Nature 241:271.

Yagi, N., and M. Nomura.

1988. Sighting survey on marine debris in the midwestern Pacific from 1977 through 1986. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, October 13-16, 1987, p. 130-142. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Yoshida, K., and N. Baba.

1985a. Results of the survey on drifting fishing gear or fish net pieces in the Bering Sea. Document submitted to the 28th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, Tokyo, 4-12 April 1985, 13 p.

- 1985b. A survey of drifting stray fishing net fragments in the northern Sea of Japan (western Pacific Ocean). Document submitted to the 28th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, Tokyo, 4-12 April 1985, 13 p.
1988. Results of research on the effects of marine debris on fur seal stocks and future research programs. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, October 13-16, 1987, p. 95-129. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

**CHARACTERIZATION OF MARINE DEBRIS IN
SELECTED HARBORS OF THE UNITED STATES**

Wayne R. Trulli, Heather K. Trulli
Battelle Memorial Institute
397 Washington Street
Duxbury, Massachusetts 02332, U.S.A.

and

David P. Redford
Office of Marine and Estuarine Protection
U.S. Environmental Protection Agency
401 M Street, S.W.
Washington, D.C. 20460, U.S.A.

ABSTRACT

As part of a program to characterize anthropogenic debris in the marine environment, the U.S. Environmental Protection Agency conducted field surveys in the harbors of nine major metropolitan areas of the United States: New York, Boston, Philadelphia, Baltimore, and Miami on the east coast, and Tacoma, Seattle, Oakland, and San Francisco on the west coast. The surveys were designed to provide information on the types, relative amounts, and distributions of marine debris in several geographic regions of the United States. Neuston net (0.3-mm mesh) tows were conducted during outgoing tides on consecutive days. After each net tow, the debris, which ranged in size from small plastic pellets to large plastic sheeting, was identified, categorized, and counted. Seven of the ten most common debris items collected were plastic or polystyrene materials. The data are being used to qualitatively characterize marine debris in coastal metropolitan areas and to examine potential regional variations and sources.

INTRODUCTION

In response to domestic and international concerns about marine plastic debris, the U.S. Congress passed the Marine Plastic Pollution Research and Control Act of 1987. Title II of this act directs the U.S. Environmental Protection Agency (EPA) to conduct a study and to issue a Report to the Congress on methods for reducing plastic pollution. One section of the comprehensive Report to the Congress discusses the types and sources of marine plastic debris, the transport and fate of this debris, and its effects on the marine environment and on human health and safety.

It also lists what EPA believes to be items of concern in the marine debris. These items are pellets, condoms, tampons, syringes and medical items, nets and traps, line and rope, six-pack yokes (or similar beverage yokes), and plastic bags and sheeting.

Because few data were available prior to the preparation of the Report to the Congress, EPA determined that field studies were necessary to collect data to adequately characterize plastic debris and its sources along the coastal United States. The Harbor Studies Program (Redford 1990), initiated by EPA in 1988, focuses on examining plastic and other floating debris in major harbors along the U.S. Atlantic and Pacific coasts. The objective of this field program was to characterize the types, relative amounts, and distributions of marine debris in representative harbors. This paper presents a summary of the results of the first nine surveys conducted under this program.

METHODS

Sample Collection

Floating debris often is observed concentrated in dense windrows, commonly referred to as debris slicks (EPA 1988), that appear to be influenced by surface currents and winds. Because the location, size, shape, and integrity of slicks were anticipated to be highly variable within each harbor, sampling was not conducted along predetermined transects but, rather, was directed toward the denser areas within the slicks.

Sampling at each harbor was conducted on 2 or 3 consecutive days between November 1988 and February 1989. Sampling dates and the total number of samples collected at each location are presented in Table 1.

The sampling plan for each location designated two or three areas within each harbor, based on criteria such as (1) presence of combined sewer overflows (CSO's) and stormwater outfalls in close proximity, (2) areas of heavy ship traffic or boating activity, (3) highly industrialized locations, and (4) areas that would represent overall debris conditions in the harbor. Because accumulated debris within the harbor is most likely to be transported out of the harbor with an outgoing tide, all sampling activities were initiated 1 to 2 h before ebb tide at each location. Selected areas within a harbor were sampled concurrently by deploying two or more small vessels.

Samples were collected by using a 0.33-mm mesh neuston net with dimensions of either $1 \times 2 \times 4$, or $0.5 \times 1 \times 4$, or $0.5 \times 1 \times 2$ m. To minimize disturbances from the wake of the vessel, the net was towed from a boom positioned abeam of the vessel.

Sampling was conducted in slicks that were observed to be generally dense with floating debris. Each tow made through a slick was considered a single sample, regardless of the tow length. Generally, tows were conducted at a speed of 2 kn for approximately 20 min, or until a sample volume of approximately 80 L was collected. If more than one tow was made

Table 1.--Summary of harbor studies sampling activities,
November 1988 through February 1989.

Location	Dates sampled	No. of samples
New York ^{a,b}	11, 12, 13 November 1988	43
Boston	2, 3, 4 December 1988	49
Philadelphia ^{a,c}	26, 27 January 1989	29
Baltimore ^a	29, 30 January 1989	29
Miami	3, 4, 5 February 1989	31
Tacoma ^b	15, 16, 17 February 1989	11
Seattle ^a	15, 16, 17 February 1989	6
Oakland	21, 22, 23 February 1989	12
San Francisco	21, 22, 23 February 1989	14
Total		224

^aCSO's observed discharging.

^bRainfall ≥ 1 in.

^cSpring high tide during sampling period.

within a slick, each tow sample was considered to be a replicate. Following each tow, the captured debris was collected and placed into labeled containers.

Meteorological conditions and the dimensions and location of each sampled slick were recorded on a sample-tracking form. Visual fixes of landmarks were used to plot tow locations on navigational charts.

Sample Processing and Analysis

All samples were processed and analyzed immediately after returning from the field. Prior to processing, all items in a sample were rinsed with tap water. Processing entailed separating all anthropogenic material from natural materials, and sorting and identifying the debris items by specific, descriptive categories (Fig. 1). Many of these categories were adapted from the national beach survey data card developed by the Center for Marine Conservation. All debris items within a category were counted and the totals were recorded on these or similar inventory or data sheets.

The data for each harbor sampling site were entered into a data base and the percent composition was calculated for each item or combination of items. Samples were photodocumented immediately upon return to the laboratory. All percentages discussed herein are calculated based on numbers of items found, not on weight or volume of the items.

RESULTS AND DISCUSSION

During this study, items were enumerated but they were not weighed or measured in any other manner. All cited percentages are based on the numbers of items found.

FLOATABLES SURVEY DATA INVENTORY SHEET

Attach Sample Labels Here

<p>PLASTICS</p> <p>Absorbent Material _____</p> <p>Bags and ties _____</p> <p>Bags ≤ 1-gal capacity _____</p> <p>Condom bag _____</p> <p>Garbage bag tie _____</p> <p>Bags > 1-gal capacity _____</p> <p>Misc. bags _____</p> <p>Misc. pieces _____</p> <p>Vegetable sack _____</p> <p>Banding Material _____</p> <p>Electrical wire tie _____</p> <p>Strapping band _____</p> <p>Brittles _____</p> <p>Bottles ≤ 1-gal capacity _____</p> <p>Bottles > 1-gal capacity _____</p> <p>Beverage bottles _____</p> <p>Misc. bottles _____</p> <p>Misc. pieces _____</p> <p>Caps and lids _____</p> <p>Caps/lids _____</p> <p>Cap/lid liners _____</p> <p>Cap/lid pieces _____</p> <p>Pull tab from plastic lid _____</p> <p>Cigarette/Cigar items _____</p> <p>Wrappers and packs _____</p> <p>Cigar tips _____</p> <p>Cigarette butts & filters _____</p> <p>Disposable lighters _____</p> <p>Containers _____</p> <p>Lemon juice dispensers _____</p> <p>Misc. containers _____</p> <p>Dishware _____</p> <p>Cups, spoons, forks, straws _____</p> <p>Dishes/plates _____</p> <p>Misc. pieces _____</p> <p>Drug Paraphernalia _____</p> <p>Crack vial caps _____</p> <p>Crack vials w/caps _____</p> <p>Crack vials w/o caps _____</p> <p>Fishing/Boating items _____</p> <p>Floats & buoys _____</p> <p>Fishing line-monofilament _____</p> <p>Netting _____</p> <p>Food wrappers--Misc. _____</p> <p>Hair Care & Cosmetic Items--Misc. _____</p> <p>Housewares & Toils--Misc. _____</p> <p>Labels--Misc. _____</p> <p>Line/Rope _____</p> <p>Filament _____</p> <p>Rope length ≤ 2 ft _____</p> <p>Rope length > 2 ft _____</p> <p>Medical _____</p> <p>Band-aids _____</p> <p>Band-aid wrappers _____</p> <p>Cough syrup bottles _____</p> <p>Cylindrical tubes _____</p> <p>Cylindrical tube pieces _____</p> <p>Lip balm & containers _____</p>	<p>Misc. _____</p> <p>Needle covers _____</p> <p>Pill vials & caps _____</p> <p>Syringes (whole) _____</p> <p>Syringes (pieces) _____</p> <p>Syringes with blood _____</p> <p>Tube ends only _____</p> <p>Vials _____</p> <p>Vial caps _____</p> <p>Miscellaneous _____</p> <p>Foil wrappers (plastic-coated) _____</p> <p>Hardhat band _____</p> <p>Misc. items _____</p> <p>Pieces _____</p> <p>Wrappers _____</p> <p>Polyvinylchloride (PVC) _____</p> <p>Toys _____</p> <p>Tubing _____</p> <p>Vials _____</p> <p>Pellets _____</p> <p>Spherules _____</p> <p>Personal Hygiene _____</p> <p>Condoms (whole) _____</p> <p>Condoms (pieces) _____</p> <p>Diapers _____</p> <p>Panty liners _____</p> <p>Q-tips (whole) _____</p> <p>Q-tips (tubes only) _____</p> <p>Sanitary items _____</p> <p>Sanitary napkins _____</p> <p>Tampon applicators _____</p> <p>Tampon wrappers _____</p> <p>Photographic items _____</p> <p>Film containers _____</p> <p>Photos _____</p> <p>Shedding _____</p> <p>≤ 2 ft _____</p> <p>> 2 ft _____</p> <p>Six-Pack Holders _____</p> <p>Whole _____</p> <p>Pieces _____</p> <p>Writing Utensils--Misc. _____</p> <p>GLASS</p> <p>Bottles _____</p> <p>Alcohol bottles _____</p> <p>Food bottles _____</p> <p>Light bulbs _____</p> <p>Misc. pieces _____</p> <p>PAPER</p> <p>Bags _____</p> <p>Whole _____</p> <p>Pieces _____</p>	<p>Cartons/Cardboard Boxes _____</p> <p>Whole _____</p> <p>Pieces _____</p> <p>Food Items & Wrappers _____</p> <p>Beverage cartons _____</p> <p>Cups & plates _____</p> <p>Fast food wrappers _____</p> <p>Food wrappers _____</p> <p>Gun wrappers _____</p> <p>Lollipop sticks _____</p> <p>Houseware Items & Toils _____</p> <p>Hand-wipes _____</p> <p>Matches _____</p> <p>Tar paper _____</p> <p>Miscellaneous _____</p> <p>Cap liners _____</p> <p>Misc. items _____</p> <p>Misc. pieces _____</p> <p>Misc. wrappers _____</p> <p>Sanitary items _____</p> <p>Tissues _____</p> <p>Toilet paper _____</p> <p>TEXTILES</p> <p>Shoes--athletic _____</p> <p>Canvas _____</p> <p>Clothing--whole & pieces _____</p> <p>Lint _____</p> <p>Medical _____</p> <p>Cotton _____</p> <p>Cotton balls _____</p> <p>Linen _____</p> <p>Rope _____</p> <p>STYROFOAM</p> <p>Band-pieces _____</p> <p>Bags & Pockets _____</p> <p>Buoys _____</p> <p>Deck float pieces _____</p> <p>Food Containers _____</p> <p>Beverage labels _____</p> <p>Cups & bowls (pieces) _____</p> <p>Cups & bowls (whole) _____</p> <p>Egg cartons _____</p> <p>Fast food containers (whole) _____</p> <p>Fast food containers (pieces) _____</p> <p>Plates & trays (pieces) _____</p> <p>Plates & trays (whole) _____</p> <p>Miscellaneous _____</p> <p>Pieces smaller than a baseball _____</p> <p>Pieces larger than a baseball _____</p> <p>Polyurethane foam _____</p> <p>Spheres _____</p> <p>Stripping (possibly rubber) _____</p> <p>Wrappers _____</p> <p>Packing Material _____</p> <p>Peanuts _____</p> <p>Misc. _____</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

(Continued on Reverse Side)

Figure 1.--Data inventory sheet showing descriptive categories. Form adapted from Center for Marine Conservation National Beach Survey Data Card.

Supplemental Sheet Yes No (Check One)

All Cities

For all cities combined, 81% of all the debris collected was plastic or polystyrene (Fig. 2). Polystyrene, a plastic material, is treated separately based on its physical properties and uses. Miscellaneous debris, composed primarily of grease balls, tar, and slag, represented 12% of all debris. The remaining major debris categories (wood, paper, metal, rubber, glass, and textile) comprised approximately 7% of all debris. A summary of debris in each major category for each city and in all cities combined is presented in Table 2 and Figure 3.

The most abundant category of debris was plastic. Of the cities sampled, Tacoma had the greatest percentage of plastic (84%), due primarily to an unusually large number of plastic pellets/spherules collected in a single sample. Baltimore ranked second highest with 70%. Debris from Seattle contained the lowest percentage of plastic (41%).




Figure 2.--Percent composition of all debris categories in all cities, November 1988 through February 1989. Percentages are based on the total number of items.

Table 2.--Summary of percent composition^a of items of Environmental Protection Agency concern and all debris categories found in marine debris samples collected in U.S. harbors, November 1988 through February 1989. All percentages are rounded to the nearest tenth.

Composition	New York		Boston		Phila- delphia		Balti- more		Miami		Tacoma		Seattle		San Francisco		Oakland		All cities	
	19.5	29.7	34.0	19.5	24.0	85.5	16.4	16.8	29.7	30.4										
Pellets	0.2	0.2	0.9	0.4	0.1	0.0	0.0	0.1	0.1	0.2										
Condoms	0.2	0.1	0.5	0.2	0.0	0.0	0.0	0.0	0.0	0.1										
Tampons	0.1	0.1	0.1	0.6	0.1	0.0	0.0	0.1	0.0	0.1										
Syringes or medical	0.0	0.0	0.0	0.0	0.1	0.0	0.4	0.0	0.0	0.0										
Nets or traps	3.8	0.3	0.6	0.5	1.1	0.2	3.4	0.7	0.3	1.6										
Line or rope	0.1	0.1	0.3	0.3	0.7	0.0	0.1	0.1	0.1	0.1										
Six-pack yokes	4.8	1.9	9.1	16.0	17.3	1.6	8.0	5.3	9.2	6.3										
Plastic bags or sheeting	28.7	32.3	45.5	37.3	43.2	87.4	28.4	23.0	39.4	38.9										
Total	59.1	61.1	64.2	69.9	47.0	84.0	41.3	43.5	56.9	61.3										
All categories	0.0	0.1	0.2	0.9	0.6	0.1	0.4	0.5	0.7	0.2										
Plastic	1.6	0.7	2.1	0.9	3.5	0.3	4.8	2.1	2.2	1.5										
Glass	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.4	0.1	0.1										
Paper	9.9	18.2	24.0	24.5	37.0	12.6	43.6	46.9	34.6	20.3										
Textile	1.2	0.8	0.9	0.5	0.4	0.0	1.1	1.1	0.6	0.8										
Polystyrene	0.2	0.3	0.3	0.7	0.9	0.1	1.1	0.5	0.9	0.4										
Rubber	6.8	0.9	1.1	1.1	6.3	1.3	5.5	3.3	2.9	3.6										
Metal	21.1	18.0	7.2	1.5	4.4	1.5	2.1	1.8	1.2	11.8										
Wood																				
Miscellaneous																				

^aBased on the total number of items found in each city.

ALL CATEGORIES





Figure 3.--Percent composition of all debris categories in each city. Percentages are based on the total number of items.

The second most abundant category of debris was polystyrene. Of the cities sampled, San Francisco had the greatest percentage of polystyrene (47%). Debris from New York contained the lowest (10%). The combined percentages of plastic and polystyrene debris found in each city are given in Figure 4.

The 10 most common items found in the study (Table 3) accounted for 75% of all debris. Four of these, plastic pellets/spherules, miscellaneous plastic pieces, polystyrene pieces smaller than a baseball, and cigarette butts were among the most abundant items found in all nine cities. Plastic pellets/spherules, the raw material, or resin, used in the manufacture of plastic products, constituted the most common item overall; it was also the most common item found in five cities (Boston, Philadelphia, Baltimore, Tacoma, and Oakland) and among the five most common items in two additional cities (New York and San Francisco).

In all, seven of the most abundant items were composed of plastic (five items) or polystyrene (two items). The two remaining categories include miscellaneous (two items) and wood (one item). The three most common types of plastic item included plastic pellets/spherules, miscellaneous plastic pieces, and plastic sheeting shorter than 0.6 m (2 ft). Plastic pellets/spherules comprised 26% of all debris collected. Miscellaneous plastic pieces (13%), and plastic sheeting shorter than 0.6 m (2 ft) (5%) ranked second and fifth overall. Polystyrene spheres (4%) and polystyrene pieces smaller than a baseball (10%) were also common.




Figure 4.--Percent composition of plastic and polystyrene debris in each city. Percentages are based on the total number of items.

Table 3.--Summary of the most common items found in marine debris samples collected from all cities sampled, November 1988 through February 1989.

Category	Debris item	Quantity	Percent ^a
Plastic	Pellets/spherules ^b	11,406	25.90
Plastic	Miscellaneous pieces	5,785	13.14
Polystyrene	Pieces smaller than a baseball	4,350	9.88
Miscellaneous	Grease balls	2,696	6.12
Plastic	Sheeting <0.6 m (2 ft) ^b	2,100	4.77
Polystyrene	Spheres ^b	1,938	4.40
Plastic	Cigarette butts and filters	1,550	3.52
Miscellaneous	Slag	1,503	3.41
Plastic	Food wrappers	959	2.18
Wood	Miscellaneous pieces	923	2.10
Total of most common items		33,210	75.41
Total of all items in all cities		44,037	100.00

^aBased on the total number of items found in each city.

^bItem of Environmental Protection Agency concern.

The EPA-designated items of concern made up 39% of all debris (Fig. 5). They included the following items enumerated in this study:

- Pellets--Plastic pellets/spherules; polystyrene spheres.
- Condoms--Condoms (whole, pieces).
- Tampons--Tampons; tampon applicators; tampon wrappers.
- Syringes or medical--Syringes (whole, pieces, with blood); needle covers; vials and vial caps; insulin bottles.
- Nets or traps--Netting; floats and lures.
- Line or rope--Plastic rope (<0.6 m (2 ft) and >0.6 m (2 ft)); filaments; strapping bands; fishing line (monofilament); textile rope.
- Six-pack yokes (or similar)--Six-pack yokes (or similar) (whole and pieces).
- Plastic bags or sheeting--Plastic bags (<3.8 L (1 gal) and >3.8 L (1 gal)); condiment bags; miscellaneous plastic bags (whole and pieces); plastic sheeting (<0.6 m (2 ft) and >0.6 m (2 ft)).




Figure 5.--Percent composition of items of Environmental Protection Agency concern in all cities. Percentages are based on the total number of items.

Items of concern constituted 87 and 46% of the total debris in Tacoma and Philadelphia, respectively. The lowest percentage of items of concern (23%) was found in San Francisco. Pellets were the most common item of concern found in all cities combined (30% of all debris and 78% of all items of concern), and ranged from 16% of all debris in Seattle to 86% of all debris in Tacoma. In all cities combined, plastic bags/sheeting was the second most common item of concern (6% of all debris; 16% of all items of concern).

Several debris items found during this study are typically associated with human sewage, medical waste, or illegal drug usage, as follows:

Sewage-related debris

Condoms (whole and pieces)	Tampons
Diapers	Tampon applicators
Panty liners	Tampon wrappers
Sanitary items	Fecal matter
Sanitary napkins	

Medical-related items

Syringes (whole and pieces)	Pill vials and caps
Syringes with blood	Cylindrical tubes (whole and pieces)
Needle covers	Tongue depressors
Vials and vial caps	Miscellaneous pills
Miscellaneous medical items	

Drug-related items

Crack vials with caps	Crack vial caps
Crack vials without caps	Illegal substances

Sewage-, medical-, and drug-related debris each comprised <1% of all items in each city (Table 4) except Philadelphia, where approximately 2% of the debris was sewage-related. Philadelphia, Baltimore, and New York had the highest percentages of all three types combined. The combination of these three types made up over 2% of the debris in Philadelphia, and more than 1% of the debris in Baltimore and New York. Exactly 1% of all debris found in all cities combined was sewage-, medical-, or drug-related.

Out of approximately 200 items identified, 26 items were common to all cities. These items were

Plastic

- Bags <3.8 L (1 gal) and >3.8 L (1 gal)
- Bottles <3.8 L (1 gal)
- Caps/lids (whole and pieces)
- Cigar/cigarette wrappers and packs
- Cigar tips
- Cigarette butts and filters
- Cups, spoons, forks, straws
- Food wrappers
- Filaments
- Rope shorter than 0.6 m (2 ft)
- Miscellaneous piece
- Pellets/spherules
- Sheeting <0.6 m (2 ft)
- Coffee stirrers

Paper

- Food wrappers
- Miscellaneous pieces

Polystyrene

- Cups and bowls (pieces)
- Pieces smaller than a baseball
- Peanuts
- Miscellaneous packing material
- Spheres

Wood

- Miscellaneous pieces

Miscellaneous

- Food items
- Grease balls

Of these items, seven items (plastic pellets/spherules, plastic bags >3.8 L (1 gal) and <3.8 L (1 gal), plastic filaments, rope shorter than 0.6 m (2 ft), and two types of polystyrene spheres) were items of EPA concern. None of these items was directly attributable to sewage-, medical-, or drug-related activities.

Table 4.--Number and percent composition^a of sewage-, medical-, and drug-related debris.^b

City	Sewage-related		Medical-related		Drug-related		Total ^c	
	Number	%	Number	%	Number	%	Number	%
New York	63	0.45	32	0.23	119	0.85	214	1.54
Boston	29	0.31	10	0.11	27	0.28	66	0.70
Philadelphia	45	1.59	2	0.07	21	0.74	68	2.40
Baltimore	40	0.92	27	0.62	6	0.14	73	1.68
Miami	4	0.14	3	0.10	1	0.03	8	0.27
Seattle	0	0.00	2	0.28	0	0.00	2	0.28
Tacoma	0	0.00	1	0.02	0	0.00	1	0.02
San Francisco	3	0.09	2	0.06	0	0.00	5	0.15
Oakland	1	0.07	1	0.07	0	0.00	2	0.02
Total	185	0.42	80	0.18	174	0.40	439	1.00

^aBased on the total number of items found in each city.

^bDefined in text.

^cSum of sewage-, medical-, and drug-related debris.

It is interesting to note that, of the 26 items listed above, 7 were related to food packaging or consumption and 3 were related to tobacco use. Food and tobacco generally are packaged in a wrapper or container when purchased; increasingly these containers or wrappers are being made of plastic. Disposal of these wrappers or containers is often careless, especially if the item is consumed during travel. In addition, the plastic wrappers are very lightweight and are easily transported by the wind. Either careless disposal by consumers or wind action over approved disposal sites such as dumpsters and trash receptacles could account for the presence of many plastic food and tobacco containers and wrappers.

East Coast Versus West Coast

Comparison of the results from east coast cities (New York, Boston, Philadelphia, Baltimore, and Miami) and west coast cities (Seattle, Tacoma, San Francisco, and Oakland), showed certain similarities and differences in debris composition (Figs. 6 and 7). Cities on both coasts had nearly the same percentages of plastic debris. Glass, paper, textile, rubber, and metal debris were found in low levels (<1%) on both coasts; these items were less common than wood debris, which was found in very similar proportions on both coasts. In contrast, east coast cities had a higher percentage of miscellaneous debris, which was primarily in the form of grease balls, tar, and slag. The contribution of polystyrene to the total debris was one to two times greater in the west coast cities than in the east coast cities.



ALL CATEGORIESALL CATEGORIES

Figure 6.--Percent composition of all debris categories.
 (A) East coast cities. (B) West coast cities. Percentages
 are based on the total number of items.



ITEMS OF CONCERN (34 % OF DEBRIS)ITEMS OF CONCERN (56 % OF DEBRIS)

Figure 7.--Percent composition of items of Environmental Protection Agency concern. (A) East coast cities. (B) West coast cities. Percentages are based on the total number of items.

Over one-half of all debris items found in west coast cities were items of EPA concern, the majority of which were represented by pellets in Commencement Bay near Tacoma. Nearly one-third of the debris in eastern cities consisted of items of EPA concern, and approximately three-fourths of the items were pellets. Another item of EPA concern, plastic bags/sheeting, was common on both coasts, although proportionally greater on the east coast. Line/rope was more common on the east coast than on the west coast.

Sewage-, medical-, and drug-related debris were uncommon on both coasts (Table 4), and combined they totaled 1% of all debris found. Most of these three debris types were found in east coast cities. The larger presence of these on the east coast is likely to be due to the greater frequency of CSO and storm sewer discharges in eastern cities. No drug-related debris was found in the west coast cities.

Medical Debris

The greatest numbers of medical-related debris items were found in three east coast cities (Table 4): New York (32 items, or 0.23% of all New York debris), Baltimore (27 items, or 0.62% of all Baltimore debris), and Boston (10 items, or 0.11% of all Boston debris). A total of 7 syringes and syringe pieces, including 1 syringe containing blood, were found in New York; 7 syringes and pieces were found in Boston, and 13 syringes and pieces were found in Baltimore. Very little medical debris, only two items of which were syringes, was found in the remaining cities.

All of the syringes found during this study were the 1-cc insulin-dispensing type. In Baltimore, the needles typically were capped at one or both ends, probably indicating that they were used and disposed of by someone who had been instructed as to safe and proper syringe disposal. However, in New York and Boston the syringes usually were in pieces and not capped at either end.

SUMMARY

Plastic debris (including polystyrene) was numerically the largest component of marine debris in surface slicks from every city sampled. Plastic pellets were a significant portion of the plastic debris and were collected in every harbor. Several sewage-, drug-, and medical-related items were found during the study, but these items were not major components of the debris.

These surveys were the first in a continuing series of surveys sponsored under EPA's Harbor Studies Program. The program is providing the first semiquantitative evaluation of marine debris in U.S. harbors. Future surveys are being planned to study additional cities along the east and Gulf coasts, and many of the cities discussed in this study will be resampled.

ACKNOWLEDGMENTS

This work was supported by the EPA. The views and conclusions represent the views of the authors and do not necessarily represent the opinions, policies, or recommendations of EPA.

The authors wish to thank all EPA and Battelle employees who assisted in sample collection and processing, tasks which were typically tedious and, at times, unpleasant. We also wish to thank Margarete Steinhauer for presenting this study at the Second International Conference on Marine Debris.

REFERENCES

- Redford, D. P.
1990. Status of the U.S. Environmental Protection Agency marine debris activities and programs. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- U.S. Environmental Protection Agency.
1988. Floatables investigation. U.S. Environmental Protection Agency, Region II, N.Y., 11 p. + appendixes.

PRELIMINARY REPORT ON THE DISTRIBUTION OF
SMALL-SIZED MARINE DEBRIS IN SURUGA BAY

Mori Yukinawa and Shin-ichi Mio*
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

*Present address: Japan Sea National Fisheries Research Institute, Fisheries Agency, the Government of Japan, 1-5939-22, Suido-cho, Niigata-shi, Niigata, 951 Japan

ABSTRACT

From 17 to 29 March 1988, a survey to collect small marine debris was conducted in Suruga Bay. Tows (10-min) were made at 24 stations using a circular tow net with a diameter of 1.4 m and mesh size of 1.7 mm at the mouth and 0.5 mm at the cod end.

A total of 665 pieces of debris were collected during the survey. Of these, terrestrial debris and plants such as wood made up 50.1% of the total, followed in decreasing order by seaweed (19.5%), Styrofoam (13.8%), plastic sheets (13.2%), and other plastic pieces (3.3%).

Plastic sheet debris was found at most stations, whereas Styrofoam showed a tendency to accumulate in specific areas.

The distribution of each type of small debris corresponded well to the distribution of large debris observed by sightings conducted during the same cruise.

INTRODUCTION

Petrochemical products flowing and thrown into the sea are causing a number of problems. It has been pointed out that marine organisms swallow debris fragments together with their food. These fragments are generated when drifting petrochemical products in the sea are, in the process of their deterioration, broken up by physical factors such as waves. It is necessary to examine these small floating objects in order to understand the changes in size as well as the distribution and movement of drifting petrochemical products over the course of time.

METHODS

This research was carried out at 24 stations in Suruga Bay from 17 to 29 March 1988 (Fig. 1). The nets used for collecting debris were 1.4 m in diameter. The first two-thirds of the net had a mesh size of 1.7 mm; the remaining third a mesh size of 0.5 mm. The net was towed for 10 min at a speed of 3 kn with the mouth of the net half submerged. Collected objects were immediately preserved in formalin and brought back to the laboratory, where they were counted and weighed and the information recorded. Those with maximal dimensions of 5 cm were excluded.

RESULTS

Major petrochemical products collected in this survey were fragments of plastic, plastic sheeting, and Styrofoam. Among the natural debris was terrestrial debris such as wood fragments and straw as well as drifting seaweed (Table 1). A total of 665 pieces of debris were collected. Debris deriving from petrochemical products accounted for 30.4% of the total (Styrofoam fragments 13.8%, fragments of plastic sheeting 13.3%, and plastic fragments 3.3%). Debris of terrestrial origin accounted for 50% of the total (the largest percentage), and seaweed for 19.5%. Plastic sheet fragments were extensively distributed, being collected at 66.7% of the 24 research stations (Fig. 1). The distribution of other plastic and Styrofoam was limited, occurring at only 29.2 and 20.8% of the stations, respectively. Natural debris of terrestrial origin was collected at 66.7% of the stations and seaweed at 62.5% of the stations.

Plankton, mainly Copepoda, sardine fry, and fish eggs were collected at all the research stations. These types of plankton numbered more than 1,000 at each station.

The highest densities of marine debris were found all across the middle of Suruga Bay. Research stations with high densities of plastic sheet fragments were continuous, as seen in the case of stations 12, 13, 22, and 17. A similar distribution pattern was observed for all drifting objects. The distribution of Styrofoam fragments was limited, but high densities were found in geographical positions similar to stations 7, 6, and 1.

Currents and winds were considered to be major factors in moving this debris. It has been reported that the surface current in Suruga Bay can be affected by the Kuroshio, which flows eastward off the bay. As shown in Figure 2, Inaba (1988) pointed out two such instances. The course of the Kuroshio during the survey period was more offshore, corresponding to Case 2 in Figure 2 (Meteorological Agency of Japan 1988). It is assumed that the flow of outer oceanic water is from west of the mouth of the bay and divides into two currents near the center of the bay, with one current moving around the mouth of the bay clockwise and the other moving counterclockwise toward the inner side of the bay. The area of the highest densities of marine debris is on the boundary between these currents, which would represent a front. Strong northeast winds were blowing for several days during the survey period, but the distribution of marine debris was considered to be affected more by surface currents than by the winds.




Figure 1.--Stations (St.) and data collected in minute marine debris survey. At each station, numbers from the top down indicate plastic pieces, plastic sheets, Styrofoam, land plants, and seaweed, in that order.

Table 1.--Sampling location and composition of minute marine debris in Suruga Bay.

Station	Time	Latitude N	Longitude E	Plastic pieces	Plastic sheets	Plastic sheets	Styrofoam	Land plants	Seaweed	Copepoda	Eggs	Larvae
1	12.50	34°43.72'	138°41.68'	1	--	--	17	39	8	42	10	--
2	7.17	34°51.19'	138°22.56'	10	--	--	--	27	7	--	--	--
3	11.56	34°24.30'	138°59.13'	--	--	--	--	--	7	--	3,230	--
4	16.42	34°36.32'	138°18.60'	--	2	--	--	2	--	58	--	7
5	7.10	34°37.30'	138°16.22'	--	--	--	--	7	1	1,120	84	--
6	11.04	34°43.58'	138°22.92'	--	--	--	9	16	14	15	--	--
7	16.40	34°54.90'	138°28.68'	--	6	--	63	39	34	--	--	--
8	7.17	34°50.75'	138°22.01'	--	3	--	--	1	--	37	--	--
9	11.35	34°47.72'	138°40.85'	--	3	--	--	3	3	25	154	--
10	14.42	35°04.05'	138°34.32'	--	--	--	--	17	5	--	--	--
11	9.11	35°02.33'	138°32.60'	--	1	--	--	15	1	26	15	--
12	10.31	35°02.70'	138°44.63'	--	22	--	--	16	--	15	46	--
13	12.45	34°57.33'	138°32.44'	--	12	--	--	7	--	25	12	--
14	17.05	34°41.56'	138°20.30'	2	1	--	--	3	--	43	82	--
15	7.44	34°34.05'	138°19.06'	--	1	--	--	--	8	--	--	--
16	11.54	35°05.48'	138°37.65'	--	--	--	--	18	--	22	57	--
17	17.03	34°45.49'	138°32.08'	4	17	--	1	1	9	121	15	--
18	12.02	34°14.81'	138°30.55'	1	3	--	--	--	26	--	19	34
19	16.32	34°51.26'	138°43.14'	2	5	--	--	--	3	12	38	--
20	7.11	34°48.68'	138°44.37'	2	1	--	--	--	2	--	--	1,540
21	7.25	34°50.28'	138°37.42'	--	1	--	--	--	--	--	3	--
22	12.08	34°49.17'	138°26.96'	--	9	--	--	--	2	--	--	6
23	7.14	34°51.66'	138°22.70'	--	--	--	2	122	--	--	--	6
24	13.03	34°05.22'	138°36.72'	--	1	--	--	--	--	--	--	5




Figure 2.--Diagram showing the relationship of the Kuroshio to the surface current of Suruga Bay (Inaba 1988).

Plankton and marine debris showed an inverse correlation. Areas where plankton were found in large quantities were influenced a great deal by the open ocean. It is reasonable to assume that plankton and marine debris showed different distributions because the plankton, being totally submerged are not influenced at all by the wind.

Small floating objects were found most frequently in areas influenced heavily by coastal currents. The distribution of drifting petrochemical product fragments corresponded well to that of the larger floating objects. It is therefore conjectured that petrochemical products drifting in the sea gradually deteriorate and are broken into small fragments.

More detailed studies will be needed to find out the distribution and abundance of Styrofoam fragments.

REFERENCES

- Inaba, H.
1988. Oceanographic conditions in Suruga Bay. Bull. Jpn. Soc. Fish. Oceanogr. 52:236-240. [In Jpn.]
- Meteorological Agency of Japan.
1988. Ten-day marine report. Meteorological Agency of Japan 1494.
-

DISTRIBUTION, ABUNDANCE, AND SOURCE OF ENTANGLEMENT
DEBRIS AND OTHER PLASTICS ON ALASKAN BEACHES, 1982-88

Scott W. Johnson
Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

ABSTRACT

Sixty kilometers of outer coast beaches at 25 locations in Alaska were surveyed from 1982 to 1988 to determine distribution, composition, quantity, deposition, and source of plastic debris washed ashore. Approximately 67% of all plastic debris found was fishing gear (e.g., net fragments, rope, floats) and 33% was packaging material (e.g., plastic bags, bottles). Debris found which could entangle marine mammals, seabirds, and fish included trawl web, rope, packing straps, and monofilament gillnet. Monofilament gillnet was not abundant (usually <5 pieces/km) on beaches, but trawl web was found on beaches throughout Alaska and exceeded 10 fragments/km at more than 50% of the locations sampled. Foreign fisheries were the source of most (98%) of the monofilament gillnet washed ashore; the source of trawl web is shifting from foreign to domestic fisheries.

Trends in composition and abundance of plastic debris were monitored at three sites: Amchitka Island, Middleton Island, and Yakutat. Amchitka Island had similar quantities (~300 items/km) of total plastics in 1982 and 1987, although the amount of trawl web at this site continued to increase. Quantities of plastic debris on Middleton Island remained similar from 1984 to 1987 (average 860 items/km), with the exception of an approximate 33% decline in 1985 from the 4-year average. Near Yakutat, the quantity of trawl web deposited ashore increased from 8.8 to 10.1 fragments/km/year from 1985 to 1988. Continuing the surveys of these benchmark beaches will help determine whether recent mitigating legislation is effective in reducing the disposal of entanglement debris and other plastics at sea.

INTRODUCTION

Marine pollution has become a major environmental concern in the 1980's. One form of marine pollution that has attained international attention is plastic debris discarded or lost in the world's oceans.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

Plastics are of particular concern because they persist in the environment for years, endangering marine animals and man. Seabirds and sea turtles can ingest pieces of plastic that block their digestive tracts (Balazs 1985; Day et al. 1985); seabirds, fish, and invertebrates can become entrapped in derelict gillnets (DeGange and Newby 1980; High 1985); marine mammals can become entangled in fragments of trawl web, packing straps, and rope (Fowler 1987; Stewart and Yochem 1987); and ships can be disabled from plastic debris which fouls props or cooling intakes (Wallace 1985).

Most plastics are lightweight, float at or near the ocean surface, and often wash ashore. Plastic debris is common on Alaskan beaches because of the loss or discard of fishing gear (e.g., trawl web, rope, and floats) and other plastic debris from large commercial fishing fleets operating in the North Pacific Ocean and Bering Sea (Merrell 1985; Uchida 1985). Plastic debris washed ashore represents, to some degree, the types and quantities lost or discarded at sea. Beach surveys may be the best method of evaluating whether recent mitigating legislation (MARPOL Annex V) to reduce the input of plastics into the sea is effective.

The National Marine Fisheries Service (NMFS) has conducted beach surveys for plastic debris on Alaskan beaches periodically since 1972. The objective of this paper is to examine recent trends in the distribution, composition, quantity, deposition, and source of plastic debris on Alaskan beaches based on surveys from 1982 to 1988; the emphasis was on entanglement debris (trawl web, gillnet, rope, and packaging straps) at study sites that were repetitively sampled since 1982. The occurrence of trawl web is discussed in detail because it is one of the most abundant entanglement debris items found on Alaskan beaches (Merrell and Johnson 1987; Johnson and Merrell 1988), and it is the principal item entangling northern fur seals, *Callorhinus ursinus*, on the Pribilof Islands (Fowler 1987). Additional information on past NMFS studies can be obtained from Merrell (1980, 1984, and 1985).

METHODS

Approximately 60 km of outer coast beaches at 25 locations in Alaska have been surveyed for plastic debris since 1982 (Fig. 1). Locations of beaches surveyed at least twice as benchmarks include: Amchitka Island in the Aleutians; Middleton Island in the central Gulf of Alaska; and beaches near Yakutat in the eastern Gulf of Alaska (Fig. 1).

Beaches were surveyed primarily during summer (June-September) in all locations with the exception of those near Yakutat. Ten beaches on Amchitka Island were surveyed once in September 1982 and again in September 1987; three beaches on Middleton Island were surveyed once in either July or early August 1984 through 1987; and eight beaches near Yakutat were surveyed once in September 1985, four times in 1986 and 1987 (January, April, July, September), and twice in 1988 (March and September). Five of the eight Yakutat beaches were surveyed once in September 1984.

Survey methods were similar for all beaches (Merrell 1985). Most beaches were 1 km in length. The survey area for each beach included the




Figure 1.--Locations of beaches surveyed for plastic debris and quantity of trawl web fragments (number per kilometer) found in Alaska, 1982-88.

intertidal zone between the water's edge and the seaward limit of terrestrial vegetation at the upper limit of normal high tide. All plastic debris visible from walking height was counted (i.e., pieces ≥ 5 mm, and trawl web and monofilament gillnet fragments with five or more complete meshes). Rope of any diameter was counted if it was ≥ 1 m in length. We did not count pieces (e.g., gillnet floats and plastic bottles) if they were less than one-half their original size. We either weighed or estimated the weight of trawl web fragments depending on size and location: whether they were loose on the beach, buried, or snarled on drift logs. Stretch mesh was measured (knot to knot inside measure) for one representative mesh of each net fragment sampled. We did not search for debris within piles of drift logs or seaweed.

Beginning in 1985, all trawl web fragments at Yakutat were tagged with a small metal tag or removed and discarded inland from the beach. Trawl web fragments that were tagged and remained onshore could therefore be distinguished from new (not tagged) fragments, making it possible to determine deposition by season and year. At Middleton and Amchitka Islands, trawl web and gillnet fragments were painted with orange dye so that they could be identified in future surveys.

To determine trends in accumulation of all types of plastic debris, a 1-km beach on Middleton Island was cleared of all surface debris annually from 1984 to 1987. Debris was moved to terrestrial areas above the high-tide zone. Debris too large to move, partially buried, or snarled on drift logs was marked with paint, flagging, or tags for identification in future surveys.

The only major change in the sampling procedure was made in 1986 and 1987 when all beaches were subdivided into ten 100-m increments, thereby providing ten different data sets for each 1-km beach. This change was designed to improve the statistical precision of debris estimates (Ribic and Bledsoe 1986).

Differences in quantities of entanglement debris items on Amchitka Island were tested by paired t-tests, where observations in 1982 and 1987 were paired for each of ten 1-km beaches. Differences in quantities of individual debris items between Amchitka and Middleton Islands in 1987 were tested by t-tests. The association between quantity of trawl web and total plastic debris found on Alaskan beaches was determined by linear correlation.

RESULTS

Derelict trawl web was found on sampled beaches throughout Alaska (Fig. 1). At over 50% of the locations sampled, trawl web exceeded 10 fragments/km of beach. Locations with the highest quantities of trawl web included Little Tanaga Island in the Aleutians (216 fragments/km), Kayak Island in the central Gulf of Alaska (92 fragments/km), Amchitka Island (55 fragments/km), and Noyes Island in southeast Alaska (53 fragments/km) (Fig. 1).

Trawl web was significantly correlated ($P < 0.05$; $r = 0.37$) with the quantity of total plastic debris (all types) found per kilometer of beach. Thus, beaches that accumulated many fragments of trawl web generally also accumulated numerous other plastics. Locations with the highest quantities of total plastics included Noyes Island (1,330/km), Kayak Island (1,142/km), and Middleton Island (988/km) (Fig. 1).

Composition of total plastic debris (based on number of individual items) on Amchitka Island beaches was similar in 1982 and 1987. Likewise, composition of plastic debris on Middleton Island was similar in all years (1984-87). At both locations in 1987, nearly two-thirds of all items found were derelict fishing gear (Table 1).

Quantities of entanglement debris changed on Amchitka Island from 1982 to 1987, but only rope increased significantly ($P < 0.05$) (Fig. 2). Trawl web, strapping, gillnet, and gillnet floats (possible indicator of quantity of gillnet lost), either increased or decreased, but not significantly (Fig. 2). Because some items increased and some decreased, total plastics were similar in 1982 and 1987 (~300 items/km).

Table 1.--Percent composition of derelict fishing gear based on number of plastic debris items found on Amchitka and Middleton Islands, Alaska, 1987.

Debris items	Percent of total	
	Amchitka	Middleton
Derelict fishing gear	68	62
Rope	31%	7%
Trawl web	26%	4%
Floats	20%	82%
Straps	16%	3%
Gillnet	1%	1%
Miscellaneous	6%	3%
Packaging material	28	35
Personal effects	2	2
Miscellaneous	2	1

The number of trawl web fragments found on Amchitka Island beaches has steadily increased since 1972 (Fig. 3); the average weight of individual fragments, however, has decreased from 11 kg in 1974 to 4 kg in 1987. The frequency of occurrence of different trawl web mesh sizes measured on Amchitka Island was similar in 1982 and 1987 (Fig. 4). In both years, the most common mesh size was 101-150 mm; approximately one-third of the fragments had mesh sizes >150 mm.

Quantities of entanglement debris remained relatively stable on Middleton Island from 1984 through 1987 (Fig. 5). During these 4 years, trawl web averaged 24 fragments/km of beach; rope, 51 pieces/km; straps, 16/km; and gillnet fragments, 4/km. Gillnet floats increased 58% from 287/km in 1984 to 454/km in 1987. Total plastics found on Middleton Island were similar in 1984, 1986, and 1987. In 1985, however, there was a 33% decline in total plastics from the 4-year average of 860 items/km (Fig. 5). Differences in quantities of debris by location were evident between Middleton and Amchitka Islands in 1987 (Table 2). Significantly ($P < 0.05$) more trawl web was found on Amchitka than on Middleton Island, whereas significantly ($P < 0.001$) more gillnet floats and total plastics were found on Middleton Island. Although not significant, twice as much gillnet was found on Middleton Island as on Amchitka Island.

A 1-km beach on Middleton Island, cleared of all plastic debris annually from 1984 to 1987, accumulated debris quickly, sometimes within 1 year (Fig. 6). Trawl web, gillnet, and rope, cleared from this beach in 1986, accumulated to previous or higher quantities by 1987. Entanglement debris accumulated in a similar proportion each year; rope was the most abundant, usually followed by trawl web, gillnet, and closed straps (Fig. 6).




Figure 2.--Quantities (mean \pm SD) of entanglement debris and total plastics found on Amchitka Island, Alaska, in 1982 and 1987. Data based on ten 1-km beaches. Asterisk denotes significant difference between years $p < 0.05$.

Total deposition of trawl web at Yakutat was similar from 1985 to 1988 (range 8.8 to 10.1 fragments/km/year) (Table 3). More fragments, however, washed ashore during the fall-winter months (Oct.-Apr.) than the spring-summer months (May-Sept.). Of the beach locations examined more than once, deposition of trawl web was greatest on Amchitka Island, followed by Middleton Island and Yakutat. Some locations, such as Little Tanaga Island, Kayak Island, and Noyes Island, accumulated more trawl web than the above or adjacent locations, probably because of their favorable orientation to major ocean currents, prevailing storm winds, or increased fishing effort and loss of gear in nearby waters.

At present, the source of trawl web washed ashore is shifting from foreign vessels to domestic vessels as U.S. trawl fisheries replace foreign trawl fisheries in the North Pacific Ocean and Bering Sea in the latter 1980's (Cotter et al. 1988) (Fig. 7). Most (98%) monofilament gillnet washed ashore, however, is from foreign high seas fisheries (Fig. 7) because monofilament nylon gillnets, with the exception of a small herring fishery, are banned in Alaska (Uchida 1985). The most common (42%) mesh size of gillnet washed ashore was 110 mm stretch mesh (Table 4). Based on mesh




Figure 3.--Number and weight of trawl web fragments found on Amchitka Island, Alaska, from 1972 to 1987. Ten 1-km beaches surveyed in each year. Data for 1972-74 from Merrell (1985).




Figure 4.--Percent occurrence of different mesh sizes of trawl web fragments found on Amchitka Island, Alaska, in 1982 (n = 333) and 1987 (n = 282). The X-axis label is upper limit of interval.

size, likely sources of foreign gillnet are high sea fisheries (Japan, Taiwan, Korea) for squid and salmon (Merrell 1985; Uchida 1985).

DISCUSSION

The widespread distribution and continual accumulation of plastic debris on outer coast beaches of Alaska are indicative of the vast quantities of debris lost or discarded into the North Pacific Ocean and Bering Sea. Annually, an estimated 1,664 metric tons of plastic debris are lost or discarded from fishing vessels in Alaskan waters (Merrell 1980). Although large quantities of plastic debris were found on many Alaskan beaches, it was not evenly distributed. Some beaches with large quantities of trawl web (>10 fragments/km) and other plastic debris were adjacent to locations with small quantities of debris (Fig. 1). Accumulation of debris on beaches depends upon the orientation of the beach to major ocean currents and prevailing winds. Even within a given location, debris abundance can differ dramatically; the windward side of Middleton Island, for example, had 15 times the amount of debris found on the leeward side of the island (Johnson and Merrell 1988). Thus, when interpreting results




Figure 5.--Quantities of entanglement debris and total plastics found on Middleton Island, Alaska, from 1984 to 1987. Data based on two 1-km beaches.

Table 2.--Quantities of entanglement debris and total plastics found on Amchitka and Middleton Island beaches, Alaska, 1987 (* $p < 0.05$; ** $p < 0.001$); n = number of 100-m sections.

Debris type	Number per 100 m	
	Amchitka n = 50	Middleton n = 18
Fishing gear		
Trawl web	5.5*	2.2
Rope	6.7	4.1
Strap	3.4	1.7
Gillnet	0.2	0.4
Gillnet floats	2.5	43.8**
Total plastics	31.0	95.0**

Table 3.--Deposition of trawl web on eight 1-km sections of beach at Yakutat, Alaska, from 1985 to 1988.

Beach	Number of trawl web fragments deposited ashore									
	1985-1986			1986-1987			1987-1988			
	Oct.-Apr.	May-Sept.	Total	Oct.-Apr.	May-Sept.	Total	Oct.-Mar.	Apr.-Sept.	Total	Total
1	8	3	11	4	2	6	7	2	9	
2	4	0	4	6	1	7	6	6	12	
3	3	4	7	8	2	10	7	3	10	
4	12	7	19	16	0	16	7	10	17	
5	12	2	14	9	0	9	8	3	11	
6	0	2	2	3	1	4	2	2	4	
7	1	0	1	4	0	4	2	4	6	
8	10	2	12	18	1	19	7	5	12	
Total	50	20	70	68	7	75	46	35	81	
Mean per kilometer per year			8.8			9.4				10.1




Figure 6.--Accumulation of entanglement debris on a 1-km beach on Middleton Island, Alaska, that was cleared of all debris annually from 1984 to 1987.

of surveys, knowledge of local ocean currents and prevailing winds is necessary.

Composition of plastic debris was nearly identical on Amchitka and Middleton Islands. In both locations in 1987, over 60% of the debris found on beaches was fishing gear. This does not seem unusual, considering that 5,500 km of trawl net and 170,000 km of gillnet are available to various fisheries in the North Pacific (Uchida 1985). Of the three benchmark locations, debris washing ashore on remote Amchitka and Middleton Islands is probably most representative of the types and quantities lost or discarded at sea.

With the exception of rope, which increased significantly, quantities of entanglement debris did not change significantly on Amchitka Island from 1982 to 1987. Trawl web fragments, however, did increase from 34 to 55 fragments/km, continuing the upward trend of earlier years. The average weight of a fragment of trawl web found on Amchitka Island in 1987 was 4 kg, and some of the fragments were rectangular in shape, indicating they




Figure 7.--Major trawl and gillnet fisheries in the North Pacific Ocean and Bering Sea. Adapted from Low et al. (1985) and Merrell (1985). Broken arrows indicate major ocean currents (Reed and Schumacher 1985).

may have been patches discarded overboard from commercial trawlers from net-mending operations. Berger and Armistead (1987) estimated that from 1982 to 1984, over 2,700 pieces of trawl web were discarded overboard into Alaskan waters from net-mending operations.

Although the number of trawl web fragments that washed ashore continued to increase on Amchitka Island, the frequency of occurrence of different mesh sizes remained stable. Approximately one-third of the fragments in both 1982 and 1987 had mesh sizes >150 mm. These are the mesh sizes most likely to entangle northern fur seals (Scordino 1985; Fowler 1987). Similar occurrences of mesh sizes have been reported for other beach locations in Alaska (Johnson 1989). Therefore, assuming trawl web washed ashore is representative of that which is floating at sea, approximately one-third of the derelict trawl web at sea could entangle fur seals.

On Middleton Island, quantities of entanglement debris remained relatively stable from 1984 through 1987, and were generally lower than quantities found on Amchitka Island. More rope and strapping and significantly

Table 4.--Mesh sizes of gillnet fragments found on Alaskan beaches from 1982 to 1988, and probable fishery sources (Chen 1985; Gong 1985; Uchida 1985; United States-Taiwan Bilateral Meeting 1988).

Mesh size (mm)	Number of fragments				Fishery ^a
	Amchitka Island	Middleton Island	Yakutat	Total	
55	0	1	0	1	Herring--US
95	0	1	0	1	Squid--T,K
100	0	1	0	1	Squid--T,K
105	0	1	0	1	Squid--T,K
110	0	20	7	27	Squid--T,K; Salmon--JL
115	8	7	4	19	Squid--T,K,J; Salmon--JL
120	1	4	9	14	Squid--T,J; Salmon--JM
130	0	0	1	1	Salmon--JM
Total	9	35	21	65	

^aUS - United States, T - Taiwan, K - Korea, J - Japan, JL - Japanese land-based, JM - Japanese mothership.

more trawl web were observed on Amchitka Island than on Middleton Island in 1987, probably because of the proximity of Amchitka Island to concentrated trawl fisheries in the North Pacific Ocean and Bering Sea (Fig. 7). Gillnet fragments and floats, however, were more abundant on Middleton Island than on Amchitka Island, even though Amchitka Island is closer to gillnet fisheries in the North Pacific Ocean and Bering Sea (Fig. 7). This may be due to the eastern direction (towards North America) of the subarctic ocean current (Reed and Schumacher 1985), which may transport debris from the high-seas squid and Japanese land-based salmon fisheries (Merrell 1985) into the Gulf of Alaska and favor deposition on Middleton Island.

The quantity of debris washed ashore is affected by frequency and intensity of storms, changes in ocean currents, winds, fishing effort, and areas fished. At Amchitka Island, total plastics remained at about 300 items/km in both 1982 and 1987. At Middleton Island, however, there was a 33% reduction in total plastics in 1985, possibly the result of a change in ocean currents or an unseasonable storm which may have redistributed debris from the beach. By 1986 and in 1987, debris had accumulated on beaches on Middleton Island to quantities near those observed in 1984 (~900 items/km). A decline in quantity of debris on beaches near Yakutat was also reported in 1985 (Merrell and Johnson 1987), supporting the concept that there may have been a change in ocean conditions affecting the accumulation of debris on beaches throughout Alaska. Thus, when monitoring trends in abundance, it is best to sample each beach location at the same time each year in

order to document the variability between years due to changes in ocean conditions or fishing effort.

Of the entanglement debris washed ashore, the reason for the scarcity of gillnet is still unclear. Gillnet is perhaps the most likely of all gear types to be lost (Uchida 1985) but it is one of the least abundant entanglement debris items found on Alaskan beaches. Approximately 1,609,000 km (1 million mi) of gillnet are fished each year in the North Pacific Ocean, of which an estimated 965 km (600 mi) are lost or abandoned each year (Eisenbud 1985). A possible explanation for the lack of gillnets on Alaskan beaches is that they may sink to the ocean bottom from the weight of marine growths (e.g., algae, barnacles) and the carcasses of marine mammals, seabirds, and fish. In some cases, gillnets may drift at mid-depths, get stranded farther offshore in intertidal areas, and never reach the beach. Because derelict gillnets tend to collapse and "roll up" relatively quickly (Gerrodette et al. 1987), they may form a better substrate for marine growths and thereby attract fish and other predators which may get entangled, ultimately causing the net to sink. Trawl web, on the other hand, usually does not "roll up" like gillnet and does not appear to form a suitable substrate for collecting marine growths. This may explain why more trawl web washes ashore than gillnet.

The short period of time (sometimes within 1 year) in which plastic debris accumulated on a beach on Middleton Island that had been cleared of all debris suggests that a substantial amount of debris is probably adrift at sea. Johnson and Merrell (1988) reported a 40% accrual of new debris (previously unseen) on an Alaskan beach in just a 4-month period. The rapid accumulation and, often times, disappearance of debris on beaches are largely controlled by storms. Storms are primarily responsible for depositing debris ashore and removing or redistributing debris already stranded; some of the debris is washed inland to terrestrial areas or buried by sand (Johnson 1989).

Frequent sampling and tagging of trawl web fragments at Yakutat indicates that most fragments are washed ashore in the fall-winter months due to storms. Shiber (1982) also reported an increased deposition of plastic debris in winter on beaches in the Mediterranean Sea. The increase in deposition of trawl web at Yakutat from 8.8 fragments/km in 1985-86 to 10.1 fragments/km in 1987-88, is consistent with the increase in trawl web observed on beaches at Amchitka Island from 1982 to 1988. The reason for the increased deposition of trawl web on Alaskan beaches is unclear; although the number of fragments has increased, the areas fished and the total number of vessels (~300) operating off Alaska have remained relatively steady since 1978 (Low et al. 1985).

Monitoring plastic debris and derelict fishing gear on beaches in Alaska and in other locations may be the best method of evaluating whether the input of plastics into the sea is decreasing because of compliance with MARPOL Annex V. Monitoring plastic debris abundance at sea by aircraft and ship surveys may work, but isn't feasible considering the cost and the immense areas to be covered.

At present, beach surveys are an effective method to determine types, sources, and composition of plastic debris that washes ashore. Trends in abundance of plastic debris may be more difficult to determine because of the variability in the accumulation of debris in different locations and years. Therefore, a better understanding is needed of the interrelationship of ocean currents, storms, and drift patterns, and their effects on the distribution of plastic debris in the North Pacific. In addition, information is needed on the length of time plastic debris remains at sea once it is lost or discarded. Some answers may be gained by releasing marked floats at specific locations in the North Pacific Ocean and Bering Sea and following their recovery.

Regardless of limitations of beach surveys, by establishing benchmarks and continuing to sample at these locations at least once a year at approximately the same time, a trend should become evident as to whether quantities of debris are increasing, decreasing, or remaining the same. Alaskan beaches, specifically Amchitka and Middleton Islands, will serve as long-term benchmarks to monitor plastic pollution because: 1) they are remote from urban sources of pollution, 2) they continually accumulate debris, and 3) a data base of several years already exists.

In summary, plastic debris is found on many outer coast beaches throughout Alaska and most is composed of fishing gear. Rope and trawl web are the two most abundant entanglement debris items found; they continue to wash ashore in some locations in an increasing number. Monitoring debris on beaches in Alaska and elsewhere in the coastal United States for the next several years may help to determine if mitigating legislation is reducing the entry of entanglement debris and other plastics into the ocean.

REFERENCES

Balazs, G. H.

1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS NOAA-TM-NMFS-SWFC-54.

Berger, J. D., and C. E. Armistead.

1987. Discarded net material in Alaskan waters, 1982-1984. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-110, 66 p.

Chen, T. F.

1985. High sea gill net fisheries of Taiwan. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 253-256. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Cotter, L., E. Eckholm, C. Blackburn, and R. Bayliss.

1988. The effects of MARPOL, Annex V., on the ports of Kodiak and Unalaska. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 88-26, 64 p.

- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- DeGange, A. R., and T. C. Newby.
1980. Mortality of seabirds and fish in a lost salmon driftnet. Mar. Pollut. Bull. 11:322-323.
- Eisenbud, R.
1985. Problems and prospects for the pelagic driftnet. Boston Coll. Environ. Aff. Law Rev. 12:473-480.
- Fowler, C. W.
1987. Marine debris and northern fur seals: A case study. Mar. Pollut. Bull. 18:326-335.
- Gerrodette, T., B. K. Choy, and L. M. Hiruki.
1987. An experimental study of derelict gill nets in the central Pacific Ocean. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-18, 12 p.
- Gong, Y.
1985. Distribution and migration of flying squid, *Ommastrephes bartrami* (LeSueur), in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 109-129. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- High, W. L.
1985. Some consequences of lost fishing gear. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 430-437. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Johnson, S. W.
1989. Deposition, fate, and characteristics of derelict trawl web on an Alaskan beach. Mar. Pollut. Bull. 20:164-168.
- Johnson, S. W., and T. R. Merrell.
1988. Entanglement debris on Alaskan beaches, 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-126, 26 p.
- Low, L.-L., R. E. Nelson, Jr., and R. E. Narita.
1985. Net loss from trawl fisheries off Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 130-153. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Merrell T. R., Jr.

1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.

1985. Fish nets and other plastic litter on Alaska beaches. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 160-182. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Merrell, T. R., and S. W. Johnson.

1987. Surveys of plastic litter on Alaskan beaches, 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-116, 21 p.

Reed, D. K., and J. D. Schumacher.

1985. On the general circulation in the subarctic Pacific. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 483-496. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Ribic, C. A., and L. J. Bledsoe.

1986. Design of surveys for density of surface marine debris in the North Pacific. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 86-12, 69 p.

Scordino, J.

1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Shiber, J. G.

1982. Plastic pellets on Spain's 'Costa del Sol' beaches. *Mar. Pollut. Bull.* 13:409-412.

Stewart, B. S., and P. K. Yochem.

1987. Entanglement of pinnipeds in synthetic debris and fishing net and fishing line fragments at San Nicolas and San Miguel Islands, California, 1978-1986. *Mar. Pollut. Bull.* 18:336-339.

Uchida, R. N.

1985. The types and estimated amounts of fish net deployed in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 37-108. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

United States-Taiwan Bilateral Meeting.

1988. Report of the 1988 U.S.-Taiwan bilateral meeting on assessment of North Pacific fisheries resources, 13-15 January 1988, Northwest and Alaska Fisheries Center, Seattle, WA.

Wallace, N.

1985. Debris entanglement in the marine environment: A review. *In* R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 259-277. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

A SURVEY OF PLASTICS ON WESTERN ALEUTIAN ISLAND
BEACHES AND RELATED WILDLIFE ENTANGLEMENT

Albert M. Manville II
Defenders of Wildlife and the
Entanglement Network Coalition
Washington, D.C. 20036, U.S.A.

ABSTRACT

A 10-day survey of 25 beaches (mean length of beach surveys = 149 m (162 yd)) on seven different islands (Attu, Agattu, Shemya, Buldir, Kiska, Little Kiska, and Adak) in the outer Aleutian Islands was conducted 12-20 July 1988, using the U.S. Fish and Wildlife Service's research vessel MV *Tiglax* as a base. Sites were randomly selected, and beaches were surveyed for all plastic from sea level to high storm tide level. Representative plastic samples were collected and all beaches photographed. Of the total 3.7 km (2.3 mi) of beach observed, 3,153 plastic objects were counted, representing 67 different finished plastic products. Debris was identified from Japan, the U.S.S.R., South Korea, People's Republic of China, Taiwan, Norway, and the United States. Most prevalent were items from Japan; of those that were identifiable, most were fishing related.

A precipitous decline in the Steller's sea lion, *Eumetopias jubatus*, was noted on Attu Island (77% decrease since 1979), where pinniped surveys were conducted. The results coincide with a reported 65% overall reduction in the western Aleutian Islands population of Steller's sea lions over the past 10 years. Plastics are suspected of contributing to their decline. An adult bull sea lion on Buldir Island was photographed with a strapping band and massive entanglement scar around its neck, with reports of two other entangled, scarred, but live sea lions on Kiska Island, and one on Agattu Island. Some two dozen dead seabirds were discovered during the beach surveys wrapped in plastic although exact cause of death could be ascertained for only one. The *Tiglax* was temporarily entangled in rope from an apparently active brown king crab, *Paralithodes camtschaticus*, pot.

There was a statistically significant difference in the amount of plastic found on beaches in protected coves versus that discovered on open, unprotected beaches. There was also a statistically significant difference in fishing-related versus

non-fishing-related plastics spotted on the beaches surveyed. If the amount of plastic located on these beaches is at all indicative of that found elsewhere on Alaska's 57,924 km (36,000 mi) of shoreline, plastic debris poses a serious potential problem for fish and wildlife.

INTRODUCTION

Worldwide, plastics in the marine environment alone have been suggested to be as great a cause of mortality to marine mammals, seabirds, and sea turtles as are oil spills, pesticide poisoning, or contaminated run-off (Schneidman 1987). It is postulated that if all dumping and discarding of plastics were to stop immediately, plastics would continue to wash ashore for at least another 100 years (R. J. Wilber, Sea Education Association, Woods Hole, Mass., pers. commun.).

Reports of the presence and impacts of plastic debris in the North Pacific Ocean are fairly common in the recent scientific, popular, and governmental literature (Manville 1988). From the standpoint of origin, plastic debris can be classified as either land-based or ocean-going. Although attempts have been made to quantify at-sea plastic debris in the North Pacific and elsewhere, these attempts are difficult and yield only rough estimates. Dahlberg and Day (1985), for example, found more than 80% of the debris sighted at sea in the North Pacific to be plastic, with over 33% of this consisting of pieces of expanded polystyrene (e.g., cups, floats, boxes). Their observations were limited to floating debris, however, which does not include plastic materials denser than seawater.

Ignell and Dahlberg (1986) surveyed 7,337 km (3,960 mi) of the central and western North Pacific Ocean, and located 1,802 man-made objects adrift on the sea surface, 61 and 26% of these plastic and Styrofoam, respectively. The proportion of plastic materials they found was consistent with that found by Venrick et al. (1973), Shaw and Mapes (1979), and Dahlberg and Day (1985).

Because of the growing concerns about the aesthetic deterioration of our nation's coastline--including beaches in the North Pacific Ocean--a number of recent beach cleanup surveys have been conducted (e.g., Centaur Associates and the Center for Environmental Education (CEE) 1986), but their findings tend to emphasize floatable plastics while often excluding those plastics denser than seawater.

Ghost nets--lost or discarded nets or net fragments, especially drift gillnets--which can continue to fish for years, were reported by Manville (1988) as among the most damaging forms of plastic debris that entangle fish and wildlife in the North Pacific Ocean and elsewhere. The nets sometimes sink from the weight of dead animals, seaweed, or barnacles, and continue to catch fish on the oceans' bottoms. They also may ball up and continue to float, or wash ashore. Also reported were packing bands, six-pack yokes, nets, net fragments, and other plastics which bind and/or strangle virtually every species of marine mammal, sea turtle, seabird,

many varieties of fish, and numerous invertebrates (such as lobsters and crabs).

Fowler (1982, 1987) and Fowler and Merrell (1986) reported that perhaps the best documentation of the results of entanglement in the North Pacific involves northern fur seal, *Callorhinus ursinus*. Extensive data, including the incidence of entanglement scars, were collected from 1967 through 1984 from young male seals killed in the annual commercial seal harvest on the Pribilof Islands, Alaska. These and other data indicated an alarming trend. The population is declining annually at 4-8%; its numbers are now less than half those of 30 years ago. Entanglement, particularly in trawl net fragments, plastic packing bands, and other plastic trash, is believed to be a contributing and perhaps even significant factor in the species' decline. Northern fur seals are presently listed as "depleted" under the Marine Mammal Protection Act, and were recently petitioned for listing as "threatened" under the Endangered Species Act.

While the studies by Fowler (1982, 1987) and others provide the best evidence of wildlife entanglement in plastic debris--especially northern fur seals--and while there is clear evidence that marine debris affects individuals of many species (Manville 1988), Heneman and CEE (1988) felt that evidence of serious population effects on marine wildlife is inconclusive. They cited the fact that few studies had been done on derelict nets or traps, and that while there was clear evidence that entanglement in marine debris kills or injures seabirds, there is no evidence that this is a significant problem for any seabird population. Heneman and CEE's research, however, was not conducted in the North Pacific Ocean.

The Japanese claim that the problem of lost driftnets in the North Pacific is negligible, estimating that only 0.05% of their net sets are lost per operation (the National Marine Fisheries Service estimate is 0.06% (Hinck 1986)). When applied to the setting of more than 32,985 km (20,500 mi) of net per night, plus an additional 16,090-32,180 km (10,000-20,000 mi) of driftnet from Taiwan, South Korea, and others (S. LaBudde, Earthtrust, Honolulu, Hawaii, pers. commun.), a 0.06% loss of net means at least 29-39 km (18-24 mi) of net are lost each night and some 1,542-2,058 km (959-1,279 mi) of net each season. These figures do not account for discarded nets or net fragments.

The northern (Steller's) sea lion, *Eumetopias jubatus*, was reported to have declined by about 50% in the eastern Aleutian Islands between 1957 and 1977 (Braham et al. 1980; King 1983), while western Aleutian populations were reported fairly stable or experiencing only moderate declines during that period (Early et al. 1980; Loughlin et al. 1984). Since 1977, declines continued in the eastern Aleutian Islands (Merrick et al. 1986), but no surveys had been conducted in the western Aleutians from 1979 until 1988. Results from five sites surveyed there in the mid-1970's compared with the 1988 study indicated a 65% reduction in sea lions in the western Aleutians (Byrd and Nysewander 1988). Entanglement was suggested as a possible contributing factor to declines in the eastern Aleutians (Loughlin et al. 1986), but few incidences were reported in the western islands.

Less well known is the status of seabird populations in the Aleutian Islands, and the role plastics may play in affecting these species. Commercial fishing continues to be the largest human activity in the Bering Sea. Factory ships with their fleets of catcher boats stay on location for months processing million of tons of seafood and dumping their wastes in the process (S. LaBudde, Earthtrust, Honolulu, Hawaii, pers. commun.). In Kotzebue Sound north of the Aleutians, data collected in 1977, 1981, and 1987 indicate that the horned puffin, *Fratercula corniculata*, may be experiencing a dramatic 75% decline on Chamisso Island (A. SOWLS, Alaska Maritime National Wildlife Refuge, Homer, Alaska, pers. commun.). The cause of the decline is as yet unknown.

While plastic debris has been reported on the beaches of southern Alaska (Cottingham 1988), on the Pribilof and eastern Aleutian chain (Byrd 1984), and as far out in the Aleutians as Amchitka Island (Merrell 1980, 1984), no plastics beach surveys were reported in the literature from the far western Aleutian Islands prior to July 1988.

METHODS

Twenty-five beach surveys were conducted on seven outer Aleutian Islands from 12 to 20 July 1988 using the U.S. Fish and Wildlife Service's (FWS) research vessel MV *Tiglax* as a base. Surveys were undertaken on beaches in the westernmost U.S. islands located in the Near Islands group (Attu, Agattu, and Shemya Islands), Buldir Island, the Rat Islands (Kiska and Little Kiska Island), and the Andreanof Islands (Adak Island, Fig. 1). Surveys were conducted on an opportunistic basis when the *Tiglax* was either at anchor or was able to stop long enough to deploy us, and when weather and seas were sufficiently favorable to allow beach landings in a motorized, Zodiac inflatable. Beach sites to be surveyed were then randomly selected, and beaches were walked and scanned for all plastic from existing sea level to the storm high tide level/upper wrack line (Wilber 1987). Representative plastic samples were collected and all beaches were photographed. No attempt was made to assess the amounts by weight or volume of plastics present on the beaches, although the numbers of complete trawl nets and relative amounts of driftnets were noted.

Attempts were made to identify the source of plastic items by linking origin of the product, item, or piece by identifiers which were often embossed, stamped, or molded into the plastic.

Five open-water plastic surveys were conducted while the *Tiglax* was steaming between islands (Fig. 1). Surveys were conducted from either the bridge of the vessel or the flying bridge, looking for floating or drifting plastic visible from the bow of the ship while it cruised at speeds of 8-10 kn. Surveys were conducted for approximately 30-min intervals.

Particular attention was paid to wildlife entangled in plastic. Where such animals were spotted, they were photographed. Carcasses were carefully examined for external evidence of plastic or for plastic entanglement scars. Rough necropsies were conducted on dead seabirds whose crops were intact to determine if plastics had been ingested.




Figure 1.--Locations of 25 beach surveys conducted on 7 outer Aleutian Islands from 12 to 20 July 1988 (B), 5 Steller's sea lion surveys conducted from late June to mid-July 1988 (S), and 5 open-water plastics surveys conducted from 12 to 20 July 1988 (W). Map after Byrd and Day (1986).

Beaches were classified as protected, located in coves, bays, or harbors; or as unprotected, located on promontories, points, or similar areas facing the open ocean. In addition to the presence or absence of protective physical barriers, beach classification also was based on the likelihood of prevailing storm tracks, waves, and weather conditions which could augment accumulation of debris.

The randomization test for two independent samples (for large samples) was used to test the statistical difference in the amount of plastic found on protected beaches versus that discovered on open, unprotected beaches (Siegel 1956). This test was also used to examine the difference between fishing-related and nonfishing-related plastics located on the beaches. Fishing-related debris consisted of material specifically used for fishing, material used in the packaging of fish and fish products, or material used by fishermen during the capture and processing of fish.

Northern sea lion counts were conducted either from land or at sea between approximately 1000 and 1800 on five islands during late June and early July (Fig. 1). This enabled peak bull, cow, and pup counts (Loughlin et al. 1986; Byrd and Nysewander 1988). When counts were made on land within the rookeries, numbers of sea lions were assessed "using spook counts" where one or two researchers drove bulls and cows into the water to facilitate counting the pups still on land. All animals were carefully assessed for signs of entanglement using binoculars and a telephoto-equipped 35-mm camera. Where haul sites and rookeries could be seen from headlands above, such as on Kiska Island, counts were made from land by one researcher using binoculars. Where counts were made from the water, three or four observers stationed 30-75 m (33-82 yd) offshore in a Zodiac inflatable counted all pinnipeds. Where counts were made by more than one observer, replicated tallies were averaged to provide the most representative value for each site. Counts were conducted on Attu, Agattu, Buldir, Kiska, and Gramp Rocks. Counts made in 1988 were compared with those made in 1977 (Day et al. 1978) and 1979 (Early et al. 1980).

RESULTS AND DISCUSSION

Beach Surveys

Twenty-five beach surveys were conducted on seven outer Aleutian Islands from 12 to 20 July 1988. Beach surveys averaged 148 m (162 yd) in length. On the 3.7 km (2.3 mi) of beach observed, 3,153 plastic objects were discovered, representing 67 different finished plastic items. No raw polyethylene pellets (nibs), or spheres or spherules of polystyrene were discovered, although due to time limitations attempts were not made to look carefully for them in the high wrack lines. On the average, 126 different plastic items were found per survey. All beaches examined, including the most protected, contained plastic; at least 15 items were deposited on the cleanest (a protected cove on the south side of Shemya Island).

Most prevalent of the plastic items found on the beaches were rope, Styrofoam driftnet buoys, fishing net (mostly trawl nets, but some driftnet segments), and bottles (Table 1). Like the beaches of Bermuda and the

Table 1.--Types and incidence of plastics found on 25 beaches of
7 islands in the outer Aleutian Islands, Alaska, July 1988.

No.	Plastic item	Count	No. of beaches with item	Type ^a
1	Rope (piece, complete coil)	706	24	F
2	Styrofoam buoy	535	15	F
3	Fishing net (mostly trawl)	360	24	F
4	Bottle (other plastic)	331	21	N
5	Hard plastic buoy	215	17	F
6	Plastic piece	157	15	N
7	Piece of Styrofoam	148	23	F/N
8	Cap and lid	111	17	N
9	Strapping band	102	21	F/N
10	Fish-sorting basket	61	12	F
11	Bottle (green plastic)	55	15	N
12	Japanese beer crate	49	8	N
13	Bag	35	15	N
14	Shoe	27	10	N
15	Cup, spoon, fork, plate	23	12	N
16	Sheeting (large plastic)	19	11	F/N
17	Sheeting (small plastic)	18	9	F/N
18	Tub	16	2	N
19	Milk jug	15	9	N
20	Jug	14	5	N
21	Glove	12	8	F/N
22	Bucket	11	9	F/N
23	Polyvinyl chloride pipe	11	7	N
24	Soda bottle	9	8	N
25	Monofilament fishing line	9	4	F
26	Hard hat	7	4	F/N
27	Packaging	7	1	N
28	Styrofoam fast food container	6	4	N
29	Insulation for cable	6	3	N
30	Disposable lighter	5	2	N
31	Styrofoam egg carton	4	3	N
32	Styrofoam cup	4	2	N
33	Cable liner	4	2	N
34	Reflector	4	2	N
35	Boot (with plastic parts)	4	1	F/N
36	Brush	3	3	N
37	Six-pack holder	3	3	N
38	Styrofoam cooler	3	2	F/N
39	Insulation	3	2	N
40	Slipper	3	2	N
41	Toy	3	2	N
42	Drift card ^b	3	1	N
43	Container top	3	1	N
44	Gas can	2	2	F/N
45	Styrofoam life ring	2	2	F/N

Table 1.--Continued.

No.	Plastic item	Count	No. of beaches with item	Type ^a
46	Electrical tape	2	2	N
47	Pen	2	1	N
48	Tooth brush	2	1	N
49	Bowl	1	1	N
50	Indoor-outdoor carpet	1	1	N
51	Caulking tube	1	1	N
52	Counter top	1	1	N
53	Dishwasher sprayer	1	1	N
54	Electrical fixture	1	1	N
55	Filter	1	1	N
56	Garbage can lid	1	1	F/N
57	Ice tray	1	1	N
58	Mylar food pouch	1	1	N
59	U.S. Navy sonabuoy container	1	1	N
60	Plug	1	1	N
61	Pump	1	1	N
62	Ring	1	1	N
63	Shower curtain	1	1	N
64	Soap dish	1	1	N
65	Thermos top	1	1	N
66	Trash can	1	1	F/N
67	Watering jug for plants	1	1	N
	Subtotal	3,153		
1	Crab buoy attached to rope ^c	2	--	F
2	Piece of floating Styrofoam ^c	4	--	F/N
	Subtotal ^c	6		
	Grand total	3,159		F = 7 N = 48 F/N = 13

^aF indicates item is fishing-related; N indicates that it is non-fishing-related; F/N indicates that it is both.

^bNational Marine Fisheries Service drift card.

^cItems discovered during open-ocean survey while departing north end of Kiska Island, 19 July 1988.

Bahamas, which are heavily littered with plastic delivered from a large Atlantic Ocean circulation pattern known as the central gyre (Wilber 1987), the Aleutian Islands act as "sieves" for plastics circulated by waters from the Japanese and Bering Sea currents. Nevertheless, if the amount of plastic located on these Aleutian Island beaches is indicative of that found elsewhere on Alaska's 57,924 km (36,000 mi) of shoreline, there is tremendous opportunity for entanglement or ingestion by wildlife.

Litter was identified from Japan, the U.S.S.R., South Korea, the People's Republic of China, Taiwan, Norway, and the United States, although most of the plastic could not be specifically related to country of origin. Most prevalent were items from Japan; those identifiable were mostly fishing related.

There was a statistically significant difference in the amount of plastic found on protected beaches versus that discovered on unprotected beaches ($P < 0.001$, $df = 23, 22,502$; Table 2). There also was a statistically significant difference in the amount of fishing-related versus non-fishing-related plastics located on beaches examined ($P < 0.001$, $df = 24, 14,083$; Table 1).

Although beaches varied considerably in composition, ranging from sandy to pebbly to rocky to boulder-covered, accumulations of plastic litter were not consistently different among the beaches (Table 2). These findings were consistent with those reported by Merrell (1980, 1984).

When comparing the total amount of plastic ($N = 2,457$ items) versus Styrofoam ($N = 696$ items) found on the 25 beaches, non-Styrofoam plastic made up 78% of the waste stream while Styrofoam consisted of about 22%.

Of particular interest was the discovery of a six-pack beverage yoke on each of three remote beaches (Table 2), since I had been asked to look for and testify about them before a joint congressional hearing held after my return to Washington, D.C., on 26 July (U.S. Government Printing Office (GPO) 1988). One of these yokes was a Hi Cone Eco photodegradable beverage ring (manufactured by Illinois Tool Works), which had not then begun to show any signs of embrittlement.

Open Water Surveys

Five open-water plastic surveys were conducted while the *Tiglax* was underway between islands. One open-water survey on 19 July off the north end of Kiska Island produced two buoys from a brown king crab, *Paralithodes camtschaticus*, pot, one rope from the pot, and four pieces of floating Styrofoam over an 8 km (5 mi) course (Table 1). Even the *Tiglax* was not immune to entanglement plastics. Her hull became ensnared in the rope from an apparently active brown king crab fishing set.

Dead Seabirds

During the 25 beach surveys, some two dozen dead seabirds were located wrapped in, lying next to, or partially entangled in plastic debris,

Table 2.--Amounts of plastic found on 9 protected and 16 unprotected beaches of 7 islands in the outer Aleutian Islands, Alaska, July 1988.

No.	Beach location ^a	No. of plastic items discovered ^b
Protected		
1	South Side Beach, Shemya Island	15 (s)
2	Scotts Cove, southwest side, Shemya Island	34 (s)
3	Casco Bay, Inlet Beach, southwest side, Attu Island	19 (r)
4	Casco Bay, small subbay, southeast side, Attu Island	18 (s)
5	Casco Bay, small subbay, southeast side, Attu Island	22 (p/r)
6	Casco Bay, another subbay, Attu Island	19 (s/r)
7	Alcan Harbor, northwest boat dock, Shemya Island	37 (r)
8	Sweeper Cove, Adak Harbor, Adak Island	22 (b)
9	Sweeper Cove, Adak Harbor, Adak Island	21 (b)
Unprotected		
1	Temnac Beach, south side inlet, Attu Island	37 (s/p)
2	Etienne Cove, southwest side, Attu Island	184 (s)
3	Wrangell Beach, Wrangell Point, Attu Island	511 (p/r)
4	Earle Cove, north side, Attu Island	45 (s/p)
5	Karab Cove, south central, Agattu Island	286 (p)
6	Karab Cove, south central, Agattu Island	48 (s/p)
7	North Bight Beach, near base camp, Buldir Island	329 (r)
8	North Bight Beach, sea lion rookery, Buldir Island	63 (r)
9	North Bight Beach, near base camp, Buldir Island	235 (r)
10	Dark Cove, Kiska Island	284 (s/r)
11	Dark Cove, Kiska Island	379 (s/r)
12	Rock beach, north side, Little Kiska Island	64 (r/b)
13	Three-Mile Beach, Kiska Island	31 (b)
14	Three-Mile Beach, Kiska Island	113 (b/s)
15	Three-Mile Beach, Kiska Island	174 (b/s)
16	North Three-Mile Beach, Kiska Island	170 (b)

^aBeaches designated as protected were located in coves, harbors, or bays, while those designated as unprotected were located on points, promontories, or areas subject to direct wave action from the open ocean, prevailing storm tracks, and weather conditions which likely augmented the accumulation of debris.

^bb = boulder beach, p = pebble beach, r = rock beach, s = sand beach.

^cA six-pack beverage yoke was discovered on each of these three beaches, but at Karab Cove, Agattu Island, the yoke was a Hi Cone Eco photodegradable carrier.

including trawl nets, a piece of driftnet, and plastic rope. With the exception of one dead sooty shearwater, *Puffinus griseus*, wrapped in a piece of trawl net that appeared to strangle it, it usually was impossible to determine the cause of death, given the decomposition of the majority of the carcasses. A Leach's storm petrel, *Oceanodroma leucorhoa*, however, was discovered in August 1988 on Buldir Island entangled in monofilament fishing line which apparently killed the bird (G. V. Byrd, Alaska Maritime National Wildlife Refuge, U.S. Fish and Wildlife Service, Nome, pers. commun.).

Field necropsies revealed no ingested plastics in the few birds (a tufted puffin, *Lunda cirrhata*, two glaucous-winged gulls, *Larus glaucescens*, a sooty shearwater, two crested auklets, *Aethia cristatella*, a least auklet, *A. pusilla*, and a common murre, *Uria aalge*, whose crops were intact. More research on seabird mortality needs to be conducted in the outer Aleutian Islands. Plastics are of special concern since seabirds tend to concentrate in areas where current upwellings reach the surface or where tidal rips occur (J. F. Piatt, Alaska Fish and Wildlife Research Center, FWS, Anchorage, Alaska, pers. commun.)--the same areas where ghost nets, drifting plastic debris, and other flotsam may also occur. Although the impacts of lost or discarded fishing gear and other plastic debris have been difficult to quantify, the few data available suggest that lost gear may be as efficient at killing birds and mammals as is active gear (DeGange and Newby 1980; Jones and Ferrero 1985; Piatt and Nettleship 1987).

Northern Sea Lion Counts

Attu Island

Although counts were made for northern (Steller's) sea lions on 8 and 14 July, the second count was made at a more appropriate hour and therefore was considered more representative. A comparison of this 1988 count with the one made in 1979 (Early et al. 1980), shows a precipitous 77% decline from about 5,700 animals to approximately 1,300 (Byrd and Nysewander 1988). Cause of the decline is unknown.

Agattu Island

Counts were made in mid-June when harem bulls were at their peak and on 9-11 July after most pups were born. The estimated 1988 count of 3,000 sea lions was less than half the number counted in 1979 (Byrd and Nysewander 1988). One bull was seen with a piece of trawl net fragment wrapped around its neck.

Buldir Island

Twenty-two areas were identified for sea lion surveys at Buldir Island, and June and July counts were made for most of these sites. Less than 1,900 sea lions were counted in 1988, 70% fewer than the 1979 survey (Byrd and Nysewander 1988). I photographed a harem bull with a massive entanglement scar around its neck and the strapping band apparently still present. The animal appeared robust and generally healthy, and maintained

a territory with one cow (but no pups) several hundred meters west of the Bull Point Beach sea lion rookery.

Kiska Island

Earlier single counts on Kiska and Tanadak Islands were followed by a mid-July count from land. The overall total for Kiska and Tanadak Islands in 1988 was 2,414 sea lions, a 64% decline from the total seen in 1979 (Byrd and Nysewander 1988). A bull and a cow were seen with deep scars around their necks from previous apparent plastic entanglement.

Gramp Rocks

In 1977, Day et al. (1978) reported sighting over 2,200 sea lions on Gramp Rocks. In late June 1988, over 900 pinnipeds were observed from land, representing a 59% decrease in the population.

Although it was certainly possible that some entangled sea lions were overlooked, those observed represented only a tiny fraction of total population examined. Sea lion populations have declined, probably drastically, in the western Aleutian Islands in the past decade--an overall 65% reduction for the five sites examined--but the reasons for this decline remain unclear. Entanglement has been suggested as a possible contributing factor, especially in the eastern Aleutian Islands (Loughlin et al. 1986; Byrd and Nysewander 1988), but it needs much closer examination in the western Aleutians.

Since pups and juvenile sea lions, like their northern fur seal counterparts, are curious, inquisitive, and playful (King 1983), they may suffer much higher mortality due to entanglement in plastic fishing debris than observed. Since so little research has been done on the sea lions in the western Aleutians, mortality due to plastic entanglement--although suspected by this author to be a contributing factor to their decline--needs more detailed study and analysis.

Presentation of Survey Data at Congressional Hearing

Using data from this study, information was presented at a joint congressional hearing on six-pack yoke legislation on 26 July 1988 (U.S. GPO 1988). Those bills, H.R. 5117 and S. 1986, requiring that six-pack beverage yokes be made degradable within 24 months, were passed by Congress and signed into law late in 1988 by President Reagan.

REFERENCES

- Braham, H. W., R. D. Everitt, and D. J. Rugh.
1980. Northern sea lion population decline in the eastern Aleutian Islands. *J. Wildl. Manage.* 44:25-33.
- Byrd, G. V.
1984. Observations of flora and fauna in the Bering Sea Unit, Alaska Maritime National Wildlife Refuge, in July 1984. Alaska Maritime National Wildlife Refuge, U.S. Fish and Wildlife Service, Homer, AK, 4 p. [Unpubl. rep.]

- Byrd, G. V., and R. H. Day.
1986. The avifauna of Buldir Island, Aleutian Islands, Alaska. *Arctic* 39(2):109-118.
- Byrd, G. V., and D. I. Nysewander.
1988. Observations of northern sea lions in the western Aleutian Islands, Alaska in 1988: Evidence of a decline. Alaska Maritime National Wildlife Refuge, U.S. Fish and Wildlife Service, Homer, AK, 9 p. [Unpubl. rep.]
- Centaur Associates and the Center for Environmental Education.
1986. Report on Contract No. 50 ABNF-6-00192 to National Marine Fisheries Service. Centaur Associates and the Center for Environmental Education.
- Cottingham, D.
1988. Persistent marine debris: Challenge and response: The Federal perspective. Sea Grant Educ. Publ. No. 1, Grant No. NA86AA-D-SG041, Proj. No. A/75-01, 41 p.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Day, R. H., T. J. Early, and E. P. Knudtson.
1978. A bird and mammal survey of the west-central Aleutians. U.S. Fish and Wildlife Service, Adak, AK. [Unpubl. rep.]
- DeGange, A. R., and T. C. Newby.
1980. Mortality of seabirds and fish in a lost salmon driftnet. *Mar. Pollut. Bull.* 11:322-323.
- Early, T. J., A. B. Taber, J. Beall, and W. Henry.
1980. Results of bird and mammal surveys of the western Aleutians. U.S. Fish and Wildlife Service, Adak, AK. [Unpubl. rep.]
- Fowler, C. W.
1982. Interactions of northern fur seals and commercial fisheries. *Trans. N. Am. Wildl. Nat. Resour. Conf.* 47:278-292.
1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.
- Fowler, C. W., and T. R. Merrell.
1986. Victims of plastic technology. *Alaska Fish Game* 18(2):34-37.
- Heneman, B., and Center for Environmental Education.
1988. Persistent marine debris in the North Sea, northwest Atlantic Ocean, wider Caribbean area, and the west coast of Baja California. Report to Marine Mammal Commission and National Ocean Pollution Program Office, NOAA, U.S. Dep. Commer. Contract MM 3309598-5.

Hinck, J.

1986. Dirge of the driftnets. *Defenders* 61(6):16-25.

Ignell, S. E., and M. L. Dahlberg.

1986. Results of 1986 cooperative research on the distribution of marine debris in the North Pacific Ocean. Document submitted to the Int. North Pac. Fish. Comm., Anchorage, AK, November 1986, NWFAC, NMFS, NOAA, Auke Bay Lab., P.O. Box 21055, Auke Bay, AK 99821, 15 p.

Jones, L. L., and R. C. Ferrero.

1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

King, J. E.

1983. *Seals of the world*. Comstock Publ. Assoc., Cornell Univ. Press, Ithaca, NY, 240 p.

Loughlin, T. R., P. J. Gearin, R. L. DeLong, and R. L. Merrick.

1986. Assessment of net entanglement on northern sea lions in the Aleutian Islands, 25 June-15 July 1985. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 86-D2, 50 p.

Loughlin, T. R., D. J. Rugh, and C. H. Fiscus.

1984. Northern sea lion distribution and abundance: 1956-1980. *J. Wildl. Manage.* 48:729-740.

Manville, A. M. II.

1988. New technologies dealing with marine plastic pollution and efforts at mitigation. *Trans. N. Am. Wildl. Nat. Resour. Conf.* 53:191-203.

Merrell, T. R., Jr.

1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.

Merrick, R. L., T. R. Loughlin, and D. G. Calkins.

1986. Decline in the abundance of the northern sea lion (*Eumetopias jubatus*) in Alaska, 1956-86. U.S. Natl. Mar. Fish. Serv., Seattle, WA.

Piatt, J. F., and D. N. Nettleship.

1987. Incidental catch of marine birds and mammals in fishing nets off Newfoundland, Canada. *Mar. Pollut. Bull.* 18:344-349.

- Schneidman, D.
1987. Plastic: Progress and peril. *Marketing News*, Dec. 18, 2 p.
- Shaw, D. G., and G. A. Mapes.
1979. Surface circulation and the distribution of pelagic tar and plastic. *Mar. Pollut. Bull.* 10:160-162.
- Siegel, S.
1956. *Nonparametric statistics for the behavioral sciences*. McGraw-Hill, N.Y., 312 p.
- U.S. Government Printing Office.
1988. Statement of Al Manville, senior staff wildlife biologist, Defenders of Wildlife, on degradable six-pack rings. Joint Hearing before the Subcommittee on Fisheries and Wildlife Conservation and the Environment of the Committee on Merchant Marine and Fisheries, and the Subcommittee on Transportation, Tourism, and Hazardous Materials of the Committee on Energy and Commerce, House of Representatives, 100th Congress, Second Session on S. 1986. A Bill to Study, Control, and Reduce the Pollution of Aquatic Environments from Plastic Materials, and for Other Purposes, July 26, 1988. Ser. No. 100-81: 9-11, 14-15, 18, 20, 66-77.
- Venrick, E. L., T. W. Backman, W. C. Bartman, C. J. Platt, M. S. Thornhill, and R. E. Yates.
1973. Man-made objects on the surface of the North Pacific Ocean. *Nature* 241:271.
- Wilber, R. J.
1987. Plastic in the North Atlantic. *Oceanus* 30(3):61-68.

ENTRAPMENT OF SEA-DEPOSITED PLASTIC ON THE
SHORE OF A GULF OF MAINE ISLAND

Richard H. Podolsky
Research and Academics
Island Institute
60 Ocean Street
Rockland, Maine 04841, U.S.A.

ABSTRACT

During 1987, 300 kg of sea-deposited plastic debris were collected and removed from the 14.45-km shoreline along a wilderness island in eastern Maine. Exactly 1 year later, 124 kg of plastic were collected along the same shoreline. The plastic debris was not uniformly distributed among shoreline habitats. Beach, boulder, marsh, and meadow shorelines were found to catch plastic debris, and ledge shores were found to repel plastic. The western half of the island, facing the prevailing wind, had twice the plastic accumulation of the eastern half.

WASHUPS OF FLOATABLE WASTE MATERIALS AND THEIR
IMPACT ON NEW YORK BIGHT BEACHES

R. Lawrence Swanson and Robert Zimmer
Waste Management Institute
Marine Sciences Research Center
State University of New York
Stony Brook, New York 11794, U.S.A.

ABSTRACT*

During the summers of 1987 and 1988, the New York Bight once again experienced a series of incidents in which waterborne, floatable, waste materials and debris were stranded on area beaches. Medically-related wastes were of particular concern. The sources of floatable wastes are identified and local climatological data are used to explain the process by which floatable material was transported.

The climatology of the summers of 1987 and 1988 are compared with that of 1976, when similar strandings of floatable wastes occurred on the south shore of Long Island. The summer wind records of these years are also compared with the historical wind record, 1959-1988. The basis of these comparisons are measures of wind persistence and relative energy. These analyses indicate the unusual nature of the conditions that prevailed in 1976, 1987 and 1988 and how they differed from each other. During unusually persistent winds, floatable debris in near surface waters can be transported in excess of 100 km in a direction opposed to the general flow over the continental shelf. While major washups of floatable wastes are unusual, we now know under what conditions they are likely to occur. Emphasis must be placed on alleviating the problem at the sources.

*Abstract from "Meteorological conditions leading to the 1987 and 1988 washups of floatable wastes on New York and New Jersey beaches and comparison of these conditions with the historical record." *Estuarine, Coastal and Shelf Science* (1990) 30:59-78. Academic Press Limited. London. By permission.

**PLASTIC DEBRIS AND DERELICT FISHING GEAR ON
SHACKLEFORD BANKS, NORTH CAROLINA**

K. V. Koski

Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

ABSTRACT

Surveys of the quantity, type, and source of plastic debris on beaches on Shackleford Banks, Cape Lookout National Seashore, North Carolina, were conducted in December 1986 and September 1987 by the National Marine Fisheries Service Beaufort Laboratory and the National Park Service. Eight 1-km beaches established as benchmarks were surveyed and measured using standard beach survey methods developed in Alaska. In 1986 an average 863 items/km of beach were found. Packaging industry items comprised 51%; fishing gear (commercial and sport), 15%; fragments, 14%; maritime industry, 8%; miscellaneous, 7%; and personal effects, 5% of the total debris. Fourteen percent of the debris items were categorized as entanglement dangers, and 55% were categorized as ingestible dangers to marine animals such as the endangered marine turtles. Fishing gear contributed significantly to the entanglement items and packaging contributed significantly to the ingestible items.

In 1987, an average of 1,073 items/km of beach were found, an increase of 25% from 1986. Four transects of beach cleared of all debris in 1986 and resurveyed in 1987 increased 117% from 526 items/km to 1,141 items/km. Composition was similar to that found in 1986. Shackleford Banks ranks high in the amount of plastic debris on its beaches when compared to beaches in Texas, Oregon, and Alaska. Because several species of endangered marine turtles utilize the barrier islands of the southeast United States for nesting, plastic debris may pose a serious threat to their well-being.

ANTHROPOGENIC AND NATURAL DEBRIS ON A
SOUTH TEXAS BARRIER ISLAND BEACH

Anthony F. Amos and Pamela T. Plotkin
Marine Science Institute
University of Texas at Austin
Port Aransas, Texas 78373, U.S.A.

ABSTRACT

The results of a long-term study to estimate the quantity of both natural and man-made debris that washed up on the gulf beach of Mustang Island, Texas, are presented. The study beach is 12 km long and has been monitored since 1978 with over 1,800 observations and 4,000 man-hours of observational effort. Four types of debris measurements are made: (1) estimates of 40 different categories of debris and litter using a ranking system (done bi-daily since 1983); (2) counts of some 70 categories of debris items (done weekly since early 1987); (3) quantity (weight) and quality of all debris at three 10-m-wide beach transects (done weekly for 1 year in 1987); (4) counts of four key anthropogenic litter items thought to be typical of four separate sources (done bi-daily for the past year). Method 1 shows seasonal tendencies but is not precise enough to indicate longer term trends. Method 2 gives much more accurate quantitative data but still reveals no trends over the 2-year period. Method 3 reveals associations of anthropogenic and natural material, quantifies "uncountable" items like tar balls, and allows examination of the world of "microtrash." Method 4 shows the short- and long-term variability of litter associated with commercial fishing and offshore oil activities, and material from south of the United States-Mexico border.

Some results of the study to date:

- There is a seasonal variation in quantity of most beach debris--highs in spring and autumn, lows in summer and especially winter.
- Most anthropogenic litter comes from offshore and is identifiable with commercial fishing, recreational boating, offshore drilling and production, and the international merchant marine (items from 60 countries have been found here).

- Short-term variability is large and is attributable to winds, storms, tides, currents, beachgoing, and beach cleaning activities.
- Despite this variability, anthropogenic debris is a permanent feature of this beach's flotsam and jetsam. Some high counts: Styrofoam 1,100/km, plastic bags 1,000/km, plastics of all kinds 1,600/km, 1-gal (3.785 L) milk jugs 50/km, laughing gull 500/km and sanderling 150/km (the two dominant bird species), Portuguese man-of-war 1,200/km, people 25/km. Some high numbers by weight: Sargassum 2,600, tar balls 2,100, driftwood 1,000, plastic debris 140 kg/km.

**SOLID WASTE ON THE ISRAELI COAST--COMPOSITION,
SOURCES, AND MANAGEMENT**

Abraham Golik and Yaron Gertner
National Institute of Oceanography
Israel Oceanographic and Limnological Research
Haifa 31080, Israel

ABSTRACT

Counts of litter pieces of six Mediterranean Sea beaches in Israel were conducted at monthly intervals between May 1988 and January 1989. Litter consisted of 71.6% plastic items, 7.9% wood, 5.7% metal pieces, 3.1% glass, and 11.7% other. Most of the litter, such as beverage bottles, food containers, cosmetics remnants, plastic bags, pieces of garments, and foam rubber mattresses, is related to recreation activity and is therefore land-based garbage. The absence of fishing gear remnants and large food packaging material indicative of ships' garbage, and the sparseness of litter with inscriptions and imprints showing foreign origin, further support the conclusion that most of this garbage is land-based.

This finding contrasts those of similar studies which were carried out on other coasts around the world such as Amchitka Island, Alaska; Helgoland Island, Germany; and the west European shores. There, most rubbish consisted of fishing gear, lavatory cleansers, household cleaners, and containers bearing inscriptions indicative of foreign origin, and is marine-based. The difference in the litter reflects the differences between Israel and the mentioned locations in coastal use, ship traffic, and winds, waves, currents, and tide conditions.

The significance of identifying the litter source is that mitigation of this problem in Israel is rather simple because it is easier to control land-based litter than sea-based. There are indications that proper publicity, education, and control will reduce the problem.

INTRODUCTION

During the last two or three decades, there has been growing concern about marine pollution by persistent litter. Most of the reports on this subject deal with litter in the oceans and only a few with coastal litter. Some reports, such as those of Carpenter et al. (1972), Gregory (1977,

1983), and Shiber (1979, 1987), provided qualitative and quantitative information on coastal garbage; others showed the deleterious effect of the coastal rubbish on seals (Merrell 1980) and seabirds (Schrey and Vauk (1987); and others investigated the sources and fate of the coastal litter (Dixon and Cooke 1977; Merrell 1980; Dixon and Dixon 1981; Vauk and Schrey 1987).

Due to the temperate climate of the Mediterranean Sea and the large number of sunshine hours, its surrounding countries are presently undergoing an intensive development of coastal-oriented tourism. Although coastal pollution is one of the deterrents to tourism, tourism may be important as a contributor of waste to the beaches. Very little information is available on litter pollution of the Mediterranean Sea and its coasts. Shiber (1979, 1982, 1987) reported on the occurrence of plastic beads on the beaches of Lebanon and Spain, Saydam et al. (1985) on floating garbage off Turkey, and Morris (1980) and McCoy (1988) on litter floating in the east Mediterranean.

The gravity of solid waste as a marine and coastal pollutant was recognized by the Convention for the Protection of the Mediterranean Sea Against Pollution (Barcelona Convention) which, in one of its protocols, prohibits the dumping into the Mediterranean Sea Area of (among others) ". . . persistent plastic and other persistent materials. . . ." In order to evaluate the magnitude of this pollutant in the Mediterranean Sea, the United Nations Environmental Programme has set up a program to monitor solid and persistent litter in the Mediterranean Sea and on some of its coasts. This report is a partial result of that effort.

THE ISRAELI COASTLINE

The Mediterranean coastline of Israel extends for 200 km from the Egyptian border in the south to Akhziv on the Lebanese border in the north (Fig. 1). It is a smooth, slightly curving coastline, with Haifa Bay being the only large indentation in it, and may be divided into two sections. The southern section, from the Egyptian border to Akko, consists of long beaches, 30-50 m wide, covered by fine to medium quartz sand. The northern part, from Akko to the Lebanese border, is mostly a rocky coastline with pocket beaches which are covered by coarse, biogenic sand.

During the winter, alternating high and low barometric pressures cross the east Mediterranean Sea from west to east, subjecting the Israeli shoreline to storms at about a 10-day frequency. During the storms, winds blow from the west and southwest. Before and after the storms, wind direction is generally from the east. During spring and autumn, winds are commonly from the east (land), and during summer, the sea breeze changes from a maximum of 18-20 km/h shoreward at noon to no wind at night and about 6 km/h seaward early in the morning. The mean significant wave height during the winter is 1.1 m, during spring and autumn 0.5 m, and in the summer 0.7 m. The alongshore current during the winter storms is from south to north; whereas during the summer it is mostly from north to south in the study area. Tidal range is approximately 30 cm, and tidal currents are insignificant.




Figure 1.--Station location map.

Some 70 beaches, declared as public swimming beaches, are present on the Israeli coastline. "Declared" beaches are operated by the various municipalities, which are responsible, among other things, for keeping these beaches clean during the swimming season. Though most of the bathers go to declared beaches, undeclared beaches are popular too and are visited by many during the summer. Once or twice a year, the Office of Environmental Quality, together with local councils, conducts a cleanup of almost the entire shoreline of Israel.

The purposes of this study were to evaluate the quantity of the coastal litter on the Israeli coastline, to find out if there is any relationship between beach morphology or beach use and litter, to determine whether the rubbish is land-based or sea-based, and to recommend means and ways to treat this problem.

METHODS

Six beaches were selected for this study. They differ in their morphology, sedimentology, and type of use; their locations are given in Figure 1.

On each beach five to eight transects were established. The locations of transects were randomly determined at each sampling date. The transect was 5 m wide, oriented normal to the beach, from the water line to the back of the beach. The back of the beach was determined as the foot of the coastal cliff, the dunes, or the vegetated area. All litter pieces larger than 2 cm found on a transect constituted a sample. Sampling started in May 1988 and continued until January 1989 at roughly monthly intervals, yielding 330 samples.

RESULTS

Because the sampling program still continues as these lines are written, the distribution of litter in space and time will be discussed in the future. Only litter composition is treated here.

The relative abundance of the various litter constituents and the types of materials identified are summarized in Figure 2 and Table 1. The most abundant components are plastic fragments. Most of these are hard plastic ranging in size from 2 cm (the smallest size counted) to 30 cm, and in most cases they could be identified as fragments of plastic containers or bottles. There were also a few straps from large packing crates. Although no special count was made, most of the plastic containers and bottles originally contained beverages, food, and cosmetics (mostly suntan lotion). Only a small fraction of the plastic containers were cleansers or various types of oil related to household or industrial activities. Plastic and metal caps were counted separately, but they came from the plastic containers and bottles and were included under the plastic category. Most of the metal components were tins used for beverages; the rest were either food cans or aerosols. In a similar way the glass fraction was dominated by soft drink bottles, with low numbers of other items such as light bulbs. The wood category included driftwood as well as crate fragments.




Figure 2.--Relative abundance of litter components.

Items of foreign origin (as indicated by inscription or imprinting) were found in low numbers. Most of these were from Lebanon but a few items with Turkish, Greek, and Spanish inscriptions were found as well. Although items were found at most of our sampling stations, the one in Akhziv, close to the border of Lebanon, contained the most.

DISCUSSION

Rubbish Composition and Sources

Examination of the litter components shows that most of the garbage on the Israeli coastline results from beach recreation activity. This is indicated by the original contents of most of the containers, bottles, and cans (beverages, food, and cosmetics), by the plastic bags used by beachgoers to carry their food and other belongings, by fragments of Styrofoam floats for children who go into the water, by remnants of rubber foam mattresses for lying on the beach, and by the various garment pieces, such as rubber sandals, which are good indicators of beachgoers. Mixed with this, and represented in smaller quantities, is garbage that originated from other sources. This fraction is represented by wood, rope, miscellaneous bottles and containers, and plastic and metal straps.

The absence of fishing gear remnants and packaging material of food in bulk, characteristic of ship litter, is considered to be evidence that most of the coastal litter is land-based. The paucity of foreign garbage pieces

Table 1.--Total litter in sampling stations during study period.

Material	Number of items	Percentage
Plastic		
Fragments	3,634	31.9
Bags	2,322	20.4
Containers and bottles	1,076	9.5
Caps and covers	1,069	9.4
Other	40	0.4
Subtotal	8,141	71.6
Wood	904	7.9
Metal		
Cans	498	4.4
Containers	90	0.8
Aerosols	66	0.6
Subtotal	654	5.8
Glass		
Bottles	308	2.7
Other	51	0.4
Subtotal	359	3.1
Other		
Cartons	358	3.1
Ropes	319	2.8
Styrofoam	292	2.6
Garments	234	2.0
Foam rubber	105	0.9
Subtotal	1,311	11.4
Total	11,366	99.8

is further evidence of this. Foreign litter on Akhziv beach is a result of the alongshore current which flows during summer mostly from the north and carries the garbage from Lebanon to Israel, thus impacting the northern beaches of Israel. Had it been ships' litter, the whole coastline would have been affected.

Comparison With Litter From Other Coastlines

Merrell (1980, 1984, 1985), who studied the coastal litter in Amchitka Island, Alaska, between 1972 and 1982, found that out of the 24 most common litter items, 12 were used in commercial fishing. In 1974, these constituted 65% by counts, and in 1982, 34%. Merrell attributes the rest of the litter to garbage which was discarded from the fishing fleets in the Pacific Ocean. Vauk and Schrey (1987), who investigated litter on

the beach of Helgoland Island in the German Bight, reported that 99.2% of the 8,539 waste items found on the beach were identifiable as ships' waste.

Dixon and Dixon (1981, 1983) report on a series of coastal litter surveys which were conducted in the United Kingdom; the Western Isles of Scotland; Cherbourg Peninsula, France; West Jutland, Denmark; and Portugal. In all instances they noted that the most abundant plastic containers found were for lavatory and household cleaners. Plastic and carton containers for milk were also abundant, but their relative abundance was not the same on all of these coasts. Bottles of mineral water, wine, and soft drinks were found in small percentages. The geographical origins of the containers indicated that many were from countries foreign to the beach on which they were found. In short, most of the litter stranded on the beaches of western Europe is seaborne and has not been brought by people coming for recreational purposes.

The difference between the coastal litter found on the Israeli shore and litter items described above is clear--most of the waste in Israel is of local origin and is related to beach recreation activity, whereas a major proportion of the discussed examples is foreign and related to commercial fisheries and ship traffic. The reasons for this difference are clear. For the islands of Amchitka and Helgoland, the litter production by local inhabitants is negligible in comparison to that which lands from the sea. On western European shores, intensive ship traffic and commercial fisheries produce much marine-based litter which is spread widely by the strong winds, currents, and tide. The input of land-based litter by beachgoers, on the other hand, is rather limited due to the short summer recreation period in these countries. The situation in Israel is the opposite. The summer beach recreation period is long (April-November), ship traffic in the east Mediterranean is much lighter than in the North Sea, the English Channel, and the eastern Atlantic, and winds and currents which may bring garbage from the sea are moderate (see above).

Mitigation of Coastal Litter in Israel

The significance of the findings of this study is that the control of solid waste on the Israeli shore is less of a problem than it is on many other coastlines. Most students of coastal and marine litter problems admit that there is little hope of controlling disposal of garbage from ships in the near future (Dixon and Dixon 1981; Bean 1987). As this source of garbage is limited in the case of the Israeli coastline, the attention there should be focused on those who pollute the beach, namely the beachgoers. This is a matter of culture, and has to be treated with the classical tools of education, legislation, and law enforcement. Indeed various programs to educate bathers to keep the beach clean are under way. Plastic refuse bags are distributed to beachgoers by youngsters, and classes of young students are called for voluntary beach cleaning operations. The idea is that these activities will not only help to clean beaches but will also educate children to keep them so in the future. There are signs that this approach will be successful. Figure 3 shows a plastic bag which was filled with garbage by beachgoers who were conscientious not to leave their refuse spread on the beach. However, due



Figure 3.--Plastic bag filled by beachgoers with their garbage to prevent it from spreading on the beach.

to the lack of a nearby trash bin, the bag was left on the beach. A program of placing trash bins on some nondeclared beaches is now under way in Israel.

ACKNOWLEDGMENTS

Y. Mart critically read the manuscript. This study was financially supported by the Intergovernmental Oceanographic Commission, United Nations Educational, Scientific and Cultural Organization under Contract No. SC/UNEP/291.006.8 relevant to the Mediterranean Action Plan - MEDPOL II.

REFERENCES

- Bean, M. J.
1987. Legal strategies for reducing persistent plastic in the marine environment. *Mar. Pollut. Bull.* 18:357-360.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Miklas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. *Science* 178:749-750.
- Dixon, T. R., and A. J. Cooke.
1977. Discarded containers on a Kent beach. *Mar. Pollut. Bull.* 8:105-109.

Dixon, T. R., and T. J. Dixon.

1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.

1983. Marine litter surveillance on the North Atlantic Ocean shores of Portugal and the Western Isles of Scotland. Marine litter research program: Stage 5, Keep Britain Tidy Group:44.

Gregory, M. R.

1977. Plastic pellets on New Zealand beaches. *Mar. Pollut. Bull.* 8:82-84.

1983. Virgin plastic granules on some beaches on eastern Canada and Bermuda. *Mar. Environ. Res.* 10:73-92.

McCoy, F. W.

1988. Floating megalitter in the eastern Mediterranean. *Mar. Pollut. Bull.* 19:25-28.

Merrell, T. R., Jr.

1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.

1985. Fish nets and other plastic litter on Alaska beaches. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 27-29 November 1984, Honolulu, Hawaii*, p. 160-182. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Morris, R. J.

1980. Floating plastic debris in the Mediterranean. *Mar. Pollut. Bull.* 11:125.

Saydam, C., I. Salihoglu, M. Sakarya, and A. Yilmaz.

1985. Dissolved/dispersed petroleum hydrocarbons, suspended sediment, plastic, pelagic tar and other litter in the north eastern Mediterranean. *The Seventh Workshop on Pollution of the Mediterranean, Lucerne, Switzerland*, p. 509-518. International Committee of the Scientific Exploration of the Mediterranean Sea, Monaco.

Schrey, E., and G. J. M. Vauk.

1987. Records of entangled gannets (*Sula bassana*) at Helgoland, German Bight. *Mar. Pollut. Bull.* 18:350-352.

Shiber, J. G.

1979. Plastic pellets on the coast of Lebanon. *Mar. Pollut. Bull.* 10:28-30.

1982. Plastic pellets on Spain's Costa del Sol beaches. Mar. Pollut. Bull. 13:409-412.

1987. Plastic pellets and tar on Spain's Mediterranean Sea. Mar. Pollut. Bull. 18:84-86.

Vauk, G. J. M., and E. Schrey.

1987. Litter pollution from ships in the German Bight. Mar. Pollut. Bull. 18:316-319.

NATIONAL MARINE DEBRIS DATA BASE: FINDINGS
ON BEACH DEBRIS REPORTED BY CITIZENS

Kathryn J. O'Hara
Center for Marine Conservation
1725 DeSales Street, NW
Washington, D.C. 20036, U.S.A.

ABSTRACT

The Center for Marine Conservation (CMC) has established a National Marine Debris Data Base to involve citizens in the collection of standardized information on marine debris. This information collected over time will serve as a means to monitor legislative and other efforts to reduce marine debris.

During the first year of this program, more than 47,500 volunteers in 25 U.S. states and territories recorded detailed information on types and quantities of debris collected during one 3-h period in the fall of 1988. All completed data cards were returned to CMC for analysis.

The data showed that approximately 62% of the 1,973,995 debris items reported were plastic. The most common debris items were fragmented pieces of plastic and foamed plastic (Styrofoam-like). More than 56% of all debris was packaging and disposable plastic products that can be generated by a diversity of ocean- and land-based sources. Using indicator items it was found that approximately 8% of all debris reported was indicative of dumping of galley wastes by vessels, 2% was operational wastes generated during activities conducted by cargo vessels and offshore petroleum operations, 6% was fishing and boating gear, and 0.4% was sewage-associated wastes indicative of inadequate sewage treatment practices. The presence of these indicator items suggests that some of the untraceable debris items may also be generated by these sources. Only 0.09% of the debris was categorized as medical wastes suspected to be from illegal dumping, storm water runoff, or inadequate sewer systems. More than 1,000 debris items from 45 countries were reported, in addition to items traceable to 10 cruise line companies. Volunteers also reported finding more than 45 cases of wildlife entanglement or ingestion of debris, most of which were birds entangled in plastic fishing line.

INTRODUCTION

More than 47,500 U.S. citizens participated in the first national volunteer effort to categorize the types and quantities of marine debris found in U.S. coastal areas. Information from this citizen monitoring effort was compiled by the Center for Marine Conservation (CMC)--formerly the Center for Environmental Education--in the National Marine Debris Data Base. The data base was established to gather and analyze information collected by citizens at beach cleanups conducted as part of the annual Coastweeks celebration each fall. Sponsored by the U.S. Environmental Protection Agency (EPA), National Oceanic and Atmospheric Administration, and the U.S. Coast Guard, the data base was established to utilize the efforts of citizen volunteers to identify specific debris problems in different parts of the country and to monitor the effectiveness of Annex V and other measures implemented to reduce debris. This paper presents information on how the data base was organized and national findings on the types of debris reported the first year. Detailed information on the types of debris reported and analyzed on the national, state, and local level is available from the CMC in a report entitled "Cleaning America's Beaches: 1988 National Beach Cleanup Results."

METHODS

Since 1986, the CMC has compiled extensive information on the types and quantities of marine debris found on the Texas coastline using data collected by citizens during volunteer beach cleanups. Based on these data findings, the CMC has published two reports on the debris problem in Texas which include documentation on the sources of debris and recommendations for Federal, state, and local governments, industry, and other groups to reduce the marine debris problem (Center for Environmental Education 1987, 1988).

In 1988, using the Texas data collection system as a model, the CMC initiated the first national data collection effort. After contacting all coordinators that planned to conduct beach cleanups during Coastweeks '88 (17 September-10 October), 25 states agreed to participate in a national data collection effort. For many of these states, 1988 would be their first cleanup effort and coordinators were eager to obtain information on the types and quantities of debris found on their coastlines. The timing of this national event was also important since the data collected would establish a baseline of information on beach debris prior to the enactment of MARPOL Annex V on 31 December 1988.

In order to produce a data card that would be representative of the types of beach debris found nationwide, the CMC requested comments from beach cleanup organizers as to what types of debris were prevalent on their coastline and what information was needed to evaluate the debris problem on the state and local levels. The CMC had previously developed a data card for use in Texas that reflected the great diversity of debris known to occur on the Texas coastline. (Due to circulation patterns in the Gulf of Mexico, Texas beaches receive the brunt of debris dumped into the Gulf.) Because of this diversity of debris, the Texas data card served as

BEACH CLEANUP DATA CARD

Thank you for completing this data card. Answer the questions and return to your area coordinator or to the address at the bottom of this card. This information will be used in the Center for Environmental Education's National Marine Debris Data Base and Report to help develop solutions to stopping marine debris.

Name _____ Affiliation _____
 Address _____ Occupation _____ Phone (_____) _____
 City _____ State _____ Zip _____ M _____ F _____ Age _____
 Today's Date Month _____ Day _____ Year _____ Name of Coordinator _____
 Location of beach cleaned _____ Nearest city _____
 How did you hear about the cleanup? _____

SAFETY TIPS

1. Do not go near any large drums.
2. Be careful with sharp objects.
3. Wear gloves.
4. Stay out of the dune areas.
5. Watch out for snakes.
6. Don't lift anything too heavy.

WE WANT YOU TO BE SAFE

Number of people working together on this data card _____ Estimated distance of beach cleaned _____ Number of bags filled _____

SOURCES OF FOREIGN DEBRIS. Please list all items that have foreign labels

Country	Item Found
Example: <i>Mexico</i>	<i>plastic bottle - "Clarisol"</i>

STRANDED AND/OR ENTANGLED ANIMALS (Please describe type of animal and type of entangling debris. Be as specific as you can.)

What was the most peculiar item you collected? _____

Comments _____

Thank you!

**PLEASE RETURN THIS CARD TO
 YOUR AREA COORDINATOR
 OR MAIL IT TO:
 Center for Environmental Education
 1725 DeSales Street, NW
 Washington, DC 20036
 A Membership Organization**



*Copyright 1988, Center for Environmental Education, Inc.

Figure 1.--Beach cleanup data card, sides 1 and 2.

ITEMS COLLECTED

You may find it helpful to work with a buddy as you clean the beach, one of you picking up trash and the other taking notes. An easy way to keep track of the items you find is by making tick marks. The box is for total items; see sample below.

egg cartons HTI HTI HTI I Total cups HTI HTI HTI HTI HTI HTI Total

PLASTIC Total number of items

bags:

trash

salt

other

bottles:

beverage, soda

bleach, cleaner

oil, lube

other

buckets

caps, lids

cups, spoons, forks, straws

diapers

disposable lighters

fishing line

fishing net:

longer than 2 feet

2 feet or shorter

floats & lures

hardhats

light sticks

milk, water gallon jugs

pieces

pipe thread protector

rope:

longer than 2 feet

2 feet or shorter

sheeting:

longer than 2 feet

2 feet or shorter

6-pack holders

strapping bands

syringes

tampon applicators

toys

vegetable sacks

"write protection" rings

other (specify)

GLASS

bottles:

beverage

food

other (specify)

fluorescent light tubes

light bulbs

pieces

other (specify)

STYROFOAM® (or other plastic foam) Total number of items

buoys

cups

egg cartons

fast-food containers

meat trays

pieces:

larger than a baseball

smaller than a baseball

other (specify)

RUBBER

balloons

gloves

tires

other (specify)

METAL

bottle caps

cans:

aerosol

beverage

food

other

crab/fish traps

55 gallon drums:

rusty

new

pieces

pull tabs

wire

other (specify)

PAPER

bags

cardboard

cartons

cups

newspaper

pieces

other (specify)

WOOD (leave driftwood on the beach)

crab/lobster traps

crates

pallets

pieces

other (specify)

CLOTH

clothing/pieces

(OVER)

Figure 1.--Continued.


A GUIDE TO GOOD DATA COLLECTION

When you help at a beach cleanup, you'll be asked not only to remove marine debris, but to record on Data Cards the kinds and amounts of trash you find.

The information you record on these cards will be used by The Center for Environmental Education (CEE) in a national marine debris study to help policy makers on the state, federal and international levels develop solutions to ending the serious marine debris problems facing all coastal states.

Data collected since 1986 and analyzed by CEE has been used in reports, in testimony on Capitol Hill and at the International Maritime Organization meetings in London to determine how plastic trash will be handled by ships at sea and at ports all around the world.

DATA COUNTS! . . . YOUR HELP WILL MAKE A DIFFERENCE!



HELPFUL TIPS FOR DATA COLLECTORS:

1. Count items in groups of five like this  , and record the total in the box.
2. Do not write the words "Lots" or "Many". Only numbers of items can be put into the computer.
3. Stranded Animals: In this section, please list animals you find stranded or dead on the beach and, if possible, any entangling debris items.
4. Sources: In this section, please list foreign items found and country, if identifiable.
5. Please leave natural items on the beach like driftwood, sea whip and seaweed. Avoid stepping on dune grass and plants. These things hold the sand and prevent erosion.
6. Work with a few people, have one person record the numbers while others collect and bag the trash.
7. Please return your data card to your area coordinator so that all your data will be added to state and national totals.



National Marine Debris Data Base Sponsored By:



Copyright 1988, Center for Environmental Education, Inc.

Return this card for future use

Figure 2.--Guide used by volunteers for data collection, sides 1 and 2.


GUIDE TO MARINE DEBRIS

The best data recording can be done if you know what the items listed on your cards look like.


Here are some examples of unusual items you may find.



Light Sticks. Listed under plastic, these clear plastic tubes about 6 inches long are mostly used by fishermen. When new, the liquid will glow in the dark and attract fish to baited hooks.



Write Protection Rings. Listed under plastic, these are used on computer tapes on ships doing seismic testing.



6-Pack Rings. Listed under plastic, these items are used to hold cans.



Wooden Pallets. Listed under wood, these items are used to help stack and transport cargo.



Strapping Bands. Listed under plastic, these strong, narrow, light-weight plastic bands are used to bind materials and boxes.



55 Gallon Drums. Listed under metal, these drums could contain dangerous chemicals. Do not go near a drum because the vapor or liquid could hurt you.



Vegetable Sacks. Listed under plastic, these large mesh bags are used to hold bulk quantities of onions, potatoes, or fruit.



Sea Whip. This yellow, orange or purple colony of animals is long, thin and has a dark string-like core. This may look like wire or rope, but it is a natural item found from North Carolina to the Gulf of Mexico. Please leave this on the beach.

FOR YOUR SAFETY

Do not approach any 55 gallon drums. They may contain dangerous liquids. Even the vapor could harm you. Leave the drum, but record it on your card.

Do not go into the dunes: snakes may be there.

Be very careful of broken glass and other sharp objects.

Wear gloves.

Don't lift anything heavy.

Copyright 1988, Center for Environmental Education, Inc.



THANK YOU for your help and interest in keeping the coast and ocean safe for all of us and for marine wildlife!

Figure 2.--Continued.

an important model for the subsequent development of a system for categorizing and classifying beach debris on a national level.

The resulting data card was divided into eight major category types-- plastic, glass, Styrofoam, metal, rubber, paper, wood, and cloth (Fig. 1). ("Styrofoam" was used instead of the more technical term "foamed plastic" because it was felt that volunteers would more readily recognize this term.) In total, the data card listed 64 types of debris items. In addition, for each major category there was a listing for "other" to ensure that items not listed on the data card were recorded. Data cards also requested specific information on the sources of foreign debris items as indicated by product labels or other markings, observations of entangled or stranded marine wildlife, observations of peculiar debris items, and comments from volunteers.

With funding from the EPA, the CMC distributed 43,000 data cards to cleanup organizers in the 25 participating states and territories. One thousand additional data cards in the Spanish language were sent to Puerto Rico.

The CMC also developed and distributed 43,000 copies of a 1-page guide for data collection (Fig. 2). The guide gave information on how to use the data card, identified certain debris items that might not be familiar to volunteers, and explained how this information would be used to compile a national assessment of beach debris. Volunteers were encouraged to work in pairs during the cleanup--one person to pick up trash while the other recorded debris.

Each state beach cleanup coordinator was responsible for distributing the data cards and guides to their volunteers and for returning all completed data cards to the CMC for data entry and analysis. All data were then entered into the CMC's National Marine Debris Data Base and analyzed on the basis of national, state, and local findings.

FINDINGS

More than 47,500 volunteers participated in beach cleanups in 1988 in 25 U.S. states and territories (Table 1). One 3-h cleanup was conducted in every coastal state. While the data collected from these cleanups provided a means to assess the debris problem in marine areas, it also provided interesting insights into the extent of the debris problem in inland waters of the United States.

Beach cleanup volunteers covered more than 5,600 km (3,500 mi) of U.S. shorelines and collected nearly 1,000 tons of debris. The methods used to weigh debris varied from state to state, and therefore the weight of debris collected is not exact. However, it is of interest to note that the greatest amount of trash per mile of beach was reported in the states bordering the Gulf of Mexico, particularly Louisiana, Mississippi, and Texas.

On analyzing the data cards, it became obvious that the number of debris items recorded was only an estimate of the true amounts. Some

Table 1.--Number of volunteers, distance cleaned, and amount of debris collected during 1988 beach cleanups (asterisk indicates information not available).

State	Number of volunteers		Distance cleaned		Debris collected		Debris per mile	
	(miles)	(kilometers)	(pounds)	(kilograms)	(pounds)	(kilograms)	(pounds)	(kilograms)
Alabama	40	64	8,340	3,786	208.50	94.66		
Alaska	10+	16+	10,300+	4,676+	*	*		
California	1,100	1,770	200,000	90,800	181.82	82.55		
Connecticut	2	3	190	86	95.00	43.13		
Delaware	54	87	6,054	2,749	112.11	50.90		
Florida	914.6	1,471.6	388,000	176,152	424.23	192.60		
Georgia	50	80	200,000	90,800	4,000.00	1,816.00		
Hawaii	102.8	165.4	100,000	45,400	972.76	441.63		
Louisiana	77	124	180,000	81,720	2,337.66	1,016.30		
Maine	114	183	15,200	6,901	133.33	60.53		
Maryland	18	29	3,750	1,702	208.33	94.58		
Massachusetts	150	241	50,000	22,700	333.33	151.33		
Mississippi	30	48	90,000	40,860	3,000.00	1,362.00		
New Jersey	15.4	24.8	10,021	4,550	652.41	296.19		
New York	4.2	6.7	4,560	2,070	1,085.71	492.91		
North Carolina	150	241	94,000	42,676	626.67	284.51		
Oregon	120	193	28,400	12,894	236.67	107.51		
Pennsylvania	7	11	2,445	1,110	349.28	158.57		
Puerto Rico	17.3	27.8	12,640	5,739	730.64	331.71		
Rhode Island	100	161	15,000	6,810	150.00	68.10		
South Carolina	198	319	30,000	13,620	151.52	68.79		
Texas	120.6	194.0	428,000	194,312	3,548.92	1,611.21		
Virginia	19.8	31.9	12,900	5,857	651.51	295.79		
Virgin Islands	3.2	5.1	*	*	*	*		
Washington	100+	161	64,000	29,056	540.00+	245.16+		
Total	3,517.84	5,660.2	1,953,800	887,025				

volunteers did not count debris items but only commented on the tremendous amounts of debris found. In cases where actual counts were not made, the cards were not added to the data base. But for the most part, volunteers made deliberate and careful efforts to record information. Some who could not identify certain debris items actually sent this trash to the CMC for identification.

Understandably, data collected during volunteer beach cleanups are highly variable and therefore cannot be interpreted exactly, but beach cleanup data can reveal important trends in the relative types, quantities, and distribution of debris. For instance, the data showed that most of the debris found on our nation's coastline is plastic (including Styrofoam). The amount of plastic debris reported surpassed all other categories, accounting for 1,222,708 of the 1,973,995 debris items reported, or approximately 62% (Fig. 3). The remaining debris items consisted of approximately 11.8% paper, 11.4% metal, 9.5% glass, 2.3% wood, 1.8% rubber, and 1.3% cloth. This abundance of plastic debris is also apparent on the state level (Table 2).

The most common debris items reported nationwide were fragmented pieces of plastic and foamed plastic (Styrofoam-like). The data indicate that these plastic pieces accounted for more than 13% of all debris reported. The 12 most common debris items recorded were plastic eating utensils, metal beverage cans, foamed plastic cups, glass beverage bottles, plastic caps and lids, paper pieces (or fragments), plastic trash bags, miscellaneous types of plastic bags (other than trash or salt bags), glass pieces (or fragments), and plastic soda bottles. Collectively, these 12 debris items constituted more than 56% of all debris items recorded (Table 3). Other debris items reported in abundance included approximately 42,700 metal bottle caps, 30,800 plastic six-pack connector rings for




Figure 3.--National composition of debris reported.

Table 2.--Composition of total debris reported
during 1988 beach cleanups.

Percent of total debris collected							
	Plastic	Metal	Paper	Glass	Wood	Rubber	Cloth
National	61.94	11.36	11.77	9.54	2.34	1.80	1.25
By state							
Alabama	63.48	10.81	10.58	10.21	1.87	1.90	1.15
Alaska	56.45	17.83	6.04	15.79	1.21	1.77	0.91
California	47.97	12.11	19.43	15.86	1.42	2.04	1.19
Connecticut	65.99	7.85	7.70	10.54	3.51	2.69	1.72
Delaware	56.82	15.16	14.02	6.78	2.60	2.88	1.76
Florida	59.67	13.31	11.35	10.27	2.72	1.34	1.33
Georgia	57.42	21.13	7.12	9.26	1.99	1.98	1.10
Hawaii	52.14	11.86	16.70	14.54	1.36	2.22	1.19
Louisiana	74.42	7.37	3.58	9.32	1.71	1.87	1.74
Maine	59.69	10.85	12.38	9.74	2.30	3.60	1.45
Maryland	55.79	21.54	7.38	9.10	2.15	2.59	1.46
Massachusetts	61.03	11.40	14.17	7.06	2.49	2.29	1.56
Mississippi	66.29	12.49	7.45	9.63	2.21	1.08	0.85
New Jersey	94.53	1.66	2.47	0.13	0.13	0.85	0.22
New York	77.63	8.99	6.59	4.15	1.02	0.74	0.88
North Carolina	51.81	13.62	20.23	7.22	4.10	1.63	1.39
Oregon	70.16	5.62	14.98	4.85	1.72	1.60	1.10
Pennsylvania	55.85	10.33	22.92	5.19	1.20	2.35	2.17
Puerto Rico	43.36	19.08	11.54	22.09	2.29	0.58	1.05
Rhode Island	60.63	13.20	13.34	6.96	1.26	3.34	1.26
South Carolina	58.86	11.82	16.71	5.81	4.00	1.69	1.11
Texas	76.54	6.73	3.87	8.58	1.63	1.60	1.05
Virginia	61.42	10.65	8.80	8.87	4.29	4.74	1.22
Virgin Islands	60.36	15.23	6.39	13.12	2.44	1.14	1.33
Washington	57.46	7.12	24.58	8.12	1.24	1.07	0.41

beverage cans, 27,600 small pieces of plastic sheeting, 25,200 paper cups, 22,500 foamed plastic fast-food containers.

This information indicates that the majority of debris items found on U.S. shorelines are packaging and disposable plastic products that can be generated by a diversity of ocean- and land-based sources. Certain items, however, are traceable to specific debris sources, and can be used as "indicators" of dumping by maritime and other groups. These indicator items were first identified by the CMC in 1986 with the assistance of the Texas Coastal Cleanup Steering Committee, which included representatives of marine industry groups familiar with the types of debris that could be generated by industry members.

Table 3.--Twelve most common debris items reported during 1988 beach cleanups.

Debris item	Total number reported	Percent of total debris collected
Plastic pieces (or fragments)	134,685	6.82
Small foamed plastic (Styrofoam) pieces	125,725	6.37
Plastic cups, spoons, forks, and straws	112,465	5.70
Metal beverage cans	99,847	5.06
Foamed plastic (Styrofoam) cups	95,807	4.85
Glass beverage bottles	95,028	4.81
Plastic caps and lids	90,998	4.61
Paper pieces	85,864	4.35
Plastic trash bags	78,025	3.95
Miscellaneous types of plastic bags	74,672	3.78
Glass pieces	65,819	3.33
Plastic soda bottles	58,116	2.94
Total	1,117,051	56.59

Using this information, 28 indicator items were identified which fall under four categories: 1) galley wastes generated by crew members on vessels, 2) operational wastes generated during activities conducted by cargo vessels and offshore petroleum operations, 3) fishing and boating gear, and 4) sewage-associated wastes indicative of inadequate sewage treatment practices. Table 4 lists the debris items included under each of these categories. A fifth category, medical wastes, was also identified using plastic syringes as the indicator item. Although the source of syringes as beach debris has not been clearly identified, syringes are suspected to be from illegal dumping, storm water runoff, or inadequate sewer systems.

These 28 indicator items accounted for more than 16% of the debris reported nationwide, with approximately 8% galley wastes, 6% attributable to recreational and commercial fishing and boating, and 2% operational-type wastes. Sewage-associated wastes and medical wastes were comparatively less common, accounting for 0.4 and 0.09% respectively. This information should not be interpreted to mean that these are the only wastes generated by specific ocean- and land-based sources. Rather, the presence of indicator items may show that some of the untraceable debris items are also generated by these same sources.

Furthermore, comparisons of indicator items on the state level showed regional differences in the amount of debris traceable to these sources. For instance, the amounts of galley and operational-type wastes found in states bordering the Gulf of Mexico were much higher than the national figures. On the other hand, while offshore-generated wastes were notably absent on inland beaches on Lake Erie, Pennsylvania, the amount of sewage-associated wastes reported from the Pennsylvania cleanup was six times

Table 4.--Categories and quantities of indicator items used for national assessment of debris reported during 1988 beach cleanups.

Category	Indicator items	Total number reported
Galley wastes	Plastic trash bags	78,025
	Plastic milk and water gallon jug	26,148
	Plastic bleach, cleaner bottles	19,300
	Foamed plastic meat trays	14,721
	Foamed plastic egg cartons	9,526
	Plastic vegetable sacks	6,770
Subtotal		154,490 (7.83%)
Fishing or boating gear	Plastic rope	47,786
	Plastic fishing line	16,563
	Plastic oil and lubricant bottles	12,002
	Plastic light sticks	9,307
	Plastic fishing nets	8,136
	Foamed plastic buoys	7,876
	Plastic floats and lures	5,980
	Rubber gloves	5,748
	Plastic salt bags	3,797
	Wooden fish and crab traps	1,309
	Metal fish and crab traps	1,281
Subtotal		119,785 (6.07%)
Operational wastes	Plastic strapping bands	11,665
	Plastic sheeting longer than 60 cm (2 ft)	7,383
	Glass light bulbs	6,905
	Plastic pipe thread protectors	5,084
	Write-enable protection rings	3,054
	Fluorescent light tubes	2,209
	Wooden pallets	1,737
	Wooden crates	1,075
	Plastic hardhats	857
Subtotal		39,969 (2.03%)
Sewage-associated wastes	Plastic tampon applicators	7,584 (0.38%)
Medical wastes	Plastic syringes	1,718 (0.09%)
Total number of indicator items		343,546 (16.39%)

greater than the national figure, indicating that inadequate sewer systems were a problem in this area.

By noting product labels and other markings, volunteers also reported more than 1,000 foreign label items from 45 countries. In addition, debris from 10 cruise line companies was reported.

Finally, during the 3-h beach cleanup, volunteers reported finding more than 45 cases of wildlife entanglement or ingestion of debris. Of these, more than 40 were birds, most of which were entangled in plastic fishing line.

DISCUSSION

Due to the diversity of debris items and their multiple uses, data collected during beach cleanups cannot realistically be used to estimate total amounts of debris found in marine areas or the exact sources of debris items. However, comparison of relative amounts of debris can reveal important national, state, and local trends in the types and distributions of beach debris. In particular, the first year of the National Marine Debris Data Base demonstrated that plastics account for the majority of waste on our nation's shorelines. Having established a baseline of information on the types and quantities of plastic waste, future beach cleanups can help to monitor legislative and other efforts to control the discharge of plastic trash into marine areas.

By monitoring the presence of indicator items, citizen beach cleanups can also serve to identify what groups are not complying with offshore dumping regulations. This type of information is especially important for developing solutions to the debris problem on the state and local levels.

The great majority of items reported during beach cleanups, however, are virtually untraceable to their specific sources. Yet this information contributes greatly to the underlying theme of a beach cleanup--increased awareness. Since much of this debris consists of items that are used by the general public, those who participate in beach cleanups learn that marine industries are not the only sources of marine debris and that the solution lies with us all. Others who do not participate in beach cleanups hear about data results in the press and media and may consider proper disposal of their trash. Finally, it is hoped that those who manufacture or distribute products that are reported as debris will realize the need to initiate and support efforts that encourage proper disposal and prevent these items from becoming debris.

REFERENCES

- Center for Environmental Education.
1987. Texas coastal cleanup report, 1986. Center for Environmental Education, Washington, D.C., 52 p.
1988. Texas coastal cleanup report, 1987. Center for Environmental Education, Washington, D.C., 105 p.

GUIDELINES FOR THE DESIGN OF BEACH DEBRIS SURVEYS

Christine A. Ribic*
Center for Quantitative Science in Forestry, Fisheries, and Wildlife
University of Washington
Seattle, Washington 98195, U.S.A.

and

Scott W. Johnson
Alaska Fisheries Science Center Auke Bay Laboratory
National Marine Fisheries Service, NOAA
Auke Bay, Alaska 99821, U.S.A.

*Present address: U.S. EPA Environmental Research Laboratory, 200 SW 35th Street, Corvallis, OR 97333, U.S.A.

ABSTRACT

Beach surveys give valuable information as to the types, quantities, and sources of marine debris floating at sea. With the passage of Annex V of MARPOL, however, a decrease in marine debris needs to be detected to demonstrate the effectiveness of mitigating legislation. In order to detect a decrease in marine debris washed ashore, beach survey methodology will need to be standardized. Standardization of survey methods based upon the authors' experience in Alaska is discussed, as is the design of beach surveys to detect between 30 and 50% decreases in the amount of marine debris washed ashore after 5 years, with 95% confidence and power of 80%. Preliminary findings suggest that the number of surveys of a given beach needed to detect a 50% change will be large (bimonthly surveys for 5 years). Annual surveys have low power for detecting a 50% decrease after 5 years, although this result depends on estimates of within-beach variability. Hopefully, this proposal will lead to a discussion of standardized methodology for marine debris beach surveys and the detection of change.

INTRODUCTION

Plastics and other synthetic materials discarded at sea constitute "marine debris" and are now recognized internationally as a form of marine pollution. There is no consensus, however, on how to monitor marine debris after it has washed ashore. Standardized protocols for monitoring other

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

pollutants have been established; examples are tar ball pollution (IOCARIBE 1980) and chemical pollution (Kullenberg 1986). Developing standardized methods will make planning easier and comparison between areas more meaningful.

One of the easiest and most cost-effective methods for arriving at an index of marine debris pollution is the beach survey (Dixon and Dixon 1981; Merrell 1985). The use of beach surveys as indices of floating marine debris, however, requires planned surveys with a clear statement of objectives and assumptions.

This paper has two objectives: (1) to outline the steps involved in planning a beach survey, and (2) to consider two different sampling designs for detecting a decrease in marine debris following the implementation of MARPOL Annex V.

METHODS

A literature review was conducted to identify all published papers on the subject of beach surveys of marine debris excluding tar. Tar pollution was excluded because of the widespread use of standardized techniques to census tar balls on beaches (IOCARIBE 1980). Studies were divided into two groups. The first group focused on describing marine debris on the study area. The second focused on using the beach survey as an indicator of floating marine debris. Next, for all studies, we checked whether the entire beach was surveyed or only portions (transects) of it. Studies were then put into a conceptual framework proposed for planning beach surveys.

The design of surveys to investigate the effect of mitigating legislation (MARPOL Annex V) to reduce the input of debris into the ocean was based on intervention analysis of time series (Hipel et al. 1978; Lettermaier et al. 1978; Barnard et al. 1985). We also considered a repeated measures 1-factor experimental design (Myers 1972) as a second design.

First, we were interested in determining the sample size (number of surveys) needed over 5 years to detect a 30 to 50% change in the amount of marine debris with power (probability of detecting the change) of 0.80 and an alpha of 0.05. An estimate of the variance between years was based on data in Merrell (1985) for Amchitka Island 1972-74. This estimate was used to translate the percentage change into trend or standard deviation ratios needed to use the graphs in Lettermaier et al. (1978). We also considered the effect on power of changing sample sizes in detecting a standardized difference of 1 standard deviation (45% change) over 5 years for alpha = 0.05 and alpha = 0.20. In all cases, gamma, the ratio of number of samples before mitigation to total number of samples, was 0.15 or 0.20. Lettermaier et al. (1978) showed that gamma should be small for the linear intervention model we used.

Secondly, we were interested in the power associated with doing annual surveys for 5 years and detecting a change between 20 and 50% and an alpha of 0.10. Power was taken from Cohen (1977). For Amchitka Island, an

estimate of within-beach variability was calculated from Merrell (1980) for 1972-74 and a mean of 361 pieces of debris per kilometer (Merrell 1984) was used to translate percent change into pieces of debris per kilometer. For the Yakutat area, an estimate of within-beach variability was calculated from data collected 1984-87, and a mean of 205.95 pieces of debris per kilometer was used to translate percent change into pieces of debris per kilometer.

RESULTS

General Beach Survey Design Considerations

The process of beach survey design is summarized in Figure 1. From the published literature, most beach surveys have been short term (one-time surveys) and focused on a single study area (individual beaches) (Cundell 1973; Gregory 1977, 1978, 1983; Bigg 1982; Gregory et al. 1984; Neilson 1985; Willoughby 1986; Henderson et al. 1987; Center for Environmental Education 1988; Marine Plastic Debris Task Force 1988). This focus on the shoreline or beach has led to massive volunteer efforts to clean beaches with little or no reporting of data. These types of studies are valuable for cleaning beaches and gathering information on the types and composition of debris on various coastlines. But quantitative analysis of such data is restricted due to small sample sizes (an annual cleanup means that the sample size is 1) and missing data (especially where data are voluntarily reported).

Few studies stated that their objective was to use the beach debris surveys as an index of floating marine debris. Most studies using beach surveys as an index of oceanic debris were European (Dixon and Cooke 1977; Dixon and Dixon 1981; Shell UK 1985; Federal Republic of Germany 1986; Vauk and Schrey 1987). In the United States, the only published program using beach debris surveys as an index of marine debris is that of Merrell and Johnson in Alaska (Merrell 1980, 1984, 1985; Merrell and Johnson 1987; Johnson 1988; Johnson and Merrell 1988).

In all cases, a precise definition of the sampling unit is needed (Fig. 1). The natural unit is the entire beach from the water's edge to the seaward limit of terrestrial vegetation. In Alaska, most beaches surveyed are 1 km long (Merrell 1985). In England, Dixon and Dixon (1981) used transects, noting that there was too much debris to be totally counted. Where possible, we propose that the entire beach be the survey unit, with results standardized to length (e.g., debris per kilometer). In all cases, the same sampling units should be surveyed over time to minimize variability between surveys.

Before the actual sampling units are chosen, it is best to survey the area of interest to determine beach characteristics and debris distribution (Fig. 1). This is where massive volunteer cleanup efforts can be utilized to help plan the study. It is important to know substrate type, beach slope, prevailing winds, ease of access, and recreational use of the area (Dixon and Dixon 1981; Merrell 1985). Depending upon one's objective, all these factors can influence the choice of beach. Preferred beaches are




Figure 1.--Steps in the design of beach surveys.

moderate-to-steep sand or gravel beaches that are exposed to open ocean. Beaches should have 1 km of similar substrate and scope and be as far as possible from urban areas to minimize bias from local refuse.

Analysis of beach survey data depends on the purpose of the study as well as on the method of data collection. For example, Henderson et al. (1987) examined the distribution of net fragments on the beaches of the Northwestern Hawaiian Islands because they were interested in the location of the debris within the beach. Comparisons between years for beaches is possible with the following caveat. The common statistical tests such as t-tests and analysis of variance (Zar 1986) are not appropriate here because the same beaches are censused over time. The more appropriate techniques are paired-t tests, repeated measures analyses of variance, and their nonparametric equivalents (Conover 1980; Zar 1986). The use of time-series analysis is not appropriate because of the extremely low sample sizes usually found in these studies.

If the focus of the study is an index of floating marine debris, changes over time are of interest but the change is for an unknown area of ocean (Fig. 1). It is not appropriate to extrapolate to other beaches. Merrell (1980) stated that he had problems in extrapolating debris abundance to other beaches.

Beach Surveys as Indicators of Oceanic Debris

There are at least two important assumptions made when using beach surveys as indicators of floating marine debris (Ribic and Bledsoe 1986). The first assumption is that the debris at time t (the first sampling period) is not the same debris as that at time $t+1$ (the second sampling period). In other words, the same debris is not counted twice. The easiest way to fulfill this assumption is by clearing the beach of all surface debris after each survey (e.g., Cundell 1973; Shell UK 1985; Federal Republic of Germany 1986; Henderson et al. 1987). Sometimes this is not practical, especially when debris (e.g., trawl web) is partially buried or snarled on drift logs. In this case, debris can be tagged (Johnson 1988) for identification on later surveys. Tagging studies can provide information on minimum time between surveys as well as information on the loss and deposition rates of beach debris.

The second assumption is that the amount of debris on a beach is related to the amount of debris floating in an unknown area of ocean, and that this area is the same between surveys. In other words, the oceanic area swept onto the beach, when integrated over time, is the same between surveys. This is an important assumption if we want to conclude that a decrease in beach debris over time is due to mitigation measures and not due to a change between years in the area swept onto a beach. We would encourage a study of this assumption if beach surveys are to be useful as indicators of marine debris.

Detecting Change Due to Implementation of Legislation

We will model the potential impact of mitigating legislation (MARPOL Annex V) on the quantity of marine debris washed ashore with the simplest

model: A gradual linear decrease in marine debris over time, after the enactment of the law.

An extremely high survey effort over years will be needed to detect any decrease between 30 and 50% (Table 1). For a 45% decrease, a sample size of 180 translates into a beach survey 3 out of 4 weeks per month per year for 5 years. Detecting a 50% change would call for almost biweekly surveys every month for 5 years. However, the probability of detecting a 45% change is nil using annual surveys and is low using quarterly surveys (Table 2), i.e., the change would have to be so drastic that no statistics would be needed to notice it.

Using a different approach, we looked at treating annual debris counts as a repeated measure on beaches. An estimate of within-beach variability for Amchitka Island was 203 pieces of debris per kilometer. For the Yakutat area, an estimate of within-beach variability was 57 pieces of debris per kilometer. We then calculated minimum and maximum power for changes between 20 and 50% (Table 3) at $\alpha = 0.10$. For Amchitka Island, the probability of detecting any change was low due to the high variability within beaches. Detecting a change of 50% with annual surveys has a power as low as 0.43-0.76 (Table 3). For the Yakutat area, however, the probability of detecting a change of 40% or more using annual surveys was between 0.50 and 0.95 (Table 3). This is due to the low within-beach variability.

DISCUSSION

We are just beginning to realize the magnitude of the marine debris problem. In order to quantitatively assess the problem, standardized beach surveys can be used. Standardization of methodology will make comparisons between areas easier and will ensure the validity of estimates.

It is important for researchers to state explicitly the objectives of their beach debris surveys. Whether or not a survey will be used as an index of floating marine debris affects the survey design from the choice of a particular sampling unit to the data collection and analysis.

A key assumption in using beach surveys to detect a difference due to mitigation is that the area of ocean swept onto the beach is constant between years. Firm conclusions about the effect of mitigating measures in decreasing floating marine debris based on beach debris surveys depend on this assumption. An attempt, therefore, should be made to evaluate its reasonableness.

Beach debris surveys are useful for determining the types and quantities of debris as well as entanglement potential. But the use of beach surveys to detect change with any degree of confidence and power will be more difficult. Preliminary sample size estimates are large for detecting a 50% decrease (power of 0.80; $\alpha = 0.05$). Whether or not annual surveys will be adequate for detecting a 50% change ($\alpha = 0.10$) depends on the estimate of within-beach variability. On Amchitka Island, variability on the same beach is large, and annual surveys will not be adequate for detecting a 50% decrease. In the Yakutat area, however,

Table 1.--Required sample sizes for detecting changes between 30 and 50% of beach debris for $\alpha = 0.05$, power = 0.80, $\gamma = 0.20$, and an estimate of variability = 103.429 pieces of debris per kilometer for a linear intervention model. n = total surveys spread over 5 years.

Percent change	Standardized difference	n
30	0.71	1,000
40	0.95	200
45	1.1	180
50	1.2	100

Table 2.--Probability of detecting a 45% change over 5 years (power) with a $\gamma = 0.15$.

n	Power	
	alpha	
	0.05	0.20
5 (annual)	<0.10	<0.10
20 (quarterly)	0.20	0.45

Table 3.--Minimum and maximum power for a one-factor repeated measures design with $\alpha = 0.10$ and $k = 5$ for detecting changes in beach debris between 20 and 50% for Amchitka Island and Yakutat beaches.

Percent change	Amchitka Island		Yakutat	
	Minimum	Maximum	Minimum	Maximum
20	0.14	0.22	0.20	0.34
30	0.22	0.38	0.33	0.63
40	0.30	0.58	0.52	0.87
45	0.46	0.67	0.61	0.94
50	0.43	0.76	0.71	>0.94

within-beach variability is lower, and annual surveys have a chance of detecting a 50% decrease over 5 years.

Designing a study to measure the impact of legislation to decrease the amount of marine debris will take more planning and a greater commitment of resources. As can be seen from our preliminary findings, sample sizes to detect a given change with stated precision will be large. An improved design could be constructed if we had better variance estimates of debris between beaches as well as within beaches, as well as consensus about the magnitude of the change we would like to detect. In addition, identifying and understanding factors that affect the deposition of debris will be critical in evaluating the success of mitigating legislation.

ACKNOWLEDGMENTS

The initial work on this problem was funded by a contract with the Northwest and Alaska Fisheries Center, National Marine Fisheries Service, NOAA, Contract No. 43-ABNF-5-2498 to the first author. We would like to thank M. B. Hanson for doing the literature search. We thank V. F. Gallucci, M. B. Hanson, and T. W. Miller for reading a draft of this manuscript.

REFERENCES

- Barnard, J., W. Myers, J. Pearce, F. Ramsey, M. Sissenwine, and W. Smith.
1985. Surveys for monitoring changes and trends in renewable resources: Forests and marine fisheries. *Am. Stat.* 39:363-373.
- Bigg, M. A.
1982. Sizes of scrap fishnet and plastic packing bands from western Vancouver Island during August-September 1982. Unpubl. manuscr.; available from Pacific Biological Station, Nanaimo, British Columbia V9R SK6, Canada. Background paper submitted to the 26th Annual Meeting of the Standing Scientific Committee, North Pacific Fur Seal Commission, March 28-29, 1983, Wash., D.C.
- Center for Environmental Education.
1988. Texas coastal cleanup report. Center for Environmental Education, Wash., D.C., 105 p.
- Cohen, J.
1977. Statistical power analysis for the behavioral sciences, revised edition. Acad. Press, N.Y., 474 p.
- Conover, W. J.
1980. Practical nonparametric statistics. 2d ed., Wiley N.Y., 493 p.
- Cundell, A. M.
1973. Plastic materials accumulating in Narragansett Bay. *Mar. Pollut. Bull.* 4:187-188.

- Dixon, T. R., and A. J. Cooke.
1977. Discarded containers on a Kent beach. *Mar. Pollut. Bull.* 8:105-109.
- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.
- Federal Republic of Germany.
1986. Degree and effects of environmental pollution of the German Bight and its coasts caused by synthetic material and other litter discarded by ships. Report submitted at the Thirteenth Meeting of the Standing Advisory Committee for Scientific Advice, Amsterdam, 10-14 March 1986, 15 p.
- Gregory, M. R.
1977. Plastic pellets on New Zealand beaches. *Mar. Pollut. Bull.* 8:82-84.
1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwater Res.* 12:399-414.
1983. Virgin plastic granules on some beaches of eastern Canada and Bermuda. *Mar. Environ. Res.* 10:73-92.
- Gregory, M. R., R. M. Kirk, and M. C. G. Mabin.
1984. Pelagic tar, oil, plastics and other litter in the surface waters of the New Zealand sector of the Southern Ocean and on Ross Dependency shores. *N.Z. Antarct. Res.* 6:12-28.
- Henderson, J. R., S. L. Austin, and M. B. Pillos.
1987. Summary of webbing and net fragments found on Northwestern Hawaiian Islands beaches, 1982-86. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-11, 15 p.
- Hipel, K. W., D. P. Lettenmaier, and A. I. McLeod.
1978. Assessment of environmental impacts, part one: Intervention analysis. *Environ. Manage.* 2:529-535.
- IOCARIBE.
1980. CARIPOL manual for petroleum monitoring, October 1980. Conservation Department File No. 13/2/6.1. February 1981.
- Johnson, S.
1988. Deposition of entanglement debris on Alaskan beaches. In D. L. Alverson and J. A. June (editors), *Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, October 13-16, 1987*, p. 64-94. Unpubl. rep. by Natural Resources Consultants, 4055 21st Ave. W., Seattle, WA 98188.
- Johnson, S. W., and T. R. Merrell.
1988. Entanglement debris on Alaskan beaches, 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-126, 26 p.

- Kullenberg, G. (editor).
1986. The IOC programme on marine pollution. *Mar. Pollut. Bull.* 17:341-352.
- Lettenmaier, D. P., K. W. Hipel, and A. I. McLeod.
1978. Assessment of environmental impacts, part two: Data collection. *Environ. Manage.* 2:537-554.
- Marine Plastic Debris Task Force.
1988. Marine plastic debris action plan for Washington State. Washington State Department of Natural Resources, 45 p.
- Merrell, T. R., Jr.
1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.
1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.
1985. Fish nets and other plastic litter on Alaska beaches. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 160-182. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFS-54.
- Merrell, T. R., and S. W. Johnson.
1987. Surveys of plastic litter on Alaskan beaches, 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-116, 21 p.
- Myers, J. L.
1972. *Fundamentals of experimental design*. 2d ed. Allyn and Bacon, Inc., Boston, 465 p.
- Neilson, J.
1985. The Oregon experience. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 154-159. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFS-54.
- Ribic, C. A., and L. J. Bledsoe.
1986. Design of surveys for density of surface marine debris in the North Pacific. Northwest and Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 267, 69 p.
- Shell UK.
1985. Braefoot Bay marine litter survey. Unpubl. rep. by Exploration and Production Environmental Affairs Department, 20 p.
- Vauk, G. J. M., and E. Schrey.
1987. Litter pollution from ships in the German Bight. *Mar. Pollut. Bull.* 18:316-318.

Willoughby, N. G.

1986. Man-made litter on the shores of the Thousand Island
Archipelago, Java. Mar. Pollut. Bull. 17:224-228.

Zar, J. H.

1986. Biostatistical analysis. Prentice Hall, Inc., Englewood
Cliffs, 2d ed., NJ.

THE COMPOSITION AND ORIGIN OF MARINE DEBRIS STRANDED
ON THE SHORES OF SUBANTARCTIC MACQUARIE ISLAND

David J. Slip and Harry R. Burton
Australian Antarctic Division
Kingston, Tasmania, 7050, Australia

ABSTRACT

The coastline of subantarctic Macquarie Island (lat. 54°35'S, long. 158°55'E) was surveyed over an 8-week period in 1988 to determine types, quantities, and possible sources of marine debris. Lost fishing gear consisted of buoys, ropes, and net fragments. Gear from both trawling and longline fishing operations were represented, with debris identified from Russian, Polish, Japanese, Taiwanese, and South American sources. Three types of litter which potentially entangle marine mammals were found: plastic packing straps, ropes, and net fragments. Plastic bottles, small plastic fragments from broken plastic bottles, and small pieces of expanded polystyrene were common. Litter accumulated in highest densities on open beaches of the west coast of the island. Overall density of marine debris was less than densities reported for islands in the South Atlantic Ocean and the Indian Ocean, or from the coast of Alaska.

INTRODUCTION

Plastic litter and other man-made debris have been recognized as major pollutants of open ocean and coastal surface waters (Shomura and Yoshida 1985; Day and Shaw 1987; Pruter 1987). The distribution of this debris is widespread. Debris has been reported from coastlines near populated, industrial regions of the world (Shiber 1979, 1982), and from open water such as the Mediterranean Sea and the North Sea, which are traversed by busy shipping lanes (Morris 1980; Dixon and Dixon 1983; McCoy 1988), as well as from remote areas such as Alaska (Merrell 1984; Merrell and Johnson 1987) and islands in the Southern Ocean (Gregory 1987; Ryan 1987b; Ryan and Watkins 1988). The slow breakdown of many artifacts allows them to disperse over long distances, often distant from their source (Carpenter and Smith 1972; Ryan 1987b).

Little is known of the impact and fate of much of this debris, although adverse effects through entanglement and ingestion have been documented in seabirds (Azzarello and van Vleet 1987) and marine mammals (Bonner and McCann 1982; Fowler 1985; Laist 1987). The long lifespan and

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

widespread dispersal of such litter make it important to monitor strandings of artifacts in remote sectors of the Southern Ocean in order to identify the major sources of debris and to determine a baseline index of distribution and relative abundance. Parties to the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) have agreed to monitor and report the occurrence of artifacts in the Southern Ocean (Morris 1985). Difficulties in monitoring floating debris, particularly small plastic artifacts, highlight the importance of beach surveys in determining densities, abundance, and sources of such debris.

The present study identifies the types and origins of marine litter stranded at Macquarie Island.

STUDY AREA AND METHODS

Macquarie Island (lat. 54°35'S, long. 158°55'E) is situated in the Southern Ocean. It lies on the Macquarie Ridge approximately 1,500 km south-southeast of Tasmania and 640 km southwest of the Auckland Islands. The island is approximately 34 km long and varies in width from 250 m at the northern isthmus to a maximum of 5 km midway along the island. Commercial sealing took place on the island from 1810 until 1919 (Cumpston 1968). Since 1948, the Australian National Antarctic Research Expeditions (ANARE) have maintained a permanent station on the island. Most of the present day land-based activity on the island is centered around the isthmus and at the six field huts on the island. The nearest ship-based fishing activities occur over the Campbell Plateau, about 700 km to the northeast, and around the Kerguelen group, 5,000 km to the west.

The entire coastline of Macquarie Island (ca. 94 km) was surveyed for stranded artifacts over a 6-week period from June to August 1988. All sizable artifacts (excluding wooden objects) were identified (where possible) from manufacturers' marks in order to determine their origins. All objects were collected and removed from the beaches. Larger objects, such as fishing buoys and floats, were recorded, marked, and stockpiled above high water level at various locations around the island. The rocky nature of much of the coastline precluded sampling of items smaller than 10 mm. Artifacts were categorized according to type of material: plastic, metal, glass, and other. Objects collected from the beach below the ANARE station refuse dump were not included in the survey. Similarly, wooden objects were not included, to avoid confounding the results with wooden remains from sealing activities and old shipwrecks.

The coastline was surveyed in seven convenient arbitrary sectors (Fig. 1), which varied in size and topography. Some sectors were dominated by rock shelf coastline with large boulders, while others had more open beach (Table 1). This is the first survey of marine debris undertaken at Macquarie Island.

RESULTS

A total of 1,034 man-made artifacts was recorded from 94 km of shoreline at Macquarie Island. This represents a density of 11




Figure 1.--Macquarie Island, showing its position in the Southern Ocean and the location of the survey areas.

Table 1.--Relative topography and density of artifacts on sectors of coastline at Macquarie Island.

Sector	Length of coastline (km)	Open beach (%)	Density of artifacts	
			(items/km)	(items/km open beach)
1	4	25	2.5	10
2	16	38	12.2	32.5
3	16	16	19.4	124
4	8	44	39.0	89.1
5	5	20	13.8	69
6	8.5	24	11.6	45.5
7	36	67	0.6	0.9

artifacts/km coastline. Densities of artifacts varied over the island, with the greatest found in sector 4 (39 items/km) and the least (0.6 item/km) on the east coast (Table 1). Plastic items were most abundant, accounting for 60% of all objects. A wide variety of plastic items was recorded. Containers and bottles of various shapes, sizes, and uses were common, while fragments from degrading bottles and polystyrene pieces of various sizes were ubiquitous. Plastic bottles and containers made up 28% of plastic artifacts, while expanded polystyrene and other foamed products contributed 13% of plastic items. Miscellaneous small plastic pieces, including fragments of a size easily ingested by birds, made up 26% of plastic objects. Metal objects contributed 34%, while glass and miscellaneous items such as cork, hemp, cloth, and wax contributed 6% (Table 2). Objects of fisheries origin (buoys, ropes, and netting) accounted for 47% of all stranded artifacts, 29% of all plastic objects, and 89% of all metal objects.

Fishery Debris

A variety of objects originating from fisheries was recorded, including artifacts from both trawl fisheries (eastern European), and longline fisheries (predominantly Japanese). Common items included plastic and metal head line floats from trawl nets, and longline tellings (marker buoys) or surface floats (Table 3). Head line floats were often found attached to fragments of trawl net, and were occasionally found strung together with polypropylene rope in groups of up to six. Three varieties of metal head line floats were recorded: an aluminium alloy float with two separate attachment lugs bearing a Polish manufacturer's mark (55% of all metal buoys), an aluminium alloy float with a single attachment lug containing two holes (26%), and a steel float (17%). Longline tellings were usually covered in rope mesh, often with fragments of rope or mesh trailing from the floats. Ten inflatable plastic floats possibly used as longline pickup buoys or fenders were recovered. Two Japanese glass longline fishery floats, both covered in mesh, were found. Eleven

Table 2.--Numbers of artifacts found on beaches along different sectors of coastline at Macquarie Island.

Artifacts	Sector							Total
	1	2	3	4	5	6	7	
Plastic objects (total)	6	141	198	175	34	52	15	621
Fisheries floats	3	33	53	46	18	18	6	162
Polypropylene ropes	--	2	3	2	--	1	--	8
Netting	--	4	--	--	--	1	--	5
Expanded polystyrene	--	20	26	7	4	11	5	73
Other foamed plastics	--	--	2	1	--	1	--	4
Bags	--	3	--	--	--	--	--	3
Packing straps	--	5	4	1	--	1	--	11
Bottles and containers	--	39	54	61	6	5	3	168
Drift cards	3	3	1	6	2	--	--	15
Miscellaneous	--	32	55	51	4	14	1	157
Metal objects (total)	6	48	79	131	30	43	5	342
Fishery floats	5	40	72	117	28	39	4	305
Containers	--	4	2	6	2	1	--	15
Aerosol cans	--	3	3	8	--	1	--	15
Satellite buoys	--	--	1	--	--	2	1	4
Miscellaneous	1	1	1	--	--	--	--	3
Other objects (total)	1	9	34	12	8	5	2	71
Glass floats	--	--	--	--	1	1	--	2
Bottles and globes	1	7	25	6	3	2	1	45
Cork objects	--	--	1	--	--	2	--	3
Hemp rope	--	--	--	4	2	--	--	6
Bamboo	--	1	1	--	--	--	--	2
Miscellaneous	--	1	7	2	2	--	1	13
Grand total	13	198	311	318	72	100	22	1,034

fragments of rope were recorded. These were of varied length up to about 30 m. Two fragments of teased nylon antichafing blankets were collected, as well as a 2-m fragment of 120-mm stretched mesh codend.

Origin of Artifacts

The country of origin of some of the artifacts could be identified. Most objects were of South American, eastern European (mainly Polish or Russian), or oriental origin (Table 4). The items of fishing debris were identified from writing on floats. Country of origin of fishing floats does not necessarily imply use by that country's fishing fleet, as several nationalities are known to use Japanese floats, although Japanese floats produced for the export market are imprinted in English (R. Burbury pers. commun.). The metal head line floats (Table 3) were of the variety used by

Table 3.--Artifacts which originated from fishing activity washed ashore at Macquarie Island.

Fishing activity debris	Number
Longline fishery debris	
Longline tellings (plastic)	72
Dahn-pole floats	1
Polystyrene surface floats	4
Inflatable pickup floats	10
Trawl fishery debris	
Plastic head line floats	79
Metal head line floats	297
Steel bobbins	8
Trawl-web netting fragments	4
Ropes	11
Antichafing blanket	2
Codend net fragment	1
Miscellaneous fishery debris	
Fender or marker buoy	5
Plastic top buoys (use unknown)	6

the Russian and Polish trawl fisheries, although 55% of these bore a Polish manufacturer's mark. Plastic head line floats included 52 of Russian origin and 2 varieties of Argentinian float (brand names Moscuza and Arex, Table 4). Plastic containers and other litter were identified to country of origin where brand names or manufacturers' names could be read. Most of these artifacts were discarded containers of household consumables such as detergents, shampoos, drinks, aerosol cans, and general food items.

Fifteen plastic drift cards were collected from the coast of Macquarie Island. Of these 13 were of South African origin (Shannon et al. 1973; Lutjeharms et al. 1988), and 2 were Australian.

State of Decay

Plastic objects were found in various stages of decay, ranging from almost pristine with easily discernible printing and little or no sign of ultraviolet degradation to brittle and disintegrating. Most plastic fishing floats had broken lugs, and 32% were in fragments. Generally, fishing floats made of aluminium alloy showed little sign of degradation, although some had broken lugs while others showed signs of pitting. All steel objects were corroded. Many plastic bottles were disintegrating into small fragments, and this type of fragment contributed 46% of miscellaneous plastic objects (Table 2). Most glass bottles were whole.

Table 4.--Countries or regions of origin of artifacts found on Macquarie Island. Where regions only are used, sources of the objects could not be identified more specifically.

Country or region	Fisheries gear	Plastics	Other litter
South America	1	3	4
Argentina	17	4	1
Orient	23	--	--
Japan	37	10	4
China	3	1	--
Scandinavia	1	4	1
Germany	--	1	4
Switzerland	--	--	1
Britain	--	2	2
France	2	--	3
Eastern European	168	--	--
U.S.S.R.	51	3	12
United States	--	2	--
Australasia	--	2	9
South Africa	--	--	13

DISCUSSION

The artifacts stranded at Macquarie Island can have either local or oceanic sources. Some objects may have resulted from the activities of the ANARE station, as rubbish has been dumped in the isthmus area in the past. However, in recent years all plastic, metal, and glass refuse generated by the station has been removed from the island. Thus, in the essential absence of a local source, the stranded artifacts were all derived from oceanic sources, that is, they drifted to the island from distant source-regions.

Macquarie Island lies just to the north of the Antarctic Polar Front (Tchernia 1980). The main oceanic drift pattern which influences Macquarie Island is the West Wind Drift, which has a slight northerly component (Shannon et al. 1973; Lutjeharms et al. 1988). Within that westerly drift, mean surface drift speeds in the Southern Ocean at the latitude of Macquarie Island have been determined at between 14.6 and 19.0 cm/sec (Shannon et al. 1973; Bye 1988; Lutjeharms et al. 1988). Drifting buoys show a tendency to accumulate in areas corresponding to the historic locations of the various frontal systems of the Southern Ocean (Lutjeharms et al. 1988), and the pattern of drift towards these fronts is also demonstrated by the amount of plastic litter which tends to accumulate there (Gregory et al. 1984; Galt 1985; Day and Shaw 1987). Most litter stranded on Macquarie Island would have come from the west (e.g., South American debris), and the influence of the West Wind Drift together with

the prevailing westerly winds would explain the high incidence of strandings on the west coast of the island and the low incidence on the east coast. The northerly component of the West Wind Drift, the tendency of floating material to accumulate at oceanic fronts, and the movement of Antarctic Water toward Macquarie Island in winter, suggest debris stranding at Macquarie Island may originate with ships operating both to the north and to the south of the Antarctic Polar Front.

The debris stranded at Macquarie Island originated in several countries. However, the country of origin of artifacts does not necessarily represent the drift tracks, as much of this debris is probably derived from the fishing fleets and other vessels operating in the Southern Ocean. It is possible that some articles with Spanish writing originated on the coast of South America or from vessels operating in that region. Driftwood originating in South America (predominantly *Nothofagus* spp.) has been reported from South Georgia (Lewis Smith 1985), and from Macquarie Island (Barber et al. 1959).

The significant contribution of fisheries-related objects to the debris on Macquarie Island is similar to that found on other islands in the Southern Ocean (Burton and Williams 1985; Ryan 1987b; Ryan and Watkins 1988), and to the situation on remote beaches in the Northern Hemisphere (Merrell 1984). Fishing gear can become a marine pollutant as a result of accidental loss or deliberate dumping. Fragments of net, line, and associated gear can be lost through snagging, but fishing gear which is worn or damaged beyond further use is often discarded at sea (Pruter 1987). Although vessels have little control over damage to fishing gear, it is possible to control the amount of gear which is actively discarded.

There are two major types of fisheries which operate in the Southern Ocean: bottom and midwater trawling and ocean longlining. The fisheries-related debris which washed ashore on Macquarie Island originated in both these fisheries. Trawl fishing operations for fish and krill in the Southern Ocean are dominated by the U.S.S.R. and Japan, with lesser amounts taken by Poland and the German Democratic Republic. For fish and krill, significant fisheries occur in the Atlantic Ocean sector around South Georgia and the Scotia Arc, and for fish only, around Iles Kerguelen in the Indian Ocean sector (Northridge 1984; Williams 1988). The high number of metal head line floats, Russian plastic head line floats, and other trawl gear (Table 3) were probably lost from the eastern European trawlers operating in these areas, with the majority of fishing gear coming from Iles Kerguelen as they are not only closer to Macquarie Island than is South Georgia, but also closer to the Antarctic Polar Front. Some of the plastic head line floats were of Argentinian origin, and probably came from offshore trawling near the coast of South America. These, along with other South American artifacts, were probably carried to Macquarie Island by the West Wind Drift. Fishing gear from longline fisheries was probably lost from Japanese, Taiwanese, or Korean vessels operating to the north of the Antarctic Polar Front, and possibly quite close to Macquarie Island at times (Robins 1985).

The density of stranded artifacts along a shoreline is determined by a number of geographical factors including beach orientation relative to

prevailing currents and winds, offshore reef structure, beach gradient and texture, and local tide and storm effects (Ryan 1987b). The overall density of artifacts on Macquarie Island (11/km) was less than the densities reported for Prince Edward Island off Africa (32/km), Gough Island (14/km), Tristan da Cunha and Inaccessible Island (from 292 to 807/km; all Ryan 1987b), or Amchitka Island in Alaska (193 to 499/km; Merrell 1980). These differences may be due, in part, to sampling methods, as the surveys above concentrated on particular beaches whereas this survey took into account the entire coastline of Macquarie Island. Surveys of stranded artifacts are often concentrated in areas of noticeably high density with the aim of determining (through repeated surveys) the accumulation rate of debris at one site over time. This may result in inflated estimates of density and make intersite comparisons of density difficult or even invalid. The almost north-south orientation of Macquarie Island results in one coast being exposed to prevailing westerly winds and currents, while the other coast is relatively sheltered. Thus, densities of artifacts in different parts of the island vary widely (Table 1). The low density of artifacts on the east coast (0.6/km) is probably due to the prevailing westerly winds, and surveys from other subantarctic islands have shown that quantities of stranded litter are greatest on the windward (westward) shores (e.g., Gregory 1987; Ryan 1987b). Despite the difficulties of interisland comparisons, the density of stranded litter on Macquarie Island appears to be less than on islands in the Atlantic and Indian sectors of the Southern Ocean (Ryan 1987b), and much less than for remote areas of Alaska (Merrell 1980). This is probably related to the proximity of fishery operations, with most fishing occurring far to the west of Macquarie Island.

The high proportion of plastics among stranded artifacts at Macquarie Island is similar to that reported on other islands of the Southern Ocean (Ryan 1987b) and elsewhere (e.g., Merrell 1980, 1984; Pruter 1987; Shiber 1987). The high incidence of plastics in stranded litter is probably due to the increased use of plastics over the last few decades, particularly for packaging, and the slow rate at which plastics degrade. Surveys in the North Sea and off the coasts of northwest Europe have shown that most marine litter is primary or secondary packaging, particularly plastic bottles, the majority of which originate with the disposal of garbage by ships at sea (Dixon and Dixon 1981, 1983). Plastics decay slowly, with little ultraviolet degradation occurring at sea (Shannon et al. 1973).

The impact of stranded artifacts on local fauna and flora is difficult to assess. Plastic particles are commonly ingested by seabirds but most ingestion probably occurs at sea (Ryan 1987a). Some seabirds at Macquarie Island ingest small fragments of plastic, some of which may have been ingested on shore (Slip pers. observ.).

The density of entanglement-type litter stranded at Macquarie Island is less than densities reported for the Northern Hemisphere (Merrell 1980, 1984). Although the southern elephant seal, *Mirounga leonina*, and fur seal, *Arctocephalus* spp., populations have been closely monitored over the last 5 years, to our knowledge there have been no recent reports of entanglements in these species at Macquarie Island, although one fur seal

was sighted with a plastic packing strap collar in 1975 (G. Copson pers. commun.). Greater concentrations of trawl debris occur in areas of concentrated fishing effort (e.g., Fowler 1987), and as the major fishing effort in the Southern Ocean is far to the west of Macquarie Island, the densities of entanglement debris may be low enough to cause little impact on the marine mammals. However, these species are wide ranging, and fatal entanglements of marine mammals at sea are believed to far outnumber those where the animal reaches shore (Fowler 1987). Thus, once debris is stranded it apparently has little impact on the marine mammals, and ingestion of small plastic fragments by some bird species is likely to be the major impact on local wildlife. The impact of these artifacts is therefore likely to be much greater prior to their stranding.

The amount and variety of stranded artifacts at Macquarie Island demonstrate the preponderance and ubiquity of plastics. The hazards posed to wildlife and shipping by marine litter demonstrate the need for active programs to prevent littering at sea and on land.

ACKNOWLEDGMENTS

We thank members of the Macquarie Island 1988 ANARE for assistance in collecting debris, particularly R. Hamilton, J. Reeve, and P. Charlesworth. R. Williams and J. Van Den Hoff provided useful comments on the manuscript.

REFERENCES


- Azzarello, M. Y., and E. S. van Vleet.
1987. Marine birds and plastic pollution. *Mar. Ecol. Prog. Ser.* 37:295-303.
- Barber, H. N., H. E. Dadswell, and H. D. Ingle.
1959. Transport of driftwood from South America to Tasmania and Macquarie Island. *Nature* 184:203-204.
- Bonner, W. N., and T. S. McCann.
1982. Neck collars on fur seals. *Br. Antarct. Surv. Bull.* 57:73-77.
- Burton, H. R., and D. L. Williams.
1985. National Antarctic Research Expeditions Heard Island, 1985 report. Australian Antarctic Division, Kingston, Tasmania. [Unpublished report.]
- Bye, J. A. T.
1988. Drift cards in the Southern Ocean and beyond (1972-1988). Flinders Institute for Atmospheric and Marine Sciences. Cruise Rep. 12; Flinders University of South Australia.
- Carpenter, E. J., and K. L. Smith, Jr.
1972. Plastics on the Sargasso Sea surface. *Science* 175:1240-1241.
- Cumpston, J. S.
1968. Macquarie Island. National Antarctic Research Expeditions Sci. Rep. Ser. A(1)93:1-380.

- Day, R. H., and D. G. Shaw.
1987. Patterns of abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-1985. *Mar. Pollut. Bull.* 18: 311-316.
- Dixon, T. J., and T. R. Dixon.
1983. Marine litter distribution and composition in the North Sea. *Mar. Pollut. Bull.* 14:145-148.
- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.
- Fowler, C. W.
1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.
- Galt, J. A.
1985. Oceanographic factors affecting the predictability of drifting objects at sea. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 497-507. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Gregory, M. R.
1987. Plastics and other seaborne litter on the shores of New Zealand's subantarctic islands. *N.Z. Antarct. Rec.* 7:32-47.
- Gregory, M. R., R. M. Kirk, and M. C. G. Mabin.
1984. Pelagic tar, oil, plastics and other litter in surface waters of the Southern Ocean and on Ross Dependency shores. *N.Z. Antarct. Res.* 6:12-28.
- Laist, D. W.
1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. *Mar. Pollut. Bull.* 18:319-326.
- Lewis Smith, R. I.
1985. Nothofagus and other trees stranded on islands in the Atlantic sector of the Southern Ocean. *Br. Antarct. Surv. Bull.* 66:47-55.
- Lutjeharms, J. R. E., L. V. Shannon, and L. J. Beekman.
1988. On the surface drift of the Southern Ocean. *J. Mar. Res.* 46:267-279.


- McCoy, F. W.
1988. Floating megalitter in the eastern Mediterranean. *Mar. Pollut. Bull.* 19:25-28.
- Merrell, T. R., Jr.
1980. Accumulation of plastic litter on beaches from Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.
1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.
- Merrell, T. R., Jr., and S. W. Johnson.
1987. Surveys of plastic litter on Alaskan beaches, 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-116, 21 p.
- Morris, R. J.
1980. Floating plastic debris in the Mediterranean. *Mar. Pollut. Bull.* 11:125.
1985. Antarctica's living resources: Are they in safe hands? *Oryx* 19:65.
- Northridge, S. P.
1984. World review of interactions between marine mammals and fisheries. *FAO Tech. Paper* 251, 189 p.
- Pruter, A. T.
1987. Sources, qualities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Robins, J. P.
1985. Study of sashimi tuna potential in Western Australia. *Fins* 18:6-19.
- Ryan, P. G.
1987a. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.
1987b. The origin and fate of artifacts stranded at islands in the African sector of the Southern Ocean. *Environ. Conserv.* 14:341-346.
- Ryan, P. G., and B. P. Watkins.
1988. Accumulation of stranded plastic objects and other artefacts at Inaccessible Island, central South Atlantic Ocean. *S. Afr. J. Antarct. Res.* 18:11-13.
- Shannon, L. V., G. H. Stander, and J. A. Campbell.
1973. Oceanic circulation deduced from plastic drift cards. *Sea Fish. Branch Invest. Rep.* 108, 31 p.
- Shiber, J. G.
1979. Plastic pellets on the coast of Lebanon. *Mar. Pollut. Bull.* 10:28-30.

1982. Plastic pellets on Spain's 'Costa del Sol' beaches. Mar. Pollut. Bull. 13:409-412.
1987. Plastic pellets and tar on Spain's Mediterranean beaches. Mar. Pollut. Bull. 18:84-86.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Tchernia, P.
1980. Descriptive regional oceanography. Pergamon Press, Oxford, 253 p.
- Williams, R.
1988. Antarctic fish. Aust. Nat. Hist. 22:518-521.
-

SESSION II



ENTANGLEMENT OF MARINE LIFE AND GHOST FISHING



DISTRIBUTION OF MARINE DEBRIS AND NORTHERN
FUR SEALS IN THE EASTERN BERING SEA

Norihiisa Baba, Masashi Kiyota, and Kazumoto Yoshida
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

ABSTRACT

To obtain basic information about entanglement rate and mortality of the northern fur seal, *Callorhinus ursinus*, at sea, we conducted sighting surveys of fur seals and marine debris along eight transect lines in 1984 and four in 1985 and 1988 in summer near the Pribilof Islands in the eastern Bering Sea. These southeast to northwest transects were approximately 300-500 km long. We observed 710 fur seals and 7 debris items of fisheries origin in 1984, 345 seals and 17 debris items in 1985, and 343 seals and 18 debris items in 1988. In 1985, one dead male fur seal was observed entangled in a trawl net fragment weighing 40 kg. Distributions of both marine debris and fur seals were concentrated in the area along the continental slope west of the Pribilof Islands. It is considered that this co-occurrence is a result of the mutual relationship between fish resources, seals' feeding, fishing grounds of trawlers in the area, and northward-flowing current.

INTRODUCTION

Japanese trawlers began operating in the eastern Bering Sea in 1933 and other nations have begun fishing there later the U.S.S.R. in 1959, South Korea in 1968, Taiwan in 1974, Poland in 1979, and West Germany in 1980. The estimated total number of trawl-fishing vessels off Alaska increased from 5 in 1933 to 432 in 1963, and dropped to 317 in 1983 (Low et al. 1985).

At the 10th meeting of the North Pacific Fur Seal Commission (NPFSC), the survival rate of fur seals that were entangled in fishing net fragments was reported (NPFSC 1967). Since then, the United States has been actively collecting data on entanglement of fur seals (Scordino 1985). Japan-United States joint research started in 1983 (Bengtson et al. 1988; Scordino et al. 1988). The fur seal population on the Pribilof Islands has steadily declined since the 1960's, and entanglement of seals has been suggested as a partial cause (Fowler 1982).

Merrell (1980) estimated that about 1,645 metric tons of plastic material were dumped into the Bering Sea and Aleutian Islands area each year in the 1970's. Dahlberg and Day (1985) encountered 0.356 trawl debris items per 1,000 km in the central North Pacific between Kodiak Island and Hawaii, whereas Jones and Ferrero (1985) found 1.349 pieces of trawl net debris per 1,000 km off the Aleutian Islands in the North Pacific. The common pelagic distribution of fur seals and marine debris has not, however, been studied at all. If the drifting routes and local accumulations of marine debris coincide with migration routes and feeding grounds of fur seals, the probability that seals will become entangled in marine debris will increase. The greater the density of marine debris, the greater will be the number of entangled seals. To properly assess the impact of entanglement on the fur seal population, the common distribution and density of fur seals and marine debris at sea must be known.

In this study, we conducted a sighting survey for fur seals and marine debris in the eastern Bering Sea in 1984, 1985, and 1988, and obtained basic information on the pelagic distribution of fur seals and marine debris.

MATERIALS AND METHODS

We conducted sighting surveys of northern fur seals and marine debris along eight transect lines from 13 July to 8 August 1984, four from 12 to 21 July 1985, and four from 10 to 23 July 1988 in the eastern Bering Sea using RV *Shunyo Maru* (Table 1). These southeast to northwest transects were approximately 300 to 500 km long. The survey areas made up of blocks measuring 30 min of latitude by 1 degree of longitude, were 248,845 km² in 1984, 152,937 km² in 1985, and 184,066 km² in 1988. In 1984, both western and eastern areas of the Pribilof Islands were surveyed, and in 1985 and 1988 only the western area was surveyed (Fig. 1).

Sightings were conducted by one or two people from the pilothouse and four or five people from the flying deck (8 m above sea level) on top of the pilothouse each day from sunrise to sunset. Observers were placed on both sides of the ship and surveyed the area on only one side. Each observer engaged in sightings for 4 h and rested for 1 h. Binoculars (7 × 50) were used only to confirm the kind and number of objects observed. We recorded the number, the time, and the location of fur seals and marine debris encountered. The speed of the ship during sightings was about 8 kn in 1984 and 1985 and about 10 kn in 1988. The ship's course was not changed except to collect debris of fishing origin such as fishing net fragments, plastic packing bands, floats, and ropes. When visibility dropped to less than about 200 m, the survey was interrupted.

In 1984 and 1985 we concentrated on sighting of fisheries-related debris; however, in 1988 we recorded all floating debris including Styrofoam, nylon bags, wood, and debris of fisheries origin. Because the debris surveys differed among years, we compared only the distributions and densities of fisheries-related debris.

Table 1.--Research period, area east or west of the Pribilof Islands, distance traveled, and number of fur seals and pieces of debris of fisheries origin observed.

Period	Days	Area	Distance traveled (km)	Number of fur seals observed	Number of pieces of debris of fisheries origin observed
13-19 July 1984	7	East	1,133	71	1
25 July-8 Aug. 1984	9	West	1,855	639	6
12-21 July 1985	9	West	1,892	345	17
10-23 July 1988	9	West	2,184	343	18

RESULTS

Distribution and Kinds of Debris of Fisheries Origin

In 1984, seven pieces of debris of fisheries origin were found: two on the continental shelf northwest of the Pribilof Islands, four near the continental slope southeast of the islands, and one to the south of St. George Island (Fig. 2A). In 1985, 17 pieces of debris of fisheries origin were found 2 on the continental shelf, 12 along the continental slope, and 3 northwest of Umnak Island (Fig. 2B). In 1988, 18 pieces of debris of fisheries origin were collected 2 on the continental shelf, 9 along the continental slope, and 7 southwest of the Pribilof Islands (Fig. 2C). Generally, debris items were found along the continental slope during the 3 years.

Debris of fisheries origin collected in 1984, 1985, and 1988 included trawl nets, gillnets, string, rope, floats, and plastic packing bands. Fifteen trawl net pieces collected ranged in weight and mesh size from 15 g, 7 cm to 40 kg, 20.5 cm; two gillnet pieces were similar at 1.75 kg, 11.7 cm and 1.8 kg, 11.5 cm. Three of four packing band pieces collected weighed 6.4 kg or more (Table 2). Trawl net accounted for 71.4% (five pieces) of all debris in 1984, 41.2% (seven pieces) in 1985 and 16.7% (three pieces) in 1988 (Table 3). Trawl net constituted the major part of the collection in 1984, floats in 1988.

Entanglement of Fur Seal

We found a dead male fur seal (110 cm long and weighing 20 kg), which we estimated to be 2 years old, entangled in a net fragment about 30 nmi southwest of St. Paul Island on 19 July 1985. The net fragment was gray trawl net weighing 40 kg (mesh size 20.5 cm; twine size 7.6 mm).

Distribution of Fur Seals

The sighting frequency of fur seals (number of fur seals sighted per 1 km) was calculated for each block measuring 30 min of latitude by 1 degree of longitude.




Figure 1.--Transect line surveyed in 1984 (A), 1985 (B), and 1988 (C) (— = 100 fathoms, --- = 1,000 fathoms, - . - = 2,000 fathoms).




Figure 2.--Location of debris of fisheries origin.

● = trawl net, ○ = gillnet, * = unidentified net,
 ⊗ = string, × = rope, Δ = float, ■ = plastic packing band.

Table 2.--Kinds (PP = plastic packing band), date, location, and characteristics of fisheries-related debris observed on the line transect surveys, 1984, 1985, and 1988, in the eastern Bering Sea.

No.	Kinds	Date	Latitude N	Longitude W	Color	Weight (kg)	Mesh (mm)	Twine (mm)	Length (m)
1	Gillnet	14 July 1984	58°20'	171°26'	Green	1.8	115	0.5	--
2	Trawl net	25 July 1984	55°13'	167°18'	Blue	17.6	140	4.0	--
3	Trawl net	29 July 1984	57°34'	172°20'	Blue	3.0	130	5.0	--
4	Trawl net	5 Aug. 1984	54°40'	167°37'	Green	0.015	70	3.0	--
5	Trawl net	5 Aug. 1984	54°48'	167°55'	Orange	0.35	200	3.5	--
6	Trawl net	5 Aug. 1984	55°10'	168°43'	Green	1.15	135	4.0	--
7	PP band	6 Aug. 1984	55°04'	170°22'	Yellow	9.4	--	--	--
8	Trawl net	12 July 1985	53°43'	170°21'	^a Green	--	--	--	--
9	Trawl net	12 July 1985	54°20'	171°45'	Green	6.4	155	3.0	--
10	Trawl net	15 July 1985	54°16'	169°31'	Gray	0.52	170	3.0	--
11	Trawl net	19 July 1985	57°02'	171°20'	Orange	2.9	125	6.0x4.5	--
12	Trawl net	19 July 1985	56°57'	171°07'	Gray	40.0	205	7.6	--
							114	3.4	--
13	Trawl net	20 July 1985	56°05'	169°12'	Green	0.07	--	4.2	--
14	Trawl net	21 July 1985	55°25'	168°01'	Orange	0.03	195	2.2x4.6	--
					Black	--	--	3.4	--
					Green	--	--	2.0x4.4	--
15	Gillnet	21 July 1985	55°25'	168°06'	Green	1.75	117	0.5	--
16	Rope	17 July 1985	56°09'	171°33'	White	0.71	--	18.5	2
17	Rope	17 July 1985	56°59'	173°18'	Yellow	0.2	--	25.0	1.3
18	Rope	20 July 1985	56°18'	169°49'	Yellow	4.8	--	18.0	20
19	Rope	20 July 1985	56°12'	169°31'	White	1.8	--	19.8	6
20	Rope	20 July 1985	56°03'	169°06'	Yellow	8.6	--	18.0	50
21	Rope	20 July 1985	55°50'	168°32'	Yellow	2.0	--	17.4	13
22	Rope	21 July 1985	55°13'	167°27'	Yellow	0.1	--	12.4	1
23	String	21 July 1985	55°16'	167°34'	Orange	0.02	--	3.0x5.0	2
24	PP band	20 July 1985	55°50'	168°32'	White	0.01	--	--	2
25	Net	17 July 1988	57°17'	171°58'	(a)	--	--	--	--
26	Trawl net	12 July 1988	55°32'	172°25'	Gray	--	--	--	--
27	Trawl net	23 July 1988	56°06'	169°19'	Orange	0.82	129	5x3	--
28	Trawl net	23 July 1988	56°13'	169°32'	Orange	0.75	195	5x3	--
29	Float	10 July 1988	54°17'	171°41'	--	--	--	--	--
30	Float	12 July 1988	56°28'	174°18'	--	--	--	--	--
31	Float	12 July 1988	56°29'	174°18'	--	--	--	--	--
32	Float	12 July 1988	55°59'	173°13'	--	--	--	--	--
33	Float	13 July 1988	55°16'	171°40'	--	--	--	--	--
34	Float	15 July 1988	55°39'	170°21'	--	--	--	--	--
35	Float	16 July 1988	56°14'	171°42'	--	--	--	--	--
36	Float	16 July 1988	56°34'	172°28'	--	--	--	--	--
37	Float	16 July 1988	56°44'	172°47'	--	--	--	--	--
38	Float	16 July 1988	56°44'	172°48'	--	--	--	--	--
39	Float	17 July 1988	57°28'	172°21'	--	--	--	--	--
40	Float	23 July 1988	55°37'	168°18'	--	--	--	--	--
41	PP band	10 July 1988	54°09'	171°24'	Yellow	6.4	--	--	^b Roll
42	PP band	15 July 1988	56°01'	171°11'	Yellow	6.4	--	--	^b Roll

^aNot collected.

^bThe roll of plastic packing band was estimated to be >100 m.

Table 3.--Kinds, number, and percent of fisheries-related debris pieces collected during line transect surveys in 1984, 1985, and 1988 in the area west of the Pribilof Islands.

Kind		1984	1985	1988	Total
Trawl net	No.	5	7	3	15
	%	(71.4)	(41.2)	(16.7)	(35.7)
Gillnet	No.	1	1	0	2
	%	(14.3)	(5.9)	(0.0)	(4.8)
Unidentified net	No.	0	0	1	1
	%	(0.0)	(0.0)	(5.6)	(2.4)
Float	No.	0	0	12	12
	%	(0.0)	(0.0)	(66.7)	(28.6)
Rope	No.	0	7	0	7
	%	(0.0)	(41.2)	(0.0)	(16.7)
String	No.	0	1	0	1
	%	(0.0)	(5.9)	(0.0)	(2.4)
Plastic packing band	No.	1	1	2	4
	%	(14.3)	(5.9)	(11.1)	(9.5)
Total	No.	7	17	18	42

In July 1984, we surveyed the area east of a line extending through St. George and St. Paul Islands. Most blocks showed fewer than 0.3 seal/km or showed no seals in this area.

In August 1984 and in July 1985 and 1988, we surveyed the area west of same line. In August 1984, more than 0.3 seal/km were seen in many blocks along the continental slope, and southwest of the islands over 0.9 seal/km were seen. No seals were found to the southeast or over the continental slope (Fig. 3A). Mean frequency of seals in all blocks west of the islands was greater than to the east, differing significantly ($t = 4.7528$, $P < 0.0001$).

In July 1985, we found fur seals mainly on the continental shelf and along the continental slope. Frequencies of over 0.6 seal/km occurred in two blocks to the northwest and southwest of St. Paul Island (Fig. 3B). Mean frequency of seals of this year was less than in August 1984, differing significantly ($t = 2.449$, $P < 0.005$).

In July 1988, we found frequencies greater than 0.3 seal/km only on the continental shelf and along the continental slope within about 200 km of the Pribilof Islands. Frequencies were greater than 0.9 seal/km in two




Figure 3.--Sighting frequency of fur seals per block measuring 30 min of latitude by 1 degree of longitude including transect line. Frequency equals the number of fur seals sighted per 1 km of research distance. (NI = not investigated.)

blocks along the continental slope south of St. George Island. Generally we saw no seals near the northwestern and southeastern ends of the continental slope or along the southwestern edges (Fig. 3C). Mean frequency of seals this year was almost the same as in July 1985, not significantly different ($t = 0.6499$, $P > 0.006$).

Coincident Sightings of Fur Seals and Fisheries-Related Debris

The sighting frequencies of fur seals in coincidental areas surveyed in 1984, 1985, and 1988 were 360.4/1,000 km in 1984, 211.0/1,000 km in 1985, and 197.8/1,000 km in 1988, whereas the densities of debris (of fisheries origin only) were 2.561/1,000 km in 1984, 7.975/1,000 km in 1985, and 9.798/1,000 km in 1988 (Table 4).

DISCUSSION

Most female fur seals at St. Paul Island deliver pups in July. After a perinatal fast of 8-10 days, they go to sea to feed for 4- to 10-day periods punctuated by 1-2 days of nursing their pups. Feeding trips lengthen as pups age until they are weaned at about 120 days postpartum (Peterson 1968). York and Kozloff (1987) reported that the number of new-born pups on St. Paul Island did not change greatly between 1981 and 1986. Therefore, we believe that the greater number of fur seals sighted in 1984 was due to the later survey period (late July-early August) and consequent greater proportion of lactating females at sea then, compared with 1985 and 1988 when surveys were in early to mid-July.

In all years, we found most seals near the continental slope in the eastern Bering Sea. Echo soundings of fish biomass, which we conducted simultaneously with transect surveys, indicated that the walleye pollock, *Theragra chalcogramma*, biomass was greatest in that area (Harada et al. 1985). Kajimura (1984) reported that fur seals in the Bering Sea ate mostly capelin, *Mallotus villosus*, and walleye pollock in July and August.

Table 4.--Sighting frequency of fur seals and fisheries-related debris (fishing nets, rope, string, plastic bands, floats) observed on the line transect surveys in 1984, 1985, and 1988 west of the Pribilof Islands.

Period	Research distance (km)	Seals	Debris	Frequency per 1,000 km	
		No.	No.	Seals ^a	Debris ^b
25 July-8 Aug. 1984	1,562	563	4	360.4	2.561
12-21 July 1985	1,630	344	13	211.0	7.975
10-23 July 1988	1,633	323	18	197.8	9.798

^aNumber of seals divided by research distance.

^bNumber of debris items divided by research distance.

Trawl-net fisheries for those species also operate primarily along the continental shelf and the continental slope in the eastern Bering Sea (Mito 1986), also suggesting that the greatest fish biomass is concentrated there. We believe that our observations showing most marine debris and most fur seals concentrated in the area from the continental shelf to the continental slope west of the Pribilof Islands are related to the concentration of prey resources and marine debris (e.g., fishing net, plastic packing bands) in that area and to the northward currents along the continental slope (Favorite et al. 1976) which act to concentrate debris there.

As fur seals migrate in winter from the Pribilof Islands to as far south as Mexico in the eastern Pacific (lat. 32°N) (Kajimura and Loughlin 1988), we feel that it is important to conduct surveys in waters off British Columbia, Washington, Oregon, and California in the future to ascertain the distribution and abundance of marine debris and fur seals there.

ACKNOWLEDGMENT

We greatly appreciate the crew of RV *Shunyo Maru* for their cooperation in the survey and the officials of the Fishing Ground Preservation Division, Fisheries Agency, Government of Japan, for their assistance and advice in planning the survey. We especially thank the National Marine Mammal Laboratory, Alaska Fisheries Science Center, National Marine Fisheries Service, NOAA, who provided us the opportunity to conduct these surveys.

REFERENCES

- Bengtson, J. L., C. W. Fowler, H. Kajimura, R. L. Merrick, K. Yoshida, and S. Nomura.
1988. Fur seal entanglement studies; juvenile males and newly weaned pups, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Favorite, F., A. J. Dodimead, and K. Nasu.
1976. Oceanography of the subarctic Pacific region, 1960-71. Int. North Pac. Fish. Comm. Bull. 33:1-187.
- Fowler, C. W.
1982. Interactions of northern fur seals and commercial fisheries. Trans. N. Am. Wildl. Nat. Resour. Conf. 47:278-292.

- Harada, Y., N. Nagai, S. Toishi, K. Yoshida and N. Baba.
 1985. Estimation of the pollock biomass in the sea around the Pribilof Islands in the Bering Sea from mid-July to mid-August, 1984. Background paper submitted to the 28th meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 18 p.
- Jones, L., and R. C. Ferrero.
 1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Kajimura, H.
 1984. Opportunistic feeding of the northern fur seals, *Callorhinus ursinus*, in the eastern North Pacific Ocean and Bering Sea. U.S. Dep. Commer., NOAA Tech. Rep. NMFS-SSRF-779:1-49.
- Kajimura, H., and T. R. Loughlin.
 1988. Marine mammals in the oceanic food web of the eastern subarctic Pacific. Bull. Ocean Res. Inst., Univ. Tokyo 26:187-223.
- Low, L.-L., R. E. Nelson, Jr., and R. E. Narita.
 1985. Net loss from trawl fisheries off Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 130-153. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Merrell, T. R.
 1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. Mar. Environ. Res. 3:171-184.
- Mito, K.
 1986. Outline of the Japanese groundfish fishery in the Bering Sea in 1985 January-December and 1986 January-July. Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Anchorage, Alaska, October 1986. Fisheries Agency of Japan, Tokyo, 100 Japan, 5 p.
- North Pacific Fur Seal Commission.
 1967. Report of the Standing Scientific Committee. In Proceedings of the Tenth Annual Meeting, North Pacific Fur Seal Commission. Issued from the Headquarters of the Commission, Washington D.C., 48 p.
- Peterson, R. S.
 1968. Social behavior in pinnipeds with particular reference to the northern fur seal. In Harrison et al. (editors), The behavior and physiology of pinnipeds, p. 3-53. Appleton-Century-Crofts, N.Y., 409 p.

Scordino, J.

1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Scordino, J., H. Kajimura, N. Baba, and A. Furuta.

1988. Fur seal entanglement studies in 1984, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), *Fur seal investigations*, 1985, p. 70-78. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.

York, A. E., and P. Kozloff.

1987. On the estimation of numbers of northern fur seal, *Callorhinus ursinus*, pups born on St. Paul Island, 1980-86. *Fish. Bull.*, U.S. 85:367-375.

POTENTIAL IMPACT OF ENTANGLEMENT IN MARINE
DEBRIS ON THE POPULATION DYNAMICS OF THE
NORTHERN FUR SEAL, *CALLORHINUS URSINUS*

Deborah P. French and Mark Reed
Applied Science Associates, Inc.
Narragansett, Rhode Island 02882, U.S.A.

ABSTRACT

A model of the population dynamics of the Pribilof Island population of the northern fur seal, *Callorhinus ursinus*, was developed using field estimates of age- and sex-specific mortality and reproductive rates. In the model, mortality rates of pups and juveniles are density dependent, while rates for older seals are constants for each age and by sex. Model-predicted pup production is compared to pup counts made in the field.

If initialized at 1912 numbers, the model population increases at the rate observed between 1912 and 1940, then oscillates around the population level of the 1940's and 1950's, thought to be the carrying capacity of the Pribilof Islands. When the female harvest (including pelagic collections) of 1956-74 is added to the simulation at the annually specified rates, the predicted pup production decreases until 1965, but recovers to preharvest levels by 1975.

Entanglement rate and resulting mortality are assumed to be proportional to the rate of entanglement of subadult males in the harvest, specified by year. When both female harvest and mortality resulting from entanglement in plastic debris are added to the simulation, the model population declines at observed rates, suggesting that entanglement mortality can account for the recent decline in the fur seal population. The model indicates that the population will continue to decline at 1% per year if mortality remains at current levels. This rate of decline is slower than the 4-8% decrease per year of the 1970's because of higher pup and juvenile survival rates at lower population density. The model indicates that a reduction in entanglement mortality rates by 20% would be enough to stop the current population decline and maintain the population at current levels.

INTRODUCTION

In the early 1900's, the population of the northern fur seal, *Callorhinus ursinus*, breeding on the Pribilof Islands in the Bering Sea had been reduced by harvesting to about 300,000 total individuals and fewer than 100,000 pups born per year. A ban on harvesting in 1911 by international agreement allowed the population to recover, such that a population level estimated at about 2 million total individuals was reached by the 1940's and continued through the 1950's. (See reviews by York and Hartley 1981; Lander and Kajimura 1982; Scheffer et al. 1984; Fowler 1985a.) In 1956, harvest of females was begun in an effort to reduce the population size to a level at which the then-predicted maximum sustainable yield in pup production could be obtained. The harvest of females was continued until 1968, and pelagic collections of females for research purposes were made between 1958 and 1974 (York and Hartley 1981). When the female harvest was ended, it was expected that the population would increase and return to the 1950's population level. However, the population has continued to decline at 4-8% per year since the late 1970's (Briggs and Fowler 1984). Evidence indicates that this decline may be the result of lethal entanglement in fishing debris and plastic packing bands (Fowler 1985a, 1985b).

In order to evaluate quantitatively the effect of the female harvest and entanglement on the Pribilof Island fur seals, we have developed a population dynamics model and have compared resulting predicted pup production against estimates made in the field. The population model consists of annual age classes of males and females with mortality and reproductive rates which are dependent on age and sex. Mortality of pups on land and of juveniles during their first 20 months at sea is density dependent, while mortality rates of older seals and pregnancy rates of females are density-independent, age-specific constants using best available estimates. Harvest mortality is considered separate from natural mortality and, when applicable, occurs only during the summer harvest season. Entanglement mortality rates are added to natural mortality when simulating the population dynamics of the last two decades.

MODEL ASSUMPTIONS

The age-specific pregnancy rates calculated by York (1979), using data from the pelagic collections conducted by the United States and Canadian Governments between 1958 and 1974, are assumed as birth rates (Table 1 and Fig. 1). Therefore, reproductive rates in the model are dependent solely on the number of adult females, assuming that enough adult males are present to impregnate those females at all population densities. The sex ratio at birth is assumed 1:1.

Natural mortality of seals age 2 or more years is assumed to be constant by age class and sex. In the model, natural mortality does not include mortality due to commercial harvest (or pelagic collection for research purposes) or recent mortality believed to be due to entanglement in fishing gear. Natural mortality rate estimates by age and sex are available from several sources (Chapman 1964; Lander 1979, 1980a, 1981; Eberhardt 1981; Smith and Polacheck 1984). Those of Lander (1980a, 1981;

Table 1.--Pregnancy rates (York 1979), age-specific natural mortality rates (Lander 1980a, 1981), and harvest rates on immature males (Lander 1980a) used in the population dynamics model. (Asterisks indicate rates which are density dependent and therefore not constants.)

Age	Percent of females pregnant	Natural survival rate (per year)		Male harvest rate (per year)
		Female	Male	
1	0	*	*	0
2	0	0.840	0.78	0.028
3	0	0.920	0.77	0.403
4	4	0.940	0.76	0.573
5	37	0.940	0.74	0.147
6	70	0.945	0.72	0
7	80	0.950	0.72	0
8	85	0.950	0.72	0
9	87	0.938	0.70	0
10	88	0.924	0.65	0
11	88	0.906	0.63	0
12	88	0.884	0.60	0
13	87	0.858	0.55	0
14	84	0.876	0.50	0
15	81	0.789	0.43	0
16	77	0.743	0.30	0
17	71	0.692	0.20	0
18	63	0.630	0.10	0
19	56	0.564	0	0
20	47	0.490		
21	37	0.411		
22	26	0.330		
23	11	0.300		
24	0	0.250		
25	0	0.200		
26	0	0.150		
27	0	0.100		
28	0	0.050		
29	0	0		

Table 1 and Fig. 2) are assumed in the present model, after correction for subadult male harvest rate was made.

The use of constant mortality rates assumes that mortality is independent of population density. While there is no evidence to date that mortality of older fur seals is density dependent, there is evidence of density dependence for the mortality of pups on land and for juveniles <2 years old (Fowler 1984, 1985a; Smith and Polacheck 1984). Thus, density-dependent relationships for these age groups are included in the model.




Figure 1.--Pregnancy rate as a function of age.




Figure 2.--Natural survival and harvest rates as functions of age and sex (after Lander 1980a, 1981).

The mortality of pups on land appears to increase with increasing number of pups counted on the rookeries (Lander 1979; Swartzman 1984). In the model, natural mortality rate of pups on land is a function of the number of pups born, assuming the functional relationship drawn by Swartzman (1984) between pups born on St. Paul Island and their estimated mortality (line SL in Fig. 3). Pups born on St. Paul are assumed to represent 80% of the total for the two islands, St. Paul and St. George (Briggs and Fowler 1984; Smith and Polacheck 1984). Pups are assumed to remain on land for 128 days (based on data of Gentry and Holt 1986) and on-land mortality is applied over that time period. Male and female pups suffer the same rate of mortality while on land.

Survival during the first 20 months at sea also appears to be related to the number of pups born (Chapman 1961; Lander 1979; Eberhardt 1981; Fowler 1985a, 1985b). Using data for the 1950 to 1970 year classes (Fig. 4, line PA), Lander (1979) found that survival of males during their first 20 months at sea is linearly related to pup survival on land. As pointed out by Fowler (1985b), when data for the years after 1970 are included, the relationship no longer holds. Fowler presented evidence that this difference in juvenile mortality is related to the increase of entanglement in fishing debris and other plastic materials, which he believed to become significant after 1965. Therefore, a linear regression for the data of 1950-65, excluding 1956, is used in the present model, as suggested by Fowler (1985b, line PB in Fig. 4).

Survival rates of female juveniles appear to be higher than those of males of the same age (Chapman 1961, 1964; York 1987). However, the magnitude and density dependence of female juvenile survival is unknown. Using a variety of techniques, Chapman (1961, 1964) estimated the ratio of female to male survival to age 3. From a simple population model using weighted-average pregnancy rates and mortality rates of females older than 3 years, Chapman (1961) calculated that this ratio should be about 2.0. In a similar analysis, Chapman (1964) calculated a ratio of 1.72 using 1920's population estimates and 1.27 using 1950's data. Based on tagging returns, Chapman (1964) estimated the ratio of female to male survival to age 3 for 10 year classes (1951-60), finding an average value of 1.64. However, he pointed out that these estimates are probably biased such that the ratio should be higher. Two of the ten tagging estimates were about 1.27, and the other eight (later) estimates averaged 1.74. Thus, two values for the ratio of female to male survival to age 3 were tried in the present model, 1.27 and 1.74. Since mortality of pups on land is assumed the same for both sexes and mortality from age 2 to 3 is 1.077 times as high for females as males (Table 1), the ratio of female to male survival during the first 20 months at sea is assumed 1.18 or 1.62.

Because the number of males over the age of 2 years has no influence on female survival rate, reproductive rate, or future population size in the model, the mortality rate of males older than 2 influences only the number of males in the population. Thus, harvest rate of subadult males is assumed to be zero in all simulations reported here, except as indicated where the 1970's harvest rates in Table 1 and Figure 3 are used.




Figure 3.--Alternate density-dependent pup mortality curves. Data are from Lander (1980b); SL is the curve drawn by Swartzman (1984) through the data; MAX and MIN are the maximum and minimum curves tested here; UEN is a sigmoid alternative to MAX; SIG is a sigmoid alternative to SL; and LOW is a lower version of SL.




Figure 4.--Alternate linear regressions of juvenile mortality rate as a function of pup survival. PA is from Lander (1979), regressed on the data for 1950-70; PC is the same, but for 1950-65 (before significant entanglement is thought to have occurred); and PB is for 1950-65, excluding 1956.

Harvest rate of females is assumed to be zero, except when specifically simulating the years 1956 to 1974. For those years, the numbers of females by age harvested or collected pelagically (as reported in York and Hartley 1981) are subtracted from the model population at the time of harvest during the specific year. As assumed by York and Hartley, the age distribution of females in the harvest where age was not determined is assumed to be the same as that for harvested females of known age. The age distribution of females grouped as older than a specified age (e.g., 10 years) is assumed to be equivalent to the age distribution in the model population for that year.

Entanglement mortality rates are the least known of the model parameters which must be estimated. Fowler (1985a, 1985b) has shown that, while there is a linear relationship between male juvenile survival the first 20 months at sea and pup survival on land using data up to 1965, following 1965 the model no longer fits the observations. The discrepancy between the model and observed juvenile mortality for the years after 1965 is linearly correlated with the rate of entanglement observed in the subadult male harvest. This suggests the discrepancy is due to entanglement mortality. The discrepancy (as percent mortality) is added to natural mortality of juveniles up to 2 years of age in the model when entanglement is added to the simulation. Because the average percent entanglement in the harvest has stabilized at 0.4% since the late 1970's, the corresponding discrepancy of 15% mortality is assumed for both male and female juveniles as a starting point in model simulations. The discrepancy has ranged from 7 to 25% since 1966 as rates of entanglement have varied (Fowler 1985b).

Mortality rate for entangled males between 2 and 3 years of age was estimated by Fowler (1985b) from the relative frequency of entangled males in the 2- and 3-year-old age classes in the harvest. His estimate is that 5.5% of 2- to 3-year-old males become entangled each year and that 90.3% of entangled males die. Thus, 5% mortality due to entanglement is used in model simulations for male seals between 2 and 3 years of age. Females of this age are assumed to suffer the same or higher rates of entanglement mortality, due to their smaller size (see Discussion).

Entanglement mortality rates of older seals are not available. Fowler (1985b) suggests that entanglement mortality of seals over 3 years of age is assumed to be between 0 and 5% per year. Different values were tried in various simulations. Both males and females are assumed to suffer the same entanglement mortality once past the age of 3.

For all simulations, the populations were initialized at 1 January sizes and ages. A daily time step was used. Mortalities and births were calculated on appropriate dates in the yearly cycle.

RESULTS

Simulation of the Population at Carrying Capacity

Assuming the above-described pregnancy rates, constant natural mortality rates for seals over age 2, no harvest on either males or females, and

no entanglement mortality, the density-dependent pup mortality relationship SL in Figure 3, and the male juvenile survival relationship PB in Figure 4, population equilibria were found for each of the two assumed values of the ratio of female to male survival up to the age of 3, 1.27 and 1.74. Assuming this ratio is 1.27, the population reaches 1.73 million total individuals (censused on 1 January), 1.15 million total females, and 442,000 pups born per year. If the female to male survival ratio is assumed to be 1.74, the total population reaches 1.95 million seals, 1.39 million females, and 540,000 pups born each year.

In the 1940's and 1950's, the fur seal population on the Pribilofs was relatively stable, with 525,000-576,000 pups born each year in the 1950's. This population is thought to have been at carrying capacity for the Pribilof Islands (Fowler 1985a). The model population, assuming a female-to-male survival ratio to age 3 of 1.74, is consistent with observations, while the lower ratio of 1.27 causes the model population to fall short of the observed levels.

If the model is initialized at 15% of the equilibrium population size, a level where pup production matches that of 1912, the model population and number of pups born increase at the same rate as observed pup counts in the 1910's and 1920's, 7.4% per year (Fig. 5). The model population overshoots the equilibrium size in the early 1940's, and afterwards oscillates around the equilibrium of 540,000 pups born per year, with the oscillation damping out over time. The agreement between the model and the observed rate of increase in the 1910's and 1920's suggests that the assumed pregnancy and mortality rates are realistic. Furthermore, a ratio of female to male survival to age 3 on the order of 1.74 is consistent with available data. This ratio was suggested by Chapman's results (1961, 1964), even though he and others have been more comfortable with a much lower ratio. When this same simulation is run with the female-to-male juvenile mortality ratio at 1.27, the projected population increase from the 1912 level is less than the observed rate of increase (Fig. 6).

To determine how sensitive the simulated carrying capacity population is to the various pregnancy and mortality rates assumed, these rates were varied individually within the range of possibilities observed to determine resulting population size. Alternate pregnancy rates by age (up to age 11) were taken from Chapman (1964) and from estimates reported in Smith and Polacheck (1984, table 11), which were based on Japanese collections of females between 1958 and 1960. The Japanese estimates represent the highest reported values for pregnancy rates of younger females. These high rates increase pup production to 583,000 per year (+8%) and decrease the total number of females by 9% (Table 2). Chapman's pregnancy rate estimates increase pup production by only 2% and have almost no effect on the total number of females in the population.

The density-dependent relationship between pups born and pup mortality on land is associated with a large amount of variation, and a number of alternate curves through the observed data were tried (Fig. 3). The resulting carrying capacity population varies from 23% higher to 25% lower than the model result using the standard assumptions (Table 2). Thus, the model is fairly sensitive to this assumed relationship.




Figure 5a.




Figure 5b.

Figure 5.--Simulated increase in (a) population level (on 1 January) and (b) number of pups born each year from the depleted population of 1912 to carrying capacity reached by the 1940's. The ratio of female to male survival to age 3 is 1.74. Harvest and entanglement-induced mortality are assumed zero in the simulation.

Table 2.--Resulting equilibrium population levels (on 1 January) and number of pups born each year (in millions of seals) when reproductive rates or mortality rates of certain age groups are varied (see text for specific rate variations) from the standard run producing the best fit to the 1950's carrying capacity population (pregnancy as in Table 1; pup mortality as curve SL in Figure 3; male juvenile survival as line PB in Figure 4; ratio of female to male survival to age 3 = 1.74; no harvest; and mortality over age 2 as in Table 1).

Reproductive or mortality rate(s) varied	Total seals	Total females	Pups born
Standard run (equilibrium of Fig. 5)	1.95	1.39	0.540
Pregnancy rates:			
As in Chapman (1964)	1.93	1.38	0.553
As in Japanese collections, 1958-60	1.79	1.27	0.583
Pup mortality curve (from Fig. 3):			
MIN	2.37	1.71	0.647
LOW	2.23	1.59	0.622
SIG	2.20	1.57	0.611
UEN	1.62	1.16	0.450
MAX	1.46	1.04	0.407
Male juvenile survival line (from Fig. 4):			
PA	1.92	1.37	0.533
PC	1.91	1.36	0.527
Ratio of female to male survival to age 3:			
2.0	2.00	1.46	0.570
1.64	1.92	1.36	0.529
1.37	1.81	1.23	0.473
1.27	1.73	1.15	0.442
1.08	1.54	0.99	0.376
Subadult male harvest rates:			
As in Table 1	1.92	1.39	0.540
Adult female mortality rates:			
As in Chapman (1964)	1.96	1.39	0.509
As in Smith and Polacheck (1984)	1.90	1.31	0.454

Male juvenile survival as a function of pup survival on land was varied to be as line PA in Figure 4, i.e., the relationship in Lander (1979), and as line PC in Figure 4, i.e., including all data from 1950 to 1965 in the regression. The resulting population size differed from the standard run using line PB (Fig. 4) by at most 2%. Thus, the results are insensitive to this amount of variation of the relationship.

The ratio of female to male survival to age 3 has been estimated by Chapman (1961, 1964) as between 1.25 and 2.0. The resulting model population sizes range from 18% less than to 5% greater than the standard population model run where the ratio is 1.74 (Table 2). Thus, the model is fairly sensitive to the assumed ratio within the range of estimates which have been made. If the ratio is assumed to be 1.0, the population reaches an equilibrium 30% lower than the standard model equilibrium (Table 2), a level well below the 1950's population size.

As mortality of males over the age of 2 has no influence on female or pup population size in the model, the model population is insensitive to variation in subadult male harvest rates or variation in natural mortality of males over the age of 2. A subadult male harvest rate equivalent to that in the 1970's (Lander 1980a) reduces the total male population by 5% (Table 2).

Mortality rates of females over the age of 3 as given by Chapman (1964) result in the same number of total females in the population as the standard model assumptions (Lander 1981), but the age distribution is such that only 509,000 pups are born per year (-6%, Table 2). The lower estimated survival rates of Smith and Polacheck (1984) reduce the female population by 6% and pup production at equilibrium by 16%. However, the estimates of Lander (1981) are based on considerably more data than the other two sets of estimates, and the error of the standard model run associated with error in adult female mortality rates is probably somewhat less than these results.

Simulation of the Pribilof Population Decline Since 1958

When the female harvest and pelagic collection of 1956-74 is removed from the simulated carrying capacity population during those years, the predicted number of pups born decreases until 1965 and subsequently recovers to the preharvest level by 1975 (Fig. 7). While the female harvest may account for some of the decline after 1958, it is clearly a minor perturbation from which the population should have rapidly recovered, assuming the carrying capacity remained unchanged from the 1950's level.

The population decline after the female harvest ceased has been a subject of much discussion in recent years, with the prevailing opinion being that lethal entanglement is the most likely causative factor (Fowler 1985a, 1985b). Assuming that entanglement became significant in 1966 (as suggested by Fowler), that an additional 15% of males and females die from entanglement by age 2, and that 5% mortality of all seals age 2 and older is due to entanglement, the resulting pup production is as in Figure 8. Under these assumptions, the model population does not decline as fast as the observed pup production indicates it should. This suggests that entanglement mortality may have been significant before 1966, or that there is some other cause of the additional mortality after 1958. If the same assumed mortality rates are initiated in 1960, the model decline fits the observed more closely (Fig. 9).




Figure 6.--Number of pups born in model population assuming the same rates as in Figure 5, except the ratio of female to male survival to age 3 is assumed to be 1.27.




Figure 7.--Number of pups born to modeled and observed populations over time, including simulation of female harvest between 1958 and 1974 but assuming no mortality due to entanglement.




Figure 8.--Number of pups born to modeled and observed populations over time, including simulation of female harvest (1958-79) and constant entanglement mortality of 15% before and 5% after 2 years of age beginning in 1966.




Figure 9.--Number of pups born to modeled and observed populations over time, including simulation of female harvest (1958-79) and constant entanglement mortality of 15% before and 5% after 2 years of age beginning in 1960.

In the model runs shown in Figures 8 and 9, all seals over age 2 are assumed to suffer 5% mortality per year due to entanglement. If seals over age 3 are assumed to suffer insignificant (zero) mortality due to entanglement, as suggested by Fowler (1985b), the model population declines to only 1.06 million total females with 401,000 pups born per year, a level much higher than observations for the last two decades (Fig. 10). Thus, adult females must be suffering mortality over and above Lander's (1981) estimated rates, whether due to entanglement or some other cause.

The rate of entanglement in the subadult male harvest was not constant between 1967 (the first year of observation) and the present. Before 1970, entanglement rate was below 0.4%. From 1973 to 1975 it increased to about 0.7% and subsequently stabilized at about 0.4% (Fowler 1985b). Before 1970, male juvenile survival was closer to 10%, rather than 15%, below the expected rate from line PB in Figure 4. Between 1973 and 1975 the discrepancy between observed and expected (line PB) was about 20%.

Assuming that entanglement varied proportionately for all age classes of seals, all entanglement mortality rates in the model were reduced by one-third over 20 months (one-fifth per year) before 1970 and increased by the same amount for 1973 to 1975. Thus, an additional 10 to 20% of juveniles are assumed to die from entanglement by age 2. Entanglement mortality of all seals over 2 years is assumed to vary between 4 and 6% per year. The resulting pup production is compared to the observed in Figure 11. The fit to the observed is improved by this inclusion of variable entanglement rate in the model (compare Figs. 9 and 11).

If the same assumptions used for the simulation of Figure 11 are made, only with 2- to 3-year-olds suffering 8 to 12% mortality per year (an annual rate equivalent to the additional 15% lost over 20 months as juveniles +20% of that value) while rates for seals over 3 remain at 4 to 6% per year, the resulting decline is as in Figure 12. This latter simulation brings the model population level down to the same level as the current pup counts on St. Paul Island indicate. Figure 13 shows associated total population numbers.

Model Predictions of Future Population Numbers

The model simulation of Figure 13 (and Figs. 8, 9, 11, and 12 as well) indicates that, while the current rate of decline is much slower than the 4 to 8% decline of the 1970's, the Pribilof fur seal population would continue to decline at about 1% per year for the next 50 years or more if mortality rates remain at current levels. If entanglement mortality were eliminated (and assuming pre-1960 survival rates still hold), the population could recover to the 1950's carrying capacity level after about 15 years, as evidenced by the increase from the late 1920's to the 1940's (Fig. 5). If current entanglement mortality rates (as assumed in Figs. 8 and 9) were halved for all age groups, the population could recover in about 50 years to a total of 1.61 million seals, 1.13 million total females, and 400,000 pups born, where the model population stabilizes (Fig. 14). This is a size equivalent to the observed population in 1960. If entanglement mortality rates were held at the levels assumed for the 1960's




Figure 10.--Number of pups born to modeled and observed populations over time, including simulation of female harvest (1958-74) and, beginning in 1966, constant entanglement mortality of 15% before age 2, 5% from age 2 to 3, and no entanglement mortality over the age of 3 years.




Figure 11.--Number of pups born to modeled and observed populations over time, including simulation of female harvest (1958-74) and variable entanglement mortality rates (10-20% before and 4-6% after age 2 years) beginning in 1960.




Figure 12.--Number of pups born to modeled and observed populations over time, including simulation of female harvest (1958-74) and variable entanglement mortality rates (10-20% before age 2, 8-12% between 2 and 3 years, and 4-6% after age 3 years) beginning in 1960.




Figure 13.--Simulated population numbers (on 1 January) for the simulation of Figure 12.




Figure 14.--Simulated population numbers (on 1 January) assuming that current entanglement rates are halved and after initializing with the estimated population size in 1987.

in the model run of Figure 12 (an additional 10% of juveniles die before age 2, 8% per year from age 2 to 3, and 4% per year over age 3), the population would increase slowly from present levels (917,000 total seals, 637,000 females, and 214,000 pups born in the 1987 simulated population) to 1.29 million seals, 896,000 females, and 301,000 pups born per year after 50 years and would stabilize at 1.34 million seals, 932,000 females, and 315,000 pups after 90 years.

DISCUSSION

The model results show that current estimates of pregnancy and natural mortality rates as outlined above are consistent with observational data from 1912 to 1958 if the ratio of female to male survival to age 3 is assumed to be 1.74. The model simulates the population increase from 1912 to the 1940's and predicts the appropriate carrying capacity level thought to have prevailed in the 1940's and 1950's. Lower estimates of the female-to-male juvenile survival ratio yield much lower equilibria and rates of population increase than have been observed on the Pribilofs. Clearly, the model is sensitive to female juvenile survival rate. However, this vital statistic is the least well known and least studied of all the vital rates involved. Tagging or other studies on female juveniles would greatly facilitate the estimation of their current survival rates.

Model sensitivity analysis shows that model results are also highly dependent upon the choice of the density-dependent relationship between pup mortality and number of pups born. The considerable variability associated with this functional relationship (Fig. 3) is undoubtedly due to the dependence of pup mortality on other factors besides total number of pups born. Actual pup density on the rookeries, climatic conditions, and competition with other species for food are other possible factors (which are not necessarily independent) which could be considered in developing more predictive functional relationships.

The model is relatively insensitive to changes in pregnancy rates and adult female natural mortality rates within the range of estimates available in the literature. Therefore, these vital statistics appear to be well enough known for modeling purposes.

The carrying capacity of the Pribilof Islands population appears to be regulated by the density of pups born on the rookeries. Subsequent survival as juveniles appears to be causally related to survival rate on land (Lander 1979, Fig. 4). This suggests that lower survival rate at higher population density is primarily due to reduced fitness of pups, whether by increased spread of disease, slower weight gain due to competition for food or space among lactating females, or other effects of crowding. The model is consistent with the view that there is no significant density-dependent control of mortality over the age of 2 years.

Owing to the weak density-dependent control of the fur seal population, the maximum pup production is at carrying capacity, not at a lower population size. This is in agreement with observations in the field during and after the period of female harvest in the late 1950's and 1960's.

The population decline after 1958 has been a subject of intense interest in recent years (e.g., Eberhardt 1981; Fowler 1984, 1985a, 1985b; Smith and Polacheck 1984; Trites 1984). The initial decline may be partially accounted for by the female harvest of 1956 to 1968 (York and Hartley 1981 and present model), but after 1960 it is evident that other factors are involved (Fig. 7). The model results suggest that entanglement mortality can account for the decline after 1960 if the following assumptions are made regarding female mortality (male mortality has no influence on pup production): An additional 15% of juveniles (less than age 3), on average, died from entanglement; an average of 5% of seals over age 3 suffered lethal entanglement; and significant entanglement mortality (at these rates) began in 1960.

The fit of the model to the observed pup counts is improved considerably if entanglement mortality rate is varied as a function of the entanglement rate observed in the subadult male harvest. A variation of $\pm 50\%$ of entanglement rate, i.e., 0.2 to 0.6%, appears to result in $\pm 33\%$ variation in the mean discrepancy between observed male juvenile survival rate and that predicted from line PB in Figure 4 (Fowler 1985a), or $\pm 20\%$ variation in entanglement mortality rate per year. If entanglement mortality rates of all age groups vary to this degree in proportion to

variation in observed entanglement rates, the appropriate curvature in the model pup curve after 1965 is produced.

The best fit is obtained if 2- to 3-year-old females are assumed to have suffered twice the entanglement mortality rates of males of the same age. If females over 3 years of age are assumed to not die at significant rates from entanglement, the decline in pups born since 1960 cannot be accounted for, and the model predicts that a stable population size of 1.49 million seals, 1.06 million females, and 401,000 pups born per year should have been reached by about 1985 (Fig. 10). Higher entanglement mortality of females as opposed to males of the same age could be accounted for if entanglement rate is a function of body size, rather than age as assumed by Fowler (1985a, 1985b). Even at 1 year of age, male fur seals are almost twice the size of females; a 3-year-old male is the same size as a 6-year-old adult female, and the largest females (up to 20 years old) are not as large as a 5-year-old male (Lander 1980a). This suggests that if entanglement is primarily a function of size, adult females could be expected to suffer the same entanglement rates as subadult males, perhaps 5% per year as estimated by Fowler (1985b), and immature females even higher rates. The assumption of 10% entanglement mortality rate for 2- to 3-year-old females is consistent with this hypothesis. Clearly, more information is needed on female entanglement rate and mortality as a function of age. The assumption that males and females of the same age suffer the same mortality rates is unlikely, especially given the disparity in size.

In the above analysis it is assumed that pups on land do not suffer entanglement mortality. In the model, pups of mothers dying from entanglement are assumed to die. Recent evidence reported by DeLong et al. (1988) suggests that mortality rates of pups may be increased by nonlethal entanglement of their mothers, since the females are less efficient at foraging and providing milk for the pup. However, as evident in Figure 3, pup mortality rates on land at a given density do not appear to have changed from 1914-65 to 1966-83, before and after the hypothesized onset of entanglement. Therefore, there is no evidence to support the hypothesis that nonlethal entanglement of lactating females has significantly affected pup mortality on land or the pup population as a whole.

As shown by the model results in Figure 14, a reduction of the current extrinsic mortality rate (due to entanglement or some other cause) by 50% would allow the fur seal population to recover at least partially. If this mortality is in fact due to entanglement, halving the current entanglement rate (to 0.2% from 0.4% of subadult males), which would reduce the annual entanglement mortality rate by 20%, would stop the current decline in pup production, according to model predictions. This suggests that it would be necessary to halve the density of plastic debris in the North Pacific and Bering Sea region to stop the decline and maintain the population at current levels.

The model results reported here suggest that the fur seal population can recover from single-event perturbations such as substantial harvest of the breeding population (females in the case of fur seals) or some other cause of mortality of finite duration. After the fur seal population was

reduced by 1912 to 15% of its carrying capacity level, the population was able to recover fully in 30 years. Recovery would be faster after smaller perturbations. However, long-term, continuous additional sources of mortality, such as lethal entanglement, form a much more serious threat to the population. While single-event perturbations are significant in the short term, chronic sources of additional mortality are much more important when considering the long-term stability of the population.

ACKNOWLEDGMENTS

This research was supported by the Minerals Management Service, Alaska Outer Continental Shelf (OCS) Region, U.S. Department of the Interior, OCS Study No. MMS 86-0045, Contract No. 14-12-0001-30145. John Calambokidis, James Cabbage, and Gretchen Steiger of Cascadia Research Collective of Olympia, Washington assisted in literature review and provided valuable comments. The authors wish to especially thank Charles Fowler of the National Marine Mammal Laboratory (NMML), National Marine Fisheries Service, Seattle, for his advice and comments. Also, comments and suggestions by Ann York, Roger Gentry, Wendy Roberts, Tom Loughlin, and Michael Goebel of NMML are gratefully acknowledged, as are those of Gordon Swartzman of the University of Washington, Brent Stewart of Hubbs-Marine Research Institute, Maxwell Dunbar of McGill University, Douglas DeMaster of the Southwest Fisheries Science Center, National Marine Fisheries Service, La Jolla, and Steve Treacy of the Minerals Management Service, Alaska OCS Region.

REFERENCES

- Briggs, L., and C. W. Fowler.
 1984. Tables and figures of the basic population data for northern fur seals of the Pribilof Islands. Background paper submitted to the 27th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, Moscow, USSR, April 1984. [Available from C. W. Fowler, National Marine Mammal Laboratory, NMFS, 7600 Sand Point Way, N.E., Seattle, WA, 98115.]
- Chapman, D. G.
 1961. Population dynamics of the Alaska fur seal herd. *Trans. N. Am. Wildl. Nat. Resour. Conf.* 26:356-369.
1964. A critical study of Pribilof fur seal population estimates. *U.S. Fish Wildl. Serv., Fish. Bull.* 63:657-669.
- DeLong, R. L., P. Dawson, and P. J. Gearin.
 1988. Incidence and impact of entanglement in netting debris on northern fur seal pups and adult females, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), *Fur seal investigations*, 1985, p. 58-68. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.
- Eberhardt, L. L.
 1981. Population dynamics of the Pribilof fur seals. In C. W. Fowler and T. D. Smith (editors), *Dynamics of large mammal populations*, p. 197-220. Wiley, N.Y.

Fowler, C. W.

1984. Density dependence in northern fur seals (*Callorhinus ursinus*) of the Pribilof Islands, Alaska. Paper presented at the International Symposium and Workshop on Fur Seal Biology, 23-27 April 1984, Cambridge, England.

1985a. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1985b. Status review: Northern fur seals (*Callorhinus ursinus*). Presented at 28th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, held in Tokyo, Japan, April 1985. Federal Register 50(44):9231-9248, Mar. 6, 1985.

Gentry, R. L., and J. R. Holt.

1986. Attendance behavior of northern fur seals. Chapter 3. In R. L. Gentry and G. L. Kooyman (editors), Fur seals: Maternal strategies on land and at sea. Princeton Univ. Press.

Lander, R. H.

1979. Role of land and ocean mortality in yield of male Alaskan fur seal (*Callorhinus ursinus*). Fish. Bull. U.S. 77:311-314.

1980a. A life table and biomass estimate for Alaskan fur seals. In H. Kajimura, B. H. Lander, M. A. Perez, A. E. York, and M. A. Bigg (editors), Further analysis of pelagic fur seal data collected by the United States and Canada during 1958-74. Report submitted to 23rd Annual Meeting of the North Pacific Fur Seal Commission. Part 1, p. 44-57. National Marine Mammal Laboratory, NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 98115.

1980b. Summary of northern fur seal data and collection procedures. Vol. 1. Land data of the United States and Soviet Union (excluding tag and recovery records). U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA, as No. PB 81-106502.

1981. A life table and biomass estimate for Alaskan fur seals. Fish. Res. (Amst.) 1:55-70.

Lander, R. H., and H. Kajimura.

1982. Status of northern fur seals. In Mammals in the sea, p. 319-345. FAO Fish. Ser. 5(IV).

Scheffer, V. B., C. H. Fiscus, and E. I. Todd.

1984. History of scientific study and management of the Alaskan fur seal, *Callorhinus ursinus*, 1786-1964. U.S. Dep. Commer., NOAA Tech. Rep. NMFS-SSRF-780, 70 p.

Smith, T. D., and T. Polacheck.

1984. The population dynamics of the Alaska fur seal: What do we really know? Report to National Marine Mammals Laboratory, NMFS, 7600 Sand Point Way N.E., Seattle, WA 98115, 122 p.

Swartzman, G. L.

1984. Factors bearing on the present status and future of the eastern Bering Sea fur seal population with special emphasis on the effect of terminating the subadult male harvest on St. Paul Island. U.S. Marine Mammal Commission Report No. MMC-83/03. [Available from U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as No. PB84-172329, 77 p.]

Trites, A. W.

1984. Stock assessment and modeling of the North Pacific fur seal population. Final report to Canadian Department of Fisheries and Oceans, Contract No. OST83-00133, 82 p.

York, A. E.

1979. Analysis of pregnancy rates of female fur seals in the combined United States--Canadian pelagic collections, 1950-74. In H. Kajimura, R. M. Lander, M. A. Perez, A. E. York, and M. A. Bigg (editors), Preliminary analysis of pelagic fur seal data collected by the United States and Canada during 1958-74. Report submitted to 22nd Annual Meeting of the North Pacific Fur Seal Commission, p. 50-122. National Marine Mammal Laboratory, NMFS, NOAA, 7600 Sand Point Way NE, Seattle, WA 98115.

1987. Northern fur seal, *Callorhinus ursinus*, eastern Pacific population (Pribilof Islands, Alaska, and San Miguel Island, California). In J. P. Croxall and R. L. Gentry (editors), Status, biology and ecology of fur seals. Proceedings of an International Symposium and Workshop, 23-27 April 1984, Cambridge, England, p. 9-21. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 51, 212 p.

York, A. E., and J. R. Hartley.

1981. Pup production following harvest of female northern fur seals. Can. J. Fish. Aquat. Sci. 38:84-90.

STUDIES OF THE POPULATION LEVEL EFFECTS
OF ENTANGLEMENT ON NORTHERN FUR SEALS

Charles W. Fowler, Richard Merrick, and Jason D. Baker
National Marine Mammal Laboratory
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
Seattle, Washington 98115, U.S.A.

ABSTRACT

Recent studies have focused on entanglement among the juvenile male northern fur seal, *Callorhinus ursinus*, as a means of evaluating the effects of entanglement at the population level. Most entanglement-related field studies were conducted on St. Paul Island, Alaska, in the 1980's but the analyses include relevant data from the late 1970's. Reported here are the results of recent studies on monitoring of entanglement, estimates of entanglement-caused mortality, and the effects entanglement may have on the chances an animal is observed on the breeding islands.

The observed proportions of seals entangled in 1985 and 1986 were consistent with those observed during the last few years of the commercial harvest (about 0.4%). The proportion observed in 1988 was 0.29%, the lowest observed since 1970. The change reflects a drop in the numbers of animals entangled in fragments of trawl webbing. The frequency of occurrence of trawl webbing among the entangling debris was about half the former levels whereas the proportion of seals entangled in other types of debris did not change.

These studies confirm earlier estimates indicating that, after 1 year, the survival of seals entangled in debris light enough to permit the animals to return once to land is about half of the survival of nonentangled seals. Data indicate that the main factor contributing to the success of entangled animals that do survive is escapement from the debris.

Rates at which entangled animals are resighted indicate that the proportion of animals resighted drops with an increase in the size (weight) of debris.

Data from radio-tagged seals confirm that entangled seals go to sea for longer periods of time than do controls.

INTRODUCTION

Entanglement in marine debris, specifically in plastics associated with the commercial fishing industry, has been documented for a number of species of seals and sea lions (Fowler 1988). The effects of entanglement in such debris have been the subject of a number of studies, especially as related to the impact on the northern fur seal, *Callorhinus ursinus*. Many of these studies have examined effects at the population level (Fowler 1982, 1985, 1987; Swartzman 1984; French and Reed 1990; Swartzman et al. 1990). Others have studied the effects at the level of the individual (Fowler 1988).

Entanglement of northern fur seals in marine debris has been a concern for several decades. The first sightings of entangled seals occurred just after World War II. Records of entanglement among young males taken in the commercial harvest or seen in juvenile male roundups have been maintained since 1967. Concern about the potential role of entanglement-caused mortality has given rise to research focused on determining as clearly as possible the extent to which entanglement contributes to a reduction in survival and to declining trends in the population (Swartzman 1984; Fowler 1985, 1987; French and Reed 1990; Swartzman et al. 1990).

This paper reports on recent field work to assess the effects of entanglement on the population of northern fur seals breeding on St. Paul Island, Alaska. The objectives of this work are: (1) continued monitoring of the proportion of seals entangled, (2) determination of the nature of entangling debris, (3) determination of the mortality caused by trawl webbing, especially as related to effects at the population level, and (4) assessment of the relative rates at which entangled and control animals are resighted. Part of the study of relative rates of resighting addresses the question of whether or not an animal's chances of being seen again are altered by being, or having been, entangled.

METHODS

Most of the data treated in this study deal with young male fur seals of the size (roughly 105 to 125 cm in total length) formerly taken in the commercial harvest on St. Paul Island. The commercial take of fur seals, which ended in 1984, was the earliest source of data on entanglement. Other data, as the main focus of this paper, were collected during 1985, 1986, and 1988 from animals of the same size (and same approximate age) to ensure comparability with historical data. Males of this size are usually between the ages of 2 and 5 years, mostly 3-year-olds.

The studies reported here involved roundups, a procedure conducted during the breeding season. A total of 63 roundups were conducted in July and early August 1985. Sixty-one were conducted in July and early August 1986; 66 were completed during July 1988.

During roundups, young males are herded together to be examined for debris or tags and for applying tags. To conduct a roundup, field biologists approach an area (called a hauling ground) near a breeding

rookery where young males come ashore in large numbers. Avoiding disturbance to the rookeries, the members of the research team position themselves between the hauling ground and the water. The males on the hauling ground are then surrounded and herded away from the rookery but close to the water's edge. Care is taken to minimize the movement required of the animals and to allow them sufficient space to prevent crowding and overheating.

Once the seals are in a controlled group, field workers then allow small numbers of animals to leave the group and file toward the water. Once one or more seals begin moving toward the water, other seals follow. This movement is controlled (to ensure that tagged flippers will be seen) by the field crew. While moving toward the water, seals pass between observers, some of whom are engaged in counting seals while others watch for tags and entangling debris. Others of the field crew remain prepared to capture seals, while the remainder work to assure that the main group of seals remains in place.

When an entangled or tagged seal is seen among those leaving, the movement of seals from the main group is stopped. If tag numbers cannot be read, if tags are to be applied, or if a detailed examination of the debris is required, the seal is captured with a wooden pole fitted with a rope noose (<2% of these seals escape to the water without being captured). If tags are to be applied, or the debris examined in detail, the seal is placed on a restraint board (Gentry and Holt 1982) for a few minutes. Tags are applied on the trailing edge of each foreflipper, about 2-3 cm distal from the hairline.

If the captured animal is entangled, the nature of the entanglement is recorded (and tags applied if not previously tagged). Data recorded at the time of tagging include the tag number; the color, size, and type of debris; mesh size (if it is a net fragment); and the extent of the wound the debris has caused. A sample of the debris is removed (if there is enough) to be used later for measuring twine size and for any analysis necessary for identification of the plastics involved.

Two control seals about the same size as the entangled animal are also tagged to compare rates of return in succeeding years. The choice of tagging two control seals is arbitrary. Tagging more controls than entangled seals ensures a larger sample of returns to be used in comparing the relative rates of return of the two groups. It also aids in the study of the frequency of resighting rates and the locations (for study of intermixture) of resighted seals.

In most cases, seals that are not handled and seals released after being tagged or examined return directly to the water. By the end of the roundup, all seals have returned to the water.

Some of the animals seen in the first roundup are seen again in later roundups. The resulting sampling scheme is one of sampling with replacement, and the data for both the control animals and the entangled animals are treated accordingly.

Other sets of the data reported in this paper are from similar studies prior to 1985 in which animals were sighted in the commercial harvest prior to 1985. During the harvests, animals were herded together and moved to special areas where they were killed. These data from harvests, therefore, are treated as samples without replacement.

In previous studies of fur seal entanglement, two approaches have been used to categorize debris on seals according to its size (weight). For continuity and comparison, both are used in this study with distinction depending on the terms used. The first approach divides the debris into "light" or "heavy" categories depending on whether it is light enough for the entangled seals to return (at least once) to the breeding islands or so heavy that they cannot return. This definition suffers from lack of precision because the two categories are not discrete; their overlap is dependent on factors such as how far the seal has to swim to haul out on land. The upper limit of the light category is about 400 g, since over 90% of the entangled seals observed on land are in debris that weighs <400 g (Fowler 1987).

The second approach uses three distinct weight categories. The debris seen on animals is either weighed (after being removed) or subjectively evaluated (when entangled animals are released with debris intact). The weight of debris is classified as small (<150 g), medium (between 150 and 500 g), and large (>500 g).

To study the behavior of entangled animals, and the influence of entanglement on the chances of being resighted, radio transmitters (weighing about 40 g) were attached to 16 control and 16 entangled animals to monitor their presence and absence in the vicinity of the hauling grounds or rookeries. A radio transmitter was attached with epoxy glue to the back of the animal's head while the animal was restrained following procedures described in Loughlin et al. (1987). Each radio-tagged seal was also marked with bright paint applied to the radio and glue. Each radio was a 3.5-V transmitter, manufactured by Advanced Telemetry Systems, Inc. All radios transmitted within the frequency range of 164 to 166 MHz.

Data on the behavioral effects of entanglement were all collected in 1988. After attaching radios early (17 to 26 July), observers, using hand-held receivers, listened for radio-tagged seals during a daily visit to each haulout site until 29 August. A computer attached to a receiver was set up at the southern end of St. Paul Island (Reef Point) to scan for and record radio signals from each of the radio-tagged animals within receiving distance (approximately 5 km).

The amount of time the seals spent on shore was estimated in two ways. Detailed data for seven animals (three control and four entangled) were available from the computer at Reef Point. The computer scanned for the presence of these animals for 10 sec every 15 min, 24 h/day. We estimated the duration of intervals spent on land or at sea to the nearest quarter hour. Because the signals occasionally were blocked by the animals lying on the transmitters, and because the animals frequently entered the near-shore water without going to sea, we considered an animal to be at sea only

when its transmitter had not been heard for at least an hour. Hence, by this definition, trips to sea could never be shorter than 1 h.

The second method for estimating the time ashore involved the use of data obtained from observers with hand-held receivers. If the radio on a given animal was heard during a survey, the seal was considered to have been on land all day. If the signal from that radio was not heard, the seal was considered to have been at sea all day. When the signal from a given animal was heard one day but not on the next day, we assumed that the animal had departed halfway between the two observations. This gave us an onshore estimate to the nearest half day for all 32 animals.

Standard methods were employed in conducting the usual statistical tests (e.g., chi-square tests) where noted. The level of significance chosen for statistical tests was $P = 0.05$, unless otherwise noted. The analysis of data resulting from the resighting of tagged animals involved both standard approaches (e.g., the Seber-Jolly method; Seber 1973) and a regression analysis specifically designed for this study. The latter was developed to make use of all the existing data to address questions unique to this study. The specifics of the procedure used in this analysis, with the assumptions involved in estimating survival from entanglement-caused mortality, are presented in the Appendix.

RESULTS

During 1985, 1986, and 1988, 22,211, 22,572, and 24,519 (respectively) male seals of the size conventionally taken in the harvest were sampled. As will be presented in more detail below, about 25% of these totals were repeated sightings. Table 1 shows the numbers of seals that were tagged each year and percentage resighted in subsequent years.

Of the 49 tagged animals released in 1985 and resighted in 1986, 12 (24%) were originally tagged as entangled animals. The change from a ratio of 85:172 ($85/172 = 0.494$, entangled to controls) tagged in 1985 to 12:37 ($12/37 = 0.324$) resighted in 1986 is not statistically significant (chi-square test). There was no field effort in 1987, so no samples were collected in that year. Of the 14 seals tagged in 1985 and resighted in 1988, 1 (7.7%) had been tagged as entangled. The change in ratio from 85:172 ($85/172 = 0.494$) to 1:13 ($1/13 = 0.077$) between 1985 and 1988, and from 12:37 to 1:13 ($12/37 = 0.324$ to $1/13 = 0.077$) between 1986 and 1988 are statistically significant (binomial probability tests).

Of the 407 animals tagged in 1986, 128 (31.4%) were entangled. Of 46 seals tagged in 1986 and resighted in 1988, 6 (13%) were tagged as entangled seals in 1986. The change from a ratio of 128:279 ($128/279 = 0.459$) to 6:40 ($6/40 = 0.150$) between 1986 and 1988 is also statistically significant (chi-square test).

Of the eight seals resighted in 1988 after having been tagged as entangled in earlier years (including one tagged prior to 1985), six had lost their entangling debris. No seals have been resighted as entangled after originally having been tagged as controls.

Table 1.--Comparison of numbers of tags applied (in parentheses) and resighted (percent resighted shown in brackets below the numbers resighted) by year for entangled and nonentangled seals, each row corresponding to the tags released in the first year for that row (from Fowler et al. 1989).

Controls	Year			
	1985	1986	1987	1988
Nonentangled	(172)	37 [21.5]	-- --	13 [7.6]
		(279)	-- --	40 [14.3]
			-- --	-- --
				(104)
Entangled	(85)	12 [14.1]	-- --	1 [1.2]
		(128)	-- --	6 [4.7]
			-- --	-- --
				(52)

Table 2 presents the percentage of juvenile male seals found entangled, by year, for 1981 to 1988 in terms of the kinds of debris in which they were entangled. More detailed presentations of the data for 1988 are available in Fowler et al. (1989). Figure 1 illustrates the percentage of entangled seals observed in the harvests since 1967 and in the roundups since 1985. Table 2 also shows the composition of the debris found on animals in terms of proportions entangled. The proportion entangled in 1988 was the lowest observed since 1970 and was about half of the mean proportion observed from 1981 to 1986.

The frequency distribution of the size of debris seen on the animals per year is shown in Table 3. The numbers and percentages of those animals resighted in subsequent years, in relation to the size of debris, are presented in Table 4. None of the seals entangled in large pieces of trawl webbing were resighted more than 1 year subsequent to their being tagged,

Table 2.--Debris found on juvenile male fur seals in 1988 compared to 6 earlier years, expressed as the observed percent of juvenile male seals entangled by debris category.

Type of debris	Entanglement (%)						
	1981	1982	1983	1984	1985	1986	1988
Trawl net fragments	0.29	0.24	0.30	0.22	0.36	0.27	0.15
Monofilament net fragments	0.00	0.01	0.01	0.02	0.01	0.01	0.00
Plastic packing bands	0.08	0.10	0.07	0.09	0.05	0.06	0.07
Cord, rope, string	0.04	0.04	0.02	0.05	0.08	0.07	0.05
Miscellaneous items	0.03	0.01	0.03	0.01	0.01	0.01	0.01
Total	0.43	0.41	0.43	0.39	0.51	0.42	0.28
Sample size	102	102	112	87	76	70	53

Table 3.--Annual percentage frequency distribution of the size of debris on entangled seals that were tagged and released.

Year	n	<150 g(%)	150-500 g(%)	>500 g(%)
1983	84	63	23	14
1984	57	81	12	7
1985	78	72	20	8
1986	128	72	21	7
1988	53	72	15	13
Total	400	71	19	10

whereas seals in small debris were resighted up to 5 years later. The resighting rate of animals in medium-size debris was intermediate to those for large and small debris.

A summary of the results of the radio tagging study using hand-held radio receivers is presented in Table 5. The table contains data from both full and partial records because the study was of insufficient length to encompass an entire long feeding trip for all of the tagged seals. Furthermore, almost no seal completed a full cycle, from departure on a trip to sea followed by a return and an on-land interval until departure for the next feeding trip. For that reason, the estimated percentages of time spent on land during the course of this study may be different from those over an entire season spanning several full cycles. However, the entangled seals spent more time at sea than did controls. Twelve of sixteen entangled




Figure 1.--The percentage of juvenile male seals found entangled in the commercial harvest from 1967 to 1984 and in research roundups from 1985 to 1988, on St. Paul Island, Alaska.

Table 4.--The numbers and percentages of tagged animals listed in Table 3 that were resighted by year in relation to size of entangling debris and year.

Year tagged	Year resighted	Size of debris		
		<150 g(%)	150-500 g(%)	>500 g(%)
1983	1984	18(34)	3(16)	2(17)
1983	1985	4(8)	1(5)	0(0)
1983	1986	3(6)	0(0)	0(0)
1983	1988	1(0)	0(0)	0(0)
1984	1985	14(30)	2(29)	0(0)
1984	1986	9(16)	0(0)	0(0)
1984	1988	0(0)	0(0)	0(0)
1985	1986	9(16)	3(19)	0(0)
1985	1988	1(2)	0(0)	0(0)
1986	1988	6(7)	0(0)	0(0)
Combined years		65(23)	9(12)	2(5)

Table 5.--Comparison of the percent of time spent on land (present) and at sea (absent) for entangled and control seals fitted with radio tags. Data are from daily surveys with hand-held receivers on all hauling areas on St. Paul Island.

Seals ^a		Percent of time	
		Present	Absent
Entangled-fr	(N = 4)	35	65
Controls-fr	(N = 13)	28	72
Entangled-pr	(N = 12)	13	87
Control-pr	(N = 3)	10	90
Entangled-t	(N = 16)	19	81
Control-t	(N = 16)	25	75

^afr = males with full records, pr = males with partial records, and t = all males combined.

seals had not returned to land by the end of the study, whereas only 3 of 16 control seals had not returned (chi-square test, $P < 0.005$, or 0.001 with continuity correction). Typically, both entangled and control seals made several short trips while in the vicinity of St. Paul, and then departed on one long feeding trip. Selecting this longest trip to sea for each seal, we found that the entangled seals had significantly longer trips (30.9 days) than did controls (24.3 days). For seals that did not return from their long trips, the time from departure until the end of the study was used. Therefore, these were actually minimum estimates of their trip lengths.

The hand-held receivers could not detect the short trips taken between daily scans. Thus, the proportion of time on land (Table 5) actually estimates the time when the seals were in the vicinity of St. Paul, but not necessarily ashore. However, the data collection computer, which was able to detect short trips for seven seals, provided estimates of the time actually spent ashore at Reef Rookery. These data indicated that the four entangled animals spent a smaller proportion (44.8%) of their visit to St. Paul on land than did the controls (55.3%), but the difference is not statistically significant. The mean time between the application of tags and the departure to sea for a long feeding trip for entangled animals was 7.59 days; that for the controls was 6.17 days (no significant difference).

In 1988, 7 of 16 entangled seals fitted with transmitters were resighted in subsequent roundups. Four of sixteen controls with transmitters were resighted. There is no significant difference between these rates of resighting (chi-square test).

Analyses of the data in Table 1 are possible through the application of two very similar methods described in Brownie et al. (1978) and the Seber-Jolly method (Seber 1973). These methods result in estimates of survival of both categories of seals (entangled and controls). The annual survival of entangled seals estimated by these two methods (the same for each) is 0.22 (0.95 confidence limits of ± 1.00 , assuming a Poisson distribution for the resightings), and 0.51 (0.95 confidence limits of ± 0.446) for controls. Although not statistically significant, the estimated survival for the entangled animals given by these results is 42% that of the controls. The estimated survival for the controls (0.51) is lower than the estimates of survival produced by Lander (1981) for juvenile males (about 0.8, including the effects of unobserved entanglement), but the difference is within the confidence limits shown above.

We also used the data in Table 1 in a regression analysis to estimate the ratio of the probabilities of being resighted for entangled and control animals and the survival factor associated with entanglement in light debris. The basis of the regression analysis is demonstrated in Figure 2, which shows the declining rate at which entangled animals were resighted relative to the controls. Each data point is corrected for the ratio of entangled to nonentangled animals, as shown in Table 6.




Figure 2.--Relative rates of return for entangled juvenile male fur seals compared to controls (nonentangled tagged seals) for varying time intervals. The relative rate of return is $F^*(C/D) = p_{i,k}(N_{c,i}/N_{e,i})$ and the time interval is $x = (k-i)$, from Table 6 and Appendix. The point at time zero, with an adjusted ratio of entangled to control animals of one, was not used in the regression analysis.

Table 6.--List of data as extracted from Table 1 for regression analysis to estimate entanglement related survival; for a linear model of $y = a + bx$. See Appendix for details.

A Year i	B Year k	C $N_{c,i}$	D $N_{e,i}$	E $\ln(N_{c,i}/N_{e,i})$	F $p_{i,k}$	G $\ln(p_{i,k})$	y E+G	x B-A
1985	1986	172	85	0.7048	0.3243	-1.1260	-0.42	1
1986	1988	279	128	0.7792	0.1500	-1.8971	-1.12	2
1985	1988	172	85	0.7048	0.0769	-2.5649	-1.86	3

The results of the regression analysis, with the assumptions involved in estimating survival from entanglement-caused mortality, are presented in the Appendix. The estimated annual survival of seals entangled in light debris is about half (0.49) that of nonentangled seals. The probability of resighting an entangled seal was estimated to be about 1.35 times as great as the probability of resighting a control (given that they are both alive). However, this estimate is not significantly different from 1.0 (the case where the probabilities of seeing a seal from either group are the same).

It should be made clear that the total annual survival among entangled animals (including the effects of other sources of mortality along with those due to entanglement) is the product of natural survival and survival from entanglement. If we use the survival for juvenile males from Lander (1981)--about 0.8--the overall survival for seals entangled in light debris would be about 0.4 (i.e., about $0.8 \times 0.5 = 0.4$ for 3-year-old males). This is a higher survival rate than that from the Seber-Jolly analyses presented above (0.22).

Table 7 contains data on the frequency of resighting tagged seals during the season when tags were applied. These data show that the fraction of resighted control animals is nearly the same as the fraction of resighted entangled animals (both being about 25%). No statistically significant differences were found between the rates of resighting for entangled and control animals for any year or for the total (chi-square tests).

DISCUSSION

Although there is insufficient information to draw conclusions, the data collected in 1988 on St. Paul Island suggest a decline in the proportion of juvenile male northern fur seals that are entangled. Most of the change seems to be associated with a reduction in entanglement in trawl webbing, possibly a reflection of reduced occurrence of trawl webbing among pelagic debris as reported in 1988 by Japanese scientists (Fowler et al. 1989). The proportion of seals entangled in other forms of debris seems to be about equal to the proportion observed in the past 7 years. The differences between 1988 and previous years may be a result of changes in the

Table 7.--Comparison of numbers of tags applied to entangled and control juvenile male fur seals in 1985, 1986, and 1988 with the numbers in each category resighted the same season. The numbers in parentheses are the percent of the tags applied that were resighted.

Year	Number of tags			
	Controls		Entangled	
	Applied	Resighted	Applied	Resighted
1985	170	35(20.6)	76	21(27.6)
1986	165	54(32.7)	70	19(27.1)
1988	104	21(20.2)	52	15(28.8)
Total	439	110(25.1)	198	55(27.8)

rate of loss and discard of net fragments. Various education programs at national and international levels have been in place for several years, and international regulations prohibit the discard of such debris.

Severe wounds caused by prolonged entanglement in light debris contribute to death. Bengtson et al. (1988), demonstrated that pups become entangled in net fragments with mesh sizes much smaller than those seen on the subadult males in the roundups. The subsequent growth of those seals caught in debris light enough for them to survive the effects of drag in the water then results in wounds and death. Seals remaining entangled in debris often suffer from wounds that increase in size as a result of the seals' growth (DeLong et al. 1990). The degree to which wounds and resulting infections contribute to mortality in comparison to other sources of mortality caused or accentuated by entanglement (such as starvation, strangulation, and predisposition to predation) cannot be determined from existing data.

Some seals survive because they escape from the debris. Escape has been reported for animals resighted in other studies (Scordino 1985; Fowler 1987), some within the season during which animals were tagged. Of the total of eight seals resighted in 1988 after having been tagged as entangled in earlier years, six had lost their entangling debris. How this affects estimates of survival of seals in light debris has not been determined; conceivably, individuals that have lost their debris would be resighted with the same probability as control animals.

All debris on entangled animals that was later lost had been judged to weigh <150 g at its first sighting; otherwise it was similar to commonly observed debris. One possible explanation for this pattern is that the animals in small debris are the most likely of the entangled animals to

return to the breeding islands. There they can come into contact with substrates (such as rocks) where the debris can abrade or otherwise wear to the point of breaking and falling off. Such wear is noted on the debris on many of the seals seen in the roundups, and on a few occasions debris has broken and fallen off during the handling of entangled animals. In view of the small numbers of animals resighted as entangled and the low survival of entangled animals, it would appear that most animals that remain entangled eventually die as a result of the debris.

The relative rate of resighting of animals originally tagged as entangled varies with the size of debris. A statistically significant (chi-square) decline in the rate at which seals are resighted with increasing size of debris is seen in Table 4. Corresponding information reported by DeLong et al. (1990) shows that of 17 females experimentally entangled in 200-g fragments of trawl net, 2 (12%) returned to the same rookery to give birth 1 year later. This is equal to the 12% resight rate of the seals entangled in medium-sized debris (Table 4). Thus, factors such as exhaustion, starvation, and drowning (likely acute factors at sea) appear to be increasingly important in the causes of death due to entanglement as debris size increases. If the survival of seals in large debris is proportional to the rate at which they are resighted, the survival of those in debris weighing just over 500 g would be about one-fourth (5/22) the survival of those in small debris. Therefore, survival resulting from the effects of entanglement alone would be about 0.11 ($(5/22) \times 0.49 = 0.11$; using the 0.49 from the Appendix). Assuming survival from natural causes is 0.8 (Lander (1981), whose results may include some mortality due to entanglement), the total survival for this large-debris group is calculated as 0.09 ($0.8 \times 0.11 = 0.088$). This implies a turnover in the population of about 2.4 times per year (turnover meaning the number of entangled seals that die for every entangled seal occurring in the population, and being equivalent to the instantaneous mortality rate, or the negative natural log of survival; $-\ln 0.09 = 2.42$). Presumably, following the trend in Table 4 to even larger debris, the turnover rate continues to increase with the size of entangling debris. If the estimated survival for controls from the Seber-Jolly analyses presented above were used (0.51 in place of 0.8), this estimated turnover would be even more rapid.

Seals that are entangled in large debris may find it impossible to return to land. Seals are seen entangled at sea in debris that is clearly large enough to prevent their returning to land (Fowler 1987). This is important in interpreting the information in Table 4. The number of seals entangled in large debris resighted on land may be small not because the seals thus entangled have died soon after entanglement, but because the debris prevents them from returning to the islands to haul out. This effect would be greater with increasing size of entangling debris. Such a trend would affect estimates of entanglement-related mortality. However, failure to return has the same effect on the population as mortality; an animal that does not return to its breeding colony is removed from the reproductive population.

Whether or not a seal is entangled may affect its chances of being seen in roundups. This is important in estimating the proportion of seals

entangled and their survival rates. Factors that may affect estimates include: 1) time spent on land and at sea, 2) entangling debris or scars attracting the attention of observers, 3) relative proportions of the two groups which remain at sea for the entire season, a factor about which nothing is known, and 4) probability of seeing seals that have lost entangling debris compared with the probability of seeing entangled seals.

Entanglement results in prolonged at-sea portions of the feeding cycles for northern fur seals. Previous work on radio-tagged entangled male seals showed that the pelagic phase of feeding cycles was about twice as long for entangled seals as for controls (Bengtson et al. 1989). The results of this study are consistent with this effect of entanglement. Similar results have been noted for females (DeLong et al. 1990). It has not been possible to produce accurate estimates of the effects of entanglement on the portion of time spent on land. As a consequence, the relative time spent on land (as a fraction of the complete feeding cycle) remains undetermined. Thus, it is not possible, with the data from radio tagging, to quantify the effect of altered feeding cycles on the chances of a seal being seen.

Other data concerning the probability of resighting a seal are inconclusive. Based on data in Table 7, it would appear that once seals return to the islands, entanglement does not significantly affect their chances of being seen at least twice. Such a comparison can also be made with the smaller sample of radio-tagged animals (these seals being more visible with the bright paint). In 1988, no significant difference was found in the rates of resighting entangled and control seals fitted with transmitters in subsequent roundups.

Based on conventional mark-recapture analyses and results presented in the Appendix, seals entangled in light debris experience an annual survival that is about half (0.41 to 0.49) that for control seals. Previous estimates are very similar (0.42, Fowler 1987; 0.46, Fowler 1985.)

Regardless of a seal's probability of being resighted, it is obvious that entangled seals suffer higher mortality than do controls (Fig. 2). We have considered whether the reduced relative rates of resighting between initial release and the first resighting (e.g., the change between the first two points in Fig. 2) could have been due only to differences between the probabilities of seeing entangled or control seals. Both groups would have experienced similar survival, and the change would have been entirely due to a higher probability of seeing control animals. If this were the case there would be no further changes in the ratios over time. A level relationship would emerge between the points for years 1-3, all of which would be lower than the ratio at year 0, the time of release. The continued decline is indicative of the predominate effect of lowered survival among entangled seals.

Combined with other factors, the mortality caused by entanglement in light debris lowers the total survival for juvenile males entangled in light debris to about 0.39, assuming independence of the causes of mortality and a natural survival of 0.8. Each year, then, the number of

seals in light debris that die would be about the same as the number of seals in light debris that are estimated to be alive in the population at the time of sampling (94% as many, based on a turnover of 0.94 from $-\ln(0.39) = 0.94$, as the instantaneous rate of mortality).

A great deal of progress has been made in understanding the extent and effects of entanglement in marine debris on northern fur seals. However, precise estimates of the contribution of entanglement to the survival and trends at the population level have yet to be produced. Several studies indicate that young fur seals are more likely to become entangled than larger seals. Pups can become entangled even before leaving land (DeLong et al. 1988). Pups have been observed entangled or becoming entangled in large fragments of debris. Groups of pups often become entangled together or in succession (Fowler 1987; DeLong et al. 1988). Experiments show that pups are susceptible to entanglement in about four times as much debris as older animals because they can pass their heads through net fragments of smaller mesh size (Fowler 1987; Bengtson et al. 1988). A greater proportion of entangled animals among the young (also less experienced) seals is also consistent with the view that immature seals are more curious than older seals and are, therefore, more likely to be attracted to debris in which they may become entangled.

Research continues to show that mortality rates are quite high for seals that become entangled in larger debris. The results of the studies reported here indicate an annual survival (from the effects of entanglement) of about 0.09 for seals in debris weighing just over 500 g. Combined with the potential that larger net fragments have a higher probability of attracting seals and the fact that seals have been observed entangled in groups in large debris (Fowler 1985; DeLong et al. 1988), entanglement in large debris obviously deserves attention. However, logistic and financial constraints have made such studies impossible.

The need for studies to examine this problem is emphasized by the implications of previous attempts to account for the effects of large debris. Trawl webbing accounts for about two-thirds of the light debris (Table 2), so the portion of the juvenile male population entangled in light pieces of trawl webbing has been (before 1988) about 0.003 ($0.66 \times 0.004 = 0.00264$; 0.004 being the proportion entangled in light debris of all kinds). On beaches, at sea, and on entangled animals seen away from the breeding colonies, the frequency of occurrence of pieces of heavy trawl webbing is about five times that of light (Fowler 1987). Assuming, that for every piece of light debris on an entangled seal there are five pieces of heavy debris also entangling seals, then entanglement in heavy debris involves about 1.5% of the juvenile male population ($0.003 \times 5 = 0.015$).

As mentioned above, pups during their first few months at sea may be four times more susceptible to entanglement than juvenile males. If so, 6% of their numbers become entangled each year ($4 \times 0.015 = 0.06$). Accounting for the turnover from mortality of seals in large debris (estimated earlier as 2.4 times per year for debris just over 500 g) produces the implication of an entanglement-caused mortality of over 14% ($2.4 \times 0.06 = 0.14$). This does not account for the mortality in light debris. Entanglement in

heavier debris has been observed to involve more than one animal per piece (Fowler 1985, 1988; DeLong et al. 1988). This, combined with the greater attraction large debris must have for seals (the larger pieces presumably being more easily seen because of their size), could result in higher rates of entanglement and mortality.

If, as indicated by field observations (Fowler 1988; DeLong et al. 1990), entanglement involves both sexes (especially among the younger age classes), entanglement and resultant mortality may have contributed significantly to the declining trends among fur seal populations (Fowler 1985, 1988). Such implications are consistent with recently observed population trends, and models consistent with such trends have been constructed (Swartzman 1984; French and Reed 1990; Swartzman et al. 1990). These observations emphasize the need for better studies to clarify our estimates of the degree to which entanglement has caused these trends. Feasible field studies to verify the role of entanglement in large debris have yet to be designed and conducted.

CONCLUSIONS

The 1988 results of field research on entanglement of northern fur seals through roundups of juvenile males on St. Paul Island, Alaska, showed:

1. A reduction of the proportions observed entangled on land from about 0.4 to 0.29%.
2. Entanglement in fragments of trawl webbing in 1988 was about half of entanglement levels observed for this kind of debris in previous years.
3. The rate of resighting for animals tagged in 1985 and 1986 and resighted in 1988 showed that entangled animals tagged in those years were seen at rates that were significantly less than the rate at which controls were resighted.
4. The pelagic portion of the feeding cycle of entangled seals is greater (and a larger portion of their time may be spent at sea) than that of control seals, but the extent of the difference is unknown.

Analysis of these data in combination with data from previous studies (data on resighted animals collected in 1986, also on St. Paul Island, and data on debris collected from 1967 through 1988) showed that:

5. The estimated survival due to being entangled in light debris ranged from 0.41 to 0.49, close to estimates of about 0.5 or less from previous work.
6. Combined with natural survival, the total survival of entangled seals is probably <0.39 , with the equivalent of nearly a complete turnover in the population of juvenile males entangled in light debris each year.

7. Mortality increases with the size of entangling debris based on the observation that survival for seals entangled in large debris is less than for those in small debris.
8. The probability of resighting entangled seals (or seals that once were entangled), compared to that of nonentangled seals, has yet to be clearly evaluated.
9. A great deal has been learned about the specifics of mortality caused by entanglement in debris weighing <500 g. Implications for effects at the general population level are serious. However, the main result of this progress is a continuing emphasis on the need to refine estimated mortality rates caused by large debris, especially pieces much larger than 500 g.
10. More studies will be required to better understand the interacting factors associated with the probability of entangled seals being resighted in the roundups.

ACKNOWLEDGMENTS

We thank all of the many individuals who have worked in the field crews conducting roundups on St. Paul Island; without their help, the data for this paper could not have been produced. Funding for the studies reported here came from the Entanglement Research Program at the Northwest and Alaska Fisheries Center, and we extend our thanks to James M. Coe, director of that program. We thank Kara Amundson, Howard Braham, Jeff Breiwick, Jean Davis, Gary Duker, Francis Fay, Sharon Giese, Mary Lynne Godfrey, Dan Kimura, Lloyd Lowry, Marcia Muto, Gary Stauffer, Gordon Swartzman, Steve Syrjala, and Anne York for their reviews, comments, and suggestions.

REFERENCES

- Bengtson, J. L., C. W. Fowler, H. Kajimura, R. Merrick, K. Yoshida, and S. Nomura.
 1988. Fur seal entanglement studies: Juvenile males and newly-weaned pups, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.
- Bengtson, J. L., B. S. Stewart, L. M. Ferm, and R. L. DeLong.
 1989. The influence of entanglement in marine debris on the diving behavior of subadult male northern fur seals. In H. Kajimura (editor), Fur seal investigations, 1986, p. 48-56. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-174.
- Brownie, C., D. R. Anderson, K. P. Burnham, and D. S. Robson.
 1978. Statistical inference from band recovery data--a handbook. U.S. Fish Wildl. Serv., Resour. Publ. 131, 212 p.

DeLong, R. L., P. Dawson, and P. J. Gearin.

1988. Incidence and impact of entanglement in netting debris on northern fur seal pups and adult females, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), *Fur seal investigations*, 1985, p. 58-68. U.S. Dep. Commer., NOAA Tech. Memo.-NMFS-F/NWC-146.

DeLong, R. L., P. J. Gearin, J. L. Bengtson, P. Dawson, and S. D. Feldkamp.

1990. Studies of the effects of entanglement on individual northern fur seals. [Abstract.] In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Fowler, C. W.

1982. Interactions of northern fur seals and commercial fisheries. *Trans. N. Am. Wildl. Nat. Resour. Conf.* 47:278-292.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.

1988. A review of seal and sea lion entanglement in marine fishing debris. In D. L. Alverson and J. A. June (editors), *Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris*, Kailua-Kona, Hawaii, 13-16 October 1987, p. 16-63. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Fowler, C. W., R. Merrick, and N. Baba.

1989. Entanglement studies, St. Paul Island, 1988; juvenile male roundups. Northwest and Alaska Fish. Cent., Natl. Mar. Mammal Lab., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115. NWAFC Processed Rep. 89-01, 23 p.

French, D., and M. Reed.

1990. Potential impact of entanglement in marine debris on the population dynamics of the northern fur seal, *Callorhinus ursinus*. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

- Gentry, R. L., and J. R. Holt.
1982. Equipment and techniques for handling northern fur seals.
U.S. Dep. Commer., NOAA Tech. Rep. NMFS-SSRF-758, 15 p. (Available
D822, User Serv. Br. Environ. Sci. Inf. Cent., NOAA, Rockville,
MD.)
- Lander, R. H.
1981. A life table and biomass estimate for Alaskan fur seals.
Fish. Res. (Amst.) 1:55-70.
- Loughlin, T. R., J. L. Bengtson, and R. L. Merrick.
1987. Characteristics of feeding trips of female northern fur seals.
Can. J. Zool. 65:2079-2084.
- Scordino, J.
1985. Studies on fur seal entanglement, 1981-84, St. Paul Island,
Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings
of the Workshop on the Fate and Impact of Marine Debris, 26-29
November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer.,
NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Seber, G. A. F.
1973. The estimation of animal abundance. Griffin, Lond., 506 p.
- Swartzman, G. L.
1984. Factors bearing on the present status and future of the
eastern Bering Sea fur seal population with special emphasis on the
effect of terminating the subadult male harvest on St. Paul Island.
U.S. Marine Mammal Commission Report No. MMC-83/03. [Available
from U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA
22161, as No. PB84-172329, 77 p.]
- Swartzman, G. L., C. A. Ribic, and C. P. Huang.
1990. Marine debris entanglement mortality and fur seal populations:
A modeling perspective. In R. S. Shomura and M. L. Godfrey
(editors), Proceedings of the Second International Conference on
Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep.
Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this
document.]

APPENDIX

ESTIMATION OF ENTANGLEMENT-RELATED SURVIVAL AND THE RELATIVE
PROBABILITIES OF ENTANGLED AND CONTROL FUR SEALS BEING RESIGHTED

To make use of the data on the returns of male fur seals (i.e., those resighted) as shown in Table 1, we make a set of assumptions and define the following terms. Let

- $N_{c,ik}$ = the number of control seals tagged in year i and resighted in year k , where $k > i$ ($i = 1985, 1986$, $k = 1986, 1988$).
- $N_{e,ik}$ = the number of seals tagged in year i as entangled animals and resighted in year k (regardless of whether or not they were entangled when resighted), where $k > i$ ($i = 1985, 1986$, $k = 1986, 1988$).
- $P_{i,k}$ = $N_{e,ik}/N_{c,ik}$, or the ratio of numbers of seals resighted in year k that were entangled when first tagged in year i to the numbers of nonentangled (control) seals tagged in year i and resighted in year k .
- $s_{c,j}$ = the annual survival of control animals, or the animals tagged without debris in year j , for j from i to k (i.e., $s_{c,j}$ = survival from j to $j+1$). This is the probability of surviving from natural causes of mortality.
- s_e = the annual survival of animals entangled in light debris (debris light enough to return to the breeding islands), and is assumed not to vary from year to year. This is the probability of surviving entanglement given that an animal has survived natural causes of mortality and is assumed to be independent of $s_{c,j}$ (so their total annual survival is $s_{c,j}s_e$).
- $N_{e,i}$ = the number of seals tagged as entangled animals in year i ($i = 1985, 1986$), and
- $N_{c,i}$ = the number of seals tagged as controls in year i ($i = 1985, 1986$).

Different proportions of entangled seals may return to the islands to be seen when compared to controls. Once in the vicinity of the islands, entangled seals may be seen at different rates than the controls for various reasons. These include the possibility of different fractions of time spent on land and entangled seals being seen more readily than controls because of their entanglement, or the effects of having been entangled. Thus we define

- f_{*k} = the probability of resighting a seal in year k given that it was entangled when tagged and that it is alive. This

probability is expressed on the basis of a unit of searching effort that is the same as applied in looking for control animals. It is assumed to vary from year to year but not in relation to f_{ck} (below), and

f_{ck} - the probability of resighting a control animal in year k given that it is alive in the population, again as based on the unit of effort spent in searching for both control and entangled seals. This is also assumed to vary from year to year but not in relation to f_{ek} (f_{ek}/f_{ck} is assumed constant).

With these terms, the expected number of seals that were entangled when tagged and sighted in year k after being tagged in year i , for one unit of effort is

$$E(N_{e,ik} | N_{e,i}) = f_{ek} R_k s_e^{(k-i)} N_{e,i}$$

($i = 1985, 1986$, $k = 1986, 1988$, and R_k is the product of $s_{c,j}$ for j from i to k), and the expected number of controls for the same circumstances is

$$E(N_{c,ik} | N_{c,i}) = f_{ck} R_k N_{c,i}$$

(R_k is the product of $s_{c,j}$ for j from i to k).

Substituting the observed for the expected values we have the following moment estimators:

$$N_{e,ik} = f_{ek} R_k s_e^{(k-i)} N_{e,i} \quad \text{and} \quad N_{c,ik} = f_{ck} R_k N_{c,i}$$

The ratio of these two equations, then, is

$$N_{e,ik}/N_{c,ik} = p_{i,k} = (f_{ek}/f_{ck})(N_{e,i}/N_{c,i})s_e^{(k-i)}$$

which can be used to estimate f_{ek}/f_{ck} and s_e .

We note that variability in natural survival (i.e., the survival of the controls and that part of the survival of entangled animals from natural effects) can occur over time and not affect the calculation since these terms cancel in the formulation of the equation above. We also note that the probability of resighting animals from each of the two groups can vary from year to year as long as their ratio remains the same, as assumed above. Effort spent in resighting entangled and control seals is the same (the same roundups) but the number of roundups can vary from year. This is because effort for each of the two groups influences the above relationships only as a ratio in f_{ek}/f_{ck} (i.e., it cancels and need not be defined).

By rearranging terms we have

$$p_{i,k}(N_{c,i}/N_{e,i}) = (f_{ek}/f_{ck})s_e^{(k-i)},$$

and taking the natural log of this equation results in the following linear equation which can be used for regression analysis and the estimation of relevant parameters as defined above:

$$\ln\{p_{i,k}(N_{c,i}/N_{e,i})\} = \ln(f_{ek}/f_{ck}) + \ln(s_e)(k-i).$$

Using this equation and the data from Table 6, the estimated parameters determined from regression analysis for the above equation are

$$\ln(f_{ek}/f_{ck}) = 0.307 \text{ and } \ln(s_e) = -0.720 \text{ (R}^2 = 1.00, p = 0.011).$$

These results imply that the ratio of the probabilities of being resighted is about 1.35 (calculated as $e^{0.307}$, with 95% confidence limits of 0.95 to 1.95). Thus, the chances of being resighted after being tagged as an entangled animal, given that the animal has survived, are estimated to be about 1.35 times that of being resighted as a control, but this does not differ significantly from 1 or an equal probability. The estimated survival of entangled animals from the effects of entanglement is 0.49 (calculated as $e^{-0.720}$ with 95% confidence limits of 0.41 to 0.57).

In addition to the small sample size, other factors prevalent in this analysis need noting. The data points for the 1- and 3-year time intervals are not independent. A random difference between the mean (here assumed constant) survival from entanglement in the first year will be seen as a bias in the same direction in the third. With this set of data, this does not affect the estimate of entanglement-related survival as much as it does the estimated ratio of probabilities of being resighted. This is because the slope of the line as seen in Figure 2 (the estimate of survival) depends more on a rotation about the point for the 2-year time interval than the distance the line is above or below the second point. The height of the line will be affected by the interdependence of the two end points.

The effect of assuming that the survival from risks caused by entanglement is independent from surviving the risks of other, natural, causes has not been explored. The same holds for the assumption that the ratio of the probabilities of being resighted for the two categories remains the same over time. However the various steps in the derivation of the linear equation used in this analysis might contain hidden assumptions, or sources of statistical error, have yet to be examined.

**A STUDY OF THE EFFECTS OF COMMERCIAL FISHING
DEBRIS ON *CALLORHINUS URSINUS* FROM
BREEDING ISLANDS IN THE WESTERN PACIFIC**

A. E. Kuzin
Pacific Research Institute of Fisheries
and Oceanography (TINRO)
4, Shevchenko Alley, Vladivostok, U.S.S.R.

ABSTRACT

In this paper, data and analyses are presented concerning the incidence of entanglement among the northern fur seal, *Callorhinus ursinus*, from the breeding islands of the western Pacific. This work was undertaken to further explore the degree to which waste disposed from fishing vessels is a source of mortality for this species. Based on the available data, estimates of the minimum proportion of various age and sex groups entangled within the population are produced. Historical data show that injuries caused by fishing nets shreds (66%), ropes (20%), fishing line (8%), and packing bands and collars made of other materials (6%) are contributing to the mortality of northern fur seals. The incidence of entanglement, and therefore the resulting mortality among the Tyuleniy (Robben) Island population, is higher than for the population on the Komandorskie (Commander) Islands. The higher incidence of entanglement on Robben Island may be related to declines in the population on that island in comparison to the relative stability on the Commander Islands, where the incidence of entanglement is less.

BACKGROUND

The intensity of exploitation of living marine resources increases annually (Moiseev 1979). The number of fishing fleets increases correspondingly (Yerukhimovich and Yefremenko 1985). Increasing intensity in fishing brings with it various negative effects on the environment. One of the aspects of this influence is the pollution of the ocean with scraps of fishing gear, packing materials, and other waste from the commercial fisheries, directly threatening many marine organisms. Fish, seabirds, reptiles, and marine mammals get entangled in such materials and die (DeGange and Newby 1980; Fowler 1982, 1985, 1987; Kuzin 1985; Shomura and Yoshida 1985).

A determination of the extent of the negative influence of commercial fishing waste and pollution in the ocean is not possible for the majority of marine organisms. But the continuous and systematic observation of

northern fur seals, returning each year to their hauling grounds (near the breeding colonies) after wintering in the ocean, provides the opportunity to determine the number of individuals entangled in marine debris. The resulting data allow for the calculation of estimates of the extent of damage caused by marine debris resulting from loss and discard by commercial fisheries as well as other human activities.

From the historical data, the kinds of debris found on entangled seals is known. Of the total number of entangled seals on Robben Island, 66% are entangled in fragments of fishing nets. Another 20% are found entangled in ropes, 8% in fishing line, and 6% in packing bands and collars made of other materials of commercial fishing waste (Kuzin 1985). Fowler's data (1982) showed that in the Pribilof Island population as many as 50,000 seals may die annually as a result of injuries caused by foreign objects. Over a life time, entanglement may cause the death of 15% of a cohort (Fowler et al. 1990). Similar estimates for other populations of fur seals, however, are unavailable.

In order to obtain estimates of entanglement-related mortality for populations of fur seals on both Robben and the Commander Islands, the statistical data presented annually to the Northern Pacific Fur Seal Commission regarding the incidence of entanglement among northern fur seals on hauling grounds are used.

METHODS

For estimating the effects of commercial fishing debris on fur seal populations, we have used the data from the commercial harvest of bachelor seals (2- to 5-year-old males, also referred to here as juvenile males). These data were collected by counting the total and the number of entangled male seals in the commercial harvest for each year from 1975 to 1986 on both Robben Island and the Commander Islands (Table 1). All entangled seals were of the same size category as the remaining seals taken in the harvest. These harvests were conducted during the months of June and July of each year, and all entangled seals were killed along with the other seals taken. Seals with scars only were not counted as entangled such that the counts included only male seals observed with entangling debris. Entangling debris found on the seals was then identified and frequencies were tabulated for each category of debris. Pieces of monofilament gillnet were included in the category of net shreds along with pieces of trawl net debris.

With the data on total numbers taken in the harvest and the counts of entangled seals among them, collected as described above, the incidence of entangled animals among the juvenile males (2 to 5 years old) was determined by dividing the number of entangled seals taken by the total in the harvest for each year and location.

For other age groups and sex groups of seals, data are not available from their primary concentrations on the breeding rookeries. However, data have been recorded for entangled individuals from these categories, as noted for seals found on hauling grounds. Thus, the percent of all entangled animals found on hauling grounds that fall into each category

Table 1.--Number of juvenile male fur seals taken and the incidence of seals entangled in debris in the commercial harvest by year and location (breeding islands in the western Pacific).

Year	Number harvested		Number entangled		Percent entangled	
	Robben	Commander	Robben	Commander	Robben	Commander
1975	2,500	1,730	27	30	1.08	1.73
1976	2,569	2,768	69	68	2.68	2.46
1977	4,069	2,766	69	66	1.69	2.39
1978	3,188	3,032	81	32	2.54	1.06
1979	2,933	2,524	33	13	1.13	0.52
1980	3,107	2,544	26	44	0.83	1.73
1981	3,613	5,117	113	35	3.12	0.76
1982	2,924	5,075	124	75	4.20	1.48
1983	2,582	5,717	24	34	0.92	0.59
1984	2,322	5,294	35	37	1.50	0.70
1985	459	5,097	4	47	0.87	0.92
1986	2,034	--	34	--	1.67	--
Total	32,300	41,664	639	481	1.97	1.15

can be determined. The categories used in this study are: mature males (older than 6 years), half-mature males (or "half bulls," 6 years of age), bachelors (younger than 6 years), females, and pups. Since exact ages are not known, the numbers in each age category are determined on the basis of experienced judgment.

To analyze the data resulting from the field work described above, a method for estimating lower bounds of the proportion of each age-sex class was developed. The following is an explanation of the procedure used.

As mentioned above, the empirical data are for a population consisting of several categories or age-sex groups. One category is represented by data for which the incidence of entanglement can be clearly determined (i.e., the bachelor males, which will be represented by subscript j). The other groups (e.g., pups, females) are those for which we wish to have estimates of the incidence of entanglement. These groups will be represented by subscript i .

To develop a procedure for estimating the proportion of seals entangled in each group, their total numbers in the population as a whole are defined as P_i . The total for the bachelor males is defined as P_j . The proportion entangled (or incidence of entanglement) is defined as C_i and C_j , respectively, for each of the two categories, so that the numbers entangled are $C_i P_i$ and $C_j P_j$. Of these, a proportion of each category is seen, a proportion defined as α (α_i and α_j , respectively) so that the actual numbers of entangled animals seen on land are $\alpha_i C_i P_i$ and $\alpha_j C_j P_j$ for

the two cases. It is assumed that the probability of seeing an entangled animal of any category is the same as the probability of seeing a non-entangled animal of the same category or that entanglement does not influence the probability of being seen. This may be summarized as follows:

<u>Category</u>	<u>Total population</u>	<u>Fraction entangled</u>	<u>Numbers entangled</u>	<u>Proportion seen</u>	<u>Entangled animals seen</u>
i	P_i	C_i	$C_i P_i$	α_i	$\alpha_i C_i P_i$
j	P_j	C_j	$C_j P_j$	α_j	$\alpha_j C_j P_j$

The desired estimate is of C_i , knowing C_j . This can be accomplished by dividing the number of entangled animals seen from category i ($\alpha_i C_i P_i$, for which there are data) by the number of animals seen in category j ($\alpha_j C_j P_j$, also for which there are data) and multiplying this ratio by $C_j(\alpha_j P_j / \alpha_i P_i)$:

$$(\alpha_i C_i P_i / \alpha_j C_j P_j) C_j (\alpha_j P_j / \alpha_i P_i) = C_i$$

For this equation to produce the correct estimate (disregarding the statistical aspects of the problem), then, it is seen that either values for $\alpha_i P_i$ and $\alpha_j P_j$ must be known (which they are not) or their ratio must be known. If this ratio were 1.0, the number of the two segments seen would be equal.

Based on observations on hauling grounds, the total number of seals in the bachelor category is always larger than the total for each of the other categories listed in Tables 2 and 3. (Here we refer to the total present, not the number of entangled seals seen.) Because of this, the equation above can be used by assuming $(\alpha_j P_j / \alpha_i P_i) > 1.0$ to determine lower bounds to the entanglement rates for the categories other than for bachelors. The value of the expression $(\alpha_j P_j / \alpha_i P_i)$ will always be greater than 1.0 since $\alpha_j P_j$ (the number of bachelor males seen on hauling grounds) is always greater than the number seen for other groups. Thus, the expression used to produce estimated lower bounds for the proportion entangled for groups other than bachelor males is:

$$C_i > N_i C_j / N_j$$

where:

C_i is the proportion of animals of the age-sex group in question that are entangled;

$N_i = \alpha_i C_i P_i$ is the number of entangled animals of the age-sex group in question as observed at the hauling grounds;

C_j is the proportion of bachelors that are entangled; and

$N_j = \alpha_j C_j P_j$ is the number of entangled bachelors observed at the hauling grounds.

Table 2.--Number and percent of entangled fur seals falling in various age and sex categories as observed at hauling grounds on Robben Island.

Year	Bulls		Half bulls		Bachelors		Females		Pups		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1974	4	1.5	10	3.8	115	43.4	136	51.3	--	--	265	100
1975	--	--	--	--	27	77.1	8	22.9	--	--	35	100
1976	2	2.2	1	1.1	69	71.2	21	22.6	--	--	93	100
1977	2	2.8	--	--	69	97.2	--	--	--	--	71	100
1978	53	18.0	47	15.9	58	19.7	135	49.9	1	0.3	294	100
1979	16	19.2	9	10.8	16	19.3	42	50.6	--	--	83	100
1980	1	0.8	4	3.5	102	88.7	8	6.9	--	--	115	100
1981	11	15.7	5	7.1	19	21.1	32	45.7	3	4.3	70	100
1982	4	3.2	5	4.3	65	52.4	50	40.3	--	--	124	100
1983	12	12.4	--	--	13	13.4	72	74.2	--	--	97	100
1984	16	18.4	5	5.7	18	20.7	47	54.0	1	1.1	87	100
1985	10	10.8	--	--	31	30.0	61	59.8	--	--	102	100
1986	5	6.8	2	2.4	19	22.9	53	63.9	4	4.8	83	100
1987	10	13.5	8	10.8	13	17.5	43	58.1	--	--	74	100
Total	146	9.2	96	6.0	634	39.8	708	44.4	9	0.6	1,593	100
Calculated minimum percent of entangled individuals:												
0.45			0.30		*1.97		2.19		0.03			

*From Table 1.

RESULTS

The numbers of seals taken in the commercial harvests on Robben Island and the Commander Islands, and the numbers of entangled seals among them are shown in Table 1 for 1975 through 1986. Also shown is the resulting incidence of entanglement expressed as a percent of the harvest. The number of entangled seals from the other categories, as observed on the hauling grounds, are shown in Table 2 for Robben Island, and Table 3 for the Commander Islands. Tables 2 and 3 also show the fraction of the total number of entangled animals observed as accounted for by seals in each category. Thus, the totals of the categories are each 100% for all observed entanglement, by year and island.

The results of calculations to determine the lower bounds to estimates of the percent entangled among each age-sex group of northern fur seals are presented in the last lines of Tables 2 and 3. The proportion of the total population which is entangled depends on the fraction of the population comprised by each of the age-sex categories. However, it can be seen that the lower bound for the overall entanglement rate must be between 0.03 and

Table 3.--Number and percent of entangled fur seals falling in various age and sex categories as observed at hauling grounds on the Commander Island.

Year	Bulls		Half bulls		Bachelors		Females		Pups		Total	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
1974	--	--	--	--	--	--	--	--	--	--	--	--
1975	1	2.9	2	5.7	13	34.1	19	54.3	--	--	35	100
1976	--	--	--	--	--	--	--	--	--	--	--	--
1977	8	9.4	11	12.9	25	29.4	30	35.3	11	12.9	85	100
1978	--	--	--	--	--	--	--	--	--	--	--	--
1979	63	26.8	2	2.0	126	53.6	41	17.4	5	2.1	237	100
1980	26	24.1	--	--	40	37.0	39	36.1	3	2.8	108	100
1981	21	15.9	7	5.3	85	64.4	19	14.4	--	--	132	100
1982	17	12.2	16	11.5	81	58.3	25	17.9	--	--	139	100
1983	36	39.1	5	5.4	20	21.7	31	33.7	--	--	92	100
1984	--	--	--	--	--	--	--	--	--	--	--	--
1985	--	--	--	--	--	--	--	--	--	--	--	--
1986	13	27.7	3	6.4	21	44.6	10	21.3	--	--	47	100
1987	--	--	--	--	--	--	--	--	--	--	--	--
Total	185	21.1	46	5.3	411	46.9	214	24.5	19	2.21	875	100
Calculated minimum percent of entangled individuals:												
		0.52		0.13		^a 1.15		0.59		0.05		

^aFrom Table 1.

2.19% for Robben Island and between 0.05 and 1.15% for the Commander Islands. These estimated lower bounds cover the period from June through August (3 months), the time during which seals are seen on the hauling grounds.

DISCUSSION

It is natural that some seals that are entangled in commercial fishing waste die at sea without being seen on land, especially in the areas where seals from the western Pacific overwinter (Pacific Ocean and Sea of Japan in the Japanese, Korean, and American exclusive economic zones). However, data concerning entanglement rates and mortality for these areas are not available. In Fowler's (1985, 1987) work and Fowler et al. (1990), information is presented as a basis for accounting for unseen mortality at sea. After accounting for debris that is too large for seals to return to land, mesh sizes that pups become entangled in, and the mortality rates observed in large debris, as many as 30 to 35 seals per year may die for each one that is observed alive. The effects of the size composition of debris at sea has not been considered in its effects on seal populations of the

western Pacific. Neither have the effects of mesh size. However, in view of the levels of entanglement presented in Tables 2 and 3, it is clear that an even lower ratio of unseen mortalities to observed entanglement would be necessary to achieve similar levels of mortality thought possible for the Pribilof population. Further data on the size composition (by weight and mesh) of debris from beaches and pelagic habitat in the western Pacific will be necessary to extend this evaluation of the potential effects of marine debris on the mortality of fur seals.

There is a very important observation to be noted in the data in Tables 2 and 3, an observation that deserves to be emphasized. At the Commander Islands, the incidence of entanglement among seals is lower than at Robben Island. Among bachelors, the incidence of entanglement is 1.7 times as high on Robben Island as on the Commander Islands. The minimum level of entanglement for females is almost four times as high. The importance of this observation comes from the fact that this may be one of the reasons why the Commander Island seal population did not decrease in recent years, while other populations declined. This emphasizes the need for further information on the composition of debris (by weight and mesh size) to determine if there are differences between the western and eastern Pacific. Such differences might explain the divergence in dynamics between the population of fur seals on the Commander Islands (no recent declines) and the Pribilof Islands (a decline in the late 1970's).

If mortality due to entanglement is as high in the western Pacific as is thought possible for the Pribilof population, as many as 3.7 to 6.7 thousand seals die from entanglement annually. In the North Pacific as a whole, then, as many as 60,000-65,000 northern fur seals may die due to the discard and loss of gear and debris from commercial fisheries. Those figures are 1.7-2 times higher than the figures of the potential annual harvest of juvenile male northern fur seals in the U.S.S.R. and the United States.

It is known that 60% of the Robben Island seals, 5-6% of the Pribilof seals, and 28-30% of the Commander Island seals winter in Japanese waters (Ashchepkov and Kuzin 1986). Cooperative efforts involving all interested countries seem necessary for studying the problem of the pollution of the ocean by commercial fishing waste. The existing fisheries-oriented scientific associations among the Pacific countries provide the opportunity for their leadership to inform fisheries organizations about the sources of debris and the volume of the damage caused by commercial fishing to marine resources. The solving of this problem depends on how soon and how completely sailors and fishermen realize the seriousness of marine pollution and that the discard of debris is contrary to international regulations.

REFERENCES

- Ashchepkov, A. T., and A. E. Kuzin.
 1986. Abundance and age-sex composition dynamics of the different reproductive seal groups at wintering places in Sea of Japan and off the Japanese Pacific coast. *Izuchenie, okhrana i ratsionalnoe ispolzovanie morskikh mlekopitayushchikh*. Archangel. 14-16. [In Russ.]

DeGange, A. R., and T. C. Newby.

1980. Mortality of seabirds and fish in a lost salmon driftnet. *Mar. Pollut. Bull.* 11:322-323.

Fowler, C. W.

1982. Interactions of northern fur seals and commercial fisheries. *Trans. N. Am. Wildl. Natur. Resour. Conf.* 47:278-292.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.

Fowler, C. W., R. Merrick, and J. D. Baker.

1990. Studies of the population level effects of entanglement on northern fur seals. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Kuzin, A. E.

1985. Injuring of fur seals with the commercial fishing waste in the northern Pacific. *Issledovanie i ratsionalnoe ispolzovanie bioresursov dalnevostochnykh i severnykh morey. Tez. Vses. Soveshch. Valdivostok*, p. 92-93. [In Russ.]

Moiseev, P. A.

1979. Biological resources of the ocean. In *Biologicheskie resursy mirovogo okeana*, p. 13-26. Nauka Publ., Moscow. [In Russ.]

Shomura, R. S., and H. O. Yoshida (editors).

1985. *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

Yerukhimovich, V. B., and S. K. Yefremenko.

1985. Tasks of the fisheries science in relation to the far-eastern fisheries development. In *Problemy dalnevostochnoy rybokhozyaystvennoy nauki*, p. 122-136. Agropromizdat Publ., Moscow. [In Russ.]

AN INDEX OF FUR SEAL ENTANGLEMENT IN FLOATING NET FRAGMENTS

Christine A. Ribic*
Center for Quantitative Science in Forestry, Fisheries and Wildlife
University of Washington
Seattle, Washington 98195, U.S.A.

and

Gordon L. Swartzman
Applied Physics Laboratory
University of Washington
Seattle, Washington 98195, U.S.A.

*Present address: Environmental Research Laboratory, U.S. Environmental Protection Agency, 200 SW 35th Street, Corvallis, OR 97333, U.S.A.

ABSTRACT

While information has been published on transect surveys based on visual sighting of floating marine debris, few attempts have been made to link the estimates of floating marine debris density to the entanglement rate observed in subadult male fur seals. Both published and unpublished survey data were used to develop a data base consisting of the location and season during which floating marine debris were observed and the estimated density of the debris. In conjunction with this data base, similar information was used for at-sea sightings of fur seals to calculate an index of potential entanglement by season (winter and breeding season, spring and fall migration). Our main conclusion is that much more information is needed to cover the known range of migrating northern fur seals. However, with these limited data, it appears that seals are most at risk during the breeding season and during the fall migration. Our conclusions are tentative due to assumptions used in calculating the index and the lack of geographical overlap between oceanic debris surveys and fur seal surveys.

INTRODUCTION

The fact of entanglement and the problems it may cause animals is not an area of debate (Center for Environmental Education 1986). But the role of entanglement in contributing to the recent decline of the northern fur seal, *Callorhinus ursinus*, on the Pribilof Islands remains to be clearly

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

defined (Fowler 1985, 1987; Scordino 1985). We (Swartzman et al. 1990) have taken a modeling approach to this problem, but the model is only as good as the estimates of the parameters and their inherent variability. In this paper, we calculate an index of potential entanglement by using information on floating marine debris and at-sea locations of fur seals. Problems with this approach are discussed.

METHODS

A review of all papers dealing with floating marine debris was used to gather information on the density of marine debris by area. All estimates from the papers were translated into number of net fragments per square nautical mile for comparison and mapped by area (using blocks of latitude and longitude as defined in the papers). We could not separate the nets into different types (trawl web or gillnet) because the data as presented in the papers were not separated by type. Unpublished data of Yoshida and Baba (Far Seas Fisheries Research Laboratory, pers. commun.) and the 1984 marine mammal observer program data (L. Jones, National Marine Mammal Laboratory, Seattle, WA, pers. commun.) were used to calculate additional density estimates by area. A strip transect density estimate (number of net fragments per square nautical mile) was calculated using a half width of 50 m (Dahlberg and Day 1985) and mapped.

For fur seal occurrence, the data in Kajimura (1980) were used. Those data are presented in summary form in terms of number of degree blocks where 0, 1-4, 5-8, or 9+ seals were seen per hour of observation for all surveys in 1958-74 by month. We calculated an index of high density of seals by 5 degree blocks from these data by month. The index of high density was chosen as the number of degree blocks with five or more seals sighted per hour of observation divided by the total number of degree blocks sampled. We calculated this index for four time periods: January-March, April-June, July-October, and November-December. These periods correspond to winter, spring migration, breeding season, and fall migration (Kajimura 1980), respectively. For each time period, the index of high density was the average of the indices for the months making up the particular period. The indices were mapped for each time period.

The two maps (debris density and fur seal occurrence) were put onto the same scale and superimposed, and areas of overlap identified by time period. Areas of no effort (indicated as blank spots on the maps) for either base set (debris or seals) automatically meant a blank spot for the combined set. For the areas of overlap, an index of potential entanglement was defined as the product of the estimated density of marine debris and the index of high fur seal density. Assumptions made because of the available data include:

- There are no age and sex differences for probability of fur seal occurrence. (Data were not divided into age and/or sex categories.)
- There have been no major changes in the pelagic distribution of fur seals. (Data were collected between 1958 and 1974.)

- Net density is constant over the seasons. (Data were not available on a seasonal basis.)
- The density and location of small and large net fragments are the same. (Data were not presented on a size basis, therefore, differences in probability of entanglement due to differences in size of net fragment cannot be factored into the index.)
- There are no seasonal changes in fragment-specific probability of entanglement. (No information was available on this point.)

The magnitude of the numbers was used to compare the relative probability of potential entanglement between areas within a time period and between time periods.

RESULTS

The papers with the most information on the density of floating nets were those by Jones and Ferrero (1985), Yoshida and Baba (1985), Baba et al. (1988), and Mio and Takehama (1988). Most of the information on floating marine debris has been collected in the North Pacific Ocean in the middle of the fishing fleets (Fig. 1). Most of the information on the presence of fur seals has been collected along the coast and in the Gulf of Alaska (Fig. 2). Fur seals have been seen offshore in the North Pacific Ocean (Kajimura 1980). However, since sighting effort is not recorded, these data cannot be used in a direct evaluation of the probability of entanglement. There are two areas where the maps overlap: around the Pribilof Islands and off the coasts of Washington and Oregon (Fig. 3).

Although there is little overlap between the two maps, for those areas where we could calculate an index of potential entanglement, the highest indices occurred during the breeding season (July-October) around the Pribilof Islands and in the fall migration (November and December) off the coasts of Washington and Oregon (Fig. 3). The index was also relatively high for some blocks off the coasts of Washington and Oregon for the winter season (January-March) and the spring migration (April-June) (Fig. 3).

DISCUSSION

The major problem in estimating the level of entanglement at sea is the lack of systematic observations of both fur seals and marine debris in the same area. The lack of data is so extreme as to present major problems in calculating indices of entanglement while seals are in their pelagic environment. As can be seen from this analysis, there was little overlap between the sets of data for the areas that were surveyed. In addition, for the existing data sets, the units for marine debris density and the probability of fur seal occurrence are not the same. There were no data for the effort behind the fur seal sightings. These circumstances lead to obvious problems in trying to use the existing empirical data to calculate




Figure 3.--Index of potential entanglement for common areas between Figures 1 and 2 for four time periods: (a) January-March, (b) April-June, (c) July-October, and (d) November-December. The coastline is outlined in black.

an index. What is needed is a study specifically designed and coordinated for three purposes. The first purpose would be to estimate the density and location of floating net fragments; the second would be to record the numbers and location of fur seals; and the third would be to collect information on the number of fur seals entangled in debris. All three parts of the study should consider the seasonal aspects of the distribution and abundance of both seals and debris. The data from such a study would have the debris and fur seal variables on the same scale (e.g., number per unit area) as well as contain information on the location and date of observed at-sea fur seal entanglement. Until such a study is done, combining results of other studies for such purposes will be highly speculative and will depend on a large number of assumptions, as evidenced by this study.

The assumptions used in this study were made to compensate for the lack of data on variables such as differences in location by age and sex for seals and differences in large and small net fragment densities and location. Changing an assumption will affect the value of the index, but whether the comparisons are changed will depend on how the assumption is changed. For example, if the probability of entanglement upon encounter changes with season, then the comparisons between seasons would be affected.

Given all the assumptions, the index calculated here indicates that the probability of potential entanglement may change depending on the season of year and, in a related fashion, on location. If the relatively high potential entanglement index around the Pribilof Islands found here is valid, then it may indicate that the entanglement problem begins as soon as the young of the year go to sea and continues as the animals migrate south for the winter. Since the first year after weaning is a stressful time for young animals, the actual impact of entanglement may be severe for fur seals going to sea for the first time.

ACKNOWLEDGMENTS

We thank M. B. Hanson for doing the literature review on floating marine debris and entanglement. We thank N. Baba, K. Yoshida, and L. Jones for allowing us to use their unpublished data. We thank C. W. Fowler for support in this project and for providing us with contacts to obtain unpublished data. We thank an anonymous reviewer for improving this manuscript. This project was funded by the National Marine Mammal Laboratory under Contract 40-ABNF803196.

REFERENCES

- Baba, N., K. Yoshida, M. Onoda, N. Nagai, and S. Toishi.
1988. Results of research on floating fishing gear and fish net fragments in the area southwest of the Pribilof Islands and off southern coasts of the Aleutian Islands, July-August 1985. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 143-164. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Center for Environmental Education.

1986. Marine wildlife entanglement in North America. Center for Environmental Education, Wash., D.C., 219 p.

Dahlberg, M. L., and R. H. Day.

1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Fowler, C. W.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. Mar. Pollut. Bull. 18:326-335.

Jones, L. L., and R. C. Ferrero.

1985. Observations of net debris and associated entanglements in the North Pacific Ocean and Bering Sea, 1978-84. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 183-196. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Kajimura, H.

1980. Distribution and migration of northern fur seals (*Callorhinus ursinus*) in the eastern Pacific. In H. Kajimura, R. H. Lander, M. A. Perez, A. E. York, and M. A. Bigg (editors), Further analysis of pelagic fur seal data collected by the United States and Canada during 1958-74, Part I, p. 6-50. Report submitted to the 23rd Annual Meeting of the Standing Scientific Committee, North Pacific Fur Seal Commission, 7-11 April 1980, Moscow, U.S.S.R.

Mio, S., and S. Takehama.

1988. Estimation of distribution of marine debris based on the 1986 sighting survey. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 64-94. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Scordino, J.

1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Swartzman, G. L., C. A. Ribic, and C. P. Huang.

1990. Simulating the role of entanglement in northern fur seal, *Callorhinus ursinus*, population dynamics. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Yoshida, K., and N. Baba.

1985. Results of a survey on drifting fishing gear or fish net pieces in the Bering Sea. Document submitted to the 28th Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 4-12 April 1985, Tokyo, 13 p.

STUDIES OF THE EFFECTS OF ENTANGLEMENT
ON INDIVIDUAL NORTHERN FUR SEALS

Robert L. DeLong, Patrick J. Gearin, John L. Bengtson, Pierre Dawson
National Marine Mammal Laboratory
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
Seattle, Washington 98115, U.S.A.

and

Steven D. Feldkamp
Long Marine Laboratory
Institute of Marine Studies
University of California at Santa Cruz
Santa Cruz, California 95064, U.S.A.

ABSTRACT

During the field seasons of 1985 and 1986, studies were conducted to determine the effects of entanglement on the northern fur seal, *Callorhinus ursinus*. These included surveys of entanglement rates among pups and adult females, an experiment on the effects of entanglement on adult females, and a study of the selectivity of mesh size in the entanglement of pups. Complementing these studies are data on the history of the development of wounds for entangled juvenile males that have been seen more than one time.

In 1985, 40 parturient females and their pups were captured at Zapadni Reef rookery on St. Paul Island. Half of the females were treated as controls and tagged with both flipper tags and radio tags and released. Pieces of trawl net weighing 200 g were placed on the other 20, simulating entanglement common to fur seals. The attendance cycles and rates of return of these animals were then compared for the two groups for several feeding cycles, and the rates of return were compared the following season. Three of the entangled females freed themselves of the debris. Of the remaining 17, 3 failed to return after their first trip to sea, 4 failed to return after their second trip, and 2 did not return after their third trip. One control did not return after her second trip to sea. The time spent at sea by the entangled animals was twice as long as for the control animals. In 1986, 2 of the 17 entangled animals were observed, whereas 12 of the 20 controls were observed.

Ground surveys for females were conducted on rookeries chosen for ease of access to observe animals. Entangled females were counted during these surveys, and the counts were converted to entanglement rates by using the numbers of pups estimated for each of the rookeries as an indication of the number of females present. Rates calculated on this basis ranged from 0.06 to 0.23% for the sample rookeries with a mean of 0.15%. This is to be compared to the 0.4% seen for the juvenile males.

Between 11 September and 16 October 1986, 39 entangled pups were observed. Of these, five were in a single piece of trawl webbing that had become wrapped around a channel marker, and another five were in a piece of blue trawl webbing that washed ashore. As with other components of the population, trawl webbing comprised the highest portion of the entangling debris (19 out of the 39 observed). Entanglement rates for these animals are not known because we have no information on the portion of the pup population that had already departed for sea. The live entangled pups were tagged and released.

During 1985, experimental studies of pup entanglement showed that pups of the size of those found on the islands in October can become entangled in trawl debris with mesh sizes as small or smaller than 16 cm (stretched). All experimental pups placed in a tank with pieces of net with mesh sizes between 18 and 22 cm became entangled within 5 h or less. Some became entangled about their face in pieces with mesh sizes as small as 14 cm.

Data on the interannual history of a small number of entangled subadult males indicate that growth in body size and abrasion brought about by movement cause wounds to increase in size.

STUDIES OF THE EFFECTS OF NET FRAGMENT ENTANGLEMENT ON NORTHERN FUR SEALS
PART 1: DAILY ACTIVITY PATTERNS OF ENTANGLED AND NONENTANGLED FUR SEALS

Kazumoto Yoshida
Seikai National Fisheries Research Institute
Nagasaki, 850 Japan

Norihisa Baba, Masashi Kiyota
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

Masayuki Nakajima, Yasutoshi Fujimaki, and Akira Furuta
Izu-Mito Sea Paradise
Numazu, Shizuoka, 410-02 Japan

ABSTRACT

Effects of net fragment entanglement on the behavior of fur seals were examined using radio telemetry. Radio transmitters were attached to three fur seals 5-8 years old kept in an aquarium. Two of the seals were entangled with 1- and 2-kg fishing net fragments, respectively, around their necks. Using radio telemetry, their activity patterns were recorded for 36 days, from 28 January to 4 March 1985. The seal entangled with the 2-kg net showed the shortest active time per day and the nonentangled seal showed the longest. Active time of the entangled individuals increased after removal of the nets. It became clear that the active time of fur seals was diminished by entanglement in net fragments.

INTRODUCTION

It has been reported that some northern fur seal, *Callorhinus ursinus*, returning to breeding islands were entangled in marine debris such as fishing net fragments and packing bands (Scordino 1985). The fur seal population of the Pribilof Islands declined to less than half of its 1940's peak, and mortality due to entanglement is suspected as a major cause of the population decline (Fowler 1982). In what period and at what rate do entangled fur seals die? This issue was examined through tag and resighting surveys of entangled fur seals on St. Paul Island (Bengtson et al. 1988; Scordino et al. 1988). The survival period of entangled animals is considered to vary according to the damage caused by entanglement. There have been only a few reports about the effects of net fragments on fur seals (Feldkamp et al. 1987). This study was intended to examine the effects of entanglement on activity patterns of fur seals using radio telemetry.

MATERIALS AND METHODS

Experimental Fur Seals and Radio Equipment

The three female fur seals used in the experiment were captured off Joban coast, north Japan, between 4 and 9 March 1982, and had been kept in captivity for 3 years. The estimated ages of the animals were 5-8 years and body weights were 29.5-36.0 kg at the beginning of the experiment (Table 1). The experiment was conducted in an aquarium, Izu-Mito Sea Paradise, Numazu, central Japan, where the animals were kept.

A radio transmitter and trawl net fragments were attached to the seals on 28 January 1985. Of the three, one (referred to hereafter as N0) was loaded only with a transmitter. A transmitter and a 1-kg net fragment were attached to one of the remaining two (N1), and a transmitter and 2-kg net fragments to the other (N2). Trawl nets used in the experiments were those commonly used in commercial fishing; they were made of polyethylene with a twine size of 3.4 mm and a mesh size of 24 cm. The transmitter was cylindrical in shape, 35 mm in diameter, 155 mm long, and 200 g in atmospheric weight. It had a life of about 6 months (Fig. 1). Receivers and recorders were installed in an observation room near the experimental area (Fig. 2).

The transmitter was attached with a harness made of nylon webbing belts sewn together with colored tapes for individual recognition. Immediately after attachment of the transmitter and net fragments, the animals were released into the experimental area (Fig. 2). The experimental area, made by partitioning an inlet with nets, was about 1,400 m² and had a natural beach. The deepest part at high tide was about 7 m; there was a tidal range of about 1.5 m. The sea was calm throughout the year. In addition to the three experimental seals, 33 other fur seals were kept in captivity in this area. Activities of N0 were recorded continuously from 4 February to 4 March. Activities of N1 and N2, entangled in net fragments, were recorded from 28 January to 26 February. The nets were intentionally removed on 27 February and their activities without nets were recorded from 28 February to 4 March.

During the entire period, behavior and health of the experimental individuals were checked carefully at a regular time each day. Moreover, in order to compare actual movements with radio records, behavior of the three animals was observed visually several times a day for 4-5 h each time.

During the experiment, fur seals were fed with defrosted mackerel in the amount of 1-4 kg (an average of 2.44 kg) per day at 1000 and 1630 on the beach. At each feeding, food was provided first to each experimental individual, and later to the herd in general.

Weather observations were made around 1400 each day. The average air temperature during the experiment period was 10.2°C (ranging from 6.0° to 18.0°C), with average water temperature at 13.3°C (between 11.1° and 15.8°C) and average humidity at 63.1% (from 38 to 88%).

Table 1.--Information on the three fur seals used in the experiment.

Seal		Capture			Age and size at start of experiment		
ID	Sex	Date 1982	Latitude	Longitude	Estimated age (year)	Body length (cm)	Body weight (kg)
N0	F	9 March	36°30'N	141°16'E	6	123	29.5
N1	F	4 March	36°42'N	141°15'E	8	123	36.0
N2	F	8 March	36°26'N	141°06'E	5	120	33.0

Analysis of Activity Records

Figure 3 shows an example of the activity records of a fur seal wearing a telemetry device. Records representing the activity of the animals are called "actograms." Waves in the figure indicate changes in intensity of electric signals due to movement of the animals. When an animal with a transmitter was on land, a continuous wave form was observed (Fig. 3A); movements of the animal on land could be recognized as fluctuations of wave form on the recording paper. The period in which the wave form was fluctuating was defined as active time on land. When the transmitter-loaded animal was in the water and dived, no signal could be received because electric waves are greatly attenuated in seawater. The recording pen then moved straight along the baseline. When the animal emerged, a sharply pointed line was recorded on the paper, corresponding to the abrupt rise of electric wave intensity. Thus, actograms for the animal moving at the sea surface showed a pectinate wave form (Fig. 3B). When the seal was resting at sea, either a flat line or a baseline could be recorded. The former means that the animal was resting with her back upward, and the latter, resting with her back under water. Therefore, the fluctuating wave form indicates activity ashore, and the pectinate wave form indicates activity at sea. We measured the length of such "active" periods in each actogram and calculated the active time and resting time in a day.

RESULTS

General Behavior

During the experiment period, other fur seals did not exhibit special behavior such as avoiding or threatening the transmitter-loaded animals or approaching them with curiosity. The fur seals with transmitters were always within the herd.

Differences in the general behavior of the three fur seals were recognized by visual observation. For several days after the experiment began, N1 and N2 tried to get rid of the attached net fragments by shaking their necks. Seal N2 moved slowly and chiefly engaged in slow swimming or




Figure 1.--A transmitter and harness.




Figure 2.--Map showing the experimental area. A. Observatory where receiving instruments were settled. B. Indoor breeding facility. C. Partitioned section where transmitters were attached.





Figure 3.--Actograms, received electronic waves representing the activity of animals. A. Resting on land. B. Swimming at sea.

grooming. Dives and headstands in water, often seen in free-swimming seals, were not observed for N2. Seal N1 moved more smoothly than N2. Sometimes N2 made shallow dives or underwater headstands, but she made no extended deep dives. Seal N0's movements were very smooth and did not differ from those of individuals without transmitters.

For several days after the start of the experiment, N1 and N2 did not draw near to men even at feeding time. Later they approached the feeder as did the other seals. When food was thrown, the nonentangled individuals often caught it before the entangled ones. Seal N0's feeding activity was no different from that of nonloaded fur seals.

Activity Pattern

Active Time

Figure 4 shows variations in active time per day from 4 to 26 February. During this period, the amount of N0's active time fluctuated greatly, while fluctuations were small for N1 and N2. The average daily active time was longest for N0 (9.6 h/day), followed by N1 (4.1 h/day) and N2 (1.4 h/day), and any pair of them differed significantly (t-test, $P < 0.01$).

Table 2 shows the daily active times of N1 and N2 before and after the removal of net fragments. The average active time of N2 after net removal was 5.4 h/day, about four times longer than before net removal. The difference was statistically significant (t-test, $P < 0.01$). The average active time of N1 after net removal was 6.7 h/day, about double that of the period of net attachment, which also differed significantly (t-test, $P < 0.01$). The active time of N0 did not show a significant change between the two corresponding periods.

Daily Cycle of Activity

Figure 5 shows the average daily cycle of activity. The ratio of activity was calculated every 3 h (activity ratio) and averaged for the experiment period. For N1 and N2, the periods of entanglement and nonentanglement were treated separately. All three seals were very active in two time periods, 0900-1200 and 1500-1800, which corresponded to feeding times. Activity patterns of N1 and N2 did not change remarkably after removal of entangled nets, though the active time increased as mentioned above.

DISCUSSION

Baba and Yoshida (1988) conducted a field experiment in which they attached transmitters to two mature female fur seals, one of which was entangled in a 120-g net fragment, off St. Paul Island and compared their activities using radio telemetry. They reported that the frequency of dives longer than 1 min was less for the entangled animal. In our study, no extended dives were observed for animals loaded with nets of 1 and 2 kg. Although there were differences in research location and the amount of nets




Figure 4.--Variations in daily activity of the three fur seals.

attached, their report is consistent with our study in that net entanglement hindered diving activities.


Daily activity cycles were the same for entangled and nonentangled animals. Baba and Yoshida (1988) also reported that no differences were observed in behavior patterns of entangled and nonentangled fur seals in the open sea. These results indicate that activity patterns of fur seals may not change even if they are entangled in net fragments.

Most of the trawl nets entangling the fur seals on St. Paul Island were <150 g in weight, although the biggest one weighed 6.75 kg (Scordino 1985). Therefore, it is also necessary to examine the effects of smaller net fragments on activity of fur seals.

It is clear that net entanglement suppressed the activity of animals because active time of the animals was short while entangled and increased


Table 2.--Daily active times of experimental seals with and without attached nets.

Seal ID	4-26 February 1985			28 February-3 March 1985		
	Net weight (kg)	Active time per day mean (min.-max.) (h/day)	S.D.	Net weight (kg)	Active time per day mean (min.-max.) (h/day)	S.D.
N0	0	9.6 (5.0-17.4)	4.0	0	11.4 (8.2-16.1)	3.4
N1	1.0	4.1 (0.4-7.5)	1.7	0	6.7 (6.3-7.3)	0.5
N2	2.0	1.4 (0.6-2.9)	0.7	0	5.4 (3.6-6.4)	1.3




N0

4 Feb - 3 Mar ; nonentangled



N1

28 Jan - 26 Feb ; entangled
28 Feb - 3 Mar ; nonentangled



N2

28 Jan - 26 Feb ; entangled
28 Feb - 3 Mar ; nonentangled

Figure 5.--Daily activity cycles of three fur seals for entangled (dotted area) and nonentangled (solid line) periods. Activity ratio is defined as the percentage of active time in each 3-h period.

after net removal. The suppression of activity might be due to either the physical burden of nets or an adaptation of animals to conserve energy. A future task should be to study the physiological impact of entanglement and relate it to energy consumption and survival.

ACKNOWLEDGMENTS

We express gratitude to the breeding technicians of Izu-Mito Sea Paradise, who collaborated in the experiment, as well as to the officials of the Fishing Ground Environment Conservation Division of the Fisheries Agency, who provided us with the opportunity to conduct this study.

REFERENCES

- Baba, N., and K. Yoshida.
 1988. Results of research on behavior of entangled and nonentangled fur seals by using radio-wave telemetry in areas around the Pribilof Islands, late July-early August, 1985. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987, p. 165-188. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Bengtson, J. L., C. W. Fowler, H. Kajimura, R. L. Merrick, K. Yoshida, and S. Nomura.
 1988. Fur seal entanglement studies: Juvenile males and newly-weaned pups, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.
- Feldkamp, S. D., D. P. Costa, and G. K. Dekrey.
 1987. Energetic and behavioral effects of net entanglement on juvenile northern fur seals, *Callorhinus ursinus*. Contract report to the National Marine Fisheries Service, Contract No. 85-ABC-00185, 44 p. (Available from Long Marine Laboratory, Institute of Marine Science, University of California, Santa Cruz, CA 95604.)
- Fowler, C. W.
 1982. Interactions of northern fur seals and commercial fisheries. Trans. N. Am. Wildl. Nat. Resour. Conf. 47:278-292.
- Scordino, J.
 1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Scordino, J., H. Kajimura, N. Baba, and A. Furuta.
 1988. Fur seal entanglement studies in 1984, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 70-78. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.

STUDIES OF THE EFFECTS OF NET FRAGMENT ENTANGLEMENT ON NORTHERN FUR SEALS
PART 2: SWIMMING BEHAVIOR OF ENTANGLED AND NONENTANGLED FUR SEALS

Kazumoto Yoshida
Seikai National Fisheries Research Institute
Nagasaki, 850 Japan

Norihisa Baba, Masashi Kiyota
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

Masayuki Nakajima, Yasutoshi Fujimaki, and Akira Furuta
Izu-Mito Sea Paradise
Numazu, Shizuoka, 410-02 Japan

ABSTRACT

The effects of net fragment entanglement on the swimming behavior of fur seals were observed. Net fragments of six different weights (0.5 to 3 kg) were attached to the necks of eight fur seals, two males and six females, 4 to 9 years old. They were released in an aquarium pool with fish, and their swimming speed and time required to capture a fish were recorded. Of the eight individuals examined, three showed active feeding behavior. As the amount of attached net was increased, swimming speed decreased and more time was required for an entangled seal to catch a fish. Decrease in swimming speed was proportional to the relative load of net fragments (net weight/body weight).

INTRODUCTION

Marine debris is known to cause problems for various animals such as fish (High 1985), marine mammals (Calkins 1985; Henderson 1985), seabirds (Tull et al. 1972), and turtles (Balazs 1985; Cawthorn 1985). Many fur seals have been found on the breeding islands entangled in fishing net fragments and packing bands (Waldichuk 1978; Scordino 1985). Fowler (1982) noticed that entanglement probably was a cause of recent decline in the Pribilof population of the northern fur seal, *Callorhinus ursinus*. In order to understand the mechanism and impact of net entanglement on northern fur seals, the National Research Institute of Far Seas Fisheries has conducted various experiments at the Izu-Mito Sea Paradise, an aquarium, since 1983. The mechanism of entangling and influence of entanglement on activity patterns of fur seals were surveyed before (Yoshida et al. 1985,

1990). In this study, experiments were conducted to understand the effect of net entanglement on swimming behavior of fur seals.

MATERIALS AND METHODS

Swimming speed and feeding behavior of net-entangled fur seals were observed in a pool of the Izu-Mito Sea Paradise in Numazu, Japan, from 27 January to 19 February 1986. The pool was 22 m wide, 10 m long, and 4 m deep (Fig. 1). The front of the pool was made of transparent plexiglass through which underwater movements of fur seals were observed. Trawl net fragments of six different weights were attached to the necks of eight fur seals, two males, and six females, estimated to be from 4 to 9 years old (Table 1). The nets used were gray trawl nets made of polyethylene, with a twine size of 3.4 mm and a mesh size of 24 cm. Specific gravity of the net was 0.77, which meant that a net fragment weighing 1 kg of air had a buoyancy of 340 g in seawater with a specific gravity of 1.03. Weights of the six fragments were 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 kg. As a control, free-swimming animals were also observed.

Experiments were conducted between 1600 and 1730 every day. Fur seals had not been fed since the previous day so that they would respond readily to food. Measurement of swimming speed was conducted in the following manner. One seal was released into the pool and lured to one corner by a display of food. Then a man showed a fish and threw it 8-10 m from the seal. When the seal started swimming after the fish, one observer recorded the time taken by the seal to swim a distance of 6 m. The distance was measured using the interval of frames supporting the glass wall. Each individual was tested using nets of two different weights per day. For each weight of net, an individual was obliged to swim eight times. One seal could make up to 16 swims in a day. If a seal would not chase a fish, it was removed from the pool and another individual was introduced.

Time to capture a fish was measured for the three seals which readily swam for a fish (M1, M2, and F1 in Table 1). Basic design of the experiment was the same as that for swimming speed measurement. Live sardine, *Sardinops melanostictus*, 12.5 to 15.0 cm in length and 15-29 g in weight, was used as bait and was thrown 8-10 m ahead of the seal. The time it took the seal to catch the fish was measured. Eight trials were made for each net weight, although the number was reduced when an animal with heavy entanglement looked tired.

To evaluate the effect of net entanglement on the basis of body weight, relative load of attached net was calculated:

$$\text{relative load} = \frac{\text{weight of attached net}}{\text{body weight of fur seal}} \times 100 (\%)$$

During the experiment period, average air temperature was 7.1°C (1.1°-11.0°C), average water temperature was 12.7°C (12°-13.4°C), and average humidity was 57.4% (36-83%).




Figure 1.--The pool used in this study. Underwater movements of fur seals were observed through the glass wall.

RESULTS

Individuals Used in the Experiment

Of the six female fur seals used, only one individual (F1) swam actively for fish. She chased fish a total of 32 times for swimming speed measurement, whether she was free-swimming or entangled in 0.5 to 1.5 kg of nets. Two other females (F3 and F5) swam only once or twice, even when they were free from nets. The remaining three made no attempt to swim after a fish. Two males (M1 and M2) tried to catch a fish in every case, whether free-swimming or entangled in up to 3.0-kg nets.

General Behavior

Fur seals often swam on their backs when they were free from entanglement, but entangled seals did not exhibit this type of swimming. All the seals tested were able to swim down to the bottom of the pool (4 m deep) in every degree of entanglement. When they chased a fish, they swam in the upper layer within 1 m of the surface. When entangled seals swam, their bodies twisted up and down. The body undulation was intensified as the amount of attached net was increased.

Swimming Speed

Swimming speed of the three fur seals (M1, M2, and F1), calculated from the amount of time it took them to pass the 6-m mark, decreased as the weight of attached nets increased (Table 2, Fig. 2). The average swimming speed of M1 without entanglement was 2.98 m/sec, but fell to 1.05 m/sec

Table 1.--Information on the fur seals used in the experiment.

Seal	Sex	Capture			Age and size at experiment		
		Date	Location		Age (year)	Body length (cm)	Body weight (kg)
			(Lat.)	(Long.)			
M1	M	July 1981	Robben Island		4	137	58.0
M2	M	4 Mar. 1982	36°34'N	141°14'E	4	132	54.0
F1	F	July 1981	Robben Island		4	120	23.0
F2	F	10 May 1980	37°57'N	142°14'E	6 ^a	120	35.5
F3	F	8 Mar. 1982	36°26'N	141°06'E	6 ^a	121	30.5
F4	F	4 Mar. 1982	36°42'N	141°15'E	9 ^a	125	36.0
F5	F	8 Mar. 1982	36°27'N	141°10'E	8 ^a	124	38.0
F6	F	9 Mar. 1982	36°30'N	141°16'E	7 ^a	122	32.5

^aEstimated age.

when 3-kg nets were attached. That of M2 decreased from 3.04 to 0.96 m/sec when 3-kg nets were loaded. For both M1 and M2, the speed was about one-third of that in a nonentangled state. The average swimming speed of F1 free from entanglement was 2.51 m/sec, but it fell to 0.73 m/sec when 1.5 kg nets were attached.

Figure 3 shows the relation between average swimming speed and relative load of net fragments. The relationship was similar for the three individuals. Swimming speed decreased in proportion to the relative load of attached nets. Linear regression of the relationship between relative load and swimming speed was

$$\text{swimming speed (m/sec)} = 2.26 - 0.25 \times \text{relative load (\%)} \quad (r = -0.97)$$

Swimming speed of free-swimming animals was excluded from the regression.

Time Required to Capture a Fish

Table 3 shows the time it took for three seals, M1, M2, and F1, to capture a fish. F1 was not tested with net fragments heavier than 1.5 kg because that much weight seemed too heavy for her. The relationship between weight of nets and time required to capture a fish is shown in Figure 4. Although there was a considerable range, all three seals required more time to catch a fish as the amount of attached net was increased. For the three weights of nets examined, average capture time was the longest for F1. It was observed that when fur seals tried to catch a live fish, they approached the fish and turned their heads quickly to snap at it. Entangled seals had difficulty with the dash and snap.




Figure 2.--Changes in swimming speed of three fur seals due to net entanglement. M1:▲ M2:■ F1:○.

Table 2.--Changes in swimming speeds (m/sec) of three fur seals due to amount of net entanglement. Speeds were calculated using the time it took the seals to swim a distance of 6 m.

Seal	Weight of net (kg)						
	0.0	0.5	1.0	1.5	2.0	2.5	3.0
	Swimming speed						
M1							
Mean	2.98	2.16	1.92	1.63	1.50	1.20	1.05
Minimum	2.72	2.06	1.71	1.50	1.39	1.11	0.95
Maximum	3.15	2.40	2.06	1.76	1.66	1.27	1.20
Standard error	0.14	0.11	0.13	0.10	0.08	0.06	0.08
Sample number	8	8	8	8	8	8	8
M2							
Mean	3.04	2.09	1.77	1.37	1.23	1.01	0.96
Minimum	2.60	1.87	1.66	1.25	1.13	0.93	0.82
Maximum	3.33	2.30	1.87	1.57	1.33	1.17	1.09
Standard error	0.29	0.15	0.07	0.10	0.07	0.07	0.11
Sample number	8	8	8	8	8	8	8
F1							
Mean	2.51	1.66	1.09	0.73	--	--	--
Minimum	2.30	1.50	0.90	0.58	--	--	--
Maximum	2.85	1.93	1.25	0.89	--	--	--
Standard error	0.23	0.16	0.15	0.12	--	--	--
Sample number	6	6	6	6	0	0	0




Figure 3.--Relationship between average swimming speed and relative load of attached net. M1:▲—▲ M2:■---■ F1:●—●

Table 3.--Time to capture a live fish, *Sardinops melanostictus*, required by the three seals carrying different weights of nets.

Seal	Weight of net (kg)						
	0.0	0.5	1.0	1.5	2.0	2.5	3.0
Capture time (seconds)							
M1							
Mean	18.1	34.1	68.1	78.8	87.8	152.8	171.6
Range	4-47	11-62	9-115	15-200	30-178	135-191	35-346
Standard error	5.2	5.9	12.1	21.0	17.3	8.0	39.4
Sample number	8	8	8	8	8	8	6
M2							
Mean	4.4	17.6	40.6	94.3	110.9	103.6	142.2
Range	3-5	9-30	15-71	12-247	58-180	67-150	76-210
Standard error	0.3	2.5	5.9	27.4	14.9	13.0	17.0
Sample number	8	8	8	8	8	6	6
F1							
Mean	25.1	46.3	79.0	115.3	--	--	--
Range	14-59	16-97	17-218	30-264	--	--	--
Standard error	5.3	10.7	23.7	27.0	--	--	--
Sample number	8	8	8	8	0	0	0




Figure 4.--Relationship between weight of nets and time required to capture a fish. Vertical lines indicate standard error. M1:▲—▲ M2:■---■ F1:●---● .

DISCUSSION

Of the eight animals used in these experiments, only three 4-year-olds, one female and two males, tried to capture a thrown fish. M1 and F1 were brought from Robben Island in 1981 as pups and were fed milk by men. All the other seals were caught pelagically. Difference in tractability of the seals might be derived from individual history as well as age, sex, hunger, and disposition.

Swimming speed of the entangled seals decreased as the weight of nets increased. Negative linear relation was observed between swimming speed and relative net load. Decrease in swimming speed might result from two physical forces of net fragments: buoyancy and drag. Buoyancy lifted the body at the neck and shifted the center of gravity. Buoyancy of nets is likely to hinder the dives of entangled seals although all the seals could dive to the bottom of the pool. Body undulation of entangled seals, which was observed when they swan, might be brought about by lifting of the neck caused by buoyancy. At the same time, swimming efficiency was reduced by the drag of the attached nets. These two forces would interfere with diving and swimming and would increase the energy expenditure of entangled seals.

Entangled seals took longer to capture a live fish as the weight of attached nets increased (Fig. 4). The increase in the capture time was derived from a decrease in swimming speed and hindrance of quick body motion. This result indicates that foraging efficiency of entangled seals will be lower than that of free-swimming seals. Heavily entangled seals should suffer from a large expenditure and a small intake of energy. Such an energy problem may be a cause of mortality of entangled seals as well as traumatic damage.

ACKNOWLEDGMENTS

We are grateful to the breeding technicians of the Izu-Mito Sea Paradise, who collaborated in the research, and to officials of the Fishing Ground Environment Conservation Division of the Fisheries Agency, the Government of Japan, who provided us with the opportunity to conduct this study.

REFERENCES

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Calkins, D. G.
1985. Steller sea lion entanglement in marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 308-314. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Cawthorn, M. W.

1985. Entanglement in, and ingestion of, plastic litter by marine mammals, sharks, and turtles in New Zealand waters. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 336-343. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Fowler, C. W.

1982. Interactions of northern fur seals and commercial fisheries. Trans. N. Am. Wildl. Nat. Resour. Conf. 47:278-292.

Henderson, J. R.

1985. A review of Hawaiian monk seal entanglements in marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 326-335. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

High, W. L.

1985. Some consequences of lost fishing gear. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 430-437. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Scordino, J.

1985. Studies in fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Tull, C. E., P. Germain, and A. W. May.

1972. Mortality of thick-billed murres in the West Greenland salmon fishery. Nature (Lond.) 237:42-44.

Waldichuk, M.

1978. Plastics and seals. Mar. Pollut. Bull. 9:197.

Yoshida, K., N. Baba, M. Kiyota, M. Nakajima, Y. Fujimaki, and A. Furuta.

1990. Studies of the effects of net fragment entanglement on northern fur seal. Part 1: Daily activity patterns of entangled and nonentangled fur seals. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154. [See this document.]

Yoshida, K., N. Baba, M. Nakajima, Y. Fujimaki, A. Furuta, S. Nomura, and K. Takahashi.

1985. Fur seal entanglement survey report: Test study at a breeding facility, 1983. Background paper submitted to the 28th meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission. National Research Institute of Far Seas Fisheries, Shimizu, Japan, 21 p.

**SIMULATING THE ROLE OF ENTANGLEMENT IN NORTHERN FUR SEAL,
CALLORHINUS URSINUS, POPULATION DYNAMICS**

Gordon L. Swartzman, Christine A. Ribic,* and Chisheng P. Huang
Center for Quantitative Science HR-20
University of Washington
Seattle, Washington 98195, U.S.A.

*Present address: Environmental Research Laboratory, U.S. Environmental Protection Agency, 200 SW 35th Street, Corvallis, OR 97333, U.S.A.

ABSTRACT

A multiage class model treating populations of both male and female fur seals was developed to examine the plausible long-term effect of their entanglement in discarded net debris. The model is based on the data available on age-specific survival and fecundity including data supporting the assumption of density-dependent survival of pups on land and of juveniles up to age 2 at sea. Also included in the model are age-specific and sex-specific harvests for the subadult male harvest as well as other pelagic and land-based commercial and scientific harvests. Entanglement in the model is linked to the observed incidence of subadult males in the harvest (or roundup). Supporting work in model development and parameter estimation has involved evaluation of various attempts to estimate both juvenile survival at sea and the mortality rate due to entanglement. This evaluation work has considered the appropriateness of assumptions and statistical tests used. Model results were evaluated by comparison with survey estimates of pup abundance and of harem bulls on the Pribilof Islands for years when these were available (between 1912 and 1960). Post-1960 survey results were used to examine the plausibility of entanglement mortality estimates in predicting the observed fur seal abundance decline. Sensitivity analysis on the model is used to indicate areas where there is a need for either further data collection or further analysis of existing data.

INTRODUCTION

The marked decline in northern fur seal, *Callorhinus ursinus*, populations on the Pribilof Islands since the mid-1970's has been attributed to a variety of causes, one of the most compelling of which is increased mortality due to seals' entanglement in discarded fishing net

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

debris (Fowler 1985, 1987) and other flotsam (e.g., packing bands). Although an earlier decline (1957-64) was probably due to pelagic scientific sampling of females at sea and harvest of females on land, later continued declines have not been explicable as being linked to repercussions of this harvest (York and Hartley 1981).

From observations of numbers of subadult male seals in the harvest entangled (about 0.4%), Fowler (1982), using a simple differential equation model and assumptions about the length of time entangled seals survive and the ratio of seals tangled in large (>0.4 kg) versus small (<0.4 kg) debris pieces (the seals entangled in large debris being assumed to die before reaching land), gave predictions for the possible effects of entanglement on the population. These predictions indicated that the entangled seals observed in the harvest could account for annual seal mortalities as high as 17%. Swartzman (1984) expanded Fowler's model to include age classes and density-dependent pup survival on land, as reflected in data from Lander (1981). Swartzman (1984) showed that the age and duration of susceptibility to entanglement can affect the annual mortality rate due to entanglement. The worst case scenarios (i.e., 2 months or less for half the entangled seals to die with only ages 1-3 susceptible to entanglement, or less than 12 months for half the entangled seals to die when all age classes are susceptible to entanglement) result in a long-term elimination of the fur seal population.

We have developed a model to investigate the effect of entanglement on fur seal population dynamics. This model separates male and female populations by age class and separates each age-sex class into entangled and unentangled animals. Sex-specific and age-specific susceptibility to mortality and entanglement mortality rates are also considered. Annual entanglement rate in small (<0.4 kg) debris is grounded in the observed fraction of entangled subadult males in the harvest. In the long-term simulation, harvests of males and scientific samples of females are removed from the population as an amount of seals (rather than as a rate).

The population dynamics of fur seals have been the object of many studies, and several models have been built in this regard. The present model is like many in that it is age-structured. Table 1 gives an overview of these previous models. Current modeling work is motivated by the need to synthesize current entanglement information and by the lack of previous treatment of the male population, no inclusion of male harvests and no previous formal sensitivity analysis having been done on previous models. Also, earlier entanglement models are equilibrium models that, while they were based on the best parameter values available at the time they were constructed, were not evaluated by being compared with historical data on pup estimates and bull counts. Finally, our model brings many of the parameter estimates up to date by including the latest available data. The large number of previous models points to the excellent long-term data base on fur seals, although, as will be shown later, assumptions must be made to fill gaps in the data when long-term model projections are made.

Table 1.--Comparison of northern fur seal population dynamics models (F = female, M = male).

Model	Age range/sex	Years	Harvest	Comments
York and Hartley 1981	2-25/F	1956-79	1956-74F	Juvenile survival higher after 1979.
Smith and Polacheck 1981	3-20/F	Not run	--	Challenged differential M/F juvenile survival.
Eberhardt 1981	1-22/F	1952-77	1952-68F	Density-dependent survival to age 3.
Fowler 1982	Pooled	Equilibrium	None	Entanglement effect with several mortalities.
Swartzman 1984	3-20F	Future, equilibrium	None	Density-dependent survival on land.
Trites 1984	1-25/M&F	1950-80	1956-74F constant M	Leslie model sensitivity analysis.
Reed and French 1987	1-29/M&F	1912-2000	1956-76F constant M	Density-dependent pup and juvenile survival.
Swartzman and Huang	2-18M 2-25F	1911-86	1956-76F Subadult male	Entanglement.

MODEL DOCUMENTATION

The model consists of 24 female and 18 male age classes. Populations in each modeled age class are updated by age-specific and sex-specific survival. Sex-specific and year-specific harvests on either (or both) land and sea are also removed from the proper age-sex classes each year. Pup numbers are computed from the adult female population based on age-specific fecundities. Running the model consists of solving a set of differential equations (one for each age-sex class) using the Runge-Kutta method with a time step of 0.25 years. The model is run from 1911 to 1986. The 0.25 time step was chosen to fit with the time period that pups are on land in the Pribilof Islands (3 months), allowing computation of pup survival on land to occur over a single time step. Survival of juveniles from the time they leave land to age 2 is also modeled as a density-dependent factor

based on regression analysis of pup counts and male survival estimates (data provided in Lander 1981). Entanglement rate depends on a year-specific susceptibility (based on the observed proportion of entangled subadult males in the harvest each year), an age-specific relative susceptibility factor, and the ratio of seals entangled in large to seals entangled in small debris parameter. Additionally, there are age-specific and sex-specific entanglement mortality and escapement (from entanglement) rates. Animals entangled in large debris (>0.4 kg) are assumed to die rapidly (at the same rate at which they are entangled).

A mnemonic notation is used to describe model equations (Swartzman and Kaluzny 1987). The first digit of a variable name denotes the variable type, with x used for state variables, k for parameters, g for intermediate variables, and z for driving variables (unaffected by system behavior and read in from a driver data file). The following letters are descriptive mnemonics such as mrt for mortality or n for numbers. Several parameters are numerically subscripted (e.g., k_1) rather than having a mnemonic name. This was done for parameters that were not easily made mnemonic.

The model is a series of differential equations for the rate of change of female and male seals by age class, with entangled seals (in small net debris) separated from unentangled seals.

$$\frac{dxn_{ij}(t)}{dt} = -(gent_i(t) + kmrt_{ij} + gmrtlg_{ij}(t))xn_{ij}(t) + kesc_{ij}xent_{ij}(t) \quad (1)$$

where $j = 1$ for male and 2 for female

i is an age index (1-24 for female; 1-18 for male.
These denote ages 2-25 and 2-19 for females and males,
respectively)

xn_{ij} = number of unentangled seals in age class i of sex j

$gent_i$ = entanglement rate in small debris for age i seals (yr^{-1})

$kmrt_{ij}$ = natural mortality rate for age i sex j seals (yr^{-1})

$gmrtlg_{ij}$ = entanglement rate in large debris for age i sex j seals
(yr^{-1})

$xent_{ij}$ = number of entangled seals in age class i of sex j

$kesc_{ij}$ = rate of escapement from entanglement for age i sex j
seals (yr^{-1}).

Population dynamics of entangled seals are:

$$\begin{aligned} \frac{dxent_{ij}(t)}{dt} = & -(kmrtent_{ij} + kmrt_{ij} + gmrtlg_{ij}(t) + kesc_{ij})xent_{ij}(t) \\ & + gent_i(t)xn_{ij}(t) \end{aligned} \quad (2)$$

where $kmrtent_{ij}$ = entanglement mortality rate for seals entangled in small debris (yr^{-1}).

These equations account for the possibility of entangled seals escaping as reported by Scordino and Fisher (1983), Fowler (1987), and Fowler et al. 1990).

In addition to these continuous time equations there are also discrete time equations that update the population at set times of the year.

- The pupping time (i.e., early July), when pups are produced, ages of the populations are updated, and harvests of subadult males (and pelagic harvests, if any) are taken;
- The time pups leave land, when the density-dependent number of surviving pups is computed.

The number of pups produced in any year are computed from age-specific fecundity

$$xpup(t) = \sum_{i=1}^{24} kfec_i xn_{i2}(t) \quad (3)$$

where $kfec_i$ = age-specific fecundity including the influence of age-specific maturity.

At the time pupping occurs, the model updates time, ages the seals by 1 year, and removes seals by harvest for that year (harvest includes the subadult male commercial harvest, any harvest of females, and any scientific samples taken that year).

$$xn_{ij}(t+1) = xn_{(i-1)j}(t) - zharv_{ij}(t+1) \quad (4)$$

where $zharv_{ij}(t)$ = the total harvest and samples of age-sex class ij in year t (data entry).

Harvests are most commonly applied to the annual subadult males on the Pribilof Islands, but involved females between 1956 and 1968 and research samples including a variety of age-sex classes in many years. Analogous harvest equations exist for entangled male and female animals (there is a data file for entangled seals as well as for unentangled seals from the harvest statistics).

The survival of 1-year-old and 2-year-old seals is treated somewhat differently from the survival of older seals. The natural mortality of seals between age 3 months (the time pups leave land) and age 2 is computed by the density-dependent function

$$gmrt_{mj}(t) = kfmrt_j \frac{-\log_e \left[\frac{ks_1(xpup(t-m+0.25) - ks_2)}{(xpup(t-m+0.25) - ks_3)} \right]}{1.75} \quad (5)$$

Here m denotes age class (1 or 2) and j sex class. The 0.25 year (3 months) adjusts time back to the time pups leave land. The ratio of female to male mortality rates is $kfmrt_2$, and $kfmrt_1 = 1$. The total number of pups leaving land (male + female) in the year for which we are computing mortality rate ($m = 1$) or in the previous year ($m = 2$) is $xpup(t-m+0.25)$. Instead of $kmrt_{mj}$, $gmrt_{mj}(t)$ is used in equations (1) and (2) for age classes 1 and 2 to denote that these are intermediate variables rather than parameters. The natural logarithm and 1.75 are used to convert the fraction of seals surviving to age 2 (excluding entanglement) to a rate. A mortality rate must be used instead of a fraction surviving (which is what we estimate from the primary data source) because entanglement mortality may also be incurred by these younger seals. We noted earlier that the model considers age classes beginning with age 2 seals. As such, the above computations for age class 1 seals are not included in the part of the code that deals with the seal age classes, but as a separate calculation. Age class 1 animals are excluded from the model because very little is known about survival rates of pups after they leave land and estimates are based solely on the male juveniles that begin showing up in the Pribilofs at age 2.

At the time pups leave land (at 3 months of age), the model computes the number of pups leaving according to a density-dependent function (Swartzman 1984) and divides them into male and female groups assuming a 1:1 sex ratio.

$$xpup_j(t+0.25) = \frac{1}{2}k_1(1.0 - k_2e^{-k_3xpup(t)}) \quad (6)$$

Parameters are k_1 , k_2 , and k_3 in this density-dependent relationship. The seal entanglement rate is assumed to be age, sex, and time specific. A year-specific driving variable, $zprop(t)$, the proportion of entangled subadult males observed in the harvest, is multiplied by an age-specific and sex-specific variable:

$$gent_i(t) = zprop(t)k_4 \cdot e^{-k_5 \cdot i} \quad (7)$$

Here k_4 is the ratio of entanglement rate for pups to the proportion of subadult males in the harvest entangled in small debris, and k_5 is a parameter controlling the age susceptibility of seals to entanglement. Entanglement is represented as an exponential function of age, with youngest seals being most susceptible. The parameter k_5 controls the rate of decline of entanglement susceptibility with age. Setting $k_5 = 0.0$ makes all ages equally susceptible. As a way of simplifying sensitivity analysis, this function was used to represent age changes in susceptibility to entanglement by a single parameter, rather than a vector of parameters. The entanglement rate of age i seals in large debris is equal to $k_6 \cdot gent_i$. The model assumes that seals entangled in large debris die rapidly enough

for the mortality rate to be equal to the entanglement rate. Thus, the entanglement rate in large debris is equated to a mortality rate $gmrtlg_{ij}$ (as shown in equations 1 and 2). The parameter k_6 is the ratio of entanglement in large versus small debris.

The mortality rate of seals entangled in small debris $kmrtent_{ij}$ was for convenience also modeled as a function of age and sex. As with entanglement rate, an exponential function was used because it gives flexibility in the change of entanglement mortality with age. The equation is:

$$kmrtent_{ij} = kentl_j e^{-kent2_j \cdot i} \quad (8)$$

Here $kent2_j$ is a sex-specific parameter for changes in the mortality rate of entangled seals with age. It is analogous to k_5 . The mortality rate for age 0 seals (i.e., pups) is $kentl_j$.

Model Parameters

Parameter values used in this model are given in Table 2, along with sources of data. A calibration process was used to improve the fit between the model and data. It consisted of changing selected parameters to produce agreement with pup counts on the Pribilof Islands. During calibration, parameters were constrained to be changed only within "reasonable" limits ("reasonable" depending upon the accuracy of the parameter estimate).

The parameters k_1 , k_2 , and k_3 were estimated using a nonlinear regression based on equation (6) of estimates of pups born against estimates of pups leaving the Pribilof Islands. The regression gave estimates of $k_1 = 1.06 \times 10^6$, $k_2 = 1.007$, and $k_3 = 1.04 \times 10^{-6}$. When the model was run with best estimates for these and other parameters (see Table 2), the fit from 1911 to 1950, the period of population growth, was very poor. We had ascertained earlier (Swartzman 1984) that the population behavior was very sensitive to these parameters (i.e., k_1 , k_2 , and k_3). This being so, we used bootstrap resampling to obtain estimates of the variance of each parameter (by redoing the regression with different resamples) and then "searched" the parameter space (1,000 Monte Carlo runs) to see which combinations of parameter values provided the best fit to the data during the population growth period. This experiment produced the values for k_1 , k_2 , and k_3 given in Table 2.

Model-Data Comparison

Our initial desire for this model was to have it replicate the female fur seal population abundance. Any model unable to do that must be judged insufficient for investigating the effect of entanglement on fur seal population dynamics. Figure 1 compares the model to pup numbers and bulls (for the model this includes all bulls 7 years or older), which are the only long-term data available. The vertical dashed lines in Figure 1 show (from left to right) the year pelagic sampling of females began, the year entanglement began, the year commercial pelagic harvest ended, and the year all female sampling ended (both scientific sampling and commercial harvests).

Table 2.--Parameter values and data sources for northern fur seal population dynamics model.

Parameter	Definition	Value	Estimate source
k_1	Maximum land pup survival	728451	
k_2	Density-dependent land pup survival	0.982037	Regression; Lander (1981) calibration.
k_3	Density-dependent land survival exponent	1.609502e-6	Regression; Lander (1981) calibration.
k_4	Ratio of age 0 seal entanglement to fraction of entangled subadult males		Regression; Lander (1981) calibration.
k_5	Change of entanglement with age	5	Fowler (1982) calibration.
k_6	Ratio of large to small net entanglement rates	0.35	Calibration.
$kent1_1$	Entanglement mortality rate for younger males	15	Fowler (1984) calibration.
$kent1_2$	Entanglement mortality rate for younger females	0.8	Fowler (1982).
$kent2_1$	Entanglement mortality age effect for males	0.8	Fowler (1982).
$kent2_2$	Entanglement mortality age effect for females	0.35	Calibration.
$kfmrt_2$	Ratio of young female to male mortality rate	0.35	Calibration.
ks_1	Density-dependent survival to age 2	0.6	Chapman (1964) calibration.
ks_2	Density-dependent sea survival parameter	0.5428	Regression; Lander (1981).
ks_3	Density-dependent sea survival parameter	0.7643	Regression; Lander (1981).
		0.7372	Regression; Lander (1981).
Male seals (age-class-specific)			
$kmrt_{11}$	Seal mortality rate	--	Lander (1981).
$kesc_{11}$	Age-specific escapement from entanglement	0.0008	Calibration.
Female seals (age-class-specific)			
$kmrt_{12}$	Seal mortality rate	--	Lander (1981).
$kesc_{12}$	Age-specific escapement from entanglement	0.0008	Calibration.
$kfec_1$	Age-specific fecundity	--	York and Hartley (1981).





Figure 1.--Model-data comparison between pup and bull counts, 1911-86.




Figure 2.--Sensitivity of 1986 pup counts to entanglement parameters.

Although the fit to the pups appears credible, the bull counts are overestimated by the model. Through many runs of the model adjusting male parameters (the female population is unaffected by these changes), it became apparent that in order for the model not to seriously underestimate the number of bulls during the late 1950's and early 1960's (a period of large subadult male harvests), bull counts in the model during the earlier period need to be significantly higher than reported counts. The possibility that the model was in error was minimized by our checking the calculations and also observing that the bull counts do not appear to respond to marked annual changes in harvests, especially during the 1940's. Possible explanations for this model-data disparity (remember that during this period the model appears to fit pup counts well) are that (1) the actual number of bulls is much higher than the number of territorial bull counted both with females (harem bulls) and as "idle bulls," (2) the pup estimates during this period are in error and pup numbers were actually significantly higher than those obtained by the tagging estimates made in the 1950's (Chapman 1973), or (3) that estimated survival parameters for males are in error.

If the first alternative is true, then many mature bulls, especially the younger ones (e.g., ages 7 and 8), are at sea much of the time and do not show up in the bull counts. If the model's bull predictions are to be believed, there must have been a very large pool of idle bulls in some years that spent either little or no time on land in the Pribilof Islands. Furthermore, the size of this pool has changed over time, being large in the 1930's and 1940's, small around 1960, large again in late 1960's, and now being drastically reduced.

SIMULATING THE EFFECT OF ENTANGLEMENT ON FUR SEAL POPULATIONS

The fit between model and data in Figure 1 was based on a calibration, where entanglement parameters were selected to best fit the population trajectory for pups after 1960. As recorded in Table 2, several of these entanglement parameters are based on limited data and others are simply based on achieving a fit of the model to the data. This is true of the age specificity of both entanglement rate and mortality rate of entangled seals, which has not been studied in the field. The ratio of the rate of seals entangled in large versus small debris is based only on the relative incidence of these two kinds of debris in land and pelagic surveys (Fowler 1987). Also, no studies have been devised to estimate susceptibility of seals to entanglement in debris. Therefore, it is to be expected that our uncertainty about the values of these parameters is great.

To investigate the sensitivity of model predictions of pups and adult males in 1986 (chosen as a measure of model performance that directly relates to the effect of entanglement) to changes in entanglement-related parameters, we performed a Monte Carlo sensitivity experiment. Parameter values were sampled from a uniform distribution over the range of values judged to be reasonable (within our expectation of what the parameter values may be). Our choice is, of course, somewhat subjective. It is to be expected that our range of acceptable parameter values will narrow considerably as a result of this experiment. This method of using a sensi-

tivity study to narrow the tolerance limits on parameter values by choosing combinations giving realistic model behavior was devised by Hornberger and Spear (1981).

Because we are primarily interested in sensitivity of the seal population to entanglement, we restricted our sensitivity study to parameters directly related to entanglement. These include parameters k_4 - k_6 , relating to the entanglement rate in both large and small debris and the change over age in susceptibility to entanglement; $kent1_j$ and $kent2_j$, the entanglement mortality rates for males and females and how they change with age; and $kesc_{ij}$, the age-specific and sex-specific rates of escape from entanglement. The pup and bull populations predicted by the model in 1986 were used as an output variable for comparing sensitivity runs. From Figure 1 it is seen that the population consistently declines after 1960, the year entanglement mortality begins to take effect, and therefore the 1986 value is a measure of the degree of decline (all sensitivity runs are at the same population level in 1960 because they differ only in entanglement-related parameters).

Initial results indicated that the escape-from-entanglement parameters $kesc_i$ are significantly less influential than the other parameters. Therefore the Monte Carlo runs were restricted to the other seven parameters. Table 3 gives the values and ranges used for each parameter in the sensitivity study. Due to uncertainty about the parameter values, we chose to sample parameter values from uniform distributions.

Ranges of parameters were set as follows: $kent1_j$, the age 0 small debris entanglement mortality rate, was set to a range of 10% on either side of the baseline run estimate value of 0.6. Considerable effort has been devoted to estimating the mortality rate of entangled seals, both through observation of marked entangled animals and by looking at the age distribution of entangled versus unentangled seals in the subadult male harvest (Fowler 1987). As such, a modest range of variability was assumed. Three parameters control the age distribution of entanglement effects. Parameter $kent2_j$ is the exponent controlling the age distribution of entanglement mortality for male ($j = 1$) and female ($j = 2$) seals, and k_5 is the same for entanglement rate (no sex distinction here). Baseline estimate for each of these parameters was 0.35. Little is known about how susceptibility to entanglement changes with age, except that significantly more young seals are observed entangled in debris on the Pribilof Islands. Having k_5 of 0.35 makes 0-age seals 20 times as susceptible to entanglement as 8-year-olds. A range of 0 (no difference in age susceptibility) to 0.5 (ratio of 55 in age 0 to age 8 susceptibility) seemed adequate to cover the plausible range of values. For $kent2$ and $kent4$ a wider range, from -0.1 (older animals die more rapidly when entangled) to 0.7, was chosen, reflecting our having no data on how long animals at different ages survive when entangled.

The last two sensitivity parameters, k_4 and k_6 , are not well known. For k_6 , the ratio of entanglement rates in large to small debris, Fowler (1984) estimated a value of 5 based on the ratio of large to small debris in beach surveys on Amchitka (Merrell 1980) and the Pribilof Islands.

Table 3.--Parameter values and ranges for sensitivity analysis.

Parameter	Monte Carlo distribution	Best estimate
$kent1_1$	U(0.54, 0.66)	0.6
$kent2_1$	U(-0.1, 0.7)	0.35
$kent1_2$	U(0.54, 0.66)	0.6
$kent2_2$	U(-0.1, 0.7)	0.35
k_4	U(4, 10)	5
k_6	U(4, 15)	15

However, our baseline estimate, which resulted in a reasonable model-data fit, was 15. We therefore chose a range of 4 to 15, putting our estimate at the high end in deference to Fowler's more data-based measure. Parameter k_4 represents the ratio of entanglement rate in small debris for 0-age seals to the fraction of observed subadult male seals entangled on the Pribilof Islands. The latter is the only data-based time series on annual entanglement available. Estimating k_4 is like trying to assess the size of an iceberg from the part above water. There is a lot unknown below the surface. For a range of values, we blanketed our baseline estimate, 5, by 4 and 10, the relatively high lower bound being due to results of preliminary experiments with the model that showed low values of k_4 leading to an overprediction of pup abundance (too weak an effect of entanglement). One caveat of the calibration approach to parameter estimation in this case is that we are assuming entanglement to be the sole cause of the additional mortality since 1960. If, in fact, there are other yet-undiscovered causes, then entanglement parameter values estimated here would be biased. This is to be borne in mind during the discussion of the sensitivity analysis, which examines parameter ranges that lead to realistic behavior, assuming that all sources of mortality are accounted for in the model (either through harvest, sampling, entanglement, or natural mortality, or through density-dependent juvenile survival).

SENSITIVITY ANALYSIS RESULTS

Figure 2 shows results of the sensitivity study for six of the parameters plotted against pup numbers. Results are omitted for $kent1_1$, which is similar to $kent1_2$. A smooth using supersmoother (Friedman and Stuetzle 1982) was fit to each plot. The three entanglement susceptibility parameters appear to have a stronger effect on 1986 pup numbers than the entanglement mortality parameters. Low values of k_5 (no age-specific or weak age-specific entanglement susceptibility) appear to lead to low 1986 pup predictions.

To get a sense of which parameter combinations led to realistic predictions, we extracted those runs (of the 1,000 runs made) that gave 1986 pup estimates between 170,000 and 200,000 (the baseline run gave 189,000 pups in 1986, close to the pup count estimate for that year).

Figure 3 illustrates which combinations of parameters give realistic model behavior. The first three plots show the entanglement rate parameters against each other for pup numbers within this range, including a super-smooth fit of the resulting scatterplots. The next three plots show each of these parameters' values against pup numbers (for the latter restricted between 170,000 and 200,000 pups). These indicate that over the range of "realistic" pup numbers, none of these parameters has a significant effect on pup numbers (there is no significant slope to the smooths). From the first three plots we deduce that k_4 and k_5 are inversely related to each other, and k_5 and k_6 are inversely related to each other. This can be interpreted to mean that there cannot be a low value of k_5 and a high value of k_4 (and contrariwise a high value of k_5 precludes a low value of k_4). This information is useful because it sets limits of parameter combinations leading to realistic behavior. Furthermore, if more information becomes available concerning any one of these parameters it further delimits the possible values of the other parameters. For example, if observation of entangled animals at sea would indicate that seals are more susceptible to entanglement in large debris than the ratio of large to small debris in beach surveys would indicate (implying a value of k_6 near the upper end of the 4 to 15 range used here), then the entanglement rate needs to be age-specific, with older animals significantly less susceptible to entanglement than younger animals. Significant benefit for research direction can be derived from these results, because they suggest that improved estimates of entanglement rates in small debris can be obtained by seemingly unrelated (and potentially less expensive) than the studies such as finding out about age-related susceptibility to entanglement.

Another interesting result of this sensitivity study is that entanglement rates are much more important for population survival than is entanglement mortality. Another way of saying this is that the rate at which seals enter the entangled animal pool is more important to long-term population trends than the rate at which they die once they are in it. Assuming mortality rate much larger than the rate of escape from entanglement assures that most animals entering the entangled pool will die before they can leave their mark on future generations through reproduction.

ENTANGLEMENT QUESTIONS AND RESEARCH

The model can be used to explore recovery scenarios such as how the population would respond to removing entanglement or reducing it. However, such an exercise is unnecessary. Except for short-term effects, the response to removing entanglement can be observed in Figure 1 in the model's pup counts during the upward cycle starting around 1923. Intermediate entanglement rates would result in less rapid recovery rates. The actual rate of recovery depends on the specific entanglement rates and combinations of parameters. These include the entanglement parameters (k_4 - k_6) as well as the non-entanglement-related parameters that were calibrated to fit time traces from 1915 to 1960 pup counts.

More important than recovery scenarios are dominant questions suggested by the model concerning entanglement. These are:




Figure 3.--Parameter interaction for 1986 pups and bulls close to nominal run levels.

1. What is the rate of entanglement in small debris and how is this influenced by debris abundance and distribution?
2. Are there age differences in susceptibility to entanglement and if so how can we measure them?
3. How relatively susceptible are seals to entanglement in large versus small debris?

This modeling exercise has demonstrated that, within the degree of uncertainty that we can answer the above three questions, entanglement is a plausible explanation for the decline in Pribilof Island fur seal populations since the late 1960's. The model's representation of male abundance has raised some questions about using the idle and harem bull counts as an index of total adult male abundance. At the very least, large fluctuations in male harvest are not reflected by subsequent appropriately large changes in bull counts. At the most, a variable, potentially large fraction of the mature males either may not be resident on the Pribilof Islands during the summer or may be resident for only part of the summer. Finally, the modeling has defined the research questions that can help reduce uncertainty about the possible past and future effects of entanglement on seal populations. The first question, about entanglement susceptibility, requires increased (and preferably simultaneous) observation of entangled seals and debris at sea and the development of a debris encounter probability estimate and an estimate of the probability of a seal's being entangled given that it has encountered debris (Ribic and Swartzman 1990). The second question requires taking a closer look at the age-sex distribution of entangled seals on land and perhaps conducting tank experiments on a larger scale than previously done. The third question requires observation of entangled animals at sea and development of statistical methods for estimation based on very infrequent encounters. Research around both the first and third questions may benefit from additional models designed to test various assumptions made in doing the estimates. For example, a Monte Carlo seal-debris encounter model, coupled with further transect observations might help clarify what the probability of entanglement is, given an encounter at sea.

ACKNOWLEDGMENT

This work was supported by the National Marine Mammal Laboratory under Contract No. 40-ABNF-803196. The authors wish to thank Charles W. Fowler for his support and especially for his strong interest and input to this work--with data, with patience, with encouragement, with ideas, and with sound critique.

REFERENCES

- Chapman, D. G.
1964. A critical study of Pribilof fur seal population estimates.
U.S. Fish. Wildl. Serv., Fish. Bull. 63:657-669.

1973. Spawner-recruit models and estimation of the levels of maximum sustainable catch. *Rapp. P.-V. Réun. Cons. Int. Explor. Mer* 164:325-332.
- Eberhardt, L.
1981. Population dynamics of the Pribilof fur seals. In C. W. Fowler and T. Smith (editors), *Dynamics of large mammal populations*, p. 197-220. Wiley, N.Y.
- Fowler, C. W.
1982. Interactions of northern fur seals and commercial fisheries. *Trans. N. Am. Wildl. Natl. Resour. Conf.* 47:278-292.

1984. Density dependence in northern fur seals (*Callorhinus ursinus*) of the Pribilof Islands, Alaska. Paper presented at the International Symposium and Workshop on Fur Seal Biology, 23-27 April 1984, Cambridge, England.

1985. An elevation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. *Mar. Pollut. Bull.* 18:326-335.
- Fowler, C. W., R. Merrick, and J. D. Baker.
1990. Studies of the population level effects of entanglement on northern fur seals. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Friedman, J. H., and W. Stuetzle.
1982. Smoothing of scatterplots. *Project Orion Tech. Rep.* 3, Dep. State Stanford Univ. Stanford, Calif., 47 p.
- Hornberger, G. M., and R. C. Spear.
1981. An approach to the preliminary analysis of environmental systems. *J. Environ. manage.* 12:7-18.
- Lander, R. H.
1981. A life table and biomass estimate for Alaskan fur seals. *Fish. Res.* 1:55-70.
- Merrell, T. R.
1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. *Mar. Environ. Res.* 3:171-184.

Reed, M., and D. French.

1987. Simulation modeling of the effects of oil spills on population dynamics of northern fur seals. Technical report to Minerals Management Service MMS 86-0045, 142 p. + appendixes.

Ribic, C. A., and G. L. Swartzman.

1990. An index of fur seal entanglement in floating net fragments. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Scordino, J., and R. Fisher.

1983. Investigations on fur seal entanglement in net fragments, plastic bands and other debris in 1981 and 1982, St. Paul Island, Alaska. Background paper submitted to the 26th Annual Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 28 March-8 April, Wash., D.C.

Smith, T. D., and T. Polacheck.

1981. Reexamination of the life table for northern fur seals with implications about population regulatory mechanisms. In C. W. Fowler and T. Smith (editors), Dynamics of large mammal populations. Wiley, N.Y., 477 p.

Swartzman, G. L.

1984. Present and future potential models for examining the effect of fisheries on marine mammal populations in the eastern Bering Sea. In Melteff and Rosenberg (editors), Proceedings of the Workshop on Biological Interactions Among Marine Mammals and Commercial Fisheries in the Southeastern Bering Sea. Alaska Sea Grant Report 84-1, Univ. Alaska.

Swartzman, G. L., and S. P. Kaluzny.

1987. Ecological simulation primer. Macmillan, N.Y., 370 p.

Trites, A. W.

1984. Stock assessment and modeling of the North Pacific fur seal population. Submitted as background paper to the 27th meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 28 March-April 6, 1984, Moscow, U.S.S.R.

York, A. E., and J. R. Hartley.

1981. Pup production following harvest of female northern fur seals. Can. J. Fish. Aquat. Sci. 38:84-90.

HISTOLOGICAL OBSERVATION OF DAMAGE TO DERMAL TISSUE
OF FUR SEAL CAUSED BY NET ENTANGLEMENT

Masayuki Nakajima
Izu-Mito Sea Paradise
3-1 Nagahama, Uchiura
Numazu, Shizuoka, 410-02 Japan

Hiromoto Yasuda
Faculty of Medicine, Teikyo University
2-11-1, Kaga, Itabasi-ku
Tokyo, 173 Japan

and

Kazumoto Yoshida
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu, Shizuoka, 424 Japan

ABSTRACT

In 1984 and 1985, experimental studies on the damage to northern fur seals by net entanglement were carried out in Izu-Mito Sea Paradise, an aquarium where fur seals are kept for the National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan (formerly the Far Seas Fisheries Research Laboratory, Fisheries Agency of Japan).

Two adult female fur seals, captured off Sanriku, northern Japan, in March 1980, were experimentally entangled in fishing net fragments in late January 1984. One seal, No. 1-7, 123 cm in body length and 39.5 kg in body weight, and an estimated 10 years of age, was entangled around the neck with a net fragment weighing 200 g. The second seal, No. 1-8, 113 cm in body length and 36.0 kg in body weight, with an estimated age of 10 years, was entangled with net fragments 300 g in weight. Both seals were kept in the same environment.

Seal No. 1-8 died in September 1984 after 226 days of entanglement, having suffered traumatic damage to pelage and skin. Net fragments were removed from No. 1-7 in March 1985. Damage to skin and pelage was not observed even after 14 months on entanglement, although the state of entanglement was similar to No. 1-8. She died in February 1986. Abnormality was not observed in the skin at the time of death. Cause of death was acute pneumonia in both cases. When they were alive, No. 1-7 was in good health and No. 1-8 was slightly unwell.

INTRODUCTION

Fur seal are known to have died from entanglement in fishing net fragments and packing bands (Scordino 1985). It was pointed out that net entanglement may constitute one of the major factors for the decrease in the Pribilof Islands populations of the northern fur seal, *Callorhinus ursinus*. In order to understand the actual state and effects of entanglement, the National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan (formerly the Far Seas Fisheries Research Laboratory, Fisheries Agency of Japan) conducted field surveys of fur seals and marine debris on breeding islands and at sea. Also, experimental studies were carried out concerning the process of entangling and the effects of entanglement on the behavior of fur seals, both of which were difficult to observe in a natural environment. Little has yet been reported about traumatic wounds to the skin of fur seals caused by net entanglement. In this study, macroscopic observation was conducted on damage inflicted by net entanglement to the pelage and skin of fur seals kept in captivity. Post-mortem histological examination was also made of the lesions of the dermal tissues.

MATERIALS AND METHODS

The experiment was conducted for about 14 months from 10 January 1984 to 31 March 1985 in a breeding facility of the Izu-Mito Sea Paradise, an aquarium in Numazu, central Japan. Two female northern fur seals, with identification numbers 1-7 and 1-8 and an estimated age of 10 years, were used. These individuals were entangled with net fragments during an experiment on the mechanism of entanglement. Seal No. 1-7 was entangled in trawl net fragments of 200 g on 20 January 1984, and No. 1-8 was entangled in a trawl net fragment weighing 100 g on 28 January 1984 and in another fragment of 200 g on 29 January 1984. Nets were removed from No. 1-7 at the end of the experiment on 31 March 1985. Behavior of the two seals and damage to pelage were observed every day in the morning and evening. The net fragments used in the experiment were commercial trawl nets made of polyethylene, with a twine size of 3.4 mm and a mesh size of 24 cm. The nets were cut into 100- and 200-g pieces.

The two individuals were kept in an open breeding facility (Fig. 1) from 10 January to 10 March 1984; afterward they were brought to an indoor breeding pool 1.6 m wide, 2.4 m long, and 1.0 m deep. No landing place was provided in the pool, in order to prevent resting on land except for the breeding season, in imitation of pelagic life. The experimental animals were fed with defrosted mackerel each day at 1000 and 1630. Average daily food consumption of No. 1-7 was 3.8 kg/day (0-5.5 kg/day) and that of No. 1-8 was 2.8 kg/day (0.4-4.0 kg/day). Body weight of No. 1-7 remained almost constant during the entire experiment period, while that of No. 1-8 declined near the time of death.

Necropsy was conducted immediately after the death of the entangled animals for macroscopic and histological inspection of the skin lesions. Histological samples of dermal tissue were fixed with 20% formaldehyde, embedded in paraffin, and cut into sections 4 microns thick with a sliding




Figure 1.--The breeding facilities at Izu-Mito Sea Paradise.

microtome. The sections were dyed using hematoxylin eosin staining or Masson's method.

Atmospheric temperature during the period ranged from -5° to 30.1°C , water temperature ranged from 12.0° to 28.3°C , and relative humidity ranged from 33 to 92%. Details of the fur seals used in the experiment are given in Table 1.

RESULTS

Behavior of No. 1-7 and Description of Net Entanglement

This individual was in quite good health. On 20 January 1984, she was entangled around the neck by a bundle of net made into a collar. There was a space between the net collar and the neck into which one finger could be inserted. No damage to the entangled part was observed even after 14 months. Net fragments were removed at the end of the experiment on 31 March 1984. The individual remained in good health with a good appetite, finally dying on 25 February 1986, 11 months after the removal of the nets. Post-mortem examination revealed small whitish nodules about 5 mm in size scattered throughout the lungs. No pus was found in the nodules. The cause of death was diagnosed as acute pneumonia. The cervical region where the net had been entangled was also inspected, but no anomaly was recognized.

Table 1.--Details of the two female northern fur seals used in this study.

Chronology of the experiment	Seal No. 1-7	Seal No. 1-8
Capture		
Date	4 March 1980	7 March 1980
Location	Lat. 36°40'N long. 141°27'E	Lat. 36°30'N long. 141°15'E
Transport to aquarium		
Date	8 March 1980	8 March 1980
Body weight (kg)	29.0	30.5
Beginning of experiment		
Date	10 January 1984	10 January 1984
Estimated age	10	10
Body length (cm)	123	113
Body weight (kg)	39.5	36.0
Net entanglement		
Starting date	20 January 1984	28 January 1984
Weight of nets (g)	200	100 29 January 1984 200
Date of removal	31 March 1985 ^a	10 September 1984 ^b
Death		
Date	25 February 1986	10 September 1984
Body length (cm)	120.5	114.5
Body weight (kg)	34.5	30.5
Cause of death	Acute pneumonia	Acute pneumonia

^aExperiment ended.

^bSeal died.

Behavior of No. 1-8 and Description of Net Entanglement

Judging from daily behavior and feeding activities, this individual was considered in somewhat poor health even before it was entangled in nets. Two net fragments were placed around its neck on 28 and 29 January 1984.

The entangled nets made up a collar consisting of 17 meshes in a bundle (Figs. 2 and 3). The collar was 14 cm in inner diameters, 44 cm in inner circumference, 100 cm in outer circumference, and 7 to 9 cm thick. No remarkable change in behavior was observed after net entanglement. On the morning of 6 June 1984, she delivered a female pup, 65 cm long and 4.2 kg in weight, but the pup was dead when it was discovered. Anatomical inspection revealed the cause of the pup's death to be drowning. Seal No.




Figure 2.--State of entanglement of fur seal No. 1-8.




Figure 3.--Collar of entangled net on fur seal No. 1-8.

1-8 ate poorly from 4 September 1984 and died on 10 September, 226 days (7.5 months) after the initiation of net entanglement. Macroscopic and histological observations of the entangled part are described below.

Diagnosis of the Lesions on No. 1-8

The nutritive condition of the body as a whole was moderate. No external scars were observed except around the neck. In macroscopic observation, hair was worn or lost all around the neck, and the epidermis was exposed in some parts. Several wounds due to abrasion by the attached nets were conspicuous: one large scar 1-3 cm wide and 5 cm long on the right side of the neck and two other wounds of 1-2 cm at the scruff. Cross sections of the lesions revealed hardened hyperdermis and a thin muscle layer. The thick adipose tissue consisted of yellow and white parts (Figs. 4 and 5).

In the histological examination, sections of the injured dermal tissue showed degeneration and loss of hair follicles and hair matrices, and degeneration of the hair itself. Connective tissues were partially worn out and necrotized. Proliferation of collagen fiber was observed in the peripheral connective tissues. In these parts, degenerated inner membranes, supposedly derived from venous sinus, were also observed. Rupture and degeneration of muscle layers were distinctive and a part of the connective tissues had been replaced. Congestion and edema of venous ducts were conspicuous and their inner membrane revealed degeneration. However, cell infiltration and inflammatory reactions were indistinctive (Figs. 6 and 7).

As for visceral organs, hyperemia was observed in the lungs and accumulation of blood was conspicuous in the heart. Other visceral organs showed no remarkable change. The cause of death was presumed to be acute pneumonia.

DISCUSSION

No injury occurred around the neck of No. 1-7, entangled in 200 g of fishing nets for 14 months. Seal No. 1-8, entangled at the neck in 300 g of nets for 7.5 months, suffered abrasion of hair and skin. These differences seemed to have been caused by such factors as physical condition, and the amount and tightness of the entangled nets. After anatomical inspection, the cause of death for both individuals was diagnosed as acute pneumonia. As a future task, it will be necessary to examine bacterial infections from the wounds caused by entanglement.

As fur seals used in this experiment were adult, no increase in body weight was observed during the experiment period of 7.5 to 14 months. But it can be assumed that when a young, growing animal becomes entangled in fishing nets, even if the net fragment is small and loose at first, it will gradually become tighter as the animal grows, causing serious damage and possibly death.



Figure 4.--Abraded wound in the pelage and skin of No. 1-8.




Figure 5.--Cross section of dermal tissues through the lesion on the neck of No. 1-8.




Figure 6.--Section through a lesion in the cervical region of No. 1-8 (hematoxylin eosin staining $\times 40$).




Figure 7.--Section through a lesion in the cervical region of No. 1-8 (Masson's staining $\times 100$).

ACKNOWLEDGMENTS

Deep appreciation is expressed to all who cooperated in the research, especially to N. Baba and M. Kiyota of the National Research Institute of the Far Seas Fisheries, the staff of the Clinical Inspection Center of Numazu Ishikai Hospital, and breeding technicians of Izu-Mito Sea Paradise. Thanks are also due to the Fishing Ground Preservation Division, the Fisheries Agency of the Government of Japan.

REFERENCES

- Scordino, J.
1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

RECENT ENTANGLEMENTS OF HAWAIIAN
MONK SEALS IN MARINE DEBRIS

John R. Henderson
Southwest Fisheries Science Center Honolulu Laboratory
National Marine Fisheries Service, NOAA
Honolulu, Hawaii 96822, U.S.A.

ABSTRACT

During field studies on the Hawaiian monk seal, *Monachus schauinslandi*, in 1985-88, 34 incidents of entanglement in marine debris were observed, including 4 known deaths, injuries to 4 seals, and recent neck scars on 2 seals. The overall entanglement rate increased eightfold, from 0.06 incidents per 100 camp days per 100 seals in 1985 to 0.48 incidents per 100 camp days per 100 seals in 1988. This increase was probably caused by increased amounts of marine debris on and around the islands where seals haul out. Weaned pups were entangled at a higher rate than their proportion in the population, while adults were entangled at a lower rate. Entanglement rates since 1981, when corrected for each island's population size, were highest at Lisianski Island: 4.44 incidents per 100 camp days per 100 seals. Lowest rates were at French Frigate Shoals: 0.37 incidents per 100 camp days per 100 seals.

INTRODUCTION

In the past 25 years, durable and resilient plastic materials have replaced natural fibers in the maritime industry. Polypropylene and nylon nets have replaced antiquated and once prevalent tarred cotton webbing, and various plastic lines are now used in place of manila or other natural hemp fiber (Pruter 1987). This use of persistent plastics has been accompanied by an increase in the impact of lost or discarded materials on wildlife in the marine environment.

Pinnipeds in particular are susceptible to entanglement in marine debris. Entanglements of the northern fur seal, *Callorhinus ursinus*, are well documented (Fowler 1982; Scordino and Fisher 1983; Scordino 1985) and appear to have contributed to a population decline in this species during 1976-81 (Fowler 1985, 1987). Although other pinnipeds may be entangled less often, the list of species known to have become entangled is large: Fowler (1988) recently stated that 16 of the 34 extant pinniped species (47%) are known to have become entangled in marine debris.

Entanglements of the Hawaiian monk seal, *Monachus schauinslandi*, in marine debris have been observed since 1974 (Balazs 1979; Andre and Ittner 1980; Alcorn 1984; Henderson 1984, 1985). Pups are particularly susceptible (Henderson 1985). Henderson (1985) documented 35 incidents of Hawaiian monk seals entangled in debris through 1984, noting that the number of observed incidents declined following the inception of a program to periodically remove hazardous debris from haul-out beaches at all Northwestern Hawaiian Islands (NWHI). Entanglements have nonetheless continued since 1984 (Alcorn et al. 1988; Johanos and Austin 1988; Johanos and Withrow 1988; Reddy and Griffith 1988; Westlake and Siepmann 1988; Henderson and Finnegan 1990). This report summarizes all of the published and unpublished reports of Hawaiian monk seal entanglements in 1985-88, thereby updating Henderson (1985). This report also examines all entanglements since 1982 for trends in number and location of occurrences, and sizes of affected animals.

METHODS

Staff biologists of the Marine Mammals and Endangered Species Program (MMESP) of the Southwest Fisheries Science Center Honolulu Laboratory, National Marine Fisheries Service, NOAA, conducted field operations from 1982 to 1988 in the NWHI to monitor the Hawaiian monk seal population. Since 1982, most Hawaiian Islands west of Necker Island (i.e., French Frigate Shoals, Laysan Island, Lisianski Island, Pearl and Hermes Reef, Midway, and Kure Atoll) have been visited and included as study sites. The number of field camp days has varied among the six locations, as has the annual total.

During these studies, all occurrences of entangled or entanglement-scarred seals were recorded. A seal was considered entangled if any part of its body was encircled by debris. Seals resting or asleep on netting or lines were not considered entangled unless their head or body was inside a loop and field personnel thought the animal would not be able to free itself. Seals with entanglement scars, which are distinguishable from scars resulting from other injuries (Henderson 1985), were documented on scar cards and tallied only in the year in which they were first documented to have acquired the scar. Seals with scars were assumed to have become entangled at the island where first observed.

Data on the number of entangled seals per year and per location were converted to number of occurrences per 100 camp days to account for any variation in the length of the field seasons, and incidents recorded while MMESP personnel were absent from an island have been excluded from most analyses. Two such incidents [documented by the U.S. Fish and Wildlife Service (FWS) and the U.S. Coast Guard] occurred in 1985-88 and have been included only in the overall listing and in data on the size of seals that become entangled.

Data prior to 1985 are from Henderson (1985) except for two entanglements in 1984 at Pearl and Hermes Reef. These were inadvertently omitted by Henderson (1985), and are included here:

<u>Seal</u>	<u>Body part</u>	<u>Type of material</u>	<u>Fate of seal</u>	<u>Comments</u>
Juvenile male	Neck	Plastic ring	Unknown	North Is.
Weaned female	Neck, foreflipper	Net	Unknown	North Is.

The relative number of incidents may be affected by the population size of seals, with more entanglements likely among larger populations. Each island's population is relatively discrete and may differ in size from another island's (Johnson et al. 1982). Furthermore, the total population, as indexed by beach counts of nonpups, increased by approximately 24% in 1983-87 (Gilmartin 1988). In the analysis among islands, the number of incidents at any one location was therefore divided by that island's mean 1983-88 beach count of pups and nonpups. To obtain annual entanglement rates, the entanglement total for each year was divided by the summed mean annual beach counts of pups and nonpups at all islands. Data on mean beach counts are incomplete for 1982; therefore, incidents occurring in 1982 were divided by data from 1983. No total beach counts were collected at Lisianski Island in 1988; hence, the 1988 total beach count includes the 1987 total for Lisianski Island.

Because pups have historically been more susceptible to entanglement (Henderson 1985), a separate analysis divided total incidents per year for 1982-88 by the number of pups known to have been born.

RESULTS

Number of Entanglements, 1985-88

Thirty-four entanglements of Hawaiian monk seals occurred from 1985 to 1988 (Table 1). The total included four known deaths, injuries to four animals, and recent neck scars on two seals. The remaining 24 animals were uninjured and were either released from the debris (20) or escaped unassisted (4). Many of the released seals were loosely entangled; it is unknown whether they might have escaped unassisted. Totals by size classes of the affected seals were 12 pups (nursing or weaned), 7 juveniles, 9 subadults, and 6 adults. When data prior to 1985 were included, the number of documented entanglements of Hawaiian monk seals totaled 71.

Entanglements per 100 camp days per 100 seals decreased to a low of 0.06 in 1985 and rose thereafter to a high of 0.48 in 1988, an eightfold increase (Fig. 1). This trend was also evident if incidents were adjusted by the number of pups born annually, with incidents per 100 camp days per 100 pups being 0.23 in 1985 and 1.34 in 1988 (Fig. 2).

Entanglements by Location, 1982-88

The seal population at Lisianski Island experienced the highest rate of entanglement, at 4.44 incidents per 100 camp days per 100 seals since 1982 (Fig. 3). Kure Atoll was next highest (2.23), followed by Pearl and

Table 1.--Summary of entangled and entanglement-scarred Hawaiian monk seals observed in 1985-88.

Year	Location	Size/sex ^a	Incident ^b	Type of material	Fate of seal	Reference ^c	Comments
1985	Kure Atoll	W/M	E, body	Monofilament line	Rescued	Reddy and Griffith 1988	--
		W/F	E, mouth, body	Fish hook and monofilament line	Rescued	Reddy and Griffith 1988	Slight injury from hook.
	French Frigate Shoals (FFS)	J/M	E, neck	Net	Rescued	L. Martin, FWS, pers. commun.	Would have escaped; mesh broke when pulled, Tern Island.
1986	Lisianski Island (LI)	W/F	E, neck	Net and line	Rescued	Westlake and Siepmann 1988	Slight injury.
		S/M	E, muzzle	Line	Rescued	Westlake and Siepmann 1988	Could have escaped.
	Laysan Island	S/M	E, neck,	Line	Rescued	NMFS unpubl. data	--
	FFS	W/M	E, abdomen	Wire	Found dead	Henderson, observation	Wire probably relic of USCG occupation, East Island.
1987	LI	P or W	E, neck	Net	Found dead	Johanos and Withrow 1988	Uncertain if nursing or weaned pup.
		A/F	S, neck	N/A	Alive in 1988	Johanos and Withrow 1988	Seen without scar in 1986.
		W/F	S, neck	N/A	Alive in 1988	Johanos and Withrow 1988	Acquired scar early in 1987.

Table 1.--Continued.

Year	Location	Size/sex ^a	Incident ^b	Type of material	Fate of seal	Reference ^c	Comments
1987	Laysan Island	J/?	E, body	Net	Rescued	NMFS unpubl. data	Net loose.
		S/F	E, neck	Line	Escaped	NMFS unpubl. data	Molting seal, line not tight.
		A/F	E, neck	Line and floats	Escaped	NMFS unpubl. data	--
		J/M	E, neck, shoulders	Line	Rescued	NMFS unpubl. data	Line loose.
		A/F	E, neck	Line	Escaped	NMFS unpubl. data	--
		J/F	E, neck, abdomen	Net and line	Found dead	Henderson, observation	Shark island.
		S/M	E, abdomen	Line	Rescued	M. Craig, SI, pers. commun.	Line loose, East Island.
1988	Kure Atoll	S/F	E, shoulders	Line	Rescued	M. Craig, SI, pers. commun.	East Island.
		S/?	E, neck	Plastic band	No rescue	NMFS unpubl. data	Band cutting into flesh, Whale-Skate Island.
		A/F	E, neck	Line	Rescued	Henderson and Finegan 1990	Line not tight.
		P/F	E, neck, shoulders	Net and line	Rescued	Henderson and Finegan 1990	On offshore reef, nursing pup, probably would have died.

Table 1.--Continued.

Year	Location	Size/sex ^a	Incident ^b	Type of material	Fate of seal	Reference ^c	Comments
1988	Kure Atoll	W/F	E, neck	Net	Rescued	Henderson and Finnegan 1990	--
		W/F	E, neck	Net	Rescued	Henderson and Finnegan 1990	--
		W/F	E, neck	Net	Rescued	Henderson and Finnegan 1990	On offshore reef, probably would have died, rescued by USCG.
	LI	W/F	E, muzzle	Plastic cup	Rescued	NMFS unpubl. data	--
		W/?	E, body	Net	Found dead	NMFS unpubl. data	Entire body wrapped in large net, decomposed. 545
	Laysan Island	J/M	E, neck	Plastic ring	Rescued	NMFS unpubl. data	Tight but no injury, probably would have been choked.
		S/F	E, neck	Net	Rescued	NMFS unpubl. data	Cutting deeply into flesh.
		S/M	E, body	Net	Rescued	NMFS unpubl. data	Net loose, seal probably would have escaped.
		S/?	E, abdomen	Line	Rescued	NMFS unpubl. data	--
		A/F	E, muzzle	Net	Rescued	NMFS unpubl. data	Net loose.

Table 1.--Continued.

Year	Location	Size/sex ^a	Incident ^b	Type of material	Fate of seal	Reference ^c	Comments
1988	Laysan Island	A/F	E, abdomen	Net	Rescued	NMFS unpubl. data	Same net as previous incident.
		J/F	E, neck	Net	Rescued	L. Hiruki, UA, pers. commun.	--
	FFS	J/F	E, neck	Plastic screen	Rescued	R. Withrow, WA, pers. commun.	Shark Island.

^aP - nursing pup, W = weaned pup, J = juvenile, S = subadult, A = adult, M = male, F = female, and ? = sex unknown.

^bE - entangled; S = scarred.

^cSI = Smithsonian Institution, FWS = U.S. Fish and Wildlife Service, USCG = U.S. Coast Guard, UA = University of Alberta, and WA = Waikiki Aquarium.




Figure 1.--Annual rates (1982-88) of Hawaiian monk seal entanglement, adjusted for camp days and average beach count (pups and nonpups).




Figure 2.--Annual rates (1982-88) of Hawaiian monk seal entanglement, adjusted for camp days and total pup production.




Figure 3.--Location of Hawaiian monk seal entanglement, 1982-88 (Kur = Kure Atoll; P&H = Pearl and Hermes Reef; Lis = Lisianski Island; Lay = Laysan Island; FFS = French Frigate Shoals).

Hermes Reef (1.62), Laysan Island (1.52), and French Frigate Shoals (0.37). One scarred seal has been observed at Midway (Henderson 1985), but MMESP presence there is minimal, beach counts are few, and the seal population is very small (probably 10-20 animals).

Size of Entangled Seals, 1982-88

Pups are most susceptible to entanglement and adults are least susceptible when percent of entanglements is considered in relation to percent of population (Fig. 4). Pups (weaned and nursing) comprise 11.0% of the population (Gerrodette 1985) and yet account for 42.1% of all entanglements from 1982 to 1988. Adults comprise 48.9% of the population and 15.8% of all entanglements. Entanglement rates for juveniles (17.5%) and subadults (24.6%) approximate their population percentages (17.7 and 22.4%, respectively).

Although pups are more susceptible to entanglement, those locations with the most births did not have the most entanglements. Nearly one-third (32.1%) of 1982-88 entanglements occurred at Lisianski Island, whereas only 10.8% of all pups were born there. Conversely, over half (58.8%) of all pups were born at French Frigate Shoals, where 25.0% of the entanglements were documented.




Figure 4.--Size classes of Hawaiian monk seals as percentage of population and percentage of entanglements, 1982-88.

DISCUSSION

The increased incidence of entanglement may result from an increased encounter rate between seals and debris. Such a higher rate could be due to a number of causes: (1) an increase in the seal population, (2) an increase in the relative number of seals in the size class or classes most likely to become entangled, (3) an increased propensity for seals to investigate debris, and (4) an increase in the amount of debris on the islands. Entanglements in 1985-88 have increased more rapidly than any increase in seal population, including the number of pups. In 1985-87, mean beach counts of seals rose approximately 9%, and the number of births also increased by 22% (Gilmartin 1988). However, these increases do not account for the sharp rise in entanglements, which is evident even when adjusted for these factors. No data exist to evaluate whether any recent behavioral changes have occurred among the population to account for increased entanglement.

The amount of debris on beaches in the NWHI has increased in recent years. The number of nets in 1987 alone increased nearly 200% over the 1985-86 average (Henderson unpubl. data). This increase probably has contributed significantly to the rise in seal entanglements. Field biologists routinely remove hazardous debris from the NWHI, an effort credited with reducing seal entanglements (Henderson 1985), yet despite this effort, entanglements have increased. With larger amounts of debris present, more

entanglements may occur (1) during the long periods when personnel are absent from these remote islands, (2) while biologists are present but before the beaches have been cleared, or (3) while seals are at sea.

The higher incidence of entanglement at Lisianski Island may be attributable to spatial coincidence of favorable pupping habitat with areas receiving most of the debris. Pups are born predominately on the island's east side (Johanos and Henderson 1986; Johanos and Kam 1986), which also receives more debris because debris is moved toward the island by trade winds from the northeast (Henderson et al. 1987).

Pups continue to become entangled at a proportionally higher rate than other size classes, a phenomenon that may have several contributing causes (Henderson 1985): (1) entangled pups are most easily observed because they remain near shore for 1-2 months after weaning; (2) pups, unlike older seals, spend proportionally more time in the vicinity of nearshore reefs, which catch and "concentrate" floating debris; (3) weaned pups are not as strong as older seals and are therefore least able to escape from debris; and (4) recently weaned pups are learning to feed and are more likely than nonpups to explore all objects in their novel environment. Bengston et al. (1988) demonstrated experimentally that recently weaned northern fur seal pups readily explore and become entangled in net fragments, and suggested that this behavior could lead to high mortality among fur seals just after weaning.

The periodic presence of biologists in the remote habitat of the Hawaiian monk seal can reduce deaths of seals from entanglement. Of the four mortalities documented here, two could likely have been prevented had personnel been present on the island. Both of these mortalities were at Lisianski Island, a location with sparse coverage by biologists in recent years.

ACKNOWLEDGMENTS

Data were collected under authority of several special use permits and Marine Mammal and Endangered Species permits issued by the FWS and the Protected Resources Division of the National Marine Fisheries Service. I am grateful to FWS staff at the Tern Island station of the Hawaiian Islands National Wildlife Refuge for their logistical support and to the officers and crew of the NOAA ship *Townsend Cromwell* for transporting field personnel. Special thanks are extended to my colleagues in the field--D. Alcorn, S. Austin, B. Becker, M. Brown, B. Choy, M. Craig, R. Forsyth, L. Hiruki, T. Johanos, and R. Morrow--who observed many of the incidents reported here.

REFERENCES

- Alcorn, D. J.
1984. The Hawaiian monk seal on Laysan Island: 1982. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-42, 37 p.
- Alcorn, D. J., R. G. Forsyth, and R. L. Westlake.
1988. Hawaiian monk seal and green turtle research on Lisianski Island, 1984 and 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-120, 22 p.

Andre, J. B., and R. Ittner.

1980. Hawaiian monk seal entangled in fishing net. 'Elepaio 41(6):51.

Balazs, G. H.

1979. Synthetic debris observed on a Hawaiian monk seal. 'Elepaio 40(3):43-44.

Bengston, J. L., C. W. Fowler, H. Kajimura, R. Merrick, K. Yoshida, and S. Nomura.

1988. Fur seal entanglement studies: Juvenile males and newly-weaned pups, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.

Fowler, C. W.

1982. Interactions of northern fur seals and commercial fisheries. Trans. N. Am. Wildl. Nat. Resour. Conf. 47:278-292.

1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals: A case study. Mar. Pollut. Bull. 18:326-335.

1988. A review of seal and sea lion entanglement in marine fishing debris. In D. L. Alverson and J. A. June (editors), Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, 13-16 October 1987, Kailua-Kona, Hawaii, p. 16-63. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.

Gerrodette, T.

1985. Estimating the 1983 population of Hawaiian monk seals from beach counts. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-85-5, 13 p.

Gilmartin, W. G.

1988. The Hawaiian monk seal: Population status and current research activities. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-88-17, 14 p.

Henderson, J. R.

1984. Encounters of Hawaiian monk seals with fishing gear at Lisianski Island, 1982. Mar. Fish. Rev. 46(3):59-61.

1985. A review of Hawaiian monk seal entanglements in marine debris. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 326-335. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Henderson, J. R., S. L. Austin, and M. B. Pillos.
1987. Summary of webbing and net fragments found on Northwestern Hawaiian Islands beaches, 1982-86. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-11, 15 p.
- Henderson, J. R., and M. R. Finnegan.
1990. Population monitoring of the Hawaiian monk seal, *Monachus schauinslandi*, and captive maintenance project at Kure Atoll, 1988. U.F. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-150, 25 p.
- Johanos, T. C., and S. L. Austin.
1988. Hawaiian monk seal population structure, reproduction, and survival on Laysan Island, 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-118, 38 p.
- Johanos, T. C., and J. R. Henderson.
1986. Hawaiian monk seal reproduction and injuries on Lisianski Island, 1982. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-64, 7 p.
- Johanos, T. C., and A. K. H. Kam.
1986. The Hawaiian monk seal on Lisianski Island: 1983. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-58, 37 p.
- Johanos, T. C., and R. P. Withrow.
1988. Hawaiian monk seal and green turtle research on Lisianski Island, 1987. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-121, 18 p.
- Johnson, A. M., R. L. DeLong, C. H. Fiscus, and K. W. Kenyon.
1982. Population status of the Hawaiian monk seal (*Monachus schauinslandi*), 1978. *J. Mammal.* 63:415-421.
- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Reddy, M. L., and C. R. Griffith.
1988. Hawaiian monk seal population monitoring, pup captive maintenance program, and incidental observations of the green turtle at Kure Atoll, 1985. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-101, 35 p.

Scordino, J.

1985. Studies on fur seal entanglement, 1981-84, St. Paul Island, Alaska. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 278-290. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Scordino, J., and R. Fisher.

1983. Investigations on fur seal entanglement in net fragments, plastic bands and other debris in 1981 and 1982, St. Paul Island, Alaska. Background paper submitted to the 26th Annual Meeting of the Standing Scientific Committee of the North Pacific Fur Seal Commission, 28 March-8 April 1983, Washington, D.C., 290 p.

Westlake, R. W., and P. J. Siepmann.

1988. Hawaiian monk seal and green turtle research on Lisianski Island, 1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-119, 18 p.

PINNIPED ENTANGLEMENT IN SYNTHETIC MATERIALS
IN THE SOUTHERN CALIFORNIA BIGHT

Brent S. Stewart and Pamela K. Yochem
Hubbs-Sea World Research Institute
San Diego, California 92109, U.S.A.

ABSTRACT

The California sea lion, *Zalophus californianus*, the northern fur seal, *Callorhinus ursinus*, the harbor seal, *Phoca vitulina richardsi*, and the northern elephant seal, *Mirounga angustirostris*, that haul out or breed on the southern California Channel Islands, become entangled in synthetic debris at various rates. The percentages of California sea lions entangled, primarily in monofilament gillnet fragments, varied from about 0.08% in 1983 to about 0.16% from 1985 through 1988, while those of northern elephant seals, primarily in packing straps, declined from about 0.15% in 1983 to about 0.10% in 1989. The entanglement rate of harbor seals has varied from 0.0% in 1983-84 to 0.06% in 1986. Entangled northern fur seals have rarely been observed.

Inter- and intraspecific differences in entanglement rates are likely the result of age, sex, and species differences in animal size, diving behavior, and foraging areas. Although entanglement in synthetic materials contributes to mortality of some animals, our studies suggest prevailing entanglement rates have not significantly influenced pinniped demography and population trends in the Southern California Bight.

INTRODUCTION AND METHODS

Pollution of marine environments with nonbiodegradable plastic debris has become an issue of increasing concern during the past several years, especially with regard to entanglement of marine mammals and seabirds in synthetic debris (e.g., see Shomura and Yoshida 1985; Wolfe 1987).

Since 1978 we have studied the incidence of entanglement of pinnipeds in synthetic materials at San Nicolas and San Miguel Islands. We have attempted to distinguish between entanglement in floating marine debris and that resulting from direct interactions of seals and sea lions with commercial fishing and sportfishing operations (Stewart and Yochem 1985, 1987).

Below we summarize the results of our surveys made between October 1986 and March 1989, and incorporate our previous surveys to assess trends in pinniped entanglement in the Southern California Bight since 1983.

We quantified rates of entanglement of the California sea lion, *Zalophus californianus*, the harbor seal, *Phoca vitulina richardsi*, the northern elephant seal, *Mirounga angustirostris*, and the northern fur seal, *Callorhinus ursinus*, using methods described earlier (Stewart and Yochem 1985, 1987). Briefly, we surveyed pinniped populations at San Nicolas Island once each month and those at San Miguel Island periodically whenever we visited there to conduct other research. Using binoculars or a spotting telescope, we systematically examined small groups of pinnipeds on rookeries and hauling grounds and recorded the number examined (by sex and relative age whenever possible), the number entangled (and the type of entangling material), and the number scarred (presumably from prior entanglement); only those animals whose bodies could be seen clearly were sampled during those entanglement surveys.

RESULTS AND DISCUSSION

From March 1988 through February 1989, 30 (0.12%) of 24,731 California sea lions surveyed at San Nicolas and San Miguel Islands were entangled and another 25 (0.10%) were scarred from previous entanglement, slightly fewer than from October 1986 through February 1988 (Tables 1, 2). Slightly more northern elephant seals were entangled but slightly fewer scarred in 1988-89 than in 1986-88 (Tables 1, 3). Relatively few (0.03%) harbor seals were entangled in 1988-89 and none was scarred (Tables 1, 4), and we observed no scarred or entangled northern fur seals (Tables 1, 5). Neither of two Guadalupe fur seal bulls that we observed at San Nicolas Island in summer 1988 was entangled or scarred.

The percentage of California sea lions observed entangled increased from 1983 through 1987 but declined in 1988, while the percentages of entangled northern elephant seals and harbor seals have declined since 1984 (Table 1). The percentages of scarred sea lions and elephant seals observed have remained relatively constant since 1983 and 1984, respectively (Table 2). We have not observed a scarred harbor seal since 1984 nor a scarred northern fur seal since July 1987.

As in previous years (Stewart and Yochem 1985, 1987), the primary material entangling California sea lions in 1988-89 was monofilament gillnet (Table 6); no floats were attached to the entangling monofilament. Since monofilament is negatively buoyant and sinks without the support of floatation devices, we believe that sea lions that were entangled in monofilament became entangled in operational gillnets and were cut out of the nets, leaving some net remaining around the animals' necks. If sea lions (especially young animals) are capable of breaking out of gillnet panels by snapping mesh lines, then some animals may have been entangled in derelict nets (i.e., debris) which were still attached to floats as well as operational, nonderelict nets. Clearly, larger animals, particularly adult males, are capable of breaking free of gillnets once they become entangled (R. DeLong pers. commun.).

Table 1.--Entanglement and scarring rates of pinnipeds in southern California waters by synthetic materials.

Years surveyed	California sea lions	Northern elephant seals	Harbor seals	Northern fur seals
1983-84				
Surveyed	13,174	6,815	1,809	--
% entangled	0.08	0.15	0.00	--
% scarred	0.10	0.09	0.06	--
1985-86				
Surveyed	35,824	17,338	3,342	826
% entangled	0.16	0.16	0.06	0.00
% scarred	0.11	0.20	0.03	0.24
1986-88				
Surveyed	27,733	12,846	3,324	353
% entangled	0.16	0.09	0.03	0.00
% scarred	0.11	0.19	0.00	0.28
1988-89				
Surveyed	24,731	9,775	2,816	422
% entangled	0.12	0.10	0.03	0.00
% scarred	0.10	0.18	0.00	0.00

The observations that we present here, as well as our earlier ones (Stewart and Yochem 1985, 1987), indicate that sea lions become entangled primarily during the first 2 or 3 years of life. Our observations of scarred juveniles and adults indicate that some animals are freed from the entangling material, presumably monofilament, and survive. Others probably die as a result of entanglement either directly through blood loss or indirectly from infection and secondary complications. The magnitude of that mortality is difficult to assess, as many may die at sea, and an insignificant number of tagged sea lions are entangled, preventing an assessment of survival.

For the following discussions we limit the use of the term synthetic marine debris to material other than monofilament. Whether entangling monofilament is obtained during interactions with active fishing gear or from floating derelict nets or net fragments remains difficult to assess. Observations during commercial fishing operations or studies of captive sea lions might clarify whether or not these cases of entanglement are actually related to floating marine debris.

Nevertheless, the increase in numbers of sea lions observed entangled in monofilament in recent years is interesting, considering the restrictions placed in 1983 on the shark and swordfish drift gillnet fishery around the southern California Channel Islands, a fishery that accounted

Table 2.--Entanglement and scarring rates of California sea lions at San Nicolas and San Miguel Islands.

Years surveyed	Adult males	Subadult males	Females/ juveniles	Yearlings	Pups
1983-84					
Surveyed	345	803	7,206	771	4,049
% entangled	0.00	0.12	0.03	0.91	0.02
% scarred	0.58	0.75	0.07	0.00	0.00
1985-86					
Surveyed	1,577	2,272	30,548	1,427	--
% entangled	0.00	0.18	0.15	0.42	--
% scarred	0.44	0.48	0.07	0.00	--
1986-88					
Surveyed	1,384	987	17,619	2,872	4,871
% entangled	0.00	0.00	0.20	0.24	0.02
% scarred	0.14	0.40	0.14	0.00	0.00
1988-89					
Surveyed	710	833	18,670	1,347	3,171
% entangled	0.00	0.12	0.12	0.30	0.13
% scarred	0.70	0.84	0.07	0.00	0.00

for most of the sea lion entanglement and mortality in recent years. Perhaps sea lions became entangled in gillnets north of Point Conception, where the fishery has recently expanded.

In 1988 and early 1989, we were able to confirm that only about 7% of the entangled sea lions observed were entangled in synthetic debris (rubber bands, Table 6). We have no information yet about the survival of sea lions entangled in such debris, and we have not observed any dead sea lions entangled in anything except monofilament gillnet fragments.

All but one entangled northern elephant seal were entangled in synthetic debris (packing bands, Table 6). Elephant seals appear to become entangled during the first 1 or 2 years of life, probably because the circumferences of most packing band debris are too small for the bands to go over the heads of older seals. Scars around the necks of older seals indicate that some seals survive entanglement, although the type of material that entangled those seals is not known. None of the seals that we have observed entangled were tagged, preventing assessment of the influences of various kinds of debris on seals' survival. Five of the adult females that we observed with severely constricting packing bands around their necks gave birth and successfully weaned their pups in 1988.

Since 1983 we have observed only four entangled harbor seals (all juveniles, each with a packing band around its neck), suggesting that they

Table 3.--Entanglement and scarring rates of northern elephant seals at San Nicolas and San Miguel Islands.

Years surveyed	Adult males	Subadult males	Females/ juveniles	Yearlings
1983-84				
Surveyed	1,019	875	4,410	511
% entangled	0.00	0.34	0.07	0.19
% scarred	0.00	0.11	0.07	0.00
1985-86				
Surveyed	1,776	1,485	13,686	391
% entangled	0.00	0.34	0.18	1.02
% scarred	0.28	0.81	0.06	0.51
1986-88				
Surveyed	1,239	1,045	9,802	760
% entangled	0.00	0.00	0.12	0.00
% scarred	0.32	0.77	0.13	0.00
1988-89				
Surveyed	989	658	7,726	402
% entangled	0.00	0.45	0.08	0.25
% scarred	0.81	1.06	0.04	0.00

rarely encounter potentially entangling debris in southern California waters. As it has been speculated that large numbers of harbor seals are incidentally killed each year in gillnet fisheries in southern California, it is surprising that we have seen no harbor seals entangled in gillnets, especially in comparison to the number of California sea lions that are. If young California sea lions that become caught in gillnets are, in fact, capable of breaking out of gillnets, the lack of observations of harbor seals entangled in gillnet fragments may suggest that they are incapable of breaking free. We speculate that such differences may be due to the different modes of propulsion of these two species and consequent differences in potential force generated to permit them to break mesh strands. Harbor seals may then simply die in active or derelict gillnets rather than break free. Clearly, additional observations are needed to sort among these speculations.

Of all cases of pinniped entanglement observed, we can only confirm that 27% were due to marine debris in 1986-88 and 22% in 1988-89, with much of the remainder (perhaps as much as 73%) evidently related to interactions of pinnipeds (especially juvenile sea lions) with commercial gillnet fisheries (see Stewart and Yochem 1987 for additional discussion).

Because relatively few pinnipeds are observed entangled in synthetic material, analysis of trends in entanglement rates (especially debris-related) is difficult. It is clear, however, that relatively large samples

Table 4.--Entanglement and scarring rates of harbor seals at San Nicolas, San Miguel, and Santa Rosa Islands.

Years surveyed	Adults	Immatures
1983-84		
Surveyed	1,445	364
% entangled	0.00	0.00
% scarred	0.07	0.00
1985-86		
Surveyed	2,757	585
% entangled	0.00	0.34
% scarred	0.00	0.17
1986-88		
Surveyed	2,021	1,303
% entangled	0.00	0.08
% scarred	0.00	0.00
1988-89		
Surveyed	1,900	916
% entangled	0.00	0.11
% scarred	0.00	0.00

Table 5.--Entanglement and scarring rates of northern fur seals at San Miguel Island.

Years surveyed	Adult males	Subadult males	Females/ juveniles
1985-86			
Surveyed	58	108	660
% entangled	0.00	0.00	0.00
% scarred	0.00	0.92	0.15
1986-88			
Surveyed	15	63	275
% entangled	0.00	0.00	0.00
% scarred	0.00	1.59	0.00
1988-89			
Surveyed	35	59	328
% entangled	0.00	0.00	0.00
% scarred	0.00	0.00	0.00

Table 6.--Types of synthetic material observed entangling pinnipeds at San Nicolas and San Miguel Islands, 1988-89.

Pinnipeds entangled	Monofilament gillnet	Packing bands	Other debris ^a	Total
California sea lions				
Adult females/juveniles	20	--	2	22
Yearlings	15	--	--	15
Pups	3	--	1	4
Total	38	--	3	41
Northern elephant seals				
Subadult males	--	2	1	3
Adult females	--	6	--	6
Juveniles	--	1	--	1
Yearlings	1	--	1	2
Total	1	9	2	12
Harbor seals				
Adults	--	1	--	1
Total	--	1	--	1

^aIncludes rubber bands, polyfilament rope and line, and items other than trawl or gillnet fragments or nylon monofilament line.

(i.e., systematic observations of large numbers of pinnipeds ashore) are necessary to evaluate properly the true rates of entanglement.

Populations of all pinnipeds have been increasing rapidly in the Southern California Bight during the past two decades (e.g., Stewart 1989; Stewart et al. 1990), indicating that entanglement of pinnipeds in marine debris has had only minor influence on population trends.

ACKNOWLEDGMENTS

We thank A. Yablokov, M. Mina, and R. DeLong for assistance on San Miguel Island, and the Naval Command at Point Mugu and on San Nicolas Island for permitting access and arranging our travel to San Nicolas Island. We also thank the command and helicopter pilots of the U.S. Coast Guard Station at Long Beach, California, for their logistic support for travel to San Miguel Island in October 1987, and J. Coe, M. L. Godfrey, W. Gordon, J. Jehl, and D. Kent for their comments on the manuscript.

The observations summarized in this report were made incidental to research on pinnipeds conducted under contract to the U.S. Air Force,

Contract No. F04701-C-0018. Analysis and reporting of these data were supported by Contract No. 43ABNF802089 to the National Marine Fisheries Service, U.S. National Oceanic and Atmospheric Administration.

REFERENCES

- Shomura, R. S., and H. O. Yoshida.
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Stewart, B. S.
1989. The ecology and population biology of the northern elephant seal, *Mirounga angustirostris* Gill, 1866, on the Southern California Channel Islands. Ph.D. Thesis, Univ. California, Los Angeles, Calif., 295 p.
- Stewart, B. S., and P. K. Yochem.
1985. Entanglement of pinnipeds in net and line fragment and other debris in the Southern California Bight. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 315-325. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Entanglement of pinnipeds in synthetic debris and fishing net and fishing line fragments at San Nicolas and San Miguel Islands, California, 1978-1986. Mar. Pollut. Bull. 18:336-339.
- Stewart, B. S., P. K. Yochem, R. L. DeLong, and G. A. Antonelis, Jr.
1990. Status and trends in abundance of pinnipeds on the southern California Channel Islands. Proceedings of the Third California Islands Symposium, Santa Barbara, California. [In press.]
- Wolfe, D. A. (editor).
1987. Plastics in the sea. Mar. Pollut. Bull. 18:303-365.

MARINE MAMMAL AND SEA TURTLE ENCOUNTERS WITH MARINE DEBRIS
IN THE NEW YORK BIGHT AND THE NORTHEAST ATLANTIC

Samuel S. Sadove and Stephen J. Morreale
Okeanos Ocean Research Foundation, Inc.
Hampton Bays, New York 11946, U.S.A.

ABSTRACT

The incidence of ingestion of synthetics by, and entanglement of, marine mammals and sea turtles in the New York Bight (1979-88) and in Iceland (1985) was documented and related to the ecology of these animals. Post mortems of 88 cetaceans, 37 pinnipeds, and 116 sea turtles in the New York Bight revealed ingestion of synthetics in 24 animals. Differences were observed among the groups of animals. Synthetics were found in 3 mysticete whales, in 7 odontocete whales (3 delphinids, 3 physterids, and 1 phocoenid), and in 14 sea turtles (10 leatherbacks, *Dermochelys coriacea*, 3 loggerheads, *Caretta caretta*, and 1 green, *Chelonia mydas*). No synthetics were found in the gut of any pinnipeds or in Kemp's ridley turtles, *Lepidochelys kempii*. Seventy-five individuals were entangled, including 4 mysticetes, 13 odontocetes, and 58 sea turtles. In Iceland, 6 of 82 examined fin whales, *Balaenoptera physalus*, contained ingested synthetics, and 5 of 95 showed signs of previous entanglement. The types of synthetics ingested and the rate of occurrence of both ingestion and entanglement were related to the feeding behavior, timing, and distribution of the species. The results indicate that certain species of marine mammals and sea turtles are more likely to interact with debris than others. In these animals ingestion of synthetics and entanglement appear to be frequent and widespread.

INTRODUCTION

Increased human use of the oceans and inshore waters has resulted in large amounts of man-made materials with which marine organisms come into contact. Organisms interact not only with waste products and floating debris but also with actively used fishing gear. Numerous efforts have been conducted worldwide to assess the amounts (Wehle and Coleman 1983; Bean 1987), types (Carpenter et al. 1972; Dixon and Dixon 1981; Dahlberg and Day 1985; Center for Environmental Education 1987a, 1987b; Henderson et al. 1987), and sources of these materials (Horsman 1982) and their impacts on marine organisms (Shomura and Yoshida 1985; Coe and Bunn 1987; O'Hara 1989). The interactions of marine organisms with these materials, and the

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154. 1990.

resulting impacts, are better understood when the ecology of the individual species is considered.

Many marine species have global distributions and occur in both populous and remote areas. An abundance and diversity of marine mammals and sea turtles are found in the New York Bight. This is one of the most heavily stressed coastal regions in the world. With New York City at its apex, the bight is a major port for shipping and fishing. This region's coastal population of over 25 million places heavy demands upon the marine environment through activities such as recreational boating, fishing, and dumping of wastes. In contrast, the Arctic region, which supports large populations of marine mammals (Remmert 1980), is one of the few remaining areas in the world where man's influence is still limited. Despite its remoteness, it has been shown that sperm whales in this region were also impacted by marine debris (Martin and Clarke 1986).

The objective of this research was to examine the incidence of ingestion of synthetics by, and entanglement of, different types of marine mammals and sea turtles in the New York Bight and to provide comparisons with whales in Iceland waters.

METHODS

The study was conducted during the period of 1979 through 1988 in the New York Bight and in Iceland during the summer of 1985. Data on ingested materials in the New York area were collected during post mortems of digestive tracts in stranded animals. Only those stranded animals for which reliable necropsies could be performed were included in this study. Animals examined included 37 pinnipeds, 88 cetaceans (19 mysticetes and 69 odontocetes), and 116 sea turtles (Table 1). Data from Iceland were collected by examining the gut contents of 82 fin whales, *Balaenoptera physalus*, at a whaling station in Hvalfjordur during the 1985 season.

Data on entanglement were also collected during the post mortems of both the New York and Iceland specimens (Fig. 1). In New York, a large number of stranded live animals were also examined for evidence of entanglement, e.g., visible scars as reported by Hare and Mead (1987) or actual attached debris, and in Iceland, 13 additional fin whales were examined for entanglement only.

RESULTS

Ingestion of Synthetics

Evidence of ingestion of synthetic materials was found in 24 animals in the New York Bight during this study (Table 2). The frequency of occurrence varied among groups. Synthetics were present in the gut of three individual mysticetes and in seven odontocetes. Among the odontocetes, 3 out of 8 physterids, 3 of 50 delphinids, and 1 of the 9 phocoenids examined contained synthetic materials. There was no evidence of ingestion of synthetics in any of the pinnipeds.

Table 1.--Stranded marine mammals and sea turtles in the New York Bight from 1979 through 1988. A total of 461 live and dead animals were found along the shores or entangled in nets in the water.

Species	Number of individuals
Cetaceans	
<i>Balaenoptera acutorostrata</i>	5
<i>Balaenoptera physalus</i>	9
<i>Delphinapterus leucas</i>	1
<i>Delphinus delphis</i>	15
<i>Eubalaena glacialis</i>	1
<i>Globicephala melaena</i>	14
<i>Grampus griseus</i>	1
<i>Kogia breviceps</i>	5
<i>Lagenorhynchus acutus</i>	4
<i>Megaptera novaeangliae</i>	4
<i>Mesoplodon densirostris</i>	1
<i>Phocoena phocoena</i>	9
<i>Physeter catodon</i>	3
<i>Stenella coeruleoalba</i>	7
<i>Stenella plagiodon</i>	3
<i>Tursiops truncatus</i>	10
<i>Ziphius cavirostris</i>	1
Unidentified	6
Pinnipeds	
<i>Halichoerus grypus</i>	2
<i>Phoca groenlandica</i>	1
<i>Phoca vitulina</i>	34
Sea turtles	
<i>Caretta caretta</i>	103
<i>Chelonia mydas</i>	15
<i>Dermochelys coriacea</i>	85
<i>Lepidochelys kempfi</i>	122
Total	461

Among the sea turtles, varying amounts of synthetics were found in 10 of the 33 leatherbacks, *Dermochelys coriacea*, in 3 of 35 loggerheads, *Caretta caretta*, and in 1 of 4 green turtles, *Chelonia mydas*. Although there were 44 Kemp's ridleys, *Lepidochelys kempfi*, examined in this study, none of these turtles contained synthetics in its gut.

In the Iceland survey during the summer of 1985, plastic material was found in 6 of the 82 fin whales examined.




Figure 1.--The incidence of entanglement in different types of gear for marine mammals (A) and sea turtles (B) in the New York Bight from 1979 through 1988.

A wide variety of debris was observed in the stomachs of the animals examined in this study. The various types of debris found in the guts of cetaceans in the New York Bight included plastic toys, cups, polypropylene line, plastic bags, plastic sheets, and some unidentifiable synthetics. Similar materials were found in the Icelandic whales as well. One of these fin whales contained plastic that unfolded to a 1 x 2 m sheet. The most prevalent types of ingested debris observed in cetaceans from both study areas were plastic bags and small pieces of plastic sheeting.

Table 2.--Gut content analysis of marine mammals and sea turtles in the New York Bight from 1979 through 1988 and of fin whales, *Balaenoptera physalus*, from Iceland during the summer of 1985.

Location	Number examined	Number with synthetics
New York Bight		
Cetaceans		
Mysticetes	19	3
Odontocetes		
Delphinidae	50	3
Phocoenidae	9	1
Physteridae	8	3
Ziphiidae	2	0
Pinnipeds		
Phocidae	37	0
Sea turtles		
Dermochelyidae	33	10
Cheloniidae		
<i>Caretta caretta</i>	35	3
<i>Chelonia mydas</i>	4	1
<i>Lepidochelys kempfi</i>	44	0
Total		24
Iceland		
Cetaceans		
Mysticetes		
<i>Balaenoptera physalus</i>	82	6

Various lengths of monofilament line, small pieces of different colored plastic, and numerous small polystyrene balls had been ingested by sea turtles. Most of the synthetic material in sea turtles, however, was clear, thin plastic. In some instances entire plastic bags were present, and these were the predominant synthetic material found in leatherback turtles.

For several stranded animals there was strong evidence that ingestion of synthetics was contributory or causative of death. In one pygmy sperm whale, *Kogia breviceps*, a hard, black plastic ball had completely blocked the pyloric valve. The surrounding tissue was hemorrhagic and there was

extensive necrosis. This animal was also severely emaciated upon death. Another whale, a pregnant sperm whale, *Physeter catadon*, was found with approximately 300 m of polypropylene line wrapped around its jaw and extending into the stomach. The esophagus and stomach were hemorrhagic and the lower jaw was gangrenous at the time of death. Five leatherback turtles had a large bolus of plastic occluding their digestive tracts. One such bolus was made up of 15 quart-size plastic bags and was blocking the pyloric opening.

Entanglements

From 1979 to 1988 there were a total of 75 individuals in the New York Bight that exhibited signs of entanglement with either debris or inactive or active fishing gear (Fig. 1). These individuals included 4 mysticetes, 13 odontocetes, and 58 sea turtles. No pinnipeds in this study were entangled in gear or debris. In Iceland, 5 of the 95 fin whales examined showed signs of previous entanglement.

Types of entanglement varied among groups of animals (Fig. 1). Three of the four mysticetes were entangled in lines from lobster pot floats, as were three sperm whales. Of the remaining odontocetes, seven exhibited evidence of the animal's having been entrapped in unidentified nets, two in trawl nets, and one in a longline. In the Icelandic fin whales, it was not possible to identify the form of entanglement gear which had made the scars.

The majority of the entanglements occurred in sea turtles, and there were clear differences among the species. The chelonid turtles (loggerheads, greens, and Kemp's ridleys) were primarily caught in pound nets (44 out of 48 turtles), while leatherbacks were entangled in other types of nets (4 of 10) and in lobster pot lines (6 of 10).

The incidence of death among entangled animals was related to the type of entrapping gear. Those types of gear which can hold an animal underwater were more frequently associated with the animal's death. The odontocetes which showed evidence of net entanglement had all died of drowning. These animals appeared healthy prior to death, exhibiting full stomachs, normal blubber thickness, and no specific disease etiology. One leatherback turtle became entangled in a lobster pot line and could not be freed. This animal also drowned. There was no mortality among the 44 turtles entrapped by pound nets, which only encircle an animal and do not confine it under water.

DISCUSSION

Between the years of 1979 and 1988, 461 stranded and entangled animals were found in the New York Bight. These strandings included 17 species of marine mammals and 4 species of sea turtles, and many of the data were collected from carcasses that had washed up along the shores of Long Island, New York. The prevailing wind and current patterns are such that most carcasses in the Long Island Sound or in the eastern bays are transported to shore, but many of those in the ocean float farther out to sea. Thus, while some areas provide an accurate account, strandings along the entire

ocean shore probably grossly under-represent the number of pelagic animals that are impacted.

The incidence of ingested synthetics varied among species. The observed patterns could be attributed to several ecological characteristics of the animals: feeding behavior, seasonal occurrence, and habitat. The type of synthetic found in 19 of 24 animals was floating or neutrally buoyant plastic. Much of this type of refuse originates on land or comes from recreational boating near shore and concentrates inshore during the summer when human activity is highest. Many of the cetaceans are deep water animals, but during the summer they often move inshore, where they have been observed to be feeding heavily. It is likely that ingestion of synthetic materials increases at this time. Animals that stranded during the winter months, such as seals and most Kemp's ridley turtles, contained no synthetic materials.

The ingestion of synthetics also corresponded to the feeding behavior of animals. The mysticetes and a few odontocetes feed throughout the water column by capturing large quantities of food at a time. Plastics and other floating materials are probably ingested along with prey species. Leatherback turtles feed almost exclusively on jellyfish (Mortimer 1981) and probably actively feed on plastic that resembles their prey. Conversely, the Kemp's ridley feeds very selectively on crabs off the bottom and seals in the New York Bight feed primarily on crabs and benthic fish and neither was found to contain debris. In many cases where synthetics were evident, it was difficult to ascertain the direct cause of death due to the decomposed state of the carcass. However, in some animals, the ingestion of synthetic debris caused serious damage and probably resulted in the death of the animal.

The entanglement data were valuable in determining the effects of different types of debris and fishing gear on the species studied. All of these animals must come to the surface to breathe. Debris in the water column or at the surface, such as floating line, can entangle these animals during their normal activities. Lobster pot float lines proved to be a major source of entanglement for pelagic animals such as fin whales, sperm whales, and leatherback turtles. These lines can be more than 100 m long and virtually undetectable below the surface. Types of active or inactive fishing gear that hold animals below the surface, such as longlines, trawlers, and gillnets, can drown marine mammals and sea turtles. Other types of gear that merely confine animals are not a problem. Most of the Kemp's ridley, loggerhead, and green turtles were caught in pound nets with no observed mortalities.

This study examined the impact of two forms of ocean debris. However, there are many other human activities that can affect marine mammals and sea turtles. Recreational boating contributes heavily to fouling the inland waters, and a large proportion of the animals in this study had been struck by boats. Other problems such as heavy metals, pesticides, and sewage runoff are epidemic in many coastal waters. While their effects on marine life may not be immediate, pollutants may result in health problems and have detrimental effects on the long-term survival of populations.

Martineau et al. (1985) showed that ingestion of toxicants drastically reduced the reproductive rate of beluga whales. It is possible that ingestion of debris and entanglement of animals have similar long-term effects, and the numbers of impacted animals are probably much higher than shown in this study (Kraus 1990)

Although the magnitude of the problems of ocean debris is not yet fully realized, this study indicates that the impact of human activity is not restricted to highly populated areas such as the New York Bight. It occurs globally and is found even in such remote areas as Iceland.

ACKNOWLEDGMENTS

Portions of this study were funded by the New York State Department of Conservation and the New York Return A Gift To Wildlife Program, Contract Nos. C000990, C001983, and C001984. We thank Colleen Coogan for help with many of the stranded animals. Gill Lankshear and Rebecca Schneider provided much assistance in manuscript preparation. We also thank the two anonymous reviewers for their helpful comments.

REFERENCES

- Bean, M. J.
1987. Legal strategies for reducing persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:357-360.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Milkas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. *Science* 178:749-750.
- Center for Environmental Education.
1987a. "Coastweek." The entanglement network newsletter. Center for Environmental Education, November 1987, (2):2.
1987b. 1986 Texas coastal cleanup report. Center for Environmental Education, Wash., D.C., 52 p.
- Coe, J. M., and A. R. Bunn.
1987. Description and status of tasks in the National Oceanic and Atmospheric Administration: Marine Entanglement Research Program for fiscal years 1985-1987. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWFAC Processed Rep. 87-15, 39 p.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.

- Hare, M. P., and J. G. Mead.
1987. Handbook for determination of adverse human--marine mammal interactions from necropsies. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWFAC Processed Report 87-06, 35 p.
- Henderson, J. R., S. L. Austin, and M. B. Pillos.
1987. Summary of webbing and net fragments found on Northwestern Hawaiian Islands beaches, 1982-86. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-11, 15 p.
- Horsman, P. V.
1982. The amount of garbage pollution from merchant ships. Mar. Pollut. Bull. 13:167-169.
- Kraus, S.
1990. Rates and potential causes of mortality in North Atlantic right whales (*Eubalaena glacialis*). Mar. Mammal Sci. 6:278-291.
- Martin, A. R., and M. R. Clarke.
1986. The diet of sperm whales (*Physeter macrocephalus*) captured between Iceland and Greenland. J. Mar. Biol. Assoc. U. K. 66:779-790.
- Martineau, D., P. Beland, C. Desjardins, and A. Vezina.
1985. Pathology, toxicology and effects of contaminants on the population of the St. Lawrence beluga (*Delphinapterus leucas*). Quebec, Canada. Report to the Marine Mammals Committee, Int. Counc. Explor. Sea 1985/N:13.
- Mortimer, J. A.
1981. Feeding ecology of sea turtles. In K. A. Bjorndal (editor), Biology and conservation of sea turtles, p. 103-109. Smithsonian Press, Wash., D.C.
- O'Hara, K. J.
1989. Plastic debris and its effects on marine wildlife. In W. J. Chandler and L. Labate (editors), Audubon Wildlife Report 1988/1989, p. 395-434. Acad. Press, N.Y.
- Remmert, H.
1980. Arctic animal ecology. Springer-Verlag, N.Y., 250 p.
- Shomura, R. S. and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Wehle, D. H. S., and F. C. Coleman.
1983. Plastics at sea. Natur. Hist. 92(2):20-26.

A REVIEW OF GHOST FISHING BY TRAPS AND GILLNETS

Paul A. Breen
Ministry of Agriculture & Fisheries
Fisheries Research Centre
P.O. Box 297
Wellington, New Zealand

ABSTRACT

Ghost fishing occurs when lost fishing gear continues to catch and kill animals. This paper reviews what is known about ghost fishing in trap and gillnet fisheries, how the information was obtained and how it has been used, how ghost fishing can be prevented, and what regulatory approaches have been taken to address the problem. Some standard terms are proposed to prevent confusion.

Ghost fishing by traps can occur through several mechanisms. The problem is serious in several fisheries, minor in at least one, and remains unexamined for the majority of trap fisheries. Timed-release devices are simple, inexpensive, and effective at preventing ghost fishing by opening the trap some time after loss. In all Dungeness crab fisheries, such devices are required in crab traps, and other regulations attempt to minimize trap loss. In the American lobster fishery, only Connecticut and Maine address ghost fishing, which is known to be a problem. Ghost fishing by traps is poorly recognized as a problem outside North America.

Ghost fishing by coastal gillnets has been documented in several locations and may persist for several years. For large pelagic gillnets the limited evidence suggests that lost nets form tangled nonfishing masses. More work, both descriptive and experimental, is required to document the nature, extent, and persistence of ghost fishing by gillnets, especially by pelagic gillnets if their use continues.

It is not clear how to prevent ghost fishing by gillnets. Preventive measures suggested to date must be examined for possible side effects.

INTRODUCTION

Ghost fishing can be defined as "the ability of fishing gear to continue fishing after all control of that gear is lost by the fisherman" (Smolowitz 1978a), i.e., when gear is lost, a common occurrence in many fishing operations. The subject was previously reviewed for trap fisheries by Smolowitz (1978a, 1978b, 1978c) and for several gear types by High (1985).

Fishing gear that requires active control, for example trawls, troll gear, and purse seines, may become virtually inert and probably catches insignificant numbers of animals after loss. By contrast, gear which normally fishes passively, such as traps, tangle nets, and gillnets, may continue to fish at significant rates after loss.

This paper looks at what is known about ghost fishing by traps and gillnets, how this knowledge was obtained and used, and what measures can be taken to reduce ghost fishing by traps and gillnets. Other fishing gear may well ghost fish--High (1980, 1987) reports Pacific halibut, *Hippoglossus stenolepis*, striking and being caught by bare longline hooks--but the literature at this stage adds little to High's (1985) review for other gear types.

Why Ghost Fishing May Be a Problem

An increasing proportion of fishing gear is now constructed from nondegradable materials such as stainless steel, other metals, fiberglass, injection-molded plastics, vinyl-coated wire, monofilament netting, and polypropylene twine. Whereas fishing gear made from natural materials deteriorated quickly in the sea--Pacific salmon, *Oncorhynchus* spp., fishing ports all featured tanks of copper sulfate for dipping nets to preserve them--gear made from modern materials lasts much longer in the sea.

The very large volumes of fishing gear now deployed translate to a large volume of lost gear even if the loss rate is small. Some crustacean trap fisheries are so overcapitalized that jurisdictions try to limit the large number of traps used. Hundreds of thousands of kilometers of pelagic gillnets are in use. If this gear ghost fishes when lost, then there is a serious biological and economic problem.

Terminology

Some standard definitions are proposed. First, I use "lost" to describe lost or discarded fishing gear. Previous authors have used "ghost" or "derelict" to describe such gear. However, using "ghost" to mean "lost" creates confusion--the lost gear may or may not actually be ghost fishing. Sutherland et al. (1983) propose a distinction between intact lost gear, still theoretically capable of fishing, and damaged or "derelict" gear that can no longer fish. "Derelict" should be limited to this sense. Where gear loss is simulated experimentally, I use the term "simulated lost" gear.

Second, two types of special openings in traps need careful differentiation. Traps can be modified by openings designed to allow animals to escape (Wilder 1945). These openings have been termed "savings gear" (Jow 1961), "escape vents" (Pecci et al. 1978; Anthony and Caddy 1980), and "escape gaps" (Brown and Caputi 1986). Traps can also be modified by openings or mechanisms designed to release animals from a lost trap. These have been termed "biodegradable sections" (Anthony and Caddy 1980), "timed-release mechanisms" (Blott 1978), "ghost panels" (Krouse pers. commun.), "escape panels" (draft Maine legislation), and "destruct panels" (Hipkins and Beardsley 1970). I suggest that the first type of special opening be called "sublegal escape gaps" and that the second type be called a "timed-release" opening.

GHOST FISHING BY TRAPS

Mechanisms

There is no single mechanism of ghost fishing by traps because traps vary widely in their design, intended mode of capture, target species, and conditions of deployment. To understand ghost fishing it is first necessary to look briefly at trap operation.

Some traps simply attract fish or crustaceans with bait. Although the animals can apparently escape at will, a number are found inside the traps: there is a temporary balance between catch and escape rates. Examples are reef fish traps (Munro et al. 1971; Munro 1974), Australian snapper traps (Dews et al. 1988), and British Columbia prawn traps (Boutillier pers. commun.). Difficult exits in fish traps reduce escapement rates to increase the number of fish in the trap (see Munro 1983).

Escape can be made more difficult by fitting "nonreturn valves" to traps (e.g., Munro 1974). Dungeness crab, *Cancer magister*, traps are fitted with hinged metal gates called "triggers" (High 1976) that permit entry but effectively block exit. Sablefish, *Anoplopoma fimbria*, traps may have similar devices (Hipkins and Beardsley 1970). Homarid lobster, *Homarus americanus* and *H. gammarus*, traps commonly have inner chambers or "parlors" to hinder escape (Pecci et al. 1978; Lovewell et al. 1988).

In the simplest form of ghost fishing, trapped animals die in lost traps and their bodies act as bait (von Brandt 1984). Hipkins and Beardsley (1970, p. 29) state: "It appears then that blackcod (sablefish) pots. . . will continue to fish with dead fish serving as bait to attract new fish which eventually die to attract more fish and so on ad infinitum until the pot deteriorates. . . ." They present indirect evidence for this mechanism. Pecci et al. (1978) suggest this mechanism may operate in lost American lobster traps. For no species has this "autorebaiting" mechanism been conclusively demonstrated.

Traps may be rebaited by species other than the target species. Alaska king crab *Paralithodes camtschatica* traps are rebaited when Pacific halibut or Pacific cod, *Gadus macrocephalus*, enter and die (High and Worlund 1979). Pecci et al. (1978) report a variety of fishes caught and perhaps acting as bait in simulated lost American lobster traps.

Some species of fish are attracted to live conspecifics in an unbaited trap (Munro 1974); for these, ghost fishing might occur without the auto-rebaiting mechanism.

In the simplest model of ghost fishing, trapped animals starve in the traps. Other forms of mortality might be important, causing death sooner. In crustaceans, cannibalism of newly molted individuals may occur. Pecci et al. (1978) observed this in simulated lost American lobster traps; Demory (1971) and Barry (pers. commun.) observed this for Dungeness crabs. Scarratt (1965) reported predation of captured American lobsters by amphipods. Ritchie (1972) and Gabites (pers. commun.) report predation on trapped New Zealand spiny lobsters, *Jasus edwardsii*, by octopus, *Octopus maorum*; Morgan (1974) describes predation by *Octopus* sp. on the Western Australian spiny lobster *Panulirus cygnus*; High (1985) describes attempts by *O. dofleini dofleini* to capture trapped Dungeness crabs. Pecci et al. (1978) reported mortality of American lobsters in simulated lost traps caused by black sea bass, *Centropristis striata*. Trapped crabs may be smothered when the trap is buried by silt (High 1985).

Even when animals manage to escape from ghost fishing traps, they may die as a result of their confinement--High and Worlund (1979) demonstrated this important effect experimentally for Alaska king crabs.

Fishes and crustaceans may enter unbaited traps. This is reported for Hawaiian spiny lobsters, *P. marginatus*, (Paul 1984) in the laboratory; New Zealand spiny lobsters in the field (Gabites pers. commun.); and American lobsters in the field (Pecci et al. 1978; Smolowitz 1978b; but cf. Karnofsky and Price 1989). Dungeness crabs (Breen 1987) and Alaska king crabs (Meyer unpubl. manuscr.) entered empty traps months after simulated trap loss. Munro (1983) describes fish traps that catch fish unbaited. Juvenile reef fishes in Florida use traps as shelter (Sutherland et al. 1983). High and Ellis (1973) found that unbaited traps caught as many reef fish as baited traps. For such traps an autorebaiting mechanism is not necessary for ghost fishing to take place.

In some cases dead crustaceans repel conspecifics. Hancock (1974) demonstrated this effect experimentally for the spiny lobster *P. cygnus*, and also presented evidence that the crabs, *Cancer pagurus*, are not attracted to traps baited with the crab *Carcinus maeanas*. Miller (1977) demonstrated in experimental trapping that the Newfoundland snow crab, *Chionoecetes opilio*, are repelled by dead conspecifics, and High (pers. commun.) also reports this for the Alaska king crab. However, Pecci et al. (1978) found that *H. americanus* are not repelled by dead conspecifics. For species repelled by dead conspecifics, the autorebaiting mechanism will not cause ghost fishing.

Thus ghost fishing can occur through a variety of mechanisms: auto-rebaiting, rebaiting by other species, attraction by living conspecifics, or attraction by the trap alone. The trap may kill through starvation or by facilitating cannibalism and predation. For some species, conspecific repellency may prevent or reduce ghost fishing. Ghost fishing may be significant on species other than the target species.

Demonstrations That Traps Ghost Fish

Recovered Lost Gear

Recovery, especially after long periods, of lost gear that contains live and dead animals is good evidence that ghost fishing occurs. Hipkins and Beardsley (1970) recovered nine sablefish traps lost for approximately 6 weeks. These contained dead fish and up to 24 live fish per trap, suggesting that the autorebaiting mechanism was operative.

In Oregon, Demory (1971) retrieved 117 Dungeness crab traps which had been abandoned for at least 6 weeks. They contained 3,629 crabs, 91% of which were legal-sized males. Dahlstrom (unpubl. manuscr.) recovered an Oregon Dungeness crab trap, lost for 10 months, containing 20 crabs and 2 empty carapaces. The trap was still in excellent condition. Meyer (unpubl. manuscr.) reports that recovered lost Alaska king crab traps "often contain as many as 100 marketable king crab."

Smolowitz (1978a) recovered 18 intact offshore American lobster traps lost for approximately 9 weeks. They contained 24 lobsters weighing a total of 70.8 kg (156 lb). High and Worlund (1979) recovered a snow crab, *Chionoecetes* spp., trap containing 12 king and 14 snow crabs 3 months after loss. Sutherland et al. (1983), using a submersible, found five undamaged fish traps in Florida, lost for an estimated 4-6 months. These held 14 fish, 14 Caribbean spiny lobsters, *Panulirus argus*, and a fish skull.

When lost traps are empty on recovery, it is often inferred that ghost fishing does not occur. For instance, Boutillier (pers. commun.) observed lost prawn, *Pandalus platyceros*, traps from a submersible in British Columbia; none contained prawns and he concluded that ghost fishing did not occur. However, simulated lost Dungeness crab traps that were empty for considerable parts of the year caught and killed crabs (Breen 1987). If traps ghost fish through other than the autorebaiting mechanism, then an empty trap may subsequently kill. Inferences made from empty traps are suspect unless made over large numbers of traps and over several seasons.

Another inference is often made from the way catch rates fall as soak time increases. Traps left to soak for too long give poor catches; the inference is that most of the catch escapes after the bait ceases to attract. Then by extension ghost fishing is inferred not to be a problem. Examples include the Tasmanian spiny lobster, *J. novaehollandiae* (Kennedy pers. commun.), British crabs and lobsters (Bannister pers. commun.), and Dungeness crabs. However, in Dungeness crab traps the catch rate declines with increased soak time, yet lost traps continue to catch and kill at a slow rate (Breen 1987). So ghost fishing may occur in the long term despite apparent short-term escapement.

Trap Loading Experiments

Ideally, all ideas about ghost fishing should be tested experimentally. Three approaches have been used: experiments in which traps are loaded and escape rates or mortality rates measured, laboratory

observations that simulate fishing, and field experiments with simulated lost traps.

Munro (1974) found that 50% of reef fishes escaped from Antillean fish traps after 14 days; this implies a 5% escapement per day and 23% retention after a month. These rates suggest that ghost fishing is likely to occur in such traps. However, Munro (1983) estimated an escapement rate of 12% per day from "Z" fish traps, a rate implying only a 2% retention after 30 days.

Sheldon and Dow (1975) loaded 98 tagged American lobsters into 35 unbaited simulated lost traps and checked the traps by diving and hauling for nearly 2 years. The traps continued to catch lobsters, of which 12-18% died in the traps, demonstrating for the first time ghost fishing for American lobsters.

Newfoundland snow crab traps were loaded and examined at intervals by diving (Miller 1977). After 3 weeks no crabs had escaped. Miller then tested the mechanism of ghost fishing in this species by fishing with four treatments. Unbaited traps and traps baited with dead crabs caught nothing; on average squid-baited traps caught 31 crabs per trap; traps baited with a mixture of dead crabs and squid caught 7 crabs per trap. Miller concluded that dead snow crabs repel conspecifics and that the only loss from ghost fishing would involve those crabs originally attracted by the bait.

High and Worlund (1979) observed a 20% retention rate for legal size Alaska king crabs and 8% for sublegal crabs in experimental traps after 18 days. Mortality in standard traps was 2-7% over this period.

Muir et al. (1984) baited Dungeness crab traps daily and observed that 35% of the captured crabs died in the traps. High (1985) placed Dungeness crabs in traps with and without triggers and sublegal escape gaps. The mortality in traps with functional triggers and sublegal escape gaps was 17% after 12 days, confirming ghost fishing as a problem with these traps.

Laboratory Observations

Paul (1984) observed that Hawaiian spiny lobsters in a large tank normally did not escape from traps. The trap lids "had to be removed to prevent them from becoming permanently trapped inside."

Behavior of reef fishes around traps was observed in a large tank by Harper and McClelland (1983, cited by Heneman and Center for Environmental Education (CEE) 1988). Most species appeared to learn to escape, leading to "an equilibrium state. . .with frequent movements in and out of the trap."

Booth (pers. commun.) used a time-lapse camera in a large tank to record the behavior of *J. edwardsii* around simple cane traps, as used in the New Zealand fishery, and parlor-type traps not used in the fishery. There was a rapid turnover of lobsters in the simple trap, but greatly

reduced escapement in the parlor traps. Booth concluded that ghost fishing is probably not a problem for the cane traps, but could be a problem if more complex traps were introduced. Plastic truncated-cone entrances on the top of the trap appear to limit escape in this species in large laboratory tanks (Breen unpubl. data; Gabites pers. commun.).

Field Experiments

Information from trap loading and laboratory studies must be treated with caution: problems with extrapolation from the laboratory to the field and from short to long term must be carefully considered. Possibly the best information comes from underwater observations of simulated lost traps. Tagging of trapped individuals by divers can be used to follow turnover.

Pecci et al. (1978) reported only 30% escapement in American lobsters entering simulated lost traps observed by divers. Mortality rate was 25%. The authors estimated that a ghost fishing trap caught at a rate near 10% that of a surface-hauled trap, confirming ghost fishing as a problem in this fishery.

Breen (1987) simulated 10 lost Dungeness crab traps in a sheltered bay for 1 year, during which approximately 100 crabs died in the traps. At the end of the study, traps were still killing crabs at a steady rate. The results cannot be generalized directly to other Dungeness crab fisheries. For instance, many traps lost off high-energy beaches are destroyed or put ashore by wave action.

Western Australian snapper, *Chrysophrys auratus*, traps were observed in the field with underwater video (Anonymous 1984; Dews et al. 1988; Moran and Jenke 1989) partly to examine possible ghost fishing (Bowen 1961). Fish seemed capable of leaving traps easily and some even swam out "in reverse." Moran and Jenke (1989) simulated lost traps for various periods from 1 to 21 days. Catches were similar to commercial catches after 15 min, indicating that cumulative catching did not occur. These workers concluded that ghost fishing is not a problem with snapper traps. However, three fish were dead in the 21-day trap, suggesting that some ghost fishing may take place.

Hawaiian spiny lobsters appear to move out of simulated lost traps once the bait has deteriorated (Okamoto pers. commun.; Parrish pers. commun.).

Rates of Trap Loss

Traps are lost for many reasons. Simple vessel traffic and towboating may sever buoy lines or drag traps into water deeper than the buoy line. Weak or chafed buoy lines may break. Buoys may become detached from the buoy line, or may be attacked by marine birds (Smolowitz 1978b) or mammals (High 1985). Storms or strong currents may "drown" traps either directly or by rolling them over the bottom, wrapping the buoy line around the trap

(Smolowitz 1978b; Sutherland et al. 1983; von Brandt 1984). Traps set on rocky ground may snag and be unrecoverable (Bowen 1961).

Traps may be carried into deep water, or buoy lines cut, by other fishing activities such as trolling, trawling and gillnetting. When traps are set on ground lines, ground lines may be intentionally cut when lines become fouled. Internecine buoy line cutting or ground line cutting may result from unresolved fishing disputes (Smolowitz 1978b; Breen 1987). In some areas vandals cut buoy lines (Sutherland et al. 1983).

Estimates of trap loss rate must be obtained through surveys or industry interviews. These give the total loss of traps, which might include stolen traps.

American Lobster Traps

For the U.S. American lobster fishery, Smolowitz (1978a) cites anecdotal estimates of the annual loss of traps as 20-30% along the Atlantic seaboard. In the offshore lobster fishery he suggests that 40,000 all-metal traps may have been lost during the period 1971-78. In the inshore fishery Krouse (pers. commun.) suggests an annual loss rate of 5-10%. Based on a 1987 estimate of 1.87 million traps fished, this leads to an annual loss estimate of 93,500-187,000 traps lost annually. An unpublished study (CEE 1987) cited by Heneman and CEE (1988) estimated an annual loss of 500,000 traps annually. In Rhode Island a logbook study led to an estimate of 10-15% annual loss (Fogarty pers. commun.).

In Newfoundland, no estimates have been made of lobster trap loss rate, but divers observe few lost traps on the fishing grounds. Many lost traps are washed ashore (Ennis pers. commun.). Losses have not been estimated in the rest of the Canadian lobster fishery.

Dungeness Crab Traps

In California, 100,000 Dungeness crab traps are estimated lost each year (Kennedy 1986). Some silt into the bottom, but others could fish for an estimated 2 years. In Washington State, Northup (1978, cited in Muir et al. 1984), estimated that 17.6% of the coastal Washington State crab traps were lost in 1975-76, considered a typical year. Barry (pers. commun.) estimated mean annual loss in the same fishery as 11.9%. He considers that ghost fishing traps are <50% of the total loss and may be as low as 10%. In the Puget Sound portion of the fishery, gear loss was estimated from a questionnaire survey to be 15% (Bumgarner pers. commun.). Breen (1987) estimated Fraser River Dungeness crab trap loss as 11%, based on a questionnaire survey. About half those surveyed thought that half the lost traps were ghost fishing.

Thus in several coastal trap fisheries, annual trap loss rates are on the order of 10-20%. American lobster and Dungeness crab fisheries are both cases where more traps than optimum are fished (and thus lost) (Bell and Fullenbaum 1986; Methot 1986). Cumulative trap losses are a cause for concern in fisheries where ghost fishing is known to occur.

The Fate of Lost Traps

Not all lost traps become ghost fishing traps even where ghost fishing is a problem. Smolowitz (1978b) reviews sources of trap destruction. Storms destroy or strand many inshore American lobster and Dungeness crab traps in exposed locales. Burial by storm action or alluvia occurs quickly in some Dungeness crab fishing areas (Hipkins 1972, cited in Smolowitz 1978b; Breen 1987).

Untreated wooden traps are destroyed by borers in a relatively short time, but treated wooden traps may last up to 2 years (Smolowitz 1978b; Fogarty pers. commun.). Twelve percent of the wooden traps used by Sheldon and Dow (1975) were so damaged by lobster chelipeds that escape became possible. Increasingly, however, traps are made from metal (Acheson 1982) or synthetic materials. Averill (pers. commun.) believes that "wooden" American lobster traps last as long as wire traps when lost. Long-term experiments are required to determine the fishing lifespan of various trap types.

High and Worlund (1979) estimate that metal-framed, synthetic mesh-covered Alaska king crab traps could have an effective longevity of 15 years after loss. Breen (1987) found that metal-framed, stainless steel-covered Dungeness crab traps were in excellent condition after a year's submersion. Electrolytic corrosion probably destroys most metal traps eventually. New designs include plastic traps (e.g., Piatt 1988) and vinyl-coated mesh (e.g., Maynard and Branch 1988), which might last for decades. The present Maine trap inventory is 50-60% vinyl-coated wire (Averill pers. commun.).

Note that much of the information just presented is based on short-term studies. The real fate of lost fishing gear has not been well studied.

Impact of Trap Ghost Fishing

How much fishing takes place by ghost-fishing traps? To answer this for a specific fishery requires 1) estimates of the number of traps fished and the loss rate, 2) an assumption about the percentage of lost traps that ghost fish, 3) an estimate of the rate of mortality in ghost-fishing traps, and 4) an estimate of the effective ghost fishing lifespan of a trap. Ideally, for requirement 3 one should also know the natural mortality rate, because some individuals killed by ghost fishing would have died before commercial capture. Many individuals would also have grown before commercial capture. However, the unavoidable imprecision of the other estimates implies that only a crude answer can be obtained in any case.

For the Newfoundland snow crab fishery, Miller (1977) used spot interviews to estimate trap loss at 8%, and combined this with commercial catch rates and experimental observations to obtain an estimate of ghost-fished catch of 10 metric tons (MT) annually. Smolowitz (1978b) estimated the impact of ghost fishing in the U.S. portion of the American lobster fishery. The estimated annual ghost fishing catch was 670 MT, worth an

estimated 1978 US\$2.5 million. From a 1976 study using different assumptions (CEE 1987, cited by Heneman and CEE 1988), the economic loss of just those lobsters within traps at the time of loss was estimated at 1976 US\$2.5 million. Krouse (pers. commun.) assumed a loss rate of 5% in the U.S. American lobster fishery and that traps last for 2 years and take two lobsters per year. This leads to an estimate of 204 MT lost to ghost fishing annually, worth 1989 US\$1.2 million. This is a conservative estimate, because it is based on the low end of the range of trap loss estimates.

Breen (1987) estimated the impact of ghost fishing in one part of the British Columbia Dungeness crab fishery, using loss rates and lifespan estimates from an industry survey and experimental ghost fishing data. He estimated the ghost-fished catch to be 7% of reported landings, worth about 1985 Can\$80,000.

For the sablefish fishery of British Columbia, Scarsbrooke et al. (1988) used trap loss rate from an industry survey, the commercial catch rate, and simple assumptions about turnover rate, trap lifespan, and timed-release device effectiveness. For traps lost from 1977 to 1983, before timed-release devices were fully employed, the estimate of ghost fishing catch was approximately 300 MT annually, compared with landings of 1,000-4,000 MT.

These cases illustrate that ghost fishing can be substantial. I can find no fishery for which the impact of ghost fishing on stocks has been determined, or where ghost fishing is addressed by stock assessments or management plans. In Oregon, where traps are required to incorporate timed-release mechanisms, biologists consider that ghost fishing, although subtracting from the potential catch, would have no stock-recruitment effect. The size limit is set so that all legal-sized males could theoretically be taken without affecting reproduction (Demory pers. commun.).

Prevention of Trap Ghost Fishing

Remedial measures may either reduce trap loss or prevent lost traps from killing. A simple way to reduce trap loss is to reduce the number of traps fished (Smolowitz 1978b). Effort is excessive in many fisheries, so this approach is often desirable for that reason alone. The extreme solution, vessel trap limits or transferable trap entitlements, is extremely expensive to enforce and therefore was rejected as a management option in the New Zealand *J. edwardsii* fishery (Anonymous 1987).

Trap designs can be improved to reduce storm and current losses caused by traps rolling on the bottom (see Smolowitz 1978b). Losses caused by vessels can be reduced by prohibiting buoyed traps in areas of heavy traffic. In Washington State, trap-free lanes for towboats have been established to minimize trap loss from that source (Bumgarner pers. commun.). The Washington Department of Fisheries also facilitates coordination between trap and net vessels to avoid gear collisions. In the Canadian sablefish fishery, ground lines must be buoyed at each end. In practice, the marking employed far exceeds the minimum standard required (McFarlane pers. commun.). In the Puget Sound recreational trap fisheries

of Washington State, regulations require solid buoys (to prevent losses from puncture) and nonfloating buoy lines (to prevent loss from vessel traffic) (Bumgarner pers. commun.).

The large literature on sublegal escape gaps shows that they greatly reduce catches of sublegal crustaceans, presumably through escapement (e.g., Cleaver 1949; Fogarty and Borden 1980; Brown and Caputi 1986; see review in Smolowitz 1978c). Because escape gaps reduce trap saturation effects (Miller 1979), they may lead to increased catches of legal animals.

Ghost fishing mortality was reduced for sublegal American lobsters by sublegal escape gaps in simulated lost traps (Pecci et al. 1978; Smolowitz 1978a). High (1985) found greatly increased sublegal escapement in simulated lost Dungeness crab traps fitted with sublegal escape gaps. Breen (1987) found that as many sublegal as legal Dungeness crabs died in simulated lost traps fitted with appropriate sublegal escape gaps, but the absolute catch rates of legal and sublegal crabs were unknown. Sublegal crabs may have had a high turnover rate in the traps.

Measures to prevent lost traps from ghost fishing usually involve some deliberate failure (timed-release) in a trap component to open the trap or create a new opening for escapement.

Natural fiber twine can be used either to make a timed-release panel or to sew a timed-release panel shut. Panels can also be made from untreated softwood. Blott (1978) tested a variety of materials with potential for use as timed-release elements in traps. Jute and manila twine and steel wire appeared to be realistic, while wool and leather were not.

In Maine, various materials have been tested for use in closing timed-release openings (Averill pers. commun.). Industry was given traps with many openings secured with test materials and asked to fish them during their regular season. Mild steel hog rings appear to last the desired time (ca. 200 days), and are consistent in their total degradation time. Cotton twine and sisal twine are also good candidates for this purpose.

Scarsbrooke et al. (1988) tested failure rates of several binding materials for timed-release openings in sablefish traps. They also fished traps with three types of opening in alternation with control traps to measure the effectiveness of timed-release openings. Triangular or square openings were more than 90% effective in allowing trapped fish to escape; simple "slashes" were less effective. They concluded that appropriately shaped timed-release openings eliminated the problem of ghost fishing in these traps.

Plastic crab and lobster traps in Florida (Piatt 1988) have a rectangular opening which the user fills with a timed-release device such as a plywood panel.

Blott (1978) describes a solid timed-release panel made from galvanized steel and held shut with natural twine or a degradable metal ring. The panel can also incorporate the sublegal escape gap, leading to the name

"catch escape panel." Blott tested various materials for suitability as catch escape panels; galvanized sheet steel seemed most appropriate. Pecci et al. (1978) tested such panels in simulated lost American lobster traps and concluded that such panels "are an effective means of releasing entrapped lobsters." Traps with this type of panel are now commercially available from a Maine manufacturer (Lazarus 1988). However, Averill (pers. commun.) considers that the combination of sublegal escape gaps and a timed-release opening leads to confusion of two separate management issues.

In California, magnesium pins are used to hold together the two halves of plastic or fiberglass traps or to attach the lids of plastic and fiberglass traps (Estrella pers. commun.).

Dungeness crab traps are serviced through the "lid," a hinged section of the top secured by a hook attached by a rubber strap from the side of the trap (High 1976). A timed-release hook, or hook attachment, would allow the trap to open. Breen (1987) unhooked 10 simulated lost traps that had ghost fished for a year. Over a week, 22 of 29 trapped crabs escaped and no new captures were observed. Thus a timed-release device that unhooked the lid would probably be effective in this type of trap.

It is possible to make plastics that are degraded by organisms, light, oxidation, other chemical reaction, and dissolution (see review by Andrady 1988). Various degradable plastic compounds designed specifically for the fishing industry are now being tested (Gonsalves et al. 1989, Gonsalves 1990). Japanese chemists are designing "bacterial co-polymers" which degrade slowly into natural chemicals in water (Doi et al. 1988).

Premature failure of timed-release elements reduces industry acceptance of the concept (Smolowitz 1978b). The early failure of a batch of hog rings used to close timed-failure panels in lobster traps resulted in industry resistance to the devices in Maine (Anonymous 1988; Averill pers. commun.). A similar experience in California led to delayed legislation (Estrella pers. commun.). Material failure rates vary widely with local conditions and probably cannot be predicted accurately. Agencies proposing timed-release regulations must conduct widespread materials testing to find a mechanism that will both fail reliably after the desired time and not fail prematurely. Studies conducted by the industry under actual fishing conditions are more likely to be accepted by the industry.

The dollar and time costs of timed-release modifications are important to acceptance by industry (High and Worlund 1979). Breen (1987) calculated the annual economic cost of Dungeness crab trap ghost fishing done in 1985 as Can\$1.46 per trap in use, and suggested that annual modifications must therefore cost less than this. This simple study appears to be the only published cost-benefit analysis of the problem. Other managers consider that "off-the-cuff cost-benefit analysis would indicate that [ghost fishing] should be addressed" (Averill pers. commun.).

Finally, Smolowitz (1978b) suggests development of "habipots" that catch animals seeking them as shelter. Such traps would not entrap animals

and thus would have only biologically positive effects when lost. Some *Octopus* traps operate on this principle (Mottet 1975).

Regulations to Prevent Trap Ghost Fishing

The American lobster and Dungeness crab fisheries are interesting to examine for regulations designed to minimize ghost fishing. In both fisheries ghost fishing is known to occur, trap losses are high, and the fisheries take place over several jurisdictions in two countries with differing management approaches.

Dungeness Crabs

California requires all traps to incorporate timed-release devices or openings. These may be trap lid hooks made of soft steel <6 mm diameter, lid hooks attached to the strap with single loops of natural fiber twine, any modification of the upper mesh secured with natural fiber to create a 125-mm diameter hole, or magnesium pins as discussed above. Testing of these materials has been carried out, and cotton twine is the preferred option (Estrella pers. commun.). All traps or ground lines of traps must be buoyed and the buoys marked with identification markings.

Oregon requires Dungeness crab traps to contain a timed-release device as in California (Demory pers. commun.). Individual traps must be buoyed and marked.

Since October 1988, Washington also requires timed-release devices as above but not including the mild steel hook; openings must be unimpeded, at least 76 × 127 mm and closed with natural fiber. Washington also has buoy and buoy line standards described earlier.

In British Columbia, Fisheries and Oceans Canada introduced a regulation in 1990 requiring a single loop of specified cotton twine in the lid strap and nonfloating buoy lines. Traps or ground lines must be buoyed with marked floats, but this regulation is often ignored (Breen 1987).

In Alaska, Dungeness crab traps are required to have timed-release devices (Koeneman pers. commun.). At least as early as 1974, Alaska sablefish traps were required to incorporate timed-release panels (Hipkins 1974, cited in High and Worlund 1979).

Alaska also requires that "traps left unattended for over 2 weeks must have bait removed and doors secured open as protection against ghost fishing." This is the only regulation dealing with ghost fishing listed by Miller's (1976) review of crab management regulations in North America, demonstrating the relatively recent recognition of the problem.

Most other major trap fisheries on the Pacific coast have similar regulations. Scarsbrooke et al. (1988) describe the requirement for a timed-release panel in the sablefish fishery. In this case the fishing industry actually included such devices before being regulated. Regulations governing a new trap fishery for hagfish, *Eptatretus* spp., require timed-failure openings in British Columbia and Oregon (Harbo pers. commun.).

American Lobsters

In the United States, Connecticut has been the only jurisdiction to require incorporation of a timed-release panel into the trap. Maine drafted legislation in 1982, which will take effect in 1990 (Krouse 1989), requiring a timed-release panel at least 95 mm square, made of untreated natural material: twine <5 mm diameter, ferrous metal less than about 2.5 mm diameter, or softwood. In the federally controlled part of the fishery, degradable fasteners closing a timed-release opening will be required in 1992 (Fogarty pers. commun.).

In the federally regulated portion of the fishery, lobster traps must be marked with the owner's identification number, and traps set on ground lines must be marked with a buoy or flagpoles and radar reflectors, depending on how many traps are set.

In the Canadian fishery, no regulations are directed at ghost fishing. Anthony and Caddy (1980) recognized the problem and recommended that timed-release panels or "links" be included in all traps and especially deepwater traps.

GHOST FISHING BY GILLNETS

Mechanisms

Gillnets work by trapping animals in the mesh of the net; ghost fishing is a simple continuation of the gillnetting process after the net is lost.

A wide variety of species are targeted with many types of gillnet worldwide (see Uchida 1985 for a comprehensive review). In comparison with the trap fisheries reviewed above, there has been little work on ghost fishing by gillnets. This may reflect failure to recognize a problem: Herrick and Hanan (1988) review problems caused, inter alia, by California gillnets without considering ghost fishing.

Pelagic or drift gillnets are used by Japan and Taiwan in the North Pacific to catch salmon and squid (Uchida 1985), and in the South Pacific by Japan, Korea, and Taiwan to catch albacore and skipjack tuna (Hinds 1984; Murray 1988). Ghost fishing in pelagic gillnets may be overshadowed by their incidental catch performance. They catch a long list of other nontarget species including fishes, birds, turtles, and marine mammals. Even reindeer have been reported caught by gillnets (Beach et al. 1976). Sloan (1984) and McKinnell et al. (1989) give extensive species lists in the incidental catch in Japanese squid gillnetting off British Columbia. In the same fishery Jamieson and Heritage (1987) estimate the catch of birds at one per 18 km of net set, the catch of mammals at one per 140 km. Harwood and Hembree (1987) estimate the incidental catch of cetaceans in pelagic gillnetting off northern Australia, 1981-85, to have been on the order of 14,000 individuals. Incidental catches of cetaceans are also a serious problem in coastal gillnet fisheries. Read and Gaskin (1988) estimated the catch of harbor porpoises, *Phocoena phocoena*, by groundfish

gillnets in the Bay of Fundy, concluding that the incidental catches threaten the population. Recreational gillnetting is a major threat to the endangered Hector's dolphin, *Cephalorhynchus hectori*, in New Zealand (Dawson 1990).

Demonstrations of Gillnet Ghost Fishing

Recovered Lost Gear

In Iceland, synthetic cod gillnets were found a "fairly long time" after loss (von Brandt 1984); they appeared to be fishing actively based on the number and appearance of fish.

Way (1977) described catches of live fishes and crabs in lost demersal Newfoundland cod gillnets retrieved with purpose-designed dragging gear. He concluded that lost gillnets continued to fish "at a declining rate."

DeGange and Newby (1980) described finding a drifting 3.5-km pelagic gillnet lost for at least a month. The net contained 99 birds and 78 fishes. Live birds appeared to be attracted to the net, perhaps by the material already caught, and many of the fish were fresh. These authors confirm the fears of Bourne (1977) that lost gillnet fragments continue to catch and kill birds.

High (pers. commun.) found a lost salmon gillnet with fish skeletons, diving ducks, and seals, *Phoca vitulina*, in varying states of decay, indicating the net continued to kill these animals.

Underwater Observations

After discovering lost salmon gillnets in Washington, High (1985) used scuba to observe them for 6 years. The nets continued to catch crabs, fishes, and birds for 3 years. One net 180 m long contained an estimated 1,000 female crabs (High pers. commun.).

In New England, Carr et al. (1985) made observations from a submersible. They describe fishes entangled in nets estimated to have been lost for at least 2 years. Observations were continued for 3 years from a submersible and remotely operated vehicle (Carr and Cooper 1987; Carr 1988). Nets lost for 3-7 years continued to catch a variety of species, including spiny dogfish, *Squalus acanthius*; American lobsters; and bluefish, *Pomatomus saltatrix*. Later observations on one net indicated that gadoid fish successfully avoided the net, but crabs, *Cancer irroratus* and *C. borealis*, continued to be killed. Carr and Cooper (1987) estimated that lost nets were fishing at approximately 15% of the rate of commercial nets.

Dennis Chalmers (pers. commun.) reported finding a British Columbia herring (*Clupea harengus pallasii*) gillnet lost for at least 4 years: "This net was all bunched and tangled up against a rock ledge in 15 ft [4.6 m] of water and, at the time, there were a few rockfish [*Sebastes* spp.] trapped inside it." Another net found in 11-12 m depth had been lost for at least

7 years. It had no cork line, but the net had enough buoyancy to sit in fishing position and contained several fresh herring.

As in crustaceans, decaying fishes of some species may repel conspecifics. It is believed in New Zealand, for instance, that dead rig, *Mustelus lenticulatus*, and rig offal near a net reduce net catches (Bradstock pers. commun.). This effect might reduce ghost fishing for some species, but no formal research appears to have been conducted.

Schrey and Vauk (1987) reported that more than 2.6% of gannets, *Sula bassana*, visiting Helgoland become entangled in lost gillnets, which caused 30% of the total gannet mortality observed.

Field Experiments

Two simulated lost demersal gillnets were observed by divers in New England (Carr et al. 1985). The nets continued to catch fishes and crabs over 2 1/2 months of observation.

Kim Walshe (pers. commun.) observed simulated lost inshore gillnets by diving for a year in New Zealand. The nets were partly disabled by algal growth and wrapping up, but continued to catch and kill some fish at intervals through the year. Rock lobsters, *J. edwardsii*, are attracted to the fish and are themselves caught by lost inshore gillnets (Anonymous 1978).

Rate of Gillnet Loss

Storms can break gillnets or break off the end markers. Vessels and trawls may run over or cut gillnets. Marine mammals and large fishes may break and carry away nets. In northern waters ice causes gillnet loss (Way 1977). Way also suspected that some nets were simply abandoned at the end of the season. Net fragments may simply be discarded (Gerrodette et al. 1987). In inshore gillnet fisheries, nets snag on obstructions and are lost.

In the New England groundfish gillnet fishery, loss of nets was investigated by CEE (1987, cited by Heneman and CEE 1988). The study examined claims for lost gear made under a U.S. Federal act providing for compensation for gear loss caused by foreign fishing activities. For 1985 and 1986, claims were made for 48 and 29 km of net, respectively. It is unknown what proportion of the total net loss this represented.

Fosnaes (1975) estimated that 5,000 Newfoundland cod, *Gadus morhua*, gillnets were lost annually. Way (1977) conducted a program of lost net retrieval on commercial grounds, finding 148 nets in 48.3 h in 1975 and 167 nets in 53.5 h in 1976.

The density of lost demersal gillnets on a commercial ground in New England was estimated from a submersible by Carr et al. (1985). They found 10 lost nets over 40.5 ha of bottom in 37.5 h search time.

For large pelagic gillnets, a major concern is the tremendous quantity of net in the water. Eisenbud (unpubl. manusc.) estimated that 5,000 km

of net were used in the Japanese North Pacific salmon net fishery alone. Uchida (1985) estimated that 170,000 km of pelagic net were used in 1984 in the North Pacific. Coe (1986) estimated that more than 1.6 million km of squid net were used by Japan, Republic of Korea, and Taiwan in 1985. Even a very small loss rate results in a very large estimate of lost net.

Pelagic gillnets are lost from most of the same causes as coastal gillnets. Because of their great length (12-15 km), these nets are vulnerable to vessel traffic. In the Japanese fishery, intact nets are easier to recover than fragments because radio buoys and lights are installed at each end; most nets recovered by Japanese observers were fragments (Morimoto pers. commun.). Additional causes of loss suggested by Eisenbud (unpubl. manusc.) are desertion of nets in prohibited areas after removal of end markers, and simple discard of old netting. A fisheries observer, Goldblatt (1989), describes a pelagic gillnet vessel entangling her own net in the propeller, then cutting away and discarding a large fragment.

Eisenbud (unpubl. manusc.) reported an estimate that 0.06% of Japanese salmon pelagic gillnet is lost at each set. Gerrodette et al. (1987) report an estimate of 0.05%. They consider this estimate to be low, but Morimoto (pers. commun.) considers that the loss rate would be lower in the squid gillnet fishery because of calmer sea conditions. Tsunoda (1989) observed a Japanese pelagic squid gillnet vessel for 4 weeks and observed no gear loss. When nets were severed by vessels, Tsunoda reports that the crew quickly recovered the subsections. Eisenbud (unpubl. manusc.) estimated annual loss from the Japanese North Pacific squid and salmon gillnet fisheries to be approximately 2,500 km of net.

The density of lost gillnet material can be estimated at sea from transect surveys (Baba et al. 1990; Day et al. 1990). However, the absolute density of lost nets is very low, net fragments cannot be seen from a significant distance, and the tendency is for drifting debris to become nonrandomly distributed by winds and currents. Assessing the impact of lost gillnets through direct surveys is therefore difficult.

Fate of Lost Gillnets

Gillnets are usually made from synthetic materials which can last for long periods of time. High (1985) observed that lost salmon gillnets continue to kill birds and fish for 3 years, and estimated that crabs may be killed for at least 6 years. The direct observations of Chalmers (pers. commun., described above) on herring gillnets tend to support these estimates.

In inshore waters, algal growth on sunken nets may stop fishing by making the nets highly visible to fishes and birds (High 1985; Dennis Chalmers pers. commun.), but Kim Walshe (pers. commun.) reports that fish are caught even in overgrown nets. Strong currents cause the net to tangle lead line over cork line (Way 1977) or end over end (High 1985). High suggests that rolled netting stops catching birds and fishes but may continue to catch crabs. Drift macrophytes and the catch of fish and crabs may cause the net to sink and stop fishing efficiently (Way 1977; Carr et

al. 1985; Millner 1985, cited in Heneman and CEE 1988). Dogfish caused twisting of the demersal gillnets observed by Carr et al. (1985). These authors found three main types of lost net configuration, and speculated that these related to how the nets were lost.

Gerrodette et al. (1987) attached radio transmitters to four sections of gillnet 50-1,000 m long, then monitored the simulated lost nets. The shortest net "collapsed" very quickly, but the largest net remained in fishing condition for at least 10 days. The authors estimated that a 1-km net would remain in a fishing configuration for several weeks.

In a similar study, Mio et al. (1990) examined five simulated lost pelagic gillnets, each 1,200 km long, for nearly 4 months. At the end of this time all nets had twisted themselves together end for end to form a large mass. One net completed this process in 20 days; the others took longer.

The wrapping up of nets may be accelerated by storms. Sloan (1984) observed that squid gillnets off British Columbia became tangled at wind speeds >65 km/h.

Merrell (1984) estimated that netting at sea survives for <10 years. This estimate is based on "aging" nets found stranded.

Prevention of Gillnet Ghost Fishing

As with traps, the most effective way to prevent ghost fishing is to prevent gear loss. In the Japanese pelagic gillnet fishery, vessels are required to mark nets with a radio buoy at one end and radar reflectors at both ends. Radio communication is used to deflect vessel traffic around the nets. Discarding of netting is prohibited, and old netting is disposed of on land (Morishita pers. commun.).

Gillnets could be hung from the cork line with natural fiber twine (Way 1977; von Brandt 1984). In theory when the net is lost the twine would rot, and the lead line would pull the net into deep water. This idea is being examined experimentally for coastal gillnets in New England (McKenzie pers. commun.). The tendency of nets to become tangled (lead line over cork line) might prevent sinking, but would also reduce ghost fishing potential. There is also a danger that sinking the net simply transfers a surface ghost fishing problem to the bottom, as suggested by the salmon gillnet observations.

In British Columbia, a proposal to require herring gillnets to be hung with cotton twine has been drafted, but is still under discussion with industry (Dennis Chalmers pers. commun.).

I am aware of no research into degradable materials for use in the web itself. The use of natural fiber for gillnets would be a backward step because of the massive effort required to maintain and preserve nets during fishing. Gillnets are commonly made from monofilament nylon (Uchida 1985), whereas the major effort in degradable plastics has been aimed at poly-

ethylene or polyolephanes (Scott 1990) or composites of polyethylene or polypropylene and natural material (Blott pers. *commun.*). A potential problem is that degradable nets would form many smaller net fragments instead of one large one.

CONCLUSIONS

Ghost fishing has not been well studied. Significant information exists for only two gear types: traps and gillnets. The importance of ghost fishing as a potential problem is underscored by very large volumes of fishing gear in use, high gear loss rates in many fisheries, and the widespread use of nondegradable materials such as plastics and stainless steel for fishing gear construction.

The fishing behavior of lost traps has been examined for only a handful of fisheries, mostly in North America. For most of the world's many trap fisheries, the impact of lost gear has simply not been addressed.

Ghost fishing by traps can operate through several mechanisms depending on trap type and the target species. Where impact has been estimated, ghost fishing sometimes emerges as only a small problem (e.g., Newfoundland snow crab and Western Australian snapper); in other cases (American lobsters, Dungeness crabs), ghost fishing is clearly an important biological and economic waste.

Modifications to stop traps from ghost fishing are simple and effective, and can be inexpensive. Such modifications are quick and easy to service once installed. Management agencies should determine whether ghost fishing is a problem in specific trap fisheries. If it is, they should conduct research into material failure rates and require timed-release devices or panels in all traps. Appropriate and properly designed research is required both to convince the industry of the problem and to develop effective timed-failure devices for specific situations.

For Dungeness crab fisheries, all jurisdictions now recognize the ghost fishing problem and attempt to control it. In the American lobster fishery, where ghost fishing was well documented much earlier, most jurisdictions have still not addressed the problem.

In the American lobster and British Columbia Dungeness crab fisheries, the amount of waste caused by ghost fishing would not have been recognized without appropriate experimentation. In no fishery should ghost fishing be rejected as a serious potential problem until proper research has been conducted.

Ghost fishing has been documented in a variety of coastal gillnet fisheries. Lost nets may kill fishes, crabs, birds, and seals for several years. Loss rates of coastal gillnets have not been estimated, but at least two studies indicate a substantial density of lost demersal gillnets on commercial fishing grounds.

The situation in pelagic gillnets is less clear. Loss rates are poorly estimated. At least one study indicates that ghost fishing and

continuing entanglement of birds occurs; other studies suggest that pelagic nets form tangled nonfishing masses in a short time. Further information is needed in two areas: documentation of lost gear encountered at sea, and direct study of the fishing behavior of lost pelagic gillnets.

Short of preventing net loss or prohibiting gillnetting, it is not clear how to prevent ghost fishing in gillnets. Studies of preventive measures such as using degradable hangings are embryonic. Preventive measures may simply change the form of the problem. Side effects of intended preventive measures must therefore be examined carefully.

ACKNOWLEDGMENTS

Special thanks to Richard Shomura. I am indebted to the following people who provided local information, unpublished material, references, or comments on the manuscript: James Acheson (University of Maine); Philip Averill (Maine Department of Marine Resources); Peter J. Auster (University of Connecticut); Colin Bannister (United Kingdom Ministry of Agriculture, Fisheries, and Food); Steve Barry (Washington State Department of Fisheries); J. Pike Bartlett (Friendship Trap Co., Friendship, Maine); A. J. Blott (United States National Marine Fisheries Service, Narragansett); John D. Booth (New Zealand Ministry of Agriculture and Fisheries); William Bourne (Aberdeen University); James A. Boutillier (Fisheries and Oceans Canada, Nanaimo); Mike Bradstock, consultant biologist (Auckland, New Zealand); Rhys Brown (Western Australian Fisheries Department); Richard H. Bumgarner (Washington State Department of Fisheries); H. Arnold Carr (Massachusetts Division of Marine Fisheries); Dennis Chalmers (Fisheries and Oceans Canada, Nanaimo); James M. Coe (U.S. National Marine Fisheries Service, Seattle); Steve Dawson (University of Otago, Dunedin, New Zealand); Darrell Demory (Oregon Department of Fish and Wildlife); Yoshiharu Doi (Tokyo Institute of Technology); G. P. Ennis (Fisheries and Oceans Canada, St. John's); Bruce T. Estrella (Massachusetts Division of Marine Fisheries); Alan R. Everson (U.S. National Marine Fisheries Service, Honolulu); Michael J. Fogarty (U.S. National Marine Fisheries Service, Woods Hole); Bruce Gabites (University of Auckland); Ken Gonsalves (Stevens Institute of Technology, Hoboken, New Jersey); Vivian Haist (Fisheries and Oceans Canada, Nanaimo); Rick Harbo (Fisheries and Oceans Canada, Nanaimo); Bill High (U.S. National Marine Fisheries Service, Seattle); Jeffrey A. June (National Resources Consultants, Seattle); Brian Kanenaka (Hawaii Department of Land and Natural Resources); Robert B. Kennedy (Tasmania Department of Sea Fisheries); Jay S. Krouse (Maine Department of Marine Resources); Wade Landsburg (Fisheries and Oceans Canada, Moncton); Skip McKinnell (Fisheries and Oceans Canada, Nanaimo); Sandy McFarlane (Fisheries and Oceans Canada, Nanaimo); Tracey McKenzie, (U.S. National Marine Fisheries Service, Narragansett); Rob Mattlin (New Zealand Ministry of Agriculture and Fisheries); Bob Miller (Fisheries and Oceans Canada, Halifax); Mike Moran (Western Australian Fisheries Department); Minoru Morimotu (Fisheries Agency, Government of Japan); J. Morishita (Fisheries Agency, Government of Japan); Margaret C. Murphy (Alaska Department of Fish and Game); Talbot Murray (New Zealand Ministry of Agriculture and Fisheries); Henry Okamoto (Hawaii Department of Land and Natural Resources); Frank A. Parrish (U.S. National Marine Fisheries Service,

Honolulu); Liz Slooten (Otago University, Dunedin, New Zealand); Larry Tsunoda (U.S. National Marine Fisheries Service, Seattle); Kim Walshe (New Zealand Ministry of Agriculture and Fisheries); Ronald W. Warner (California Department of Fish and Game); and Edward R. Zyblut (Fisheries and Oceans Canada, Vancouver).

REFERENCES

- Acheson, J. M.
 1982. Metal traps: A key innovation in the Maine lobster industry. In J. R. Maiolo and M. K. Orbach (editors), *Modernization and marine fisheries policy*, p. 279-312. Ann Arbor Science.
- Andrady, A.
 1988. The use of enhanced degradable plastics for control of plastic debris in the marine environment. In D. L. Alverson and J. A. June (editors), *Proceedings of the North Pacific Rim Fishermen's Conference on Marine Debris, 13-16 October 1987, Kailua-Kona, Hawaii*, p. 384-403. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199.
- Anonymous.
 1978. Ghost fishing under study. *Catch* (New Zealand Ministry of Agriculture and Fisheries) 5(8):3.
 1984. Progress in the snapper study. *Fish. Ind. News* (Western Australian Department of Fisheries) 17(1):3-6.
 1987. Proposed management policy for rock lobster. *Fish. Bull.* (New Zealand Ministry of Agriculture and Fisheries) 2(2):1-4.
 1988. Ghost panels delayed. *Commer. Fish. News* [March 1988]:17.
- Anthony, V. C., and J. F. Caddy.
 1980. Scientific recommendations for management. In V. C. Anthony and J. F. Caddy (editors), *Proceedings of the Canada-U.S. Workshop on Status of Assessment Science for N.W. Atlantic Lobster (*Homarus americanus*) Stocks*, p. 180-182. *Can. Tech. Rep. Fish. Aquat. Sci.* 932.
- Baba, N., M. Kiyota, and K. Yoshida.
 1990. Distribution of marine debris and northern fur seals in the eastern Bering Sea. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Beach, R. J., T. C. Newby, R. O. Larson, M. Pedersen, and J. Juris.
 1976. Entanglement of an Aleutian reindeer in a Japanese fish net. *Murrelet* 57:66.

- Bell, F. W., and R. F. Fullenbaum.
1986. The American lobster fishery: Economic analysis of alternative management strategies. *Mar. Fish. Rev.* 10(5):1-6.
- Blott, A. J.
1978. A preliminary study of timed release mechanisms for lobster traps. *Mar. Fish. Rev.* 40(5-6):44-49.
- Bourne, W. R. P.
1977. Nylon netting as a hazard to birds. *Mar. Pollut. Bull.* 8:75-76.
- Bowen, B. K.
1961. The Shark Bay fishery on snapper (*Chrysophrys unicolor*). Fisheries Department, Western Australia, Rep. 1:1-15.
- Breen, P. A.
1987. Mortality of Dungeness crabs caused by lost traps in the Fraser River estuary, British Columbia. *N. Am. J. Fish. Manage.* 7:429-435.
- Brown, R. S., and N. Caputi.
1986. Conservation of recruitment of the western rock lobster (*Panulirus cygnus*) by improving survival and growth of undersize rock lobsters captured and returned by fishermen to the sea. *Can. J. Fish. Aquat. Sci.* 43:2236-2242.
- Carr, H. A.
1988. Long term assessment of a derelict gill net found in the Gulf of Maine. *Oceans '88 (Proceedings of the Marine Technology Society):*984-986.
- Carr, H. A., E. H. Amaral, A. W. Hulbert, and R. Cooper.
1985. Underwater survey of simulated lost demersal and lost commercial gill nets off New England. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 438-447. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Carr, H. A., and R. A. Cooper.
1987. Manned submersible and ROV assessment of ghost gill nets in the Gulf of Maine. *Oceans '87 (Proceedings of the Marine Technology Society):*622-624.
- Center for Environmental Education.
1987. *Plastics in the ocean: More than a litter problem.* Center for Environmental Education, 1725 DeSales St., Wash., D.C., 128 p.
- Cleaver, F. C.
1949. Preliminary results of the coastal crab (*Cancer magister*) investigation. *Wash. Dep. Fish. Biol. Bull.* 49A:47-82.
-

- Coe, J. M.
1986. Derelict fishing gear: Disaster or nuisance? Master of Marine Affairs Thesis, University of Washington, Seattle, 79 p.
- Dahlstrom, W. A.
MS. Report of lost crab trap recovery. Unpubl. manuscript. Marine Resources Region, California Department of Fish and Game, unpagged.
- Dawson, S. M.
1990. Sounds, acoustic behaviour, and gillnet entanglement in Hector's dolphin. Ph.D. Thesis, Univ. Canterbury, Christchurch, New Zealand, 136 p.
- Day, R. H., D. G. Shaw, and S. E. Ignell.
1990. The quantitative distribution and characteristics of marine debris in the North Pacific Ocean, 1984-88. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- DeGange, A. R., and T. C. Newby.
1980. Mortality of seabirds and fish in a lost salmon driftnet. Mar. Pollut. Bull. 11:322-323.
- Demory, D.
1971. Abandoned crab pots near Cannon Beach, Oregon. Fish Comm. Oregon, Res. Div., Shellfish Invest. Inf. Rep. 70-6:1-5.
- Dews, G., K. Sainsbury, W. Whitelaw, and M. Moran.
1988. CSIRO goes fish trapping on NW Shelf. Aust. Fish. 47(9):28-29.
- Doi, Y., M. Kunioka, Y. Nakamura, and K. Soga.
1988. Nuclear magnetic resonance studies on unusual bacterial copolyesters of 3-hydroxybutyrate and 4-hydroxybutyrate. Macromolecules 21:2722-2727.
- Eisenbud, R.
MS. The pelagic driftnet. Unpubl. manuscript, submitted to the FAO World Conference on Fisheries Management and Development, Rome. Greenpeace International, Lewes, East Sussex, U.K., unpagged.
- Fogarty, M. J., and D. V. D. Borden.
1980. Effects of trap venting on gear selectivity in the inshore Rhode Island American lobster, *Homarus americanus*, fishery. Fish. Bull., U.S. 77:925-933.
- Fosnaes, T.
1975. Newfoundland cod war over use of gill nets. Fish. News Int. 14(6):40-43.

- Gerrodette, T., B. K. Choy, and L. M. Hiruki.
1987. An experimental study of derelict gill nets in the central Pacific Ocean. Southwest Fish. Cent. Honolulu Lab., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI. Southwest Fish. Cent. Admin. Rep. H-87-18, 12 p.
- Goldblatt, R.
1989. Drift netting [off the] Federated States of Micronesia, February 3rd - March 3rd 1989. N.Z. Professional Fisherman 3(2):27-29.
- Gonsalves, K. E.
1990. Degradable materials for the marine environment. Submitted to the Proceedings, Fisheries Conservation Engineering Special Topic Workshop. Kingston, Rhode Island, April 1990.
- Gonsalves, K. E., S. H. Patel, and X. Chen.
1989. Development of degradable materials for fishing ropes and traps. Unpubl. manusc., Polymer Processing Institute, Stevens Institute of Technology, Hoboken, N.J.
- Hancock, D. A.
1974. Attraction and avoidance in marine invertebrates--their possible role in developing an artificial bait. J. Cons. 35:328-331.
- Harper, D., and D. McClelland.
1983. Observations on the behavior and survival of trap-caught reef fish. Unpubl. manusc., U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., Miami, Fla.
- Harwood, M. B., and D. Hembree.
1987. Incidental catch of small cetaceans in the offshore gill net fishery in northern Australian waters: 1981-1985. Rep. Int. Whal. Comm. 37:363-367.
- Heneman, B., and the Center for Environmental Education.
1988. Persistent marine debris in the North Sea, northwest Atlantic Ocean, wider Caribbean area, and the west coast of Baja California. Unpublished report to the Marine Commission and the National Ocean Pollution Program Office. Center for Environmental Education, 1725 DeSales St., Wash. D.C.
- Herrick, S. F. Jr., and D. Hanan.
1988. A review of California entangling net fisheries, 1981-1986. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-108, 38 p.
- High, W. L.
1976. Escape of Dungeness crabs from pots. Mar. Fish. Rev. 38(4):19-23.

1980. Bait loss from halibut longline gear observed from a submersible. *Mar. Fish. Rev.* 42(2):26-29.
1985. Some consequences of lost fishing gear. In R. S. Shomura, and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 430-437. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
1987. Biological and technical observations of halibut longline gear from a submersible. *NOAA Symp. Ser. for Undersea Res.* 2(2):93-100.
- High, W. L., and I. E. Ellis.
1973. Underwater observations of fish behavior in traps. *Helgol. Wiss. Meeresunters.* 24:341-347.
- High, W. L., and D. D. Worlund.
1979. Escape of king crab, *Paralithodes camtschatica*, from derelict pots. U.S. Dep. Commer., NOAA Tech. Rep. NMFS-SSRF-734, 11 p.
- Hinds, V.
1984. Gill-netters in port. *Catch (New Zealand Ministry of Agriculture and Fisheries)* 11(2):6-7.
- Hipkins, F. W.
1972. Dungeness crab pots. U.S. Dep. Commer., Natl. Mar. Fish. Serv., *Fish. Facts* 3:1-13.

1974. A trapping system for harvesting sablefish. U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., *Fish. Facts* 7:1-20.
- Hipkins, F. W., and A. J. Beardsley.
1970. Development of a pot system for harvesting blackcod (*Anoplopoma fimbria*). Unpubl. manuscript, U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., Seattle, Wash., 31 p.
- Jamieson, G. S., and G. D. Heritage.
1987. Experimental flying squid fishing off British Columbia, 1985 and 1986. *Can. Ind. Rep. Fish. Aquat. Sci.* 179:1-103.
- Jow, T.
1961. Crab trap escape-opening studies. *Bull. Pac. Mar. Fish. Comm.* 5:49-71.
- Karnofsky, E. B., and H. J. Price.
1989. Behavioural response of the lobster *Homarus americanus* to traps. *Can. J. Fish. Aquat. Sci.* 46:1625-1632.
- Kennedy, R.
1986. The Dungeness crab of North America. *Safish (South Australian Department of Fisheries)* 9(6):35.

- Krouse, J. S.
1989. Performance and selectivity of trap fisheries for crustaceans. In J. F. Caddy (editor), *Marine invertebrate fisheries: Their assessment and management*, p. 307-325. Wiley, N.Y.
- Lazarus, P.
1988. Resistance to lobster escape panels slows down legislation. *Natl. Fisherm.* 69(4):69-70.
- Lovewell, S. A., A. E. Howard, and D. B. Bennett.
1988. The effectiveness of parlour pots for catching lobsters (*Homarus gammarus* (L.)) and crabs (*Cancer pagurus* L.). *J. Cons.* 44:247-252.
- Maynard, D. R., and N. Branch.
1988. Retention of lobster (*Homarus americanus*) in three compartment wire traps as a function of escape mechanisms. *CAFSAC Res. Doc.* 88/40:1-10.
- McKinnell, S. M., T. Gjernes, W. Shaw, and S. Whiteaker.
1989. Canadian North Pacific pelagic study, *Arctic Harvester*, July 12-August 22, 1989. Department of Fisheries and Oceans, Pacific Biological Station, Nanaimo, B.V., Canada V9R 5K6, 13 p. Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Seattle, Washington, U.S.A., October 1989.
- Merrell, T. R., Jr.
1984. A decade of change in nets and plastic litter from fisheries off Alaska. *Mar. Pollut. Bull.* 15:378-384.
- Methot, R. D.
1986. Management of Dungeness crab fisheries. In G. S. Jamieson and N. F. Bourne (editors), *North Pacific Workshop on Stock Assessment and Management of Invertebrates*, p. 326-344. *Can. Spec. Publ. Fish. Aquat. Sci.* 92.
- Meyer, R. M.
MS. A study concerning the problem of derelict pots in the king crab fishery. Unpubl. manuscr. *Natl. Mar. Fish. Serv.*, NOAA, Kodiak, Alaska, 8 p.
- Miller, R. J.
1976. North American crab fisheries: Regulations and their rationales. *Fish. Bull.*, U.S. 74:623-633.
1977. Resource underutilization in a spider crab industry. *Fisheries* (Bethesda) 2(3):9-13.
1979. Saturation of crab traps: Reduced entry and escapement. *J. Cons.* 38:338-345.

- Millner, R. S.
1985. The use of anchored gill and tangle nets in the sea fisheries of England and Wales. Ministry Agriculture, Fisheries and Food, Lowestoft, Suffolk, U.K., Laboratory Leaflet 57:1-22.
- Mio, S., T. Domon, K. Yoshida, and S. Matsumura.
1990. Preliminary study on change in shape of drifting nets experimentally placed in the sea. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Moran, M. J., and J. Jenke.
1989. Effects of fish trapping on the Shark Bay snapper industry. Western Australian Department of Fisheries, Fish. Rep. 82:1-29.
- Morgan, G. R.
1974. Aspects of the population dynamics of the western rock lobster, *Panulirus cygnus* George. II. Seasonal changes in the catchability coefficient. Aust. J. Mar. Freshwater Res. 25:249-259.
- Mottet, M. G.
1975. The fishery biology of *Octopus dofleini* (Wulker). Wash. Dep. Fish. Tech. Rep. 16:1-39.
- Muir, W. D., J. T. Durkin, T. C. Coley, and G. T. McCabe, Jr.
1984. Escape of captured Dungeness crabs from commercial crab pots in the Columbia River estuary. N. Am. J. Fish. Manage. 4:552-555.
- Munro, J. L.
1974. The mode of operation of Antillean fish traps and the relationships between ingress, escapement, catch and soak. J. Cons. 35:337-350.

1983. The composition and magnitude of trap catches in Jamaican waters. In J. L. Munro (editor), Caribbean coral reef fishery resources, p. 33-49. ICLARM Studies and Reviews 7.
- Munro, J. L., P. H. Reeson, and V. C. Gaut.
1971. Dynamic factors affecting the performance of the Antillean fish trap. Proc. Gulf Caribb. Fish. Inst. 23:184-194.
- Murray, T.
1988. Drift netters threaten South Pacific fisheries. Catch (New Zealand Ministry of Agriculture and Fisheries) 15(11):13-14.
- Northup, T.
1978. Development of management information for coastal Dungeness crab fishery. Project Completion Report, Project No. 1-114-R.

- Paul, L. M. B.
1984. Investigations into escape vent effectiveness and ghost fishing in captive populations of the spiny lobster, *Panulirus marginatus*. In R. W. Grigg and K. Y. Tanoue (editors), Proceedings of the Second Symposium on Resource Investigations in the Northwestern Hawaiian Islands, Vol. 2, p. 283-295. University of Hawaii Sea Grant Report MR-84-01.
- Pecci, K. J., R. A. Cooper, C. D. Newell, R. A. Clifford, and R. J. Smolowitz.
1978. Ghost fishing of vented and unvented lobster, *Homarus americanus*, traps. Mar. Fish. Rev. 40(5-6):9-24.
- Piatt, C.
1988. Florida fishermen are slow to adopt plastic traps. Natl. Fisherm. 69(4):66-70.
- Read, A. J., and D. E. Gaskin.
1988. Incidental catch of harbor porpoises by gill nets. J. Wildl. Manage. 52:517-523.
- Ritchie, L. D.
1972. Octopus predation on pot-caught rock lobster, Hokianga area, New Zealand. N.Z. Fish. Tech. Rep. 81:1-44.
- Scarratt, D. J.
1965. Predation of lobsters (*Homarus americanus*) by *Anonyx* sp. (Crustacea, Amphipoda). J. Fish. Res. Board Can. 30:1370-1373.
- Scarsbrooke, J. R., G. A. McFarlane, and W. Shaw.
1988. Effectiveness of experimental escape mechanisms in sablefish traps. N. Am. J. Fish. Manage. 8:158-161.
- Schrey, E., and G. J. M. Vauk.
1987. Records of entangled gannets (*Sula bassana*) at Helgoland, German Bight. Mar. Pollut. Bull. 18(B):350-352.
- Scott, G.
1990. The philosophy and practice of degradable plastics. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Sheldon, W. W., and R. L. Dow.
1975. Trap contributions to losses in the American lobster fishery. Fish. Bull., U.S. 73:449-451.
- Sloan, N. A.
1984. Canadian-Japanese experimental fishery for oceanic squid off British Columbia, summer 1983. Can. Ind. Rep. Fish. Aquat. Sci. 152:1-42.

- Smolowitz, R. J.
1978a. Trap design and ghost fishing: An overview. *Mar. Fish. Rev.* 40(5-6):2-8.
1978b. Trap design and ghost fishing: Discussion. *Mar. Fish. Rev.* 40(5-6):59-67.
1978c. An annotated bibliography on lobster trapping and related subjects. *Mar. Fish. Rev.* 40(5-6):68-77.
- Sutherland, D. L., G. L. Beardsley, and R. S. Jones.
1983. Results of a survey of the south Florida fish-trap fishing grounds using a manned submersible. *Northeast Gulf Sci.* 6(2):179-183.
- Tsunoda, L. M.
1989. Observations on board a Japanese, high-seas, squid gill net vessel in the North Pacific Ocean July 1-August 14, 1986. U.S. Dep. Commer., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWAFC Processed Rep. 89-02.
- Uchida, R. N.
1985. The types and estimated amounts of fish net deployed in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 37-108. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Von Brandt, A.
1984. *Fish catching methods of the world.* Fishing News Books, U.K., 418 p.
- Way, E. W.
1977. Lost gill net (ghost net) retrieval project 1976. Environment Canada, Fisheries and Marine Service, Industrial Development Branch, St. John's, Newfoundland, 30 p.
- Wilder, D. G.
1945. Wider lath spaces protect shorter lobsters. *Fish. Res. Board Can., Atl. Biol. Stn. Circ.* 4:1.

AN EXPERIMENTAL STUDY OF DERELICT GILLNET
FRAGMENTS IN THE CENTRAL PACIFIC OCEAN

Tim Gerrodette,* Barry K. Choy,** and Lisa M. Hiruki***
Southwest Fisheries Science Center Honolulu Laboratory
National Marine Fisheries Service, NOAA
Honolulu, Hawaii 96822, U.S.A.

*Present address: Southwest Fisheries Science Center La Jolla Laboratory,
National Marine Fisheries Service, NOAA, La Jolla, California 92038, U.S.A.

**Present address: c/o NOAA Ship *John N. Cobb*, 1801 Fairview Avenue E,
Seattle, Washington 98107, U.S.A.

***Present address: Department of Zoology, University of Alberta, Edmonton,
Alberta T6G 2E9, Canada.

ABSTRACT

An experiment designed to investigate the behavior and fate of derelict gillnet fragments was initiated in August 1986 in the central Pacific Ocean. Four fragments of high-seas squid gillnet, varying in length from 50 to 1,000 m, were observed closely for 3 days and subsequently tracked for up to 10 months by satellite. The net fragments changed length, shape, heading, and location under the influence of wind and current. The time a net remained open in a fishing configuration varied from hours to weeks, depending on its initial length. The nets drifted at an average speed of 15 km/day, but with frequent changes in direction, they remained in the general vicinity of the Hawaiian Archipelago. The complex movement of the net fragments means that predicting the drift of marine debris is an oceanographic problem that requires detailed knowledge of surface currents and wind.

INTRODUCTION

The amount of debris in the world ocean is a matter of increasing concern, both to the scientific community and the public at large. The Workshop on the Fate and Impact of Marine Debris (Shomura and Yoshida 1985) focused attention on the problem, and the National Marine Fisheries Service (NMFS) shortly afterwards established a program (Coe and Bunn 1987) to coordinate research, public awareness, and mitigation efforts.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

One source of marine debris is fishing operations. The amount of fishing gear in use is staggering. For example, considering only gillnets in the North Pacific, at least 180,000 km of net is available to the various gillnet fisheries (Chen 1985; Gong 1985; Shima 1985; Uchida 1985), an amount that would stretch 4.5 times around the Earth. (This figure is conservative; it does not include gillnet in the coastal fisheries of Korea and Taiwan.) Even if only a small percentage, for example, 0.05% (Komatsu 1986), is lost in the course of these fishing operations, 90 km of gillnet would enter the North Pacific Ocean every time these nets are used. Gillnets may be lost as a result of storms, cut adrift by a ship crossing the float line, or discarded overboard after being damaged. Such derelict gillnets are a cause for concern because they may 1) continue to catch fish, leading to waste of marine resources and inaccurate estimates of fishing mortality; 2) present a hazard to navigation by fouling ships' propellers; and 3) ensnare and kill such nontarget species as seals, dolphins, whales, turtles, and seabirds.

The impact a piece of derelict gillnet will have depends on its size, shape, location, and length of time in the ocean. This paper reports the results of experiments designed to investigate some of these questions. Specifically, the objectives of the study were to measure the change in shape of derelict gillnet fragments of various sizes over time, to determine the fishing ability of derelict gillnets of known age, and to track the movement of drifting net fragments for periods up to 1 year.

METHODS

Thirty sections ("tans") of used, 113-mm monofilament gillnet of the type used in the Japanese high-seas squid fishery were purchased from Kyoei Unyu Company, Ltd., Hakodate, Japan. Each section measured 50 m long and 9 m deep, with floats at 1-m intervals. Sections were joined together to make four nets, 50, 100, 350, and 1,000 m long.

Attached to each of the four nets was a small, dual-frequency, radio-satellite transmitter buoy (Fig. 1), designed by Telonics of Mesa, Arizona. The buoy, 90 cm in length and 9 cm in diameter, allowed tracking and potential recovery of each net. The UHF satellite transmitter portion of each buoy used the Argos system (Argos 1984) to give a location on the Earth's surface accurate to within several hundred meters. The satellite transmitter broadcast on a schedule of 24 h on, 72 h off; a series of locations was, therefore, available once every 4 days. The VHF radio transmitter portion of each buoy allowed close-range directional tracking and recovery within a radius of approximately 10 km. The radio transmitter broadcast once a second without interruption. The combination of long- and short-range location systems was designed to allow physical recovery of the buoy and net after drifting freely in the ocean for up to 18 months.

To reduce windage and to avoid accidental discovery by fishermen or others, the buoy also was designed to be as inconspicuous, both visually and electronically, as possible. The buoy projected only 25 cm above the ocean's surface. Further, the megahertz frequencies transmitted by the buoy's location systems were beyond the kilohertz frequencies commonly used in ships' radio direction finders (RDF's) for locating buoys.




Figure 1.--Dual-frequency transmitter buoy used to track and recover experimental gillnets.

The nets and their associated buoys were deployed on 12 August 1986 from the NOAA ship *Townsend Cromwell* about 10 km east of Southeast Hancock Seamount, northwest of the Hawaiian Archipelago (Table 1). The nets were deployed by letting the ship drift downwind; hence, all nets were initially set parallel to the wind with the transmitter buoy at the downwind end. Measurements of net heading, length, and catch were made three times a day for 3 days from an inflatable boat. A temporary buoy was attached to one end of the net to serve as a visual target; from the other end, heading was then measured with a hand-held bearing compass, and length with an optical range finder. The configuration of each net was sketched. The longest (1,000-m) net was surveyed in a similar way, except that several visual targets were placed along its length, and measurements made in sections. Catch and fish aggregation around the nets were monitored by snorkel or scuba diving and documented photographically with video and 35-mm cameras.

Observations and measurements of the nets were confined to daylight hours. To track the nets during the night, larger buoy systems were attached to some nets each night and removed the next morning. Such a system consisted of a large RDF transmitter buoy; a long bamboo pole buoy with strobe light on top and large inflatable float; and a small, round, plastic buoy at the end of a tag line. The whole system had considerable windage. Because such a system probably affected a net's dynamics, the periods during which a large transmitter buoy system was attached to a net were considered in the interpretation of the results. The small radio-satellite transmitter buoys (Fig. 1), which were attached to the nets at all times, were considered to have negligible effects.

On the 10th day after deployment, the *Townsend Cromwell* returned and relocated all four nets. Rough seas, however, prevented launching a small boat, and the nets had to be observed from the deck of the *Townsend Cromwell*. After the cruise, the buoys were tracked by satellite until each buoy was either recovered or the signal from the satellite transmitter was lost. Positions were determined from monthly reports of Service Argos, Toulouse, France.

Table 1.--Experimental gillnet deployment on 12 August 1986 and tracking in the central Pacific Ocean. Dates and times are Midway standard time.

Buoy No.	Net length (m)	Deployment			Recovery			Days tracked (No.)
		Time (h)	Latitude N	Longitude E	Date	Latitude N	Longitude	
10013	50	0841	29°46.7'	179°10.3'	11/3/86	28°48.9'	176°57.6'W	83
10010 ^a	100	0945	29°46.8'	179°09.8'	1/7/87	28°14.2'	178°13.2'W	148
10011 ^a	350	1226	29°47.0'	179°08.6'	10/8/86	29°35.2'	176°18.9'W	57
10012	1,000	1330	29°47.6'	179°08.0'	6/17/87	23°00.1'	178°00.0'E	309

^aBuoy was not recovered; recovery data reflect time and location at which the signal was lost.

RESULTS

Shape and Heading

During the first 3 days, fair weather and calm seas greatly aided tracking and observation of the nets. Wind was east-southeast during this initial period but shifted to east-northeast on the second and third days and rose slightly in strength (Table 2). The nets first drifted north-northwest, then north-northeast, traveling about 9 km/day.

The 50- and 100-m nets shortened soon after deployment (Fig. 2). The 50-m net, in fact, had already collapsed by the time of the first observation, 30 min after deployment. "Collapsed" means that the net was folded like an accordion and all floats were close together. The net, however, was still hanging freely in the water; it was not tangled with itself.

The 350-m net contracted to about 40% of its original length during the first few hours, but then contracted more slowly (Fig. 2). By the 10th day, it had collapsed completely. The rate of collapse of this net may have been affected by the large transmitter buoy attached to the downwind end during the first night. The net was slightly longer the next morning (observation at 21 h). After removal of the large buoy system, the net further contracted (25 h) but was longer in the evening (29 h). The next day, the net followed a similar pattern, contracting between morning and afternoon and lengthening by evening. Greater detail of the changes in configuration of this net is shown in Figure 3A. Interestingly, the net rotated so that its heading 50 h after deployment was approximately 140° from its original heading.

Table 2.--Summary of wind and swell observations on 12-14 August 1986, the first 3 days of the gillnet experiment. Data are means calculated from the ship's hourly weather log.

Date	Time (h)	Wind		Swell	
		Speed (kn)	Heading	Height (ft)	Heading
12 August	0100-1200	9	116°	3	121°
	1300-2400	11	115°	3	116°
13 August	0100-1200	10	100°	3	118°
	1300-2400	14	70°	4	98°
14 August	0100-1200	13	74°	3	88°
	1300-2400	13	68°	4	83°




Figure 2.--Percentage of original lengths of four experimental derelict gillnets over time. During the first and second nights of observation, large transmitter buoys attached to certain nets may have affected net lengths.




Figure 3.--Shape and heading of two experimental derelict gillnets at various points in time after deployment. The original windward end of the net (without the transmitter buoy) was placed at the origin (★) of each sketch. A) 350-m net. B) 1,000-m net.

The 1,000-m net followed a pattern of contraction and expansion similar to the 350-m net, except that its relative size 52 h after deployment was only about 50% of original length instead of 25% (Fig. 2). The 1,000-m net also rotated in the same direction as the 350-m net, but only by about 30° during the same period. By the 10th day (218 h), its heading had changed completely (Fig. 3B).

The time required for a gillnet to collapse was related to its original length. Figure 4 shows the time required for the various fragments of net to collapse to 10% of their original length as a function of their original length.

Catch

Very little was caught in any of the gillnets during the initial 3 days of observation. On the morning of the second day, a small marlin (*Makaira* sp.), about 1 m total length, was entangled in the 1,000-m net at the surface. On the third day, a large flyingfish (Exocetidae) was similarly caught in the same net. None of the other nets had any animals entangled in them by the end of the third day. Three small kahala (*Seriola* sp.) were observed swimming around the 350-m net on the second day. One opelu (*Decapterus* sp.) and three small kahala were seen around the 1,000-m net on the third day. Soon after the nets were deployed, several albatross (*Diomedea* sp.) landed on the water near the float line, but each left after a short investigation.

After 10 days, nothing was visible in the 50-, 100-, or 1,000-m nets, although one mahimahi, *Coryphaena hippurus*, was swimming near the latter. The floats of the 350-m net were in a tight group with numerous small kahala swimming nearby. A rotting, 2-m shark of undetermined species was entangled in the net, together with several bony fish too rotten to identify.

Movement

The location of each net during the entire course of the study, plotted once every 4 days, is shown in Figure 5. The number of days the buoys were tracked ranged from 57 to 309 (Table 1). The buoys and nets stayed in the general vicinity of the northwestern end of the Hawaiian Archipelago. For several months they remained north of Midway, then moved south. After 83 days at sea, the 50-m net and buoy 10013 were recovered by the *Townsend Cromwell*. Several species of fish were swimming near the net, and two pilotfish, *Naucrates ductor*, were caught in it (Table 3). No large animals were entangled in the net.

Buoy 10012, which was tracked the longest, traveled as far south as lat. 17°37.8'N, then returned north and west (Fig. 5). It was recovered after 309 days at sea by the chartered fishing vessel *Feresa*. The 1,000-m net was no longer attached to the buoy at that time. It is not known when the net became separated from the buoy, but the absence of barnacles, together with damage to the buoy, suggested that separation may have occurred only a short time before recovery. The buoy failed to transmit a




Figure 4.--Time required for a fragment of gillnet to collapse to 10% of original length as a function of its original length.

position on 8 June 1987, 9 days before recovery (Fig. 5), and was possibly entangled in the net at that time.

The two remaining nets and buoys were not recovered because their signals were lost. Signals stopped after 57 days for the 350-m net and 148 days for the 100-m net (Table 1). The reason for signal loss is not known, but the most likely explanation is that the buoys became entangled in the nets and submerged. Buoy 10010 on the 100-m net stopped transmitting about 15 km southeast of Kure Atoll (Fig. 5). Possibly the net became caught on the reef, but searches by plane and boat in August 1987 failed to find it.

For each 4-day interval, the mean speed of each buoy was computed. Mean speed per 4-day interval varied widely, from less than 1 km/day to nearly 50 km/day (Fig. 6). The 4-day mean speeds reflect several types of water movement: advection, inertial movement, and other eddies of various scales. Overall mean speed was 14.8 km/day or about 0.3 kn. The frequent and abrupt changes in speed and direction, however, meant that the distance




Figure 5.--Positions of four experimental derelict gillnets in the central Pacific Ocean, plotted at 4-day intervals. Dotted lines connect positions more than 4 days apart.

from the point of release did not bear any simple relation to time (Fig. 7). The four buoys traveled more or less together as long as they were tracked; the buoy tracked the longest (buoy 10012) drifted nearly 1,500 km from the point of deployment, then returned (Figs. 5, 7).

DISCUSSION

The amount of gillnet that becomes lost, detached, or discarded in the course of gillnet fishing operations is not known with any precision. Based on fishing activity, however, the total amount is undoubtedly large. The loss rate of 0.05%, mentioned earlier, is an unsubstantiated estimate given by a Japanese Government official (Komatsu 1986) during public hearings on the incidental catch of marine mammals during high-seas driftnet salmon fishing. Eisenbud (1985), citing a 1982 letter from Richard B. Roe,




Figure 6.--Mean speed in each 4-day interval for the four gillnets, plotted as a function of time from deployment.




Figure 7.--Distance of each fragment of gillnet from the point of deployment, plotted as a function of time from deployment.

Table 3.--Biological observations near the 50-m gillnet recovered on 3 November 1986 north of Midway. The net had been drifting for 83 days. *N* = number of individuals sighted.

Species sighted	<i>N</i>	Sighting location
Pilotfish, <i>Naucrates ductor</i>	2	Caught in net.
Barracuda, <i>Sphyraena helleri</i>	20	Swimming near net.
Mahimahi, <i>Coryphaena hippurus</i>	2	Swimming near net.
<i>Alutera scripta</i>	11	Swimming near net.
<i>Naucrates ductor</i> juveniles	14	Swimming near net.
Unidentified fish, possibly <i>Kyphosus</i> sp.	7-12	Swimming near net.
Black-footed albatross, <i>Diomedea nigripes</i>	7	On surface near net.

NMFS, mentions an estimated 0.06% loss rate for the same fishery. Such a low loss rate, even if accurate for this fishery, is unlikely to apply to other types of gillnet fishing. For example, coastal gillnetting operations are likely to have a higher rate of net loss because of more boat traffic and a greater chance of nets becoming hung up on the bottom. Even so, applying this minimum loss rate to the total amount of available gillnet means that thousands of kilometers of derelict gillnet enter the North Pacific every year.

The fishing ability of a net depends on its size and configuration in the water (see also Mio et al. 1990). Left alone, a drifting gillnet will eventually collapse and become entangled with itself. The rate at which this happens depends, among other things, on the original length of the net. Rapid collapse of short sections of net is expected because of the weight of the lead line and local turbulence; longer nets have greater resistance to these small-scale effects. Over the range of sizes of gillnet fragments used in this study, it appeared that the rate of collapse was approximately 100 m/day (Fig. 4). Thus, net fragments less than 100 m long collapsed in less than a day, those several hundred meters long in several days, and those 1 km or longer in several weeks. These rates give a first approximation of the length of time a derelict gillnet would remain in an active fishing configuration. Note that these estimates apply to *intact* fragments of gillnet--that is, with both float and lead lines attached. The absence of a lead line, in particular, might affect the rate at which a net fragment collapses.

The rate of collapse may also depend on other factors. High wind and swell may make the net collapse faster. If a large animal, such as a shark or seal, is caught in a net, its struggling may also hasten the collapse of the net. If a buoy is attached to one end of the net, the force of wind on the buoy may keep the net open much longer, as demonstrated by the effect of the large transmitter buoy system on the 100-m net in this study. After 1 day, the net was completely collapsed, but after the large buoy system was attached to it overnight, the net lengthened (Fig. 2). The nearby 50-m net, which did not have a large buoy system attached to it, did not

lengthen during the same period. The force of the wind on the large buoy, which was at the downwind end of the net, caused a constant pull on one end of the net and was the likely cause of its lengthening.

Once collapsed, a gillnet is still capable of catching fish, though much less effectively. The rapid collapse of the nets in this study suggests that the catch rate of a lost or discarded gillnet will, for the target species, decline rapidly. Whether the hazard of a derelict gillnet also declines rapidly for nontarget species, however, is not resolved by this study. A floating mass of net will attract fish that may, in turn, attract predators like birds, sharks, seals, and dolphins.

The movement of debris on the ocean's surface is controlled by a combination of wind and surface currents. The gillnets used in this study have a large surface area in the water and little above it. Hence, their movement over a period of months (Fig. 5) reflects mainly the movement of the upper 10 m of water rather than wind drift. Currents in the Hawaiian Archipelago are complex and irregular (Wyrcki et al. 1969). Eddies of various sizes are common in Hawaiian waters (Seckel 1955; Patzert 1969), and the loops executed by buoys 10010 and 10012 may indicate such eddies. Movements on smaller space and time scales, such as inertial motion, are not resolved by the 4-day interval between buoy positions in this study. Inertial circling was observed in the finer scale measurements of Matsumura et al. (1990).

The abrupt changes in speed and direction of the nets in this study illustrate that predicting the movement of marine debris is a difficult problem (Galt 1985; Seckel 1985). The movement of marine debris can be approached both experimentally and through simulation modeling (Matsumura et al. 1990). At least around the Hawaiian Archipelago, simple models of linear motion (distance proportional to time) or diffusion (distance proportional to the square root of time) will not predict the movement of derelict gill nets (Fig. 7). The general problem of predicting the movement and fate of debris in the ocean requires greater knowledge of factors affecting the "birth" and "death" rates of the various "species" of marine debris (Gerrodette 1985).

ACKNOWLEDGMENTS

Thanks to the captains and crews of the NOAA ship *Townsend Cromwell* and fishing vessel *Feresia* for their assistance in gillnet deployment and recovery. Michael Seki, NMFS, Honolulu, contributed the observations in Table 3. Alan Reichman, Greenpeace, Seattle, provided several references on gillnet entanglement. Partial financial support from the Marine Entanglement Research Program is gratefully acknowledged.

REFERENCES

- Argos.
1984. Location and data collection satellite system user's guide. Service Argos Inc., Landover, MD 20785, 36 p.

- Chen, T. F.
 1985. High sea gill net fisheries of Taiwan. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 253-256. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Coe, J. M., and A. R. Bunn.
 1987. Description and status of tasks in the National Oceanic and Atmospheric Administration's Marine Entanglement Research Program for fiscal years 1985-1987. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. Processed Rep. 87-15, 39 p.
- Eisenbud, R.
 1985. Problems and prospects for the pelagic driftnet. Boston Coll. Environ. Aff. Law Rev. 12:473-490.
- Galt, J. A.
 1985. Oceanographic factors affecting the predictability of drifting objects at sea. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 497-507. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Gerrodette, T.
 1985. Toward a population dynamics of marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 508-518. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Gong, Y.
 1985. Distribution and migration of flying squid, *Ommastrephes bartrami* (Lesueur), in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 109-129. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Komatsu, M.
 1986. Statement of Masayuki Komatsu, International Affairs Division, Ministry of Agriculture, Japan Fisheries Agency. [Presented at the public hearings before the National Marine Fisheries Service on the take of marine mammals incidental to commercial salmon fisheries operation, MMPAH-1986-01, 7 p.]
- Matsumura, S., Y. Wakata, and Y. Sugimori.
 1990. Movements of floating debris in the North Pacific. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

- Mio, S., T. Domon, K. Yoshida, and S. Matsumura.
1990. Preliminary study on change in shape of drifting nets experimentally placed in the sea. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154. [See this document.]
- Patzert, W. C.
1969. Eddies in Hawaiian waters. Hawaii Inst. Geophys. HIG-69-8, 57 p.
- Seckel, G. R.
1955. Mid-Pacific oceanography, Part VII, Hawaiian offshore waters, September 1952-August 1953. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 164, 250 p.

1985. Currents of the tropical and subtropical North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 461-482. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Shima, K.
1985. Summary of Japanese net fisheries in the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 252. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.
- Uchida, R. N.
1985. The types and estimated amounts of fish net deployed in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 37-108. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Wyrteki, K., V. Graefe, and W. Patzert.
1969. Current observations in the Hawaiian Archipelago. Hawaii Inst. Geophys. HIG-69-15, 97 p.

PRELIMINARY STUDY ON CHANGE IN SHAPE OF DRIFTING
NETS EXPERIMENTALLY PLACED IN THE SEA

Shin-ichi Mio, Takashi Domon, Kazumoto Yoshida, and Satsuki Matsumura
National Research Institute of Far Seas Fisheries
Fisheries Agency, the Government of Japan
Shimizu-shi, Shizuoka, 424 Japan

ABSTRACT

Research activities to understand the impact of lost drifting nets on marine organisms were initiated in 1988, and an experiment to clarify how the lost nets change their shape at sea was conducted as the first stage of the activities.

Five driftnet sets (40 tans each) were placed in the water in the area around lat. 38°N, long. 158°E, and their shapes were observed from 5 to 25 May. After this, the net sets were allowed to drift, and about 3 months later (in early September) were again observed and subsequently retrieved.

The observations were visual, and recordings were made using a camera-equipped balloon and a video camera from a research vessel.

Three days after setting, one of the nets began twisting into a mass near each end of the net. As time passed, the mass grew larger: the ends of the net approached each other and the net folded in half. Each mass continued to grow, and several small masses also were formed in portions of the long, overlapped net. Twenty days after setting, the net had become one large mass.

All nets observed became masses in the same way, although the speed of formation varied. In September, when the research vessel visited these nets again, each was found floating in a mass.

INTRODUCTION

It has been noted that fishing nets, especially gillnet fragments (hereafter referred to as drifting nets), drift in the sea out of man's control and continue catching marine organisms such as fish while drifting. However, there has been only fragmentary information concerning movements of drifting nets and the actual damage done to marine organisms. Therefore,

since 1988 we have been conducting research on the movement and changes in shape of drifting nets as well as their impact on marine organisms. This paper examines the changes in the shape of drifting nets in the course of time. Gillnets are thought to be the most effective means of catching fish, as they are set in the sea in a straight line. Once freed from man's control, gillnets are believed to change shape as time passes, with their fishing efficiency gradually declining. We conducted a survey on changes in the shape of drifting nets over time in order to establish a basis for research on related changes in fishing efficiency.

METHODS OF SURVEY

The research was conducted in the North Pacific Ocean on the salmon driftnet fishing ground at lat. 35° - 45° N, long. 150° E- 180° . The first survey was conducted from 1 to 30 May 1988, and the second from 17 August to 30 September. Five nets were used for the experiments, and each consisted of 40 tans of nylon monofilament gillnet with a mesh size of 115 mm.

An Argos buoy and a "self-call" buoy were attached to the net, one at each end. The location of each net was recorded an average of six times a day using information from the Argos buoy.

On the first cruise, experimental drifting nets were observed by sighting from on board the vessel and photographing from above with a remote control camera attached to a balloon. On the second cruise, experimental drifting nets were located using information from the buoy. They were retrieved after visual confirmation as well as confirmation through a remote control television attached to a balloon.

RESULTS

The six experimental nets, stretched tight, were set in the area lat. $39^{\circ}20'$ - $38^{\circ}43'$ N, long. $154^{\circ}33'$ - $155^{\circ}44'$ E from 1406 on 6 May to 1710 on 7 May. The experimental nets were observed a total of only 16 times, since they moved in two different directions after setting, with two drifting northeast and the other three southeast. Three nets were observed four times and two were observed twice before being retrieved (Table 1).

Except for net No. 1, each experimental net showed generally the same pattern of changing although they differed in pace. First, each end of the net twisted and formed a small mass (Fig. 1A). Second, each net folded in half and its two ends approached each other. The two ends formed a mass, twisting with each other, and the rest of the net stretched long, overlapping more and more (Fig. 1B and C). Third, as time passed, the stretched part wound around the mass. After reaching the third stage, the stretched part of the net formed a mass slowly, becoming entangled and disentangled. Observed 15 and 18 days after release, it was 50 to 60 m long compared with its original length of 2 km, indicating that it did not need many days to become a complete mass. When the five experimental nets were all collected after drifting for a long time, each net had formed a complete mass (Fig. 1D).

Table 1.--Trajectory and width ($\leftarrow \rightarrow$) change of six floating nets in 1988.

Net No.	Date	Width (m)	Latitude N	Longitude E
1	25 May	2,000	40°04'	153°06'
	15 Sept.	5	40°20'	161°08'
2	7 May	2,000	39°19'	154°50'
	10 May	1,250	39°10'	155°12'
	12 May	120	39°19'	155°25'
	18 May	250	38°12'	159°04'
	23 May	60	38°31'	158°15'
	3 Sept.	5	33°56'	169°15'
3	7 May	2,000	39°05'	155°28'
	11 May	160	38°14'	156°52'
	19 May	130	37°18'	160°14'
	28 Aug.	5	40°51'	171°28'
4	7 May	2,000	38°42'	155°19'
	12 May	310	38°14'	154°55'
	18 May	120	37°29'	158°49'
	31 Aug.	5	35°34'	179°46'
5	7 May	2,000	38°48'	154°04'
	13 May	800	39°25'	153°58'
	16 May	500	39°23'	153°56'
	21 May	250	40°06'	153°45'
	25 May	50	40°12'	153°05'
	11 Sept.	5	39°57'	158°10'
6	6 May	2,000	39°06'	154°33'
	8 May	1,150	39°20'	154°28'
	10 May	1,080	39°34'	153°23'
	16 May	600	39°19'	158°07'
	22 May	180	39°32'	153°19'

Note: Measurement of width ($\leftarrow \rightarrow$) refers to Figure 1.

As for the time required to reach each stage, the five nets (excluding net No. 1) remaining in this experiment can be divided into two groups (Fig. 2): One group needed 4 to 5 days after release to reach the third stage; the other needed 14 to 16 days to reach the third stage.

The approach of both ends of a drifting net is the basic process of changing the shape. The structure and arrangement of nets and accompanying buoys also seem to affect the changes. In this experiment, buoys were




Figure 1.--Schematic diagram showing formation of a mass of floating net after setting ((← →) denotes width of floating net).





Figure 2.--Relationship between floating period and longest width of net.


attached at both ends of a net. It may be assumed that the difference in resistance between the buoys and the net helped the two ends approach each other. The Argos buoy is a cylinder 160 mm in diameter and 790 mm in height, with a float 440 mm in diameter and 200 mm in height. It is quite small compared to the size of the net. Eight days into the experiment, the Argos buoy temporarily attached to No. 4 net dropped off, and one end of the net was subject to the same resistance as the net alone. However, both ends of the No. 4 net approached each other in the same way as the other nets.

As both ends behaved in a manner similar to other nets with buoys attached on both ends, it is suggested that there would be no changes in the basic configuration even though the presence of buoys may affect the speed of shape change. We plan to conduct further experiments to study effects of different conditions on net shape changes.

SESSION III



INGESTION BY MARINE LIFE



THE EFFECTS OF INGESTED PLASTIC AND OTHER MARINE DEBRIS ON SEABIRDS

Peter G. Ryan
Percy FitzPatrick Institute of African Ornithology
University of Cape Town
Rondebosch 7700, South Africa

ABSTRACT

Seabirds ingest plastic particles and other marine debris more frequently than do any other taxon. Despite considerable speculation as to the adverse effects ingested debris has on seabirds, there have been few experiments designed to test these hypotheses. Initial attempts to demonstrate adverse effects were based on correlations between plastic load and bird condition. However, unless the influence of season and breeding status are removed, negative correlations cannot be used to infer an adverse effect from ingesting plastic. Few statistically significant negative correlations have been found among adequately controlled samples, suggesting that the effects of ingestion are either relatively minor or that they frequently are masked by other variables. To avoid ambiguous results, carefully designed experiments are required to assess the severity of the specific adverse effects that have been hypothesized to result from debris ingestion.

Ingested debris may have three specific effects on seabirds: physical damage and blocking of the digestive tract, impairment of foraging efficiency, and the release of toxic chemicals. The severity of these effects depends upon the types of debris ingested and their retention time within seabirds.

At present, severe physical damage and obstruction of the digestive tract is infrequent in seabirds, and probably affects only a small proportion of populations. Virgin (raw) plastic particles were not found to affect the assimilation efficiency of the white-chinned petrel, *Procellaria aequinoctialis*, but seabird feeding may be affected by large plastic loads that reduce the food storage volume of the stomach, causing reduced meal size and, consequently, the ability to accumulate energy reserves. Experiments on free-ranging seabirds are required to confirm this, potentially the most serious consequence of plastic ingestion by seabirds. An estimate of the critical load size is required to determine the proportion of populations likely to be affected by reduced food intake.

Little is known about the transfer of toxic compounds from ingested plastic to seabirds, but a significant positive correlation between polychlorinated biphenyls and plastic loads in the great shearwater, *Puffinus gravis*, independent of other organochlorines, suggests that the pathway exists. This warrants confirming experimentally, and the toxicity of various additives such as plasticizers and colorants needs to be determined. Other types of debris such as tar balls and paint also are sources of toxic chemicals to seabirds; the incidence of their ingestion needs to be investigated.

Not all birds are equally vulnerable to the effects of ingested debris. Species that seldom regurgitate indigestible stomach contents, and thus accumulate large debris loads, are most prone to adverse effects. Immature petrels apparently are particularly vulnerable because they cannot unload their accumulated debris by feeding chicks.

INTRODUCTION

On a global scale, seabirds ingest plastic particles and other marine debris more frequently than do any other taxon. At least 82 seabird species out of 140 examined have been found to contain ingested plastic and other debris, and the incidence of ingestion exceeds 80% of individuals in several species (e.g., Day et al. 1985; Fry et al. 1987; Ryan 1987b; Sileo, Sievert, Samuel, and Fefer 1990). Most work to date on plastic and other debris ingestion by seabirds has focused on recording the incidence of ingestion in various taxa (e.g., Day et al. 1985; Furness 1985a, 1985b; van Franeker 1985; Ryan 1987b; van Franeker and Bell 1988; Sileo, Sievert, Samuel and Fefer 1990). This has proved valuable in several ways; in addition to providing baseline data on the temporal and spatial increase in debris at sea (e.g., Rothstein 1973; Baltz and Morejohn 1976; Harper and Fowler 1987), it has helped raise public awareness of the marine debris problem, and has provided an insight into the dynamics of ingested plastic in seabird populations (e.g., Day et al. 1985; Ryan 1988b). However, despite being the most important question arising from plastic ingestion, there have been few studies on the severity of adverse effects resulting from plastic ingestion by seabirds (Day et al. 1985; Azzarello and van Vleet 1987; Ryan 1987a). This paper reviews what is known of the impacts of ingested plastic and other debris in seabirds, and identifies key areas for future research.

GENERAL INDICATORS OF ADVERSE EFFECTS

Most attempts to demonstrate adverse effects resulting from plastic and other debris ingestion by seabirds have been based on correlations between debris load and indicators of bird condition (Day et al. 1985; Ryan 1987a). Weak negative correlations between plastic loads and either body mass or the mass of fat deposits have been detected (Day 1980; Connors and Smith 1982; Furness 1985a, 1985b; Ryan 1987a), but in most cases the lack of adequately controlled sampling (for factors such as age, reproductive status, and time of year) seriously hampers the interpretation of results (Ryan 1987a). The poor relationship between indicators of bird condition

and plastic load suggest that the effects of ingestion are either relatively minor or that they frequently are masked by other variables.

The main drawback to using correlations to demonstrate adverse effects from plastic ingestion is the inability to separate cause from effect. Ingested plastic may cause poor bird condition, or a bird in poor condition may be more prone to ingest plastic (Conners and Smith 1982), assuming at least some plastic is ingested as a result of "misdirected foraging" (i.e., plastic eaten directly, not incidentally with prey; Ryan 1987b). Similarly, stranded birds may have higher-than-average plastic loads because the ingested plastic has affected the birds' ability to survive adverse weather conditions, or because starving birds are less discriminating and eat more plastic immediately prior to stranding (Bourne and Imber 1982; Ryan 1987b). Day (1980) recorded larger plastic loads in nonbreeding than in the breeding parakeet auklet, *Cyclorhynchus psittacula*, but this could also be due to age-related foraging differences (Day et al. 1985). The only way to avoid these ambiguous results is to perform experiments designed to test the specific adverse effects postulated to result from debris ingestion.

SPECIFIC EFFECTS OF PLASTIC AND OTHER DEBRIS INGESTION

The specific effects of ingested debris on seabirds can be divided into three categories; physical damage and blocking of the digestive tract, impairment of foraging efficiency, and the release of toxic chemicals (Day et al. 1985; Ryan 1987a). The severity of these different categories of effects varies according to the types of debris ingested and their retention time within seabirds.

Physical Damage and Obstruction of the Digestive Tract

Physical damage and blocking of the digestive tract is the most obvious effect of ingested debris on seabirds, resulting in starvation in extreme cases of gastrointestinal obstruction (e.g., Parslow and Jefferies 1972; Dickerman and Goelet 1987; Fry et al. 1987). However, obstruction of the digestive tract currently is infrequent in seabirds, and probably affects only a small proportion of populations (Ryan and Jackson 1987). Gastrointestinal obstruction by plastic has been suggested to be an important cause of chick mortality among albatross chicks in the North Pacific (Pettit et al. 1981; Fry et al. 1987), but this is not supported by recent observations (Sileo, Sievert, and Samuel 1990) which found only occasional instances of obstruction.

Threads and fibers may result in obstruction more frequently than other debris types because they form dense, intertwined balls in seabird gizzards, blocking the entrance to the intestine (Parslow and Jefferies 1972; Day et al. 1985). However, intestinal obstruction was not found in any of the more than 200 white-chinned petrels, *Procellaria aequinoctialis*, sampled off southern Africa, despite fibers comprising almost half the mass of ingested plastics (Ryan 1987b; Ryan and Jackson 1987).

To test whether ingested debris interferes with digestion, the assimilation efficiency (digestive efficiency) of white-chinned petrels fed large loads (1.4 g, more than twice the maximum load recorded for the

species; Ryan 1987b) of virgin polyethylene pellets was compared with that of control birds (Ryan and Jackson 1987). No significant difference was detected, suggesting that at least virgin pellets have little effect on seabird digestive efficiency. However, similar experiments with other types of plastics are warranted.

Cuts and ulcerations of the stomach lining caused by ingesting sharp objects are more frequent than is intestinal obstruction (e.g., Day et al. 1985; Zonfrillo 1985; Fry et al. 1987; Ryan and Jackson 1987). These lesions are seldom likely to be lethal, because seabirds tolerate similar injuries from sharp prey items (Baltz and Morejohn 1976; Bourne and Imber 1982; Fry and Lowenstein 1982). However, lesions may have sublethal effects, reducing disease resistance and thus influencing survival (Fry et al. 1987).

Impaired Foraging Efficiency

Debris accumulated in the stomachs of seabirds has been postulated to impair foraging efficiency as a result of mechanical distension of the stomach. This has two effects: it induces a false feeling of satiation and reduces the food-storage volume of the stomach (Day et al. 1985; Ryan 1988a). Both these mechanisms would tend to reduce foraging efficiency and consequently the ability to accumulate energy reserves essential for reproduction, molting and the survival of adverse weather conditions (Ryan 1988a). However, there have been no direct tests of this effect of ingested debris on seabirds.

Ryan (1988a) showed that chickens fed 10 virgin plastic pellets ate smaller meals and grew more slowly than did control birds, although production (growth per unit food eaten) was not affected by plastic loads (which is to be expected if plastic pellets have little or no influence on digestive efficiency, see above). This provides empirical evidence that plastic loads comparable to those found in similarly-sized seabirds affect foraging efficiency in birds. However, experiments on free-ranging seabirds are required to assess the severity of this problem. One possible test would be to monitor the breeding success of birds whose chicks are fed additional plastic loads.

Given the very large frequency of occurrence of ingested plastic and other debris in some seabird populations, it is essential to estimate the critical load size (relative to bird size) beyond which stomach distension caused by accumulated debris has a deleterious effect. A few small particles are unlikely to have an adverse effect, because many seabirds store quantities of squid beaks and naturally occurring indigestible debris such as pumice in their ventriculi (e.g., Furness 1985a; Ryan 1988b). Fortunately, the distributions of total plastic loads in individual birds are strongly skewed, with most birds having very small plastic loads (Fig. 1), and this probably results in a fairly small proportion of seabird populations being adversely affected by stomach distension caused by ingested debris. Even the species with the greatest occurrence of ingested plastic off southern Africa, the blue petrel, *Halobaena caerulea* (92% of birds containing plastic; Ryan 1987b), has 85% of birds containing plastic loads <25% of the maximum load recorded (Fig. 1).




Figure 1.--The total plastic loads (by mass) of individual Antarctic prions, *Pachyptila desolata*, and blue petrels, *Halobaena caerulea*. See Ryan (1987b) for sampling procedures. Hatching depicts birds containing no plastic particles. Note the highly skewed distributions, with relatively few birds having large plastic loads.

Ingested Debris as a Source of Toxic Chemicals

It has been suggested that plastics and other debris ingested by seabirds are a source of toxic chemicals (e.g., Pettit et al. 1981; Bourne and Imber 1982; Day et al. 1985; van Franeker 1985). Plastics contain a variety of toxic additives including colorants, plasticizers, and heat and ultraviolet stabilizers (Gregory 1978; van Franeker 1985; Wirka 1988), and at sea the surface of plastic particles adsorb certain organochlorine compounds, notably polychlorinated biphenyls (PCB's) (Carpenter et al. 1972). The only direct evidence to indicate that seabirds receive toxic chemicals from ingested plastic is the positive correlation found between plastic and PCB loads in female great shearwaters, *Puffinus gravis*, immediately after egg-laying, independent of other organochlorine loads (Ryan et al. 1988). More circumstantial evidence is that both interspecific and intraspecific (geographical) variations in PCB concentrations in eggs of Hawaiian seabirds (Ohlendorf and Harrison 1986) correlate with the prevalence of ingested plastic (Sileo, Sievert, Samuel, and Fefer 1990), although this pattern may result from foraging differences (Ryan et al. 1988). The uptake of toxic compounds from ingested plastics needs to be confirmed by examining seabird tissues for traces of plastic-specific additives (Ryan 1988b). Although the toxicity of plastic additives to seabirds (both singly and synergistically) needs to be determined, it is likely that plastics contribute only a small proportion of the total toxic chemical load borne by seabirds (Bourne 1976; Ohlendorf et al. 1978; Fry et al. 1987; Ryan 1988b).

Other types of debris ingested by seabirds are potentially more serious sources of toxic chemicals. Although not ingested at sea, Laysan albatross, *Diomedea immutabilis*, chicks are killed by lead and perhaps mercury poisoning from ingestion of paint peeling off buildings on Midway (Fry et al. 1987; Sileo, Sievert, and Samuel 1990). This presumably is a localized problem, and can be readily alleviated. However, paint flakes have also been found in the stomach of a pintado petrel, *Daption capense*, collected at sea off southern Africa (Ryan 1990). Birds that frequently scavenge from vessels are likely to ingest some paint, particularly when ship scraping and repainting occurs at sea. Seabirds also ingest tar balls (Brown et al. 1981; van Franeker 1985; Ryan 1986), and petroleum products are known to have adverse toxicological effects on seabirds (e.g., Fry et al. 1986; Koth and Vauk-Hentzelt 1988). More information is required on the incidence of paint and tar ball ingestion by seabirds before an estimate of impacts on seabird populations can be made. Particular attention should be paid to the lifespan of paint and tar balls after ingestion; if they are rapidly broken down in seabirds' stomachs, the incidence of ingestion may be greater than these scattered records indicate.

VULNERABILITY TO THE EFFECTS OF INGESTED DEBRIS

The vulnerability of a given species or age-class of seabirds to the effects of debris ingestion is determined by the type of debris ingested: the sizes and shapes of pieces of debris presumably are important in determining the degree of physical damage to the digestive tract, and different compounds vary as regards toxicity. However, probably the major factor affecting vulnerability is the dynamics of debris ingestion and loss. The magnitude of debris loads in birds are a function of the balance between the rate of ingestion and the rate of loss of ingested debris.

Virtually all debris ingested by seabirds floats in seawater, and is eaten when mistaken for food items, or in association with prey (Day et al. 1985; Ryan 1987b; but see Fry et al. 1987). Inter- and intraspecific comparisons of plastic loads illustrate that the rate of ingestion is related to foraging technique (greatest incidence in surface feeders), foraging niche width (greatest in generalists), and the local density of debris at sea (Ryan 1987b). The rate of loss is related to the maximum size of particles passed through the digestive tract, the rate of erosion within the stomach, and the frequency of regurgitation of indigestible objects. Of these factors, the frequency of regurgitation appears to be the primary determinant of whether or not seabirds accumulate plastic particles and other debris in their stomachs (Ryan 1987b, 1988b).

There are three patterns of regurgitation among seabirds (Fig. 2). Some birds, including giant-petrels, cormorants, skuas, gulls, terns, and albatrosses, frequently regurgitate indigestible stomach contents, preventing any accumulation of ingested plastic or other debris (Ryan 1988b). These birds are unlikely to suffer many serious effects from the ingestion of persistent debris. The main problems are likely to be ulcerations and lesions caused by sharp objects (e.g., glass in gulls feeding at refuse dumps) or the release of toxic chemicals (either those rapidly absorbed from the surface of particles, or those associated with debris that is not easily regurgitated, such as tar balls that adhere to the stomach lining).

Other seabird taxa apparently seldom regurgitate indigestible stomach contents (Furness 1985a, 1985b; Ryan 1987b), and it is these accumulators of ingested debris that are likely to show adverse effects resulting from debris causing stomach distension and from obstruction of the digestive tract. For the majority of procellariiform seabirds (petrels, shearwaters, storm-petrels and diving-petrels), the main avenue for removing ingested debris occurs during the chick-feeding period, when plastic particles accumulated throughout the nonbreeding season are fed to the single chick along with the chick's meals that are stored in the parents' stomachs (Fry et al. 1987; Ryan 1988b). This pattern of annual regurgitation results in an annual cycle in the amount of plastic and other debris in breeding adult petrels (Ryan 1988b, fig. 2). Other taxa that seldom regurgitate indigestible stomach contents, such as auks and phalaropes, but do not feed their chicks food stored in the stomach (e.g., Bedard 1969), apparently lack this avenue for plastic loss. These taxa accumulate ingested plastic and other debris (Connors and Smith 1982; Day et al. 1985; Ryan 1988b), but the dynamics of debris loads are poorly understood, and particle loss may depend almost entirely on erosion and subsequent excretion.

Not all seabirds that accumulate ingested plastic and other debris in their stomachs are equally vulnerable to the effects of ingested debris. A consequence of the intergeneration transfer of ingested debris is that chicks fledge with a debris load approximately equal to 2 years' accumulation by adult birds. This initial loading is exacerbated subsequently by the greater ingestion rate of young, naive birds (Day et al. 1985; Ryan 1988b) and the lack of an effective loss mechanism during the protracted immature period (up to 10 years; Croxall 1984). This suggests that immature petrels have the largest debris loads stored in their stomachs, and are most likely to show adverse effects from debris ingestion.




Figure 2.--Diagrammatic representation of the effect of different regurgitation frequencies on the pattern of persistent debris accumulation in seabird stomachs.

FUTURE RESEARCH DIRECTIONS

The preceding review has indicated several areas where both the effects and the dynamics of debris ingestion are poorly understood. The following points summarize key problems that warrant special attention.

1. Verification that stomach distension resulting from accumulated debris causes reduced meal size and thus reduces the foraging efficiency of seabirds.
2. Assuming that 1) holds, an estimate is needed of the critical load size beyond which stomach distension has serious effects.
3. Examination of seabird tissues for plastic-specific additives is required to test whether toxic compounds are assimilated from ingested plastic particles.
4. Assuming that 3) holds, tests of the toxicity of plastics additives to seabirds (both singly and synergistically) are needed.
5. Experimental assessment of the lifespans of different types of debris in seabird stomachs is essential to interpret correctly the dynamics of ingestion.
6. Continued monitoring of debris loads in seabirds is warranted to detect major changes in ingestion patterns or effects, such as an increase in the incidence of obstruction of the digestive tract.

Ingestion by seabirds is a function of the density of debris at sea, and it is only by reducing this density that the incidence of ingestion can be reduced. However, a thorough understanding of both the dynamics and effects of plastic ingestion may enable implementation of measures specifically targeted to lessen the effects on seabird populations.

ACKNOWLEDGMENTS

I am grateful to the organizers of the Second International Conference on Marine Debris for the opportunity to attend the conference. Financial and logistical support for my work was received from the South African Council for Scientific and Industrial Research, South African Department of Environment Affairs, and the South African Scientific Committee for Antarctic Research.

REFERENCES

- Azzarello, M. Y., and E. S. van Vleet.
1987. Marine birds and plastic pollution. *Mar. Ecol. Progr. Ser.*
37:295-303.

- Baltz, D. M., and G. V. Morejohn.
1976. Evidence from seabirds of plastic particle pollution off central California. *West. Birds* 7:111-112.
- Bedard, J.
1969. Adaptive radiation in Alcidae. *Ibis* 111:189-198.
- Bourne, W. R. P.
1976. Seabirds and pollution. In R. Johnston (editor), *Marine pollution*, p. 403-502. Acad. Press, London.
- Bourne, W. R. P., and M. J. Imber.
1982. Plastic pellets collected by a prion on Gough Island, central South Atlantic Ocean. *Mar. Pollut. Bull.* 13:20-21.
- Brown, R. G. B., S. P. Barker, D. E. Gaskin, and M. R. Sandeman.
1981. The foods of great and sooty shearwaters, *Puffinus gravis* and *P. griseus*, in eastern Canadian waters. *Ibis* 123:19-30.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Miklas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. *Science* 178:749-750.
- Connors, P. G., and K. G. Smith.
1982. Oceanic plastic particle pollution: Suspected effect on fat deposition in red phalaropes. *Mar. Pollut. Bull.* 13:18-20.
- Croxall, J. P.
1984. Seabirds. In R. M. Laws (editor), *Antarctic ecology*, p. 533-618. Acad. Press, London, 2.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Dickerman, R. W., and R. G. Goelet.
1987. Northern gannet starvation after swallowing Styrofoam. *Mar. Pollut. Bull.* 18:293.
- Fry, D. M., S. I. Fefer, and L. Sileo.
1987. Ingestion of plastic debris by Laysan albatrosses and wedge-tailed shearwaters in the Hawaiian Islands. *Mar. Pollut. Bull.* 18:339-343.
- Fry, D. M., and L. J. Lowenstein.
1982. Insults to alcids: Injuries caused by food, by burrowing and by oil contamination. *Pac. Seabird Group Bull.* 9:78-79.
-

- Fry, D. M., J. Swenson, L. A. Addiego, C. R. Grau, and A. Kang.
1986. Reduced reproduction of wedge-tailed shearwaters exposed to single doses of weathered Santa Barbara crude oil. *Arch. Environ. Contam. Toxicol.* 15:453-463.
- Furness, R. W.
1985a. Ingestion of plastic particles by seabirds at Gough Island, South Atlantic Ocean. *Environ. Pollut. (Ser. A)* 38:261-272.
1985b. Plastic particle pollution: Accumulation by procellariiform seabirds at Scottish colonies. *Mar. Pollut. Bull.* 16:103-106.
- Gregory, M. R.
1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwater Res.* 12:399-414.
- Harper, P. C., and J. A. Fowler.
1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptila* spp.) 1958-1977. *Notornis* 34:65-70.
- Koth, T., and E. Vauk-Hentzelt.
1988. Influence of plumage and oiling on body and organ growth in young Kittiwakes. *Mar. Pollut. Bull.* 19:71-73.
- Ohlendorf, H. M., and C. S. Harrison.
1986. Mercury, selenium, cadmium and organochlorines in eggs of three Hawaiian seabird species. *Environ. Pollut. (Ser. B)* 11:169-191.
- Ohlendorf, H. M., R. W. Risebrough and K. Vermeer.
1978. Exposure of marine birds to environmental pollutants. U.S. Fish Wildl. Serv., Wildl. Res. Rep. 9.
- Parslow, J. L. F., and D. J. Jefferies.
1972. Elastic thread pollution of puffins. *Mar. Pollut. Bull.* 3:43-45.
- Pettit, T. N., G. S. Grant, and G. C. Whittow.
1981. Ingestion of plastics by Laysan albatross. *Auk* 98:839-841.
- Rothstein, S. I.
1973. Plastic particle pollution of the surface of the Atlantic Ocean: Evidence from a seabird. *Condor* 75:344-345.
- Ryan, P. G.
1986. Seabirds as water quality monitors. *Mar. Pollut. Bull.* 17:435-436.
1987a. The effects of ingested plastic on seabirds: Correlations between plastic load and body condition. *Environ. Pollut.* 46:119-125.
1987b. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.

- 1988a. Effects of ingested plastic on seabird feeding: Evidence from chickens. *Mar. Pollut. Bull.* 19:125-128.
- 1988b. Intraspecific variation in plastic ingestion by seabirds and the flux of plastic through seabird populations. *Condor* 90:446-452.
1990. The marine plastic debris problem off southern Africa: Types of debris, their environmental effects, and control measures. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Ryan, P. G., A. D. Connell, and B. D. Gardner.
1988. Plastic ingestion and PCBs in seabirds: Is there a relationship? *Mar. Pollut. Bull.* 19:174-176.
- Ryan, P. G., and S. Jackson.
1987. The lifespan of ingested plastic particles in seabirds and their effect on digestive efficiency. *Mar. Pollut. Bull.* 18:217-219.
- Sileo, L., P. R. Sievert, and M. D. Samuel.
1990. Causes of mortality of albatross chicks at Midway Atoll. *J. Wildl. Dis.* 26:329-338.
- Sileo, L., P. R. Sievert, M. D. Samuel, and S. I. Fefer.
1990. Prevalence and characteristics of plastic ingested by Hawaiian seabirds. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Van Franeker, J. A.
1985. Plastic ingestion in the North Atlantic fulmar. *Mar. Pollut. Bull.* 16:367-369.
- Van Franeker, J. A., and P. J. Bell.
1988. Plastic ingestion by petrels breeding in Antarctica. *Mar. Pollut. Bull.* 19:672-674.
- Wirka, J.
1988. *Wrapped in plastics: The environmental case for reducing plastics packaging*. Environmental Action Foundation, Wash., D.C., 159 p.
- Zonfrillo, B.
1985. Petrels eating contraceptives, polythene and plastic beads. *Br. Birds* 78:350-351.
-

INGESTION OF PLASTIC PARTICLES BY SOOTY AND SHORT-TAILED SHEARWATERS IN THE NORTH PACIFIC

Haruo Ogi

Research Institute of the North Pacific Fisheries
Faculty of Fisheries, Hokkaido University
Hakodate, Hokkaido, 041 Japan

ABSTRACT

Differences in rates of ingestion and types of plastic particles ingested by 218 sooty shearwaters, *Puffinus griseus*, and 324 short-tailed shearwaters, *P. tenuirostris*, obtained between 1970 and 1987 were examined. Of these seabirds, 193 sooty shearwaters (88.5%) and 265 short-tailed shearwaters (81.8%) were found to have ingested plastic particles. Significant differences in ingestion rates by year and area of collection were observed for short-tailed shearwaters. However, only one case of significant difference was observed for sooty shearwaters in the northern North Pacific.

After analyzing plastic particles ingested by these two species of seabirds on the basis of shape and color, plastic molding materials ingested by short-tailed shearwaters were found to account for 67.2% of all particles. On the other hand, sooty shearwaters mainly ingested particles of plastic products, with plastic molding materials accounting for only 38.4%. These differences were believed to reflect the differences in food habits of the two species.

INTRODUCTION

The behavior of actively ingesting objects with no nutritional value, called Pica Phenomenon, is commonly observed among birds, including seabirds (Day 1980). Since the second half of the 1960's, plastic production has increased sharply, and plastic has become a major pollutant of the marine environment. However, the impact of plastic ingestion by seabirds has not been made sufficiently clear.

This study analyzes characteristics of plastic particle ingestion by sooty, *Puffinus griseus*, and short-tailed shearwaters, *P. tenuirostris*, based on records of the number, shape, and color of plastic particles found in gastric contents of the two species of shearwaters prevalent in the subarctic North Pacific Ocean at the same time in summer.

MATERIALS AND METHODS

All of the shearwaters from whose stomachs plastic particles were extracted were individuals killed incidentally in the course of driftnet fishing. Dates and sites of collection and the number of specimens are shown in Figures 1 and 2 and Tables 1 and 2.

A total of 218 sooty shearwaters were examined. They were obtained at 62 driftnet fishing sites between April and October for 7 years during the period between 1975 and 1987 (Fig. 1, Table 1). A total of 324 short-tailed shearwaters were collected at 62 driftnet fishing sites between April and August for 8 years between 1975 and 1987.

Geographically, collection sites of sooty shearwaters were distributed in the North Pacific in an area bounded by lat. 31°-51°N and long. 143°E-143°W (Fig. 1). Those of short-tailed shearwaters were limited to the northwestern part of the Pacific, two sites in Bristol Bay in the Bering Sea, and all of the Aleutian Basin except for five sites around Cape Navarin (Fig. 2).

Shape and color of plastic particles taken from the stomachs of the two species of shearwaters were examined on an individual basis. Colors were classified into 11 types: white, yellow, brown, yellow-brown, blue, green, red, dark blue, dark green, dark red, and black/gray. Shapes and forms were classified into 13 types: cylinder, pill, dome, sphere, box, asymmetrical molding materials, string, cone, fragments of asymmetrical plastic products, vinyl, rubber, unidentifiable particles, and other. Of these, vinyl and rubber, whose shapes and forms are difficult to determine, are dealt with as independent categories because of their high ingestion rates by seabirds. On the basis of these data, frequencies of appearance of plastics were compared and examined by year, month, latitude, and longitude. The cases in which the number of bird individuals were 10 or fewer per item were excluded from statistical testing.

Classification and recognition of plastic particles in this study followed the method of Day (1980).

RESULTS

Interspecies Differences in Plastic Ingestion Rates

Tables 1 and 2 show the location and number of sooty and short-tailed shearwaters with and without plastic particles. The numbers of individuals ingesting plastic particles were 193 (88.5%) for sooty shearwaters (Table 1) and 265 (81.8%) for short-tailed shearwaters (Table 2). The difference in plastic particles ingestion rates between the two species was $\chi^2 = 4.023$ (df = 1, $0.025 < P < 0.05$), indicating slight interspecies differences, although they were not so obvious.




Figure 1.--Stations where sooty shearwaters were sampled.




Figure 2.---Stations where short-tailed shearwaters were sampled.

Table 1.--Frequency of occurrence of plastics in sooty shearwaters.

Date	Location		With plastic		Without plastic		Total number
	Latitude	Longitude	No.	Frequency (%)	No.	Frequency (%)	
19 Apr. 1975	40°35'N	167°31'E	1	100	0	0	1
13 June 1977	47°38'N	178°01'E	1	50.0	1	50.0	2
14 June 1977	47°37'N	179°44'E	0	0	2	100	2
14 July 1978	44°28'N	167°25'E	8	100	0	0	8
17 July 1978	45°30'N	172°25'E	3	75.0	1	25.0	4
26 July 1978	48°47'N	177°28'E	2	66.7	1	33.3	3
28 July 1978	44°58'N	179°58'E	1	100	0	0	1
2 Aug. 1978	48°31'N	177°34'W	3	75.0	1	25.0	4
2 Aug. 1978	46°00'N	175°33'E	1	100	0	0	1
3 Aug. 1978	47°30'N	175°30'E	2	100	0	0	2
5 Aug. 1978	50°25'N	177°32'W	1	100	0	0	1
8 Aug. 1978	49°29'N	164°12'E	1	100	0	0	1
14 June 1979	45°00'N	180°00'	1	100	0	0	1
19 June 1979	40°06'N	179°59'W	5	100	0	0	5
22 June 1979	39°00'N	175°00'W	1	100	0	0	1
23 June 1979	40°00'N	175°00'W	1	100	0	0	1
24 June 1979	41°00'N	174°59'W	9	100	0	0	9
25 June 1979	42°00'N	175°00'W	6	100	0	0	6
26 June 1979	43°00'N	175°00'W	7	100	0	0	7
27 June 1979	43°59'N	174°59'W	4	100	0	0	4
26 July 1979	49°33'N	177°32'E	0	0	1	100	1
29 July 1979	43°15'N	175°24'E	5	100	0	0	5
30 July 1979	44°32'N	175°29'E	14	77.8	4	22.2	18
31 July 1979	43°58'N	179°58'W	4	100	0	0	4
31 July 1979	46°00'N	175°30'E	3	75.0	1	25.0	4
1 Aug. 1979	43°01'N	180°00'	1	100	0	0	1
1 Aug. 1979	47°30'N	175°29'E	2	100	0	0	2
2 Aug. 1979	48°00'N	173°28'E	9	81.8	2	18.2	11
3 Aug. 1979	48°29'N	171°19'E	2	66.7	1	33.3	3
4 Aug. 1979	48°59'N	169°09'E	4	100	0	0	4
7 Aug. 1979	42°48'N	153°02'E	4	100	0	0	4
5 Sept. 1979	43°54'N	172°40'E	1	100	0	0	1
11 Sept. 1979	46°46'N	178°19'E	1	100	0	0	1
16 Sept. 1979	45°35'N	178°24'W	1	100	0	0	1
2 Oct. 1979	40°40'N	178°03'W	1	100	0	0	1
3 Oct. 1979	40°40'N	178°03'W	6	100	0	0	6
9 Oct. 1979	43°56'N	171°41'E	1	100	0	0	1
10 Oct. 1979	44°02'N	170°37'E	10	90.9	1	9.1	11
11 Oct. 1979	43°53'N	169°22'E	0	0	1	100	1
12 Oct. 1979	44°01'N	169°19'E	1	100	0	0	1
13 Oct. 1979	45°56'N	168°58'E	2	100	0	0	2
14 Oct. 1979	43°51'N	169°38'E	1	100	0	0	1
15 Oct. 1979	43°27'N	167°53'E	5	100	0	0	5
16 Oct. 1979	43°22'N	167°35'E	2	100	0	0	2

Table 1.--Continued.

Date	Location		With plastic		Without plastic		Total number
	Latitude	Longitude	No.	Frequency (%)	No.	Frequency (%)	
13 June 1980	40°02'N	179°58'E	2	66.7	1	33.3	3
16 June 1980	42°58'N	179°59'W	0	0	1	100	1
20 June 1980	46°04'N	179°34'W	1	50.0	1	50.0	2
29 July 1980	43°00'N	175°28'E	4	100	0	0	4
21 Apr. 1986	38°35'N	145°30'E	3	100	0	0	3
27 Apr. 1986	39°30'N	144°30'E	1	50.0	1	50.0	2
29 Apr. 1986	39°37'N	145°38'E	10	90.9	1	9.1	11
30 Apr. 1986	38°32'N	144°23'E	1	100	0	0	1
1 May 1986	38°31'N	143°32'E	1	100	0	0	1
2 May 1986	39°00'N	145°30'E	1	50.0	1	50.0	2
18 July 1986	45°00'N	150°02'W	1	100	0	0	1
5 May 1987	39°30'N	143°29'E	1	100	0	0	1
6 May 1987	39°30'N	144°30'E	2	100	0	0	2
7 May 1987	39°29'N	145°30'E	0	0	1	100	1
18 Aug. 1987	42°28'N	170°33'W	7	100	0	0	7
21 Aug. 1987	43°59'N	163°33'W	6	100	0	0	6
23 Aug. 1987	43°59'N	163°57'W	4	80.0	1	20.0	5
24 Aug. 1987	43°58'N	163°40'W	1	100	0	0	1
3 Sept. 1987	45°40'N	153°24'W	9	100	0	0	9
Total			193	88.5	25	11.5	218

Yearly Differences in Plastic Ingestion Rates

Plastic ingestion rates by year and by species are shown in Tables 3 and 4. Based on comparisons of data for the period (excluding 1975, 1977, and 1980), the plastic ingestion rate of sooty shearwaters was $\chi^2 = 1.034$ (df = 3, $0.5 < P < 0.75$), indicating no significant differences (Table 3). On the other hand, the ingestion rate for short-tailed shearwaters for the entire period excluding 1973 was $\chi^2 = 74.757$ (df = 6, $P < 0.005$), showing significant differences (Table 4). The plastic particle ingestion rate of short-tailed shearwaters for 1970-73 was conspicuously lower than rates for other years, although the number of specimens was small for all the year. Comparing 1970-72 with 1975-79, an extremely significant difference of $\chi^2 = 67.822$ (df = 1, $P < 0.005$) was observed. It was suggested that the increase in plastic particle ingestion by short-tailed shearwaters reflected the increased production of synthetic resins.

Monthly Differences in Plastic Ingestion Rates

Plastic ingestion rates of two species of shearwaters by month were shown in Tables 5 and 6. No significant difference was found in plastic

Table 2.--Frequency of occurrence of plastics in short-tailed shearwaters.

Date	Location		With plastic		Without plastic		Total number
	Latitude	Longitude	No.	Frequency (%)	No.	Frequency (%)	
5 July 1970	55°21'N	164°00'W	13	46.4	15	53.6	28
18 July 1972	56°23'N	161°03'W	27	52.9	24	47.1	51
29 Apr. 1973	49°30'N	170°30'E	2	50.0	2	50.0	4
30 Apr. 1973	50°30'N	170°30'E	2	100	0	0	2
21 Apr. 1975	42°30'N	167°30'E	1	100	0	0	1
22 Apr. 1975	43°33'N	167°30'E	4	100	0	0	4
26 Apr. 1975	46°30'N	167°30'E	2	50.0	2	50.0	4
28 Apr. 1975	47°30'N	167°30'E	1	50.0	1	50.0	2
4 May 1975	51°30'N	167°30'E	1	100	0	0	1
9 May 1975	50°30'N	171°00'E	1	100	0	0	1
11 May 1975	49°30'N	171°00'E	2	66.7	1	33.3	3
16 May 1975	47°30'N	169°01'E	4	100	0	0	4
17 May 1975	48°30'N	169°00'E	3	100	0	0	3
27 May 1975	50°25'N	172°30'E	2	100	0	0	2
10 June 1976	55°00'N	171°00'E	11	100	0	0	11
11 June 1976	55°00'N	172°00'E	12	100	0	0	12
13 June 1976	55°00'N	173°00'E	12	92.3	1	7.7	13
15 June 1976	55°00'N	176°00'E	2	100	0	0	2
16 June 1976	55°10'N	177°17'E	2	66.7	1	33.3	3
17 June 1976	55°00'N	178°00'E	1	100	0	0	1
18 June 1976	55°03'N	179°20'E	1	100	0	0	1
19 June 1976	55°00'N	179°00'W	2	100	0	0	2
20 June 1976	55°00'N	178°00'W	1	100	0	0	1
21 June 1976	55°03'N	176°50'W	1	100	0	0	1
24 June 1976	56°55'N	176°00'W	1	100	0	0	1
26 June 1976	57°00'N	177°00'W	5	100	0	0	5
27 June 1976	57°00'N	178°00'W	2	100	0	0	2
29 June 1976	58°56'N	178°08'E	2	100	0	0	2
30 June 1976	59°00'N	178°00'E	1	100	0	0	1
6 July 1976	59°00'N	171°00'E	2	100	0	0	2
9 July 1976	61°44'N	177°40'E	4	100	0	0	4
11 July 1976	61°54'N	179°59'W	2	100	0	0	2
12 July 1976	61°48'N	179°39'W	1	50.0	1	50.0	2
13 July 1976	62°02'N	179°40'E	2	66.7	1	33.3	3
17 July 1976	62°05'N	179°36'E	2	100	0	0	2
2 June 1977	49°30'N	165°01'E	1	100	0	0	1
3 June 1977	49°22'N	165°43'E	4	100	0	0	4
4 June 1977	50°03'N	166°59'E	4	100	0	0	4
5 June 1977	49°26'N	168°48'E	5	100	0	0	5
7 June 1977	50°16'N	169°00'E	4	100	0	0	4
8 June 1977	50°35'N	168°31'E	3	100	0	0	3
9 June 1977	50°54'N	168°10'E	2	100	0	0	2
11 June 1977	50°08'N	175°21'E	16	94.1	1	5.9	17
13 June 1977	47°38'N	178°01'E	2	100	0	0	2

Table 2.--Continued.

Date	Location		With plastic		Without plastic		Total number
	Latitude	Longitude	No.	Frequency (%)	No.	Frequency (%)	
14 June 1977	47°37'N	179°44'E	3	100	0	0	3
23 June 1977	56°04'N	176°14'W	2	100	0	0	2
25 June 1977	56°00'N	175°14'W	5	100	0	0	5
30 June 1977	57°02'N	178°02'W	1	100	0	0	1
2 July 1977	57°58'N	176°18'W	5	100	0	0	5
4 July 1977	58°10'N	175°37'W	4	100	0	0	4
5 July 1977	59°03'N	175°32'W	1	100	0	0	1
8 July 1977	58°45'N	178°54'W	1	100	0	0	1
9 July 1977	58°30'N	179°58'E	1	100	0	0	1
11 July 1977	59°13'N	177°51'W	4	80.0	1	20.0	5
12 July 1977	58°52'N	177°28'W	2	100	0	0	2
14 June 1978	52°55'N	176°59'W	14	93.3	1	6.7	15
19 June 1978	54°00'N	178°02'W	10	83.3	2	16.7	12
26 July 1978	48°47'N	177°28'E	1	50.0	1	50.0	2
28 July 1978	44°58'N	179°58'E	1	100	0	0	1
14 June 1979	51°02'N	177°31'W	33	91.7	3	8.3	36
4 Aug. 1979	46°24'N	158°18'E	1	50.0	1	50.0	2
4 Aug. 1979	48°59'N	169°19'E	1	100	0	0	1
Total			265	81.8	59	18.2	324

particle ingestion rates for either species. The rate for sooty shearwaters was $\chi^2 = 3.517$ (df = 5, $0.5 < P < 0.75$), and the rate for short-tailed shearwaters was $\chi^2 = 4.060$ (df = 3, $0.25 < P < 0.5$).

More than 85% of the sooty shearwaters were found to have ingested plastic particles in all the months between April and October except May, when there were few specimens collected (Table 5). Slightly lower values were obtained for short-tailed shearwaters in April, although no significant differences were shown in ingestion rates (Table 6). Day (1980) made clear that there are seasonal differences in plastic particle ingestion by short-tailed shearwaters observed near Kodiak Island, and showed that the birds there actively ingest in June and August. In this study, no seasonal trend in plastic particle ingestion was identified, as collection sites of short-tailed shearwaters were scattered in outer waters and collection was not made at regular monthly intervals in areas where this species stays for a long time.

Latitudinal Differences in Plastic Particle Ingestion

Tables 7 and 8 show the plastic particle ingestion rates by 5° latitudinal belts. A significant difference of $\chi^2 = 7.248$ (df = 2,

Table 3.--Frequency of occurrence of plastics in sooty shearwaters by year.

Year	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
1975	1	100	0	0	1
1977	1	25.0	3	75.0	4
1978	22	88.0	3	12.0	25
1979	114	91.2	11	8.8	125
1980	7	70.0	3	30.0	10
1986	18	85.7	3	14.3	21
1987	30	93.8	2	6.3	32
Total	193	88.5	25	11.5	218

Table 4.--Frequency of occurrence of plastics in short-tailed shearwaters by year.

Year	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
1970	13	46.4	15	53.6	28
1972	27	52.9	24	47.1	51
1973	4	66.7	2	33.3	6
1975	21	84.0	4	16.0	25
1976	69	94.5	4	5.5	73
1977	70	97.2	2	2.8	72
1978	26	86.7	4	13.3	30
1979	35	89.7	4	10.3	39
Total	265	81.8	59	18.2	324

0.025 < P < 0.05) was found for sooty shearwaters as a result of comparisons of the first three belts in Table 7. Further, examination by belt presented a rate of $\chi^2 = 5.750$ (df = 1, 0.01 < P < 0.025) between lat. 40°-45°N and 45°-50°N, indicating a difference. The plastic ingestion rate for the belt of lat. 40°-45°N was higher at 93.1%, while that for lat. 45°-50°N was slightly lower at 80.3%.

Table 5.--Frequency of occurrence of plastics in sooty shearwaters by month.

Month	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
April	16	88.9	2	11.1	18
May	5	71.4	2	28.6	7
June	38	86.4	6	13.6	44
July	45	84.9	8	15.1	53
August	48	90.6	5	9.4	53
September	12	100	0	0	12
October	29	93.5	2	6.5	31
Total	193	88.5	25	11.5	218

Table 6.--Frequency of occurrence of plastics in short-tailed shearwaters by year.

Month	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
April	12	70.6	5	29.4	17
May	13	92.9	1	7.1	14
June	205	81.0	48	19.0	253
July	33	89.2	4	10.8	37
August	2	66.7	1	33.3	3
Total	265	81.8	59	18.2	324

Table 7.--Frequency of occurrence of plastics in sooty shearwaters by 5° latitudinal belts.

Latitudinal range	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
35°-40°N	21	84.0	4	16.0	25
40°-45°N	122	93.1	9	6.9	131
45°-50°N	49	80.3	12	19.7	61
50°-55°N	1	100	0	0	1
Total	193	88.5	25	11.5	218

Table 8.--Frequency of occurrence of plastics in short-tailed shearwaters by 5° latitudinal belts.

Latitudinal range	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
40°-45°N	6	100	0	0	6
45°-50°N	31	81.6	7	18.4	38
50°-55°N	93	92.1	8	7.9	101
55°-60°N	124	74.7	42	25.3	166
60°-65°N	11	84.6	2	15.4	13
Total	265	81.8	59	18.2	324

As regards short-tailed shearwaters, a significant difference of $\chi^2 = 12.645$ (df = 3, $0.005 < P < 0.01$) was found after comparing the last four belts in Table 8. Further, in the comparison of all areas, there were significant differences, with the rates standing at $\chi^2 = 0.030$ (df = 1, $0.01 < P < 0.05$) between lat. 45°-50°N and 60°-65°N and $\chi^2 = 11.347$ (df = 1, $P < 0.005$) between lat. 50°-55°N and 50°-60°N. This suggests the presence of differences in plastic ingestion rates for short-tailed shearwaters between the North Pacific and the north Bering Sea. Comparing lat. 45°-50°N and 55°-60°N, a rate of $\chi^2 = 8.954$ (df = 1, $P < 0.005$) was obtained. Thus visible differences were found in the plastic ingestion rates between north and south.

These were believed to reflect both differences in distribution, migration, and summer residence of both species of shearwaters in the subarctic North Pacific region, and differences in abundance of plastic particles by area.

Longitudinal Differences in Plastic Ingestion Rates

Tables 9 and 10 show the plastic particle ingestion rates for two species of shearwaters by 5° longitudinal strips.

With regard to sooty shearwaters, no significant difference in ingestion rates was found between strips ($\chi^2 = 13.559$, df = 7, $0.05 < P < 0.1$). On the other hand, a significant difference was observed for short-tailed shearwaters, with $\chi^2 = 69.748$ (df = 4, $P < 0.005$). This significant difference occurred because the plastic ingestion rate for 40 individuals obtained in Bristol Bay in Alaska was only 50.6%, and the collection sites for these seabirds were limited to long. 165°-160°W. It was not evident why the plastic particle ingestion rate for short-tailed shearwaters in this strip was conspicuously low. These 40 short-tailed shearwaters were full of *Thysanoessa raschii*, a species of euphausiids, suggesting that the abundance of plastic particles in the area was low and plastic particles in the stomach and gizzards had moved to the intestine.

Table 9.--Frequency of occurrence of plastics in sooty shearwaters by 5° longitudinal belts.

Longitudinal range	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
140°-145°E	6	85.7	1	14.3	7
145°-150°E	14	82.4	3	17.6	17
150°-155°E	4	100	0	0	4
155°-160°E	--	--	--	--	--
160°-165°E	1	100	0	0	1
165°-170°E	24	96.0	1	4.0	25
170°-175°E	26	83.9	5	16.1	31
175°E-180°	40	78.4	11	21.6	51
180°-175°W	22	88.0	3	12.0	25
175°-170°W	35	100	0	0	35
170°-165°W	--	--	--	--	--
165°-160°W	11	91.7	1	8.3	12
160°-155°W	--	--	--	--	--
155°-150°W	10	100	0	0	10
Total	193	88.5	25	11.5	218

Table 10.--Frequency of occurrence of plastics in short-tailed shearwaters by 5° longitudinal belts.

Longitudinal range	With plastic		Without plastic		Total number
	No.	Frequency (%)	No.	Frequency (%)	
155°-160°E	1	50.0	1	50.0	2
165°-170°E	40	93.0	3	7.0	43
170°-175°E	46	92.0	4	8.0	50
175°E-180°	41	91.1	4	8.9	45
180°-175°W	97	92.4	8	7.6	105
175°-170°W	--	--	--	--	--
170°-165°W	--	--	--	--	--
165°-160°W	40	50.6	39	49.4	79
Total	265	81.8	59	18.2	324

Number of Shapes and Colors of Plastic Particles Ingested by Two Species of Shearwaters

Tables 11 and 12 show the number of plastic particles classified by shape and color found in the stomachs of 192 sooty shearwaters and 265 short-tailed shearwaters. For both species, 13 shapes and 11 colors were used.

Such shapes as cylinder, pill, dome, sphere, box, and asymmetrical plastic pellets were apparently molding materials for making plastic products. All the other particles were judged to be fragments of plastic products. Ingestion rates of these molding materials were 38.4% for sooty shearwaters (Table 11) and 67.2% for short-tailed shearwaters (Table 12). A significant difference was found in the interspecies comparison of ingestion rates ($\chi^2 = 298.7$, $df = 1$, $P < 0.005$), showing that sooty shearwaters have a strong tendency to ingest plastic particles with no preference as to shape. On the other hand, short-tailed shearwaters have a strong tendency to ingest molding material particles having consistent shapes. Molding materials were 39.4% of the total weight for sooty shearwaters, whereas for short-tailed shearwaters they were 74.6%. It is interesting to note that a highly significant difference of $\chi^2 = 624.830$ ($df = 12$, $P < 0.005$) was found after comparing the rates of total number of particles by shape for the two species. However, in terms of order of ingestion by shape, $r_s = 0.9272$ ($n = 13$, $P < 0.01$), indicating similarity between the two species.

As for the color of plastic particles ingested by the two species of seabirds, a highly significant difference ($\chi^2 = 515.588$, $df = 10$, $P < 0.005$) was found when comparing the numbers of particles by color, but the order of ingestion was similar for the two species ($r_s = 0.8727$, $n = 11$, $P < 0.01$).

The average number and weight of plastic particles ingested per individual were 8.45 particles weighing 134 mg for sooty shearwaters and 8.79 particles weighing 140 mg for short-tailed shearwaters.

DISCUSSION

Day (1980) examined plastics ingested by seabirds in Alaska and discussed the occurrence and characteristics of plastic pollution in them. Much remains unknown about the impact of plastic particle ingestion on seabirds, but Carpenter et al. (1972) found that polychlorinated biphenyls (PCB's) are concentrated on the surface of spherical polystyrene molding materials. Further, as regards great shearwater, *P. gravis*, Ryan et al. (1988) found that a high-level positive interrelation of $r_s = 0.700$ existed between the amount of ingested plastic particles and PCB in adult birds. Day (1980) suggested that hydrocarbon pollutants arising from plastic particle ingestion are not only affecting breeding capability but also causing abnormal behavior.

For short-tailed shearwaters, a species subject to global-scale migration, it is believed that the variation in amounts of body fat in

Table 11.--Number of shapes and colors of plastic ingested by sooty shearwaters.

Shape	Color										Total Percent		
	White	Yellow	Brown	Yellow brown	Blue	Green	Red	Dark blue	Dark green	Dark red		Black/gray	
Cylinder	41	6	0	70	0	2	1	0	1	0	117	238	14.6
Pill	14	6	29	26	0	0	1	0	1	1	28	105	6.4
Dome	2	0	2	4	0	0	0	0	0	0	1	9	0.6
Sphere	0	0	64	2	0	0	1	0	0	0	8	75	4.6
Box	1	0	2	6	0	0	0	0	0	0	3	12	0.7
Asymmetrical molding material	53	10	21	40	4	40	1	1	3	0	14	187	11.5
String	52	0	5	0	2	4	0	0	1	0	50	114	7.0
Cone	1	0	0	0	0	0	0	0	0	0	1	2	0.1
Asymmetrical plastic products	168	27	28	67	12	42	8	1	6	1	41	401	24.6
Vinyl	90	0	16	0	9	3	0	0	0	0	180	298	18.3
Rubber	0	0	1	0	0	0	0	0	0	0	3	4	0.3
Unidentifiable particles	27	3	3	17	2	10	0	1	2	1	21	87	5.3
Other	17	7	12	36	0	15	1	2	1	0	8	99	6.1
Total	466	59	183	268	29	116	12	5	15	3	475	1,631	
Percent	28.6	3.6	11.2	16.4	1.8	7.1	0.7	0.3	0.9	0.2	29.1		

Table 12.--Number of shapes and colors of plastic ingested by short-tailed shearwaters.

Shape	Color											Total Percent	
	White	Yellow	Brown	Yellow brown	Blue	Green	Red	Dark blue	Dark green	Dark red	Black/gray		
Cylinder	232	21	225	407	3	7	9	0	0	4	68	976	41.9
Pill	47	1	56	111	0	0	1	0	0	3	14	233	10.0
Dome	3	6	7	18	0	0	0	0	0	0	0	34	1.5
Sphere	4	0	2	11	0	10	0	0	0	0	2	29	1.2
Box	6	0	10	10	0	1	0	0	0	0	1	28	1.2
Asymmetrical molding material	58	13	49	92	12	26	2	0	1	0	12	265	11.4
String	20	0	3	0	6	15	0	0	0	0	4	48	2.1
Cone	1	0	1	1	0	1	0	0	0	0	0	4	0.2
Asymmetrical plastic products	97	16	18	126	10	42	49	3	8	1	10	380	16.3
Vinyl	102	0	5	0	19	5	1	0	0	0	9	141	6.1
Rubber	29	0	1	7	0	0	0	0	0	0	6	43	1.9
Unidentifiable particles	20	0	12	28	4	10	5	2	3	0	12	96	4.1
Other	19	1	4	14	2	9	2	1	0	0	1	53	2.3
Total	638	58	393	825	56	126	69	6	12	8	139	2,330	
Percent	27.4	2.5	16.9	35.4	2.4	5.4	3.0	0.3	0.5	0.3	6.0		

connection with long-distance migration between Northern and Southern Hemispheres causes changes in PCB density in internal organs and tissues (Tanaka et al. 1986). It will be necessary to examine the impact of plastic particle ingestion on seabirds throughout their entire life history.

In this study, it is suggested that differences in plastic particle ingestion by sooty and short-tailed shearwaters are related not only to food habits of the two species but also to their distribution and migration in the subarctic North Pacific region. Sooty shearwaters arrive in the area just south of the Subarctic Boundary at lat. 38°-40°N in early April. These adult birds have just completed breeding in the Southern Hemisphere. Sooty shearwaters migrate northward, keeping pace with northward migration of Pacific saury, *Cololabis saira*, and the Japanese sardine, *Sardinops melanostictus*. In June and July, subadults and hatching year birds arrive also from the Southern Hemisphere to the area just south of the Subarctic Boundary, and then migrate even farther north in the same manner as adult birds. Therefore, as Day and Shaw (1987) showed, life of the sooty shearwaters in the Northern Hemisphere starts in areas where plastic pollution is intense. Further, they range in summer from lat. 41°-42°N to the Aleutian Islands, extending to around lat. 55°N in the Okhotsk Sea. Sooty shearwaters tend to stay in this area longer than short-tailed shearwaters. Especially to be noted is that sooty shearwaters migrate only in a nomadic mode after arriving in the North Pacific Subarctic Zone, and their movement is determined by food distribution. This causes emergence of flocks composed of individuals in different stages of growth. Therefore it may be difficult to determine changes in plastic particle ingestion rates in terms of year, month, latitude, and longitude.

Short-tailed shearwater parent birds, which complete breeding in Tasmania in the Southern Hemisphere, fly toward the Northern Hemisphere in mid- and late March and arrive in the Bering Sea in late April (Shuntov 1961). These short-tailed shearwaters continue northward migration without resting around the Subarctic Boundary. After staying in the summer resident area, they migrate directly southward to the Southern Hemisphere for breeding. They therefore live in a condition immune from the plastic-polluted areas in the North Pacific. Lower plastic particle ingestion rates for 40 short-tailed shearwater individuals collected at Bristol Bay in the Bering Sea were probably due to the fact that they were breeding adult birds. Subadult and hatching year birds arrive in areas around lat. 40°N in mid-April and early June, continuing northward to around lat. 53°N in June-July. The northern limit for adult birds is north of lat. 70°N, while subadult and hatching year birds live in a nomadic mode between lat. 45°-60°N. This means that younger birds are exposed to plastic pollution for a longer period of time, with the likelihood of increased plastic particle ingestion rates over both time and space. The results of this study coincide with this finding.

This study showed that short-tailed shearwaters had a tendency to ingest more plastic molding materials, while sooty shearwaters ingested more plastic product fragments. This reflects the food habits of the two species. Short-tailed shearwaters feed mainly on small, low-class

organisms originating from the biological production process in the surface layers of the ocean, these include juvenile fish, euphausiids, copepods, larval and juvenile squid, and amphipods (Ogi et al. 1980). They are particularly fond of euphausiids, which appear in the surface not as single individuals but usually as patches or swarms. The body size of euphausiids is relatively uniform. Short-tailed shearwaters must catch them quickly and continuously. They probably actively ingest plastic molding materials because these resemble organisms forming patches or swarms.

Sooty shearwaters, on the other hand, feed on Japanese sardines (Ogi unpubl. data) and Pacific sauries (Ogi 1984) in large quantities. They are also known to consume large quantities of larval and juvenile squid and euphausiids (Brown et al. 1981), but regardless, it is obvious that the food niche of sooty shearwaters is higher than that of short-tailed shearwaters. It is assumed, therefore, that they select less specifically plastic molding materials resembling small-sized organisms.

ACKNOWLEDGMENTS

I thank H. Yoshida for help with measurement and identification of plastic particles.

This study was supported by the Fisheries Agency, the Government of Japan the Nippon Life Insurance Foundation (C87110-206-13), the Toyota Foundation (89-III-042), and a Grant-in-Aid for Scientific Research on Priority Areas (From Asia to America: Prehistoric Mongoloid Dispersals) from the Japan Ministry of Education, Science and Culture.

This is Contribution No. 227 from the Research Institute of North Pacific Fisheries, Faculty of Fisheries, Hokkaido University.

REFERENCES

- Brown, R. G. B., S. P. Barker, D. E. Gaskin, and M. R. Sandeman.
1981. The foods of great and sooty shearwaters (*Puffinus gravis* and *P. griseus*) in eastern Canadian waters. *Ibis* 123:19-30.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Miklas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. *Science* 178:749-750.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., and D. G. Shaw.
1987. Patterns in the abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-1985. *Mar. Pollut. Bull.* 18:311-316.

- Ogi, H.
1984. Feeding ecology of the sooty shearwater in the western subarctic North Pacific Ocean. In D. N. Nettleship, G. A. Sanger, and P. F. Springer (editors), *Marine birds: Their feeding ecology and commercial fisheries relationships*, p. 78-84. Canadian Wildl. Serv., Ottawa.
- Ogi, H., T. Kubodera, and K. Nakamura.
1980. The pelagic feeding ecology of the short-tailed shearwater *Puffinus tenuirostris* in the subarctic Pacific region. *J. Yamashina Inst. Ornith.* 12:157-182.
- Ryan, P. G., A. D. Connell, and B. D. Gardner.
1988. Plastic ingestion and PCBs in seabirds: Is there a relationship? *Mar. Pollut. Bull.* 19:174-176.
- Shuntov, V. P.
1961. Migration and distribution of marine birds in southeastern Bering Sea during spring-summer season. *Zool. Zh.* 40:1058-1069. [In Russ. with Engl. abstr.]
- Tanaka, H., H. Ogi, S. Tanabe, R. Tatsukawa, and N. Oka.
1986. Bioaccumulation and metabolism of PCBs and DDE in short-tailed shearwater *Puffinus tenuirostris* during its transequatorial migration and in the wintering and breeding grounds. *Mem. Natl. Inst. Polar Res., Spec. Issue* 40:434-442.

THE INCIDENCE OF PLASTIC IN THE DIETS OF PELAGIC SEABIRDS
IN THE EASTERN EQUATORIAL PACIFIC REGION

David G. Ainley, Larry B. Spear
Point Reyes Bird Observatory
Stinson Beach, California 94970, U.S.A.

and

Christine A. Ribic*
Center for Quantitative Science in Forestry, Fisheries and Wildlife
University of Washington
Seattle, Washington 98195, U.S.A.

*Present address: U.S. EPA Environmental Research Laboratory, 200 SW 35th
Street, Corvallis, OR 97333, U.S.A.

ABSTRACT

Between 1984 and 1988, 921 seabirds of 39 species were collected during cruises in the eastern equatorial Pacific. Cruises were centered in the areas of the South Equatorial Current, Equatorial Countercurrent, and the northern Peru Current. The majority of species, mostly gadfly petrels and storm-petrels, had not previously been checked for plastic ingestion. Ingestion was a function of feeding behavior, area of the ocean frequented, and the amount of time passed since birds frequented polluted areas. Species that resided year-round in the equatorial region had eaten little plastic, but those species or populations that had recently come from the area of the southern Peru Current (off Chile), the North Pacific (off Japan or California), or the Tasman Sea/northern New Zealand area, had high plastic loads in their digestive tracts. Results suggest that the residency time of plastic in the digestive tract of petrels is less than 1 year.

INTRODUCTION

On the basis of research and literature reviews by Day (1980; Day et al. 1985), who worked mostly in the northern Pacific, and by Ryan (1987a, 1987b, 1988a, 1988b; Ryan and Jackson 1987) in the southern Atlantic, we now recognize that the ingestion of plastic is a pervasive phenomenon in seabirds. At present, 69 seabird species are known to ingest plastic while feeding at sea, and the incidence of ingestion has risen steadily since the

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

1960's. The species most likely to eat plastic are members of the order Procellariiformes. The factors that make these species so susceptible are 1) their feeding behavior (many feed at the surface, are omnivorous, and scavenge frequently); 2) their morphology (petrels have muscular gizzards in which indigestible items become trapped, whereas most other seabirds do not); and 3) their inability (albatrosses are an exception) to eject indigestible matter through the formation of pellets. It is believed that seabirds eat the plastic, usually in the form of industrial pellets and fragments of larger items, either because they mistake it for food or because edible organisms are attached to the plastic.

Geographic variation in the incidence of plastic indicates that seabirds feeding near to industrialized areas are the most likely to eat it. The plastic remains in digestive tracts until it degrades from abrasion and digestive processes or until adult birds feed it to their chicks. Day et al. (1985) estimated that the degradation process for an individual plastic pellet requires on the order of 6 months, but Ryan and Jackson (1987) proposed a time scale of 1 to 2 years. Ryan (1988b) further proposed that it is through feeding plastic to chicks that adults rid themselves of most of the plastic that has accumulated in their gut.

As yet, little work has been done on the incidence of ingestion in seabirds that forage far at sea, i.e., away from pollution sources. (See Ainley et al. 1990, who looked at plastic in the diets of Antarctic seabirds.) In this paper we present information on the incidence of plastic in the diets of seabirds that frequent Pacific equatorial waters. The sampling efforts closest spatially to ours were those by Harrison et al. (1983) and Sileo et al. (1990), who noted plastic in the diets of several species breeding in the Northwestern Hawaiian Islands, in the North Pacific a few thousand kilometers away from our study area. A number of species pass through our study area from various directions, and the incidence of plastic in them offers some clues as to where they are encountering it, and, on the basis of the time passed since visiting polluted areas, whether or not the degradation process and that of off-loading through chick feeding are important processes by which birds relieve themselves of plastic. In addition, several of the species we sampled had not been inspected before for evidence of plastic ingestion.

METHODS

In conjunction with the Equatorial Pacific Ocean Climate Study (EPOCS) of the National Oceanic and Atmospheric Administration (NOAA), we have been characterizing the community structure of open-ocean seabirds using the eastern equatorial Pacific as our study area. Included in our studies are the analyses of prey items. On six cruises (boreal autumn 1984, spring 1986, spring and autumn 1987, and spring and autumn 1988) we collected 921 seabirds of 39 species using a shotgun (Table 1). During each collection, we attempted to obtain up to five individuals of each species present. We noted which individuals had ingested plastic and, except for autumn 1984, counted the number of plastic pieces. Based on necropsy, we determined whether individuals were adults or subadults. The study area is between lat. 15°N and 15°S and long. 170° and 85°W (Fig. 1). Collections were made




Figure 1.--The study area in the eastern equatorial Pacific (shading), and sites where seabirds were collected (open symbols, on one cruise only; closed symbols, on four or more cruises); arrows indicate ocean currents.

Table 1.--The incidence of plastic in the diets of Pacific equatorial seabirds, and a comparison to other studies.

Species ^b	Cruise ^a						Total ^a		Other studies	
	A84 ^c	S86	S87	A87	S88	A88	No.	%	%	Source ^d
Petrels										
<i>Bulweria bulwerii</i>			0 (7)		0 (10)	0 (2)	19	0	0	3
<i>Daption capense</i> *			88 (8)				8	88	14-57	1,5
<i>Pachyptilla belcheri</i> *			83 (6)				6	83	0-57	5
<i>Procellaria aequinoctialis</i> *			67 (3)				3	67	0-57	1,5
<i>Pterodroma alba</i>			0 (2)			0 (9)	11	0		
<i>Pterodroma arminjoniana</i>			0 (8)			0 (3)	11	0		
<i>Pterodroma cookii defilippiana</i> *		0 (3)	67 (6)				9	44	10	4
<i>Pterodroma e. externa</i> *	0 (3)	0 (5)	0 (27)	0 (21)	0 (13)	0 (35)	104	1		
<i>Pterodroma e. cervicalis</i>		100 (1)	0 (2)		0 (7)		10	10		
<i>Pterodroma inexpectata</i>						0 (1)	1	0	0	1
<i>Pterodroma leucoptera</i>	0 (6)	75 (4)	20 (15)	7 (28)	26 (19)	5 (20)	92	15		
<i>Pterodroma l. longirostris</i> *		100 (1)	50 (6)	33 (6)	50 (10)	100 (2)	25	52		
<i>Pterodroma l. pycrofti</i>					50 (2)		2	50		
<i>Pterodroma neglecta</i>				0 (2)	0 (5)	0 (1)	8	0		
<i>Pterodroma nigripennis</i>		0 (1)	0 (4)	0 (3)	0 (14)	7 (15)	37	3		
<i>Pterodroma rostrata</i>	0 (2)		0 (21)	0 (13)	7 (15)	0 (16)	67	2		
<i>Pterodroma ultima</i>			0 (1)			0 (4)	5	0		
<i>Puffinus bulleri</i>					100 (1)		1	100		
<i>Puffinus griseus</i>			50 (2)		0 (6)	100 (17)	25	72	10-67	2,5
<i>Puffinus nativitatus</i>			50 (2)			0 (1)	3	33	0	3
<i>Puffinus pacificus</i>			0 (15)		0 (7)	8 (13)	35	3	0	2
<i>Puffinus tenuirostris</i>						100 (1)	1	100	47-100	2

Table 1.--Continued.

Species ^b	Cruise ^a					Total ^a		Other studies		
	A84 ^c	S86	S87	A87	S88	A88	No.	%	%	Source ^d
			Storm-Petrels							
<i>Fregata grallaria</i>		0 (2)	0 (3)		0 (1)	0 (2)	8	0		
<i>Nesofregata fuliginosa</i>	0 (2)	0 (1)	0 (1)	0 (3)	0 (1)		8	0		
<i>Oceanites gracilis</i>			0 (2)				2	0		
<i>Oceanites oceanicus*</i>			0 (8)				8	0	19-75	1,5
<i>Oceanodroma castro</i>			0 (2)		0 (3)	0 (2)	7	0		
<i>Oceanodroma hornbyi*</i>		0 (1)					1	0		
<i>Oceanodroma leucorhoa</i>	6 (18)	0 (1)	9 (22)	13 (45)	0 (16)	16 (49)	151	11	25	2
<i>Oceanodroma markhami*</i>		0 (3)	0 (7)		50 (2)		12	8		
<i>Oceanodroma melania</i>			0 (2)				2	0		
<i>Oceanodroma tethys</i>	7 (29)	0 (12)	0 (32)	0 (45)	0 (29)	0 (29)	176	1		
<i>Pelagodroma marina</i>		0 (1)	88 (8)		50 (4)		13	69	50-88	4,5
			Other							
<i>Fregata minor</i>			0 (3)				3	0		
<i>Gygis alba</i>			0 (2)				2	0	0	3
<i>Phaethon lepturus</i>			0 (1)				1	0		
<i>Phaethon rubricauda</i>			0 (2)				2	0	0	3
<i>Stercorarius parasiticus</i>					50 (2)		2	0	0-50	2,5
<i>Sterna fuscata</i>			0 (19)	0 (2)	0 (5)	0 (9)	35	0	0	3
<i>Sterna lunata</i>			0 (4)		0 (1)		5	0	0	3

^aIn each column is given the number of seabirds with plastic followed by the total number inspected.

^bSpecies marked by an asterisk (*) were collected at the outer edge of the Peru Current, or breed in that area; see text.

^cThe cruises are designated as A (autumn) or S (spring) by year (e.g., A84 = autumn 1984).

^dSource: 1, Ainley et al. 1990; 2, Day et al. 1985; 3, Harrison et al. 1983; 4, Imber in Day et al. 1985; 5, Ryan 1987b.

while the ship tended permanently moored oceanographic buoys, and thus were mostly from the same sites on each cruise. During spring 1987, however, the ship worked in the outer reaches of the Peru Current off Peru, and birds were collected there as well.

In our analyses, we statistically compared proportions using t-tests following angular transformations (Sokal and Rohlf 1969). Regressions were by Spearman rank correlation (r_s).

RESULTS

We looked for plastic in the stomachs of 39 species, 21 of which had not been previously inspected for this. Highest rates of plastic ingestion were evident in species that frequented the periphery of the study area or moved through it. Most obvious were the nine species that came from the edge of the Peru Current (Table 1). Plastic was found in all but one of these, and in seven the incidence equaled or exceeded 45% of the birds inspected. Incidence was very low in the Juan Fernandez petrel, *Pterodroma e. externa*, which nests on islands off Chile, but in the cape petrel, *Daption capense*, and the narrow-billed prion, *Pachyptila belcheri*, both of which only winter in the Peru Current, incidence exceeded 80%.

Various patterns of ingestion were apparent in several longer distance migrants that either pass through or spend the nonbreeding portion of their annual cycle in the study area. Incidence of plastic was very high in the sooty shearwater, *Puffinus griseus*, a species that migrates across the study area between its breeding islands in the southwestern Pacific (New Zealand) and its wintering grounds in the North Pacific. Incidence was 0% in shearwaters collected on their northward postbreeding journey during spring ($n = 6$) but was 95% among those individuals flying south in the autumn ($n = 19$). In the Leach's storm-petrel, *Oceanodroma leucorhoa*, which breeds on islands ringing the eastern and central North Pacific but which otherwise resides in the study area, frequency of plastic relative to its annual cycle was the opposite of the shearwater. Plastic occurred in 11% overall. During autumn, just after the breeding season, 13.4% contained plastic ($n = 112$) compared to 5.1% during spring ($n = 39$; $t = 3.261$, $P = 0.002$), after breeders had departed. The same relative seasonal pattern was also evident in the white-winged petrel, *Pterodroma leucoptera*, which breeds in the Tasman Sea but otherwise resides in the study area. Plastic occurred in 15% of these birds overall, with a seasonal breakdown of 5.6% ($n = 54$) during autumn, just before the nesting period, and 28.9% ($n = 38$) during spring, just after the nesting period ($t = 3.104$, $P = 0.003$). Finally, in the Stejneger's petrel, *Pt. l. longirostris*, which breeds off Chile and then migrates to North Pacific waters off Japan, plastic occurred in the same proportion of birds regardless of season (spring, coming from the nesting grounds off Chile: 52.9%, $n = 17$; autumn, coming from waters off Japan: 50.0%; $n = 8$).

Among species that reside in equatorial waters year-round, the incidence of plastic ingestion was very low, and in most it was nil. An instructive comparison is that between the migratory Leach's storm-petrel and the year-round resident wedge-rumped storm-petrel, *O. tethys*, two

species that are closely associated in the study area (unpubl. data). In Leach's, ingestion was 11%, but in wedge-rumped storm-petrel, it was only 1% ($t = 4.289$, $P < 0.001$).

Ingestion frequencies for each species were tracked by the number of plastic particles per bird. Individuals of species in which ingestion frequency was high also had large numbers of pieces in their gizzards and vice versa ($r_s = 0.7385$, $df = 14$, $P < 0.001$; for species where $n > 4$ individuals; Table 2). On the basis of that relationship, we might project for species in which samples were small, that if the one or two individuals inspected had eaten a large number of pieces, a larger sample should show a high proportion of individuals with plastic. For example, we inspected only one short-tailed shearwater, *Pu. tenuirostris*, but it contained 14 pieces of plastic. This species is known to eat large amounts of plastic (Day et al. 1985). The pattern should hold for the Buller's shearwater, *Pu. bulleri*, for which there are no other published data on plastic ingestion (the one individual in our sample contained seven pieces). One exception would be Juan Fernandez petrel, in which only 1 individual of 104 inspected contained plastic: 4 pieces, a relatively large number of pieces in our data set.

We compared the frequency of plastic as a function of age (subadult versus adult) in species where $n > 10$ birds. Compared to adults, a larger proportion of subadults should have ingested plastic (Day et al. 1985; Ryan 1987b). We found, though, that the proportion of breeding adults, nonbreeding adults, subadults, and unknowns that had ingested plastic (5.9, 35.6, 56.4, and 2.0%, respectively) did not differ from the proportion of these age classes in the total samples (7.9, 37.4, 51.8, and 3%, respectively; $\chi^2 = 8.78$, $df = 3$, $P > 0.05$). Plastic was not more prevalent in subadults.

DISCUSSION

We found little evidence that terns, tropicbirds, or frigatebirds ate plastic, as also noted by Harrison et al. (1983), Day et al. (1985), and Sileo et al. (1990). These species eat active prey, and even if they did ingest plastic, they lack a gizzard where, in Procellariiformes, plastic accumulates.

Among the petrels we sampled, ingestion was mainly a function of feeding behavior, what parts of the Pacific Ocean they frequented, and the amount of time passed since leaving polluted areas. Petrels such as Juan Fernandez petrel or wedge-tailed shearwater, *Pu. pacificus*, which chase airborne flyingfish and squid or prey being driven to the surface by predatory fish (i.e., very active prey), exhibited low rates of plastic ingestion. The majority of other petrel species are less specialized feeders, and they eat prey both live and dead. Some species, such as white-faced storm-petrel, *Pelagodroma marina*, and prions, genus *Pachyptila*, almost always contain very high loads of plastic (Day et al. 1985; Ryan 1987b). It is probable that their prey search-images (including particle size) and their propensity to associate with convergences account for their high susceptibility to plastic as food. On the other hand, some species

Table 2.--The average number of pieces of plastic, and standard error, in the gizzards of seabirds collected in the study area.

Species	Average number of pieces	SE	Number of birds
<i>Daption capense</i>	8.4	2.3	7
<i>Pachyptila belcheri</i>	8.2	3.4	5
<i>Procellaria aequinoctialis</i>	3.5	1.5	2
<i>Pterodroma cooki defillipiana</i>	9.8	4.8	4
<i>Pterodroma e. externa</i>	4		1
<i>Pterodroma e. cervicallis</i>	1		1
<i>Pterodroma leucoptera</i>	2.1	0.8	14
<i>Pterodroma l. longirostris</i>	3.1	0.6	14
<i>Pterodroma l. pycrofti</i>	2		1
<i>Pterodroma nigripennis</i>	1		1
<i>Pterodroma rostrata</i>	1		1
<i>Puffinus bulleri</i>	7		1
<i>Puffinus nativitatus</i>	1		1
<i>Puffinus griseus</i>	10.5	2.7	18
<i>Puffinus tenuirostris</i>	14		1
<i>Puffinus pacificus</i>	1		1
<i>Oceanodroma leucorhoa</i>	3.3	0.4	16
<i>Oceanodroma markhami</i>	2		1
<i>Pelagodroma marina</i>	12.2	3.5	9

that are obligate scavengers, such as Tahiti petrel, *Pt. rostrata*, do not eat much plastic (unpubl. data). They, too, might have a particular search-image.

Temporal and geographic aspects of plastic ingestion were also evident, and these patterns bear on the question of how long plastic is retained in seabird guts. Ryan (1988b) proposed that most of the observed seasonal change in plastic loads is a function of the transfer of plastic from breeding adults to chicks rather than degradation within the digestive tracts of adults. If this is correct, we should have found highest incidence in subadults (who have never bred and thus have never had an opportunity to disgorge plastic), lowest incidence in postbreeding adults, and intermediate values in nonbreeding adults. Our data do not indicate such a pattern. First, in species for which samples were sufficiently large to make comparisons, we found no indication that subadults had more or less plastic than adults. Second, the seasonal patterns we observed suggest that degradation is an important process, and that the time scale is on the order of 6 months or slightly longer. Ryan and Jackson (1987) assumed that degradation rate is constant throughout the time that plastic resides in a seabird digestive tract, and they based their extrapolation on plastic fed to birds and then inspected after retention for only 12 days.

However, plastic subjected constantly to digestive acids and abrasion could degrade at a faster rate as time passes, especially if its surface-to-volume ratio increases. Van Franeker and Bell (1988) found that the mass of plastic particles in cape petrels collected in the Antarctic (where there is no plastic to replenish ingested loads) decreased by 50% over a 3-month period. This rate supports a shorter degradation period than that proposed by Ryan and Jackson (1987; see Day et al. 1985).

We interpret our seasonal and geographic data as follows. Those birds that reside year-round in the region of the South Equatorial Current and the Equatorial Countercurrent (Wyrteki 1967), i.e., our study area, exhibited only incidental ingestion of plastic. We observed little floating debris during our censuses, probably because there are few, if any, large industrial source areas in the central Pacific for plastic and there are no large human population centers except for Oahu. Among petrel species of the Northwestern Hawaiian Islands, which lie in the North Equatorial Current, Harrison et al. (1983) also found only low levels of plastic ingestion. We did not collect any samples in the Panamanian Bight, where we observed much floating plastic and other flotsam on our bird censuses.

Compared to year-round equatorial residents, results were much different for petrels that were either passing through the study area or that came a long way from polluted areas to spend their nonbreeding period there. Those species that had just come from areas where industrial plastic pellets are common, for example, off Japan and California (Day et al. 1985; Pruter 1987), had ingested significant amounts of pellets. Several of these species, including adults and subadults, had higher plastic loads after frequenting polluted waters (where adults bred). This suggests that loss of plastic while in plastic-free waters (because erosion outpaces ingestion) is important. Since the migrations are annual, degradation over a 6- to 12-month period would best fit the patterns. As examples, species or populations that had come from the eastern and central North Pacific had high plastic loads (i.e., Buller's, short-tailed, and sooty shearwater and Leach's storm-petrel). The first three were in the prebreeding portion of the annual cycle, and the last was postbreeding. This inconsistency is contrary to the off-loading through chick-feeding hypothesis. When specimens of these species have been inspected in California waters they contain much plastic (Balz and Morejohn 1976). The fact that sooty shearwaters migrating north from New Zealand or Leach's storm-petrel not newly arrived from the north have low plastic loads confirms that it is in the eastern North Pacific that these species are ingesting plastic. Sooty shearwaters, probably of the South American breeding population, also frequent the Peru Current, but we did not collect any. A portion of the New Zealand population of this species migrates to polluted waters off Japan, but the eastern Pacific position of our study area and the flight directions observed indicate that the birds we sampled were moving between New Zealand and the eastern North Pacific.

The incidence of plastic ingestion was also high in species collected at the periphery of the Peru Current or that came to the study area from waters off Chile (i.e., cape petrels, prions, white-chinned petrel,

Procellaria aequinoctialis, Cook's petrel, *Pt. cooki defilippiana*, and Stejneger's petrel). Cape and white-chinned petrels collected south of South America have much lower plastic loads than those inspected in this study (Ainley et al. 1990). This supports the suggestion that the specimens inspected in this study encountered the plastic in the Peru Current. Stejneger's petrel moves through the study area between Chilean waters and waters off Japan, and thus its plastic could have come from either area. Unlike the other migrants discussed above, these petrels had similar plastic loads regardless of whether they were moving south or north. This supports the suggestion that they were ingesting plastic off Chile and off Japan. Bourne and Clark (1984) noted specific areas of nearshore Chile where significant amounts of plastic had accumulated, but little information is available on a larger geographic scale. One species they observed in association with scum and flotsam lines was *Pa. belcheri*, which in our samples contained much plastic. Species from the northern part of the Peru Current (i.e., several species of the genus *Oceanodroma*) had low plastic loads.

Though not as high as in the birds noted above, the occurrence frequency of plastic was high in white-winged petrel, which as far as we know nests in the subtropical Tasman Sea area. The pattern of low plastic incidence before breeding and after wintering away from polluted areas, but high incidence after breeding, again is contrary to the chick off-loading hypothesis. Along the Equator, one of two individuals of Pycroft's petrel, *Pt. longirostris pycrofti*, which breeds in northern New Zealand, contained plastic, and four of six white-faced storm-petrels, that could also have come from northern New Zealand, contained much plastic. Imber (in Day et al. 1985) reported plastic in 50% of specimens of the latter species from Chatham Island (NZ); Harper and Fowler (1987) found that plastic was prevalent in prions washed ashore in northern New Zealand; van Franeker and Bell (1988) noted significant loads of plastic in Antarctic-breeding species that wintered in the Indian Ocean and Tasman Sea; and Gregory (1978) has noted significant amounts of plastic on most New Zealand beaches, north and south. Thus, it is not surprising that petrels that spend time in New Zealand waters show high plastic ingestion. It is surprising that those sooty shearwaters that were newly arrived to our study area from New Zealand had no plastic, although the large majority of sooty shearwaters nest away from and south of the main islands of New Zealand, and thus away from industrialized areas. This may account for the lower rates of plastic ingestion in these birds.

ACKNOWLEDGMENTS

We wish to thank the EPOCS Council for the invitation to participate in their cruises, and the officers and crews of the NOAA ships *Discoverer*, *Oceanographer*, and *Researcher* for their logistic support. Our research was funded by the National Science Foundation (Grant OCE 8515637), the National Geographic Society, and Point Reyes Bird Observatory (PRBO). This is Contribution No. 421 of PRBO.

REFERENCES

- Ainley, D. G., W. R. Fraser, and L. B. Spear.
1990. The incidence of plastic in the diets of Antarctic seabirds. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Balz, D. M., and G. V. Morejohn.
1976. Evidence from seabirds of plastic particle pollution off central California. *West. Birds* 7:111-112.
- Bourne, W. R. P., and G. C. Clark.
1984. The occurrence of birds and garbage at the Humboldt Front off Valparaiso, Chile. *Mar. Pollut. Bull.* 15:343-344.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Gregory, M.
1978. Accumulation and distribution of virgin plastic granules on New Zealand beaches. *N.Z. J. Mar. Freshwater Res.* 12:399-414.
- Harper, P. C., and J. A. Fowler.
1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptilla* spp.). *Notornis* 34:65-70.
- Harrison, C. S., T. S. Hida, and M. P. Seki.
1983. Hawaiian seabird feeding ecology. *Wildl. Monogr.* 85:1-71.
- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Ryan, P. G.
1987a. The effects of ingested plastic on seabirds: Correlations between plastic load and body condition. *Environ. Pollut.* 46: 119-125.
1987b. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.
1988a. Effects of ingested plastic on seabird feeding: Evidence from chickens. *Mar. Pollut. Bull.* 19:125-128.

- 1988b. Intraspecific variation in plastic ingestion by seabirds and the flux of plastic through seabird populations. *Condor* 90:446-452.
- Ryan, P. G., and S. Jackson.
1987. The lifespan of ingested plastic particles in seabirds and their effect on digestive efficiency. *Mar. Pollut. Bull.* 18: 217-219.
- Sileo, L., P. R. Sievert, M. D. Samuel, and S. I. Fefer.
1990. Prevalence and characteristics of plastic ingested by Hawaiian seabirds. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Sokal, R. R., and F. J. Rohlf.
1969. *Biometry*. W. H. Freeman, San Francisco.
- Van Franeker, J. A., and P. J. Bell.
1988. Plastic ingestion by petrels breeding in Antarctica. *Mar. Pollut. Bull.* 19:672-674.
- Wyrтки, K.
1967. Circulation and water masses in the eastern equatorial Pacific Ocean. *Int. J. Oceanol. Limnol.* 1:117-147.
-

PREVALENCE AND CHARACTERISTICS OF PLASTIC
INGESTED BY HAWAIIAN SEABIRDS

Louis Sileo, Paul R. Sievert,* Michael D. Samuel
National Wildlife Health Research Center
U.S. Fish and Wildlife Service
Madison, Wisconsin 53711, U.S.A.

and

Stewart I. Fefer**
Hawaiian and Pacific Complex National Wildlife Refuge
U.S. Fish and Wildlife Service
Honolulu, Hawaii 96850, U.S.A.

*Present address: Department of Biology, University of Pennsylvania,
Philadelphia, Pennsylvania 19104.

**Present address: U.S. Fish and Wildlife Service, 1 Gateway Center, Suite
700, Newton Corner, Massachusetts 02158.

ABSTRACT

The prevalence of plastic in 18 species of seabirds at seven study sites in the Hawaiian Islands and Johnston Atoll was studied during 1986 and 1987. Stomach samples were collected by induced emesis from 1,803 live birds of 15 species and during necropsy of 277 dead birds of 5 species. The prevalence of ingested plastic varied greatly among species; age, year of collection, and location of the study site had less pronounced but significant effects. Ingested plastic was absent or uncommon in terns and noddies. Plastic was not found at all in samples from gray-backed terns, *Sterna lunata*, or white terns, *Gygis alba*. The prevalence was low in sooty terns, *Sterna fuscata* (0 to 8%), and brown, *Anous stolidus*, and black noddies, *A. minutus* (0 to 3%). Plastic was much more prevalent (67 to 100%) in chicks of black-footed, *Diomedea nigripes*, and Laysan albatrosses, *D. immutabilis*. In the 11 other species prevalence ranged from 0 to 44% depending on the age, year of collection, and location. The mean volume of plastic in samples collected at necropsy from Laysan albatross chicks was higher in 1986 (46 cc) than in 1987 (5 cc). Prevalence was generally higher in seabirds which fed at the surface. Fragments of manufactured articles were the most common type of plastic found. Other plastics included pellets, Styrofoam, bottle caps, bags, and sponges. The largest individual item (200 cc) and the greatest diversity of plastic items were found in albatross chicks.

INTRODUCTION

Production of plastic products and dumping of plastic garbage in the ocean have increased dramatically in the past 25 years. From 1960 to 1985 the United States increased plastic production from 2.9 to 20.7 billion kg (6.3 to 47.9 billion lb) per year (Iudicello and O'Hara 1986). The prevalence of plastic ingestion by seabirds also has increased. Plastic was first reported in the stomachs of seabirds in 1960, when it was discovered in broad-billed prions, *Pachyptila vittata*, that were killed during storms on New Zealand (Harper and Fowler 1987). By 1985, at least 70 species of seabirds from all the world's oceans were known to ingest plastic (Day et al. 1985; Ryan 1987a).

Seabirds ingest a wide variety of plastic types, but 2- to 5-mm diameter plastic pellets and fragments of manufactured plastic products are most common (Day et al. 1985; Ryan 1987a). Other types of plastic found in seabird stomachs include Styrofoam, fibers, bags, bottle caps, and toys (Kenyon and Kridler 1969; Harrison et al. 1983; Dickerman and Goelet 1987; Ryan 1987a). Plastic particles may be eaten intentionally, accidentally mistaken for prey, or ingested secondarily when hidden within a food item (Day et al. 1985). For example, small plastic particles may be confused with fish eggs or small invertebrates, while larger pieces can be mistaken for squid, jellyfish, fish, or other large prey. Flyingfish often deposit their egg masses on floating debris such as plastic, which may then be ingested by albatrosses (Pettit et al. 1981; Harrison et al. 1983).

Plastic ingestion has not been reported as a significant health problem in seabird populations, but several studies have indicated negative effects on individual birds. Ingested plastic may distend or block the proventriculus or intestine, causing erosions or ulcers (Pettit et al. 1981; Sileo and Fefer 1987) or possibly starvation (Dickerman and Goelet 1987). The presence of ingested plastic was correlated with elevated (but nonthreatening) concentrations of polychlorinated biphenyls in great shearwaters, *Puffinus gravis* (Ryan et al. 1988), and with reduced body weight in red phalaropes, *Phalaropus fulicarius* (Connors and Smith 1982) and domestic chickens, *Gallus domesticus* (Ryan 1988a). Other studies have shown no correlation between body weight and plastic presence (Day et al. 1985; Ryan 1987b). Ryan and Jackson (1987) found no reduction in assimilation efficiency in white-chinned petrels, *Procellaria aequinoctialis*, artificially fed large quantities of plastic particles.

Studies of the prevalence of plastic ingestion in seabird communities, based on examination of regurgitations or carcasses, have been conducted in the North Pacific (Day et al. 1985) and South Atlantic Oceans (Furness 1985; Ryan 1987a). Reports of plastic in Laysan and black-footed albatrosses have shown that these birds ingest the widest variety of plastic items (Kenyon and Kridler 1969) and ingest the largest volumes of plastic more frequently than other seabirds (Fry et al. 1987; Sileo and Fefer 1987). Harrison et al. (1983) reported cursory data for other Hawaiian seabirds, but systematic study of the prevalence of plastic ingestion in the Hawaiian avifauna has not been done. Our objective was to determine the prevalence of ingested plastic in three orders of Hawaiian

seabirds (Procellariiformes, Pelecaniformes, and Charadriiformes) from different islands by examining stomach contents. We evaluated the prevalence of ingested plastic for associations with species, location of study site, year of collection, and ages of birds sampled.

STUDY AREA AND METHODS

Stomach samples were collected from 18 species of seabirds on Kauai, Maui, Nihoa, Tern, and Laysan Islands, Pearl and Hermes Reef, and Midway in the Hawaiian Islands. Approximately 5 million seabirds of 22 species nest on these islands (Harrison et al. 1984). Samples were also collected on Johnston Atoll, 1,400 km southwest of Honolulu. Seabirds from remote, uninhabited islands were sampled by the authors or other biologists as travel circumstances permitted. One of us (Sievert) spent 4 months in 1986 and 6 months in 1987 on Midway.

To avoid killing large numbers of seabirds, a stomach pumping method was used to recover the proventricular contents (Wilson 1984; Ryan and Jackson 1986). The method reportedly recovers 89 to 100% of the proventricular contents, although Ryan and Jackson (1986) found that the proportion (by mass) of food recovered by a single pumping was negatively correlated with total stomach content in white-chinned petrels. Seawater was pumped through a 10-mm (outside diameter) plastic tube into the proventriculus of larger species with a manual pump (Black and Decker Model No. JSO-1500). Smaller species, such as terns and petrels, were given seawater using a 6-mm (outside diameter) plastic tube attached to a 140-cc syringe. Seawater was pumped into the proventriculus until water became visible in the esophagus. The tube then was quickly removed and the bird inverted over a container. The bill was held open with one hand and large objects were massaged through the esophagus if the bird was having difficulty regurgitating. A 1-mm diameter mesh net was used to skim plastic items from the surface of the regurgitant for later characterization.

The samples were collected from April 1986 to November 1987 by induced emesis from live seabirds or during necropsy of carcasses. Necropsy samples were obtained from carcasses found dead of natural causes and from six euthanized Laysan albatross chicks. All necropsies were performed by the authors. Dehydration was the most common natural cause of death in 1987 (Sileo et al. 1990). Seabird stomach samples were collected by different biologists in different locations, which resulted in statistical confounding of investigator effect with location. The confounding was unavoidable because the remoteness and inaccessibility of the study sites precluded replicative sampling by different biologists. Investigator effect was minimized by having each biologist complete a training session for the stomach pumping method. For most species, stomach samples were collected from chicks because they were more accessible and more easily captured. All live birds were approached on foot and captured by hand. Age of the seabirds was determined by plumage characters. When it was not possible to differentiate between free-flying juveniles and adults, all were assigned to an arbitrary "either" category. Birds were sampled once and banded with a U.S. Fish and Wildlife Service leg band to prevent repetitive sampling.

We tested the effectiveness of the sampling technique by pumping the stomachs of six Laysan albatross chicks, then euthanizing and examining them. The volume of proventricular content not removed by the stomach pump was determined. We necropsied birds that were found dead and inspected their digestive tracts for the presence of plastic. This was the only method used for 58 Bonin petrels, *Pterodroma hypoleuca*, 3 dark-rumped petrels, *Pterodroma phaeopygia*, and 18 Newell's shearwaters, *Puffinus auricularis* (newelli). The proventriculus and ventriculus of all necropsied birds were examined, and the intestines also were examined for a sample of 39 black-footed albatrosses, 57 Laysan albatrosses, and 12 Bonin petrels. Contents of the entire intestine from pylorus to cloaca were stripped. All plastic was separated from the stomach content by fresh water flotation and saved for later characterization.

Plastic was characterized by type, volume, and color. Types were fiber, pellet, Styrofoam, fragment, bottle cap, bag, sponge, and other. Fiber included fishing line, net fragments, and rope. Pellets were 2 to 5 mm diameter spherical polyethylene particles primarily used as the raw material for manufacturing plastic products (Carpenter et al. 1972). Bottle caps included all types of plastic lids, and bags were defined as plastic film less than 1 mm thick in the form of a bag or sheet. Other included children's toys, cigarette lighters, hair combs, balloons, gloves, condoms, tubing, sandals, wrappers, and bandages. The number of plastic items of each type per bird was counted except for fiber, which was recorded as present or absent. Plastic fragments were assigned to one of five color groups: white, which included white, yellow, tan, and brown; black, which included gray and black; blue, which included purple and blue; green, which included all shades of green; and red, which included pink and red. One of us (Sievert) was studying growth rates of a marked population of albatross chicks at a study site on Midway. Albatross carcasses found dead at this study site were dissected and the total volume of recovered plastic was determined by water displacement. The buoyant contents were placed in a wire screen bag (1.5-mm diameter mesh) and immersed in a 2,000-ml beaker filled with tap water. The volume of water displaced was measured. In 1986, all measurements were made in 1-cc increments, and in 1987, in 5-cc increments. The volume of the largest plastic fragment from each bird was calculated from the linear dimensions of the fragment.

Statistical analysis of the plastic prevalence data followed the general recommendations of Freeman (1987). Chi-square tests for association were used for 2×2 tables of plastic prevalence and other independent variables. Fisher's exact test was used when cell expectations were small (<5). Several 2×2 tables were combined using the Cochran-Mantel-Haenszel procedure to test hypotheses of an overall association between plastic prevalence and the independent variables. This procedure provided a means for stratifying the plastic prevalence data from each unique combination of island, age, species, and year so that there was only a single independent variable in each test. Hypothesis tests were considered significant at the 0.05 level, however, P values for each test are also reported. Samples from wedge-tailed shearwaters, *Puffinus pacificus*, and red-tailed tropicbirds, *Phaethon rubricauda*, from different islands permitted assessment of the effect of geographic location on

plastic ingestion. The influence of year of collection was determined by comparing prevalence in 1986 and 1987 in 11 combinations of species, location, and age. The prevalence in chicks was compared to adults in nine combinations of year, species, and location. If the prevalences of two or more closely related species were not significantly different, data from the species were combined to make comparisons with broader taxonomic groups.

RESULTS

In 1987 stomach pumping removed at least 50% of the proventricular plastic from five (83%) of six Laysan albatross chicks, each of which contained 5 to 10 cc of plastic (Table 1). Stomach pumping failed to remove any plastic from the remaining chick, which contained 5 cc of plastic as well as 100 cc of rock. The rock was primarily buoyant volcanic pumice. Induced emesis recovered half of the proventricular plastic as long as the pumice content was less than 55% of the total proventricular content. For comparison, the prevalence of proventricular plastic in the carcasses of Laysan albatross chicks examined by dissection at Midway was 94% in 1986 and 98% in 1987 (Table 2).

Stomach-pumped samples from both 1986 and 1987 were collected from 11 unique combinations of species, location, and age (Table 2). Significant differences between years of collection were found for red-tailed tropicbird chicks at Midway and adult wedge-tailed shearwaters at Kauai; however, the overall test for a significant association between year of collection and plastic prevalence was not quite significant ($P = 0.053$). To remove potential investigator effect, combinations where all the birds were sampled by only one of us (Sievert) were analyzed. These included the five combinations sampled at Midway, and the overall P value (0.029)

Table 1.--Rock, plastic, and other proventricular contents removed by stomach pumping from six Laysan albatross chicks that were then euthanized and necropsied.

Bird No.	Total content		Rock content		Plastic content	
	Volume present ^a (cc)	Percent removed	Volume present ^a (cc)	Percent removed	Volume present ^a (cc)	Percent removed
1	140	3	100	5	5	0
2	100	20	55	18	10	50
3	60	17	20	25	10	50
4	45	44	15	67	5	100
5	20	50	10	50	10	50
6	20	50	5	100	10	50

^aVolumes were measured in 5 cc increments.

Table 2.--Statistical comparisons of the prevalence of plastic in stomach samples collected from Pacific seabirds.^a

Method ^b	Year	Species ^c	Site ^d	Age ^e	N	%	P Values				Species
							1986-87	Age	Islands		
N	1986	BFAL	M	C	28	89	--	--	--	--	ab 0.433
N	1986	LAAL	M	A	31	35	--	i 0.000	--	--	--
N	1986	LAAL	M	C	78	94	--	i 0.000	--	--	ab 0.433
N	1986	BOPE	M	B	58	29	--	--	--	--	--
N	1987	BFAL	M	C	18	100	--	--	--	--	ac 1.00
N	1987	LAAL	M	C	43	98	--	--	--	--	ac 1.00
N	1987	NESH	K	I	18	11	--	--	--	--	--
P	1986	BFAL	L	C	56	79	a 0.097	--	--	--	--
P	1986	BFAL	T	C	1	0	--	--	--	--	--
P	1986	LAAL	L	C	24	92	c 0.161	--	--	--	--
P	1986	LAAL	T	C	12	92	d 0.245	--	--	--	--
P	1986	WTSB	K	A	150	18	i 0.024	--	ba 1.00	ca 0.452	--
P	1986	WTSB	M	A	15	13	j 0.304	h 0.620	ba 1.00	da 0.716	dc 1.000
P	1986	WTSB	M	C	11	27	--	h 0.620	--	--	--
P	1986	WTSB	T	A	48	23	k 0.272	--	ca 0.452	da 0.716	--
P	1986	CHSH	M	A	1	0	--	--	--	--	dc 1.000
P	1986	RTTB	M	A	8	0	--	c 0.054	--	--	--
P	1986	RTTB	M	C	16	44	g 0.001	c 0.054	--	--	--
P	1986	RFBO	L	A	34	0	--	--	--	--	--
P	1986	RFBO	L	C	7	0	e 1.00	--	gb 1.00	--	--
P	1986	RFBO	M	A	4	0	--	b 1.00	--	--	--
P	1986	RFBO	M	C	19	11	f 0.119	b 1.00	gb 1.00	--	--
P	1986	SOTE	M	A	26	8	--	f 0.172	--	--	ba 1.00
P	1986	SOTE	M	C	36	0	h 0.493	f 0.172	--	--	bb 1.00
P	1986	GBTE	M	A	7	0	--	--	--	--	ba 1.00
P	1986	GBTE	M	C	29	0	--	--	--	--	bb 1.00
P	1986	BRNO	M	A	17	0	--	a 1.00	--	--	ba 1.00
P	1986	BRNO	M	C	86	1	b 1.00	a 1.00	--	--	bb 1.00
P	1986	BLNO	L	A	18	0	--	--	--	--	--

Table 2. ---Continued.

P Values										
Method ^b	Year	Species ^c	Site ^d	Age ^e	N	#	1986-87	Age	Islands	Species
P	1987	BFAL	L	C	36	92	a 0.097	--	eb 0.710	aa 0.239
P	1987	BFAL	N	C	21	86	--	--	fb 1.00	--
P	1987	BFAL	P	C	35	97	--	--	--	ad 0.614
P	1987	BFAL	T	C	35	89	--	--	fb 1.00	ae 0.206
P	1987	LAAL	L	C	35	100	c 0.161	--	eb 0.710	aa 0.239
P	1987	LAAL	P	C	35	91	--	--	--	ad 0.614
P	1987	LAAL	T	C	6	67	d 0.245	--	--	ae 0.206
P	1987	BUPE	N	A	38	5	--	--	--	--
P	1987	WTSH	J	A	60	5	--	--	ab 0.004	--
P	1987	WTSH	K	A	35	3	i 0.024	g 0.067	bb 0.003	cb 0.064
P	1987	WTSH	K	C	7	29	--	g 0.067	--	--
P	1987	WTSH	L	A	35	14	--	--	ga 0.145	ea 0.888
P	1987	WTSH	M	A	35	29	j 0.304	--	ab 0.004	dd 0.093
P	1987	WTSH	N	A	60	3	--	--	bb 0.003	ga 0.145
P	1987	WTSH	T	A	85	15	k 0.272	--	fa 0.033	fa 0.033
P	1987	CHSH	L	A	36	17	--	--	cb 0.064	ea 0.888
P	1987	CHSH	N	A	2	50	--	--	db 0.093	da 0.782
P	1987	SOSP	L	A	18	33	--	e 0.903	--	db 0.522
P	1987	SOSP	L	C	17	35	--	e 0.903	--	--
P	1987	RTTB	J	C	50	4	--	--	aa 1.00	--
P	1987	RTTB	M	A	39	5	--	d 1.00	--	--
P	1987	RTTB	M	C	48	2	g 0.0001	d 1.00	aa 1.00	--
P	1987	RTTB	T	C	50	14	--	--	--	--
P	1987	MABO	L	C	20	5	--	--	--	--
P	1987	RFBO	L	C	35	3	e 1.00	--	ec 1.00	gc 1.00
P	1987	RFBO	M	C	35	0	f 0.119	--	dc 1.00	gc 1.00
P	1987	RFBO	T	C	35	0	--	--	dc 1.00	ec 1.00
P	1987	GRFR	M	C	45	18	--	--	--	--

Table 2.--Continued.

Method ^b	Year	Species ^c	Site ^d	Age ^e	N	%	P Values		Islands	Species	
							1986-87	Age			
P	1987	SOTE	M	C	35	3	h	0.493	--	--	bc 1.00
P	1987	GBTE	M	A	10	0	--	--	--	--	--
P	1987	GBTE	M	C	25	0	--	--	--	--	bc 1.00
P	1987	BRNO	M	C	35	0	b	1.00	--	--	--
P	1987	BRNO	T	C	15	0	--	--	--	--	ca 1.00
P	1987	BLNO	T	C	35	3	--	--	--	--	ca 1.00
P	1987	WHITE	M	C	35	0	--	--	--	--	--

^aRows having the same letter codes and the same P value were compared by the Cochran-Mantel-Haenszel procedure (Freeman 1987). The first letter of a double letter P value code indicates a species group comparison; the second letter of a double letter code indicates the two members of a paired comparison within the designated group.

^bN = sample collected at necropsy, P = sample collected by stomach pumping.

^cBFAL = black-footed albatross, BLNO = black noddy, BRNO = brown noddy, BOPE = Bonin's petrel, BUPE = Bulwer's petrel, CHSH = Christmas shearwater, GBTE = gray-backed tern, GRFR = great frigatebird, LAAL = Laysan albatross, NESH = Newell's shearwater, MABO = masked booby, RFBO = red-footed booby, RTTB = red-tailed tropicbird, SOSP = sooty storm petrel, SOTE = sooty tern, WHITE = white tern, WTSH = wedge-tailed shearwater.

^dJ = Johnston Atoll, K = Kauai, L = Laysan Island, M = Midway, N = Nihoa Island, P = Pearl and Hermes Reef, T = Tern Island.

^eA = adult, C = chick, B = both adults and free-flying immatures, and I = free-flying immatures.

indicated a significantly higher prevalence in 1986. Thus year of collection had a significant effect in species and age combinations from Midway.

To test age, prevalence data based on stomach-pumped samples collected from both adults and chicks of eight combinations of species, location, and year of collection were compared (Table 2). None of the individual results were significantly different; however, in all species except the sooty tern, the trend was for higher prevalence among chicks. This caused a significant overall association between age and prevalence ($P = 0.05$). A second analysis of age was done which included the 1986 necropsy samples from the Laysan albatrosses at Midway. There was a significant difference between the Laysan albatross chicks and adults ($P < 0.00$), and the overall association between age and prevalence was again significant ($P < 0.00$).

In 1987 the adult wedge-tailed shearwaters of Johnston Atoll had significantly lower prevalence (5%) of plastic in pumped stomach samples than at Midway (29%) in individual tests (Table 2), and the overall test of association also showed a significantly ($P = 0.01$) lower prevalence at Johnston. In 1987, adult wedge-tailed shearwaters had higher prevalence (29%) at Midway than at Kauai (3%), and at Tern Island (15%) than at Nihoa (3%). The individual tests for these particular combinations were significant (Table 2), but the overall tests of association were not ($P = 0.064$, and $P = 0.078$, respectively), suggesting the absence of a consistent pattern in differences in prevalence among these island pairs. No other significant differences were detected between locations.

It was possible to compare prevalence between black-footed and Laysan albatrosses using both stomach-pumped and necropsy samples for five combinations of location, age, and year of collection (Table 2). Neither the individual results nor the overall test of association ($P = 0.98$) were significant. Consequently, data from both species of albatrosses were combined for later comparisons with other taxa. Stomach pumped samples from three combinations of location, age, and year were used to compare gray-backed terns and sooty terns. Again, neither individual nor overall ($P = 0.264$) tests of association were significant, and both species of terns were combined for comparison with other taxa. Brown noddies and black noddies were also not significantly different ($P = 1.00$) and were combined. Finally, the combined terns were compared to the combined noddies, and again neither individual ($P = 0.542$, 1.00 , or 1.00) nor the overall ($P = 0.519$) tests of association were significant, so that all tern and noddy species were combined for comparisons with other taxa. Stomach-pumped samples from Christmas shearwaters, *Puffinus nativitatis*, and wedge-tailed shearwaters were compared for three combinations of location, age, and year (Table 2). There were no significant individual or overall ($P = 0.69$) association, and data from these two species of shearwaters were combined for comparisons with other taxa. In comparisons between taxonomic groups, the albatrosses had significantly higher prevalences, the terns and noddies the lowest, and the shearwaters, tropicbirds, and boobies were intermediate (Table 3).

Table 3.--Comparisons of the prevalence of ingested plastic in stomach samples of five taxonomic groups of Hawaiian seabirds. A "-" means that the taxa in the row had significantly lower prevalence than the taxa in the column, "ns" means that there was no significant difference between the taxa in the row and column, "+" means that the taxa in the row had significantly higher prevalence than the taxa in the column, and "nc" means no comparison was done.

Taxa ^a	LAAL/BFAL	GBTE/SOTE BLNO/BRNO	RFBO	CHSH/WTSH	RTTB
LAAL/BFAL		+	+	nc	+
GBTE/SOTE BLNO/BRNO	-		ns	-	-
RFBO	-	ns		ns	-
CHSH/WTSH	nc	+	ns		+
RTTB	-	+	+	-	
Range of prevalence in chicks (%)	67-100	0-36	0-11	27-29	2-44
Range of prevalence in adults (%)	35	0-8	0-34	3-29	0-5

^aBFAL - black-footed albatross, BLNO - black noddy, BRNO - brown noddy, CHSH - Christmas shearwater, GBTE - gray-backed tern, LAAL - Laysan albatross, RFBO - red-footed booby, RTTB - red-tailed tropicbird, SOTE - sooty tern, WTSH - wedge-tailed shearwater.

Necropsy examinations of carcasses revealed plastic in the intestines of many of the albatrosses (Table 4). While plastic was usually present in both the proventriculi and ventriculi of albatrosses, it was more common in the ventriculi of petrels and shearwaters (Table 4). The mean volume of plastic in the Laysan carcasses was about nine times greater in 1986 than 1987 (Table 5).

Albatrosses held the widest diversity of plastic types and were the only species to ingest bottle caps, bags, sponges, and a variety of other items. Styrofoam, fibers, sponges, bags, and bottle caps were more common in black-footed albatrosses than Laysan albatrosses, whereas plastic pellets were more common in the latter. Fragments were the most common type of plastic ingested by all the species except Bonin petrels, which contained fibers most frequently. Albatrosses ate the largest individual plastic items; the volume of the largest single item (plastic sheet) recovered was 200 cc. Albatrosses also ate the largest fragments (up to 25.2 cc); petrels, shearwaters, tropicbirds, boobies, and frigatebirds ingested moderate-sized ones; and storm-petrels and terns consumed the smallest (up to 2.0 cc, Table 6). Most of the fragments recovered were white regardless of the species of bird (Table 7). Samples from sooty storm-petrels, *Oceanodroma tristrami*, contained red fragments more frequently than samples from any of the other species. Great frigatebirds, *Fregata minor*, contained only white or black fragments.

DISCUSSION

Although we never recovered more than 50% of the volumes of plastic greater than 5 cc, or over 50% of the total stomach content of the euthanized albatross chicks, the reliability of stomach pumping seemed acceptable for simply detecting the presence of plastic in Laysan albatross chicks, and thus for prevalence information. Efficacy in the other species was not determined. Ryan and Jackson (1986) got higher (89-100%) yields of the total stomach content of petrels than we did for albatross chicks, although Ryan and Jackson were studying the removal of all dietary items and we were interested only in detecting the presence of plastic. Large volumes of proventricular rock seemed to interfere with removal of small volumes of plastic. Other reports also state that stomach pumping is less effective in birds that have full stomachs with tightly packed contents (Ryan and Jackson 1986). Stomach pumping removes only the proventricular content of petrels, and may or may not remove gizzard content from Pelecaniformes, Charadriiformes, and some Procellariiformes (Diomedidae) (Ryan and Jackson 1986). Our stomach pumping results possibly underestimated prevalence in our study and this may have been more significant in individuals with small plastic loads. But stomach pumping precluded killing 1,803 seabirds, simply to learn what they were eating.

There are several reports that the prevalence of ingested plastic in seabirds is increasing over the long term. Day et al. (1985) reported increases in short-tailed shearwaters, *Puffinis tenuirostris*, in the 1970's. Van Franeker (1985) noted an increase in the number of plastic particles in fulmars, *Fulmaris glacialis*, in the North Sea through the early 1980's. Harper and Fowler (1987) reported long-term interannual increases in plastic prevalence based on large samples of prions, *Pachyptila* spp., found dead on New Zealand beaches from <5% in 1960 to >20% in 1970. Fry et al. (1987) show that the prevalence (80-90%) in the Laysan albatrosses of the Northwestern Hawaiian Islands in the late 1980's was higher than the 74% reported by Kenyon and Kridler (1969), although the sampling methods were different. Frequency of occurrence increased in Antarctic prions, *P. desolata*, in the Southern Ocean during the early 1980's (Ryan 1988b). It is widely assumed that increasing prevalence in seabirds reflects increasing pollution of the marine environment, and this is probably correct; however, the marked short-term interannual difference we found (44% in red-tailed tropicbird chicks at Midway in 1986 versus 2% in 1987) in the same species at the same nesting colony examined by the same scientist suggests that long-term relationships must be interpreted carefully. Such interannual variation might be caused by changes in the amount of plastic dumped in the ocean, movement of floating plastic by winds and currents, seabird foraging areas, or feeding behavior. The volumes of plastic we recovered from the necropsied albatross chicks are 10 to 100 times greater than the volumes reported for other species (Day et al. 1985; Furness 1985; Ryan 1987a, Bayer and Olson 1988; van Franeker and Bell 1988). The mean 18.3 cc volume (estimated as $0.9053 \times \text{weight}$) recovered from Laysan albatross chicks by Kenyon and Kridler (1969) compares well with our data.

Table 4.--The prevalence of plastic in the gastrointestinal tracts of seabird carcasses examined by necropsy in the Hawaiian Islands in 1986 and 1987 [percent containing plastic (number examined)].

Species	Age ^a	Plastic location		
		Proventriculus	Ventriculus	Intestines
Black-footed albatross	Chick	95 (44)	93 (44)	30 (20)
Laysan albatross	Chick	100 (131)	95 (131)	39 (57)
Bonin petrel	Both	27 (99)	82 (99)	0 (12)
Dark-rumped petrel	Both	33 (3)	100 (3)	--
Newell's shearwater	Both	8 (36)	17 (36)	--

^aBoth = Juveniles plus adults.

Table 5.--Volume of plastic removed from the proventriculi of black-footed and Laysan albatross carcasses found dead of natural causes on Midway.

Year	Black-footed albatross			Laysan albatross		
	N	Range	Mean	N	Range	Mean
1986	25	0-198	39	45	1-186	46
1987	18	5-165	33	76	5-20	5

With the inclusion of the results from this study, 80 species, or approximately 25% of the world's seabird species, have been shown to ingest plastic. The pronounced differences in the prevalence of ingested plastic between different taxa of Hawaiian seabirds was expected. Seabird biologists have reported interspecific differences for seabird communities in the North Pacific and Southern Ocean and have attributed them primarily to differences in feeding behavior (Day et al. 1985; Ryan 1987a). Species feeding primarily at the ocean surface ingested more plastic, possibly because of increased exposure to it.

Geographic variations in prevalence in a given species are usually attributed to differences in the environmental availability of plastic, although no study has attempted to test this assumption (Day et al. 1985; Ryan 1988b). The mean density of plastic particles in Alaskan waters was 910 particles/km² (calculated from Day and Shaw 1987), in the South Atlantic it was 2,080 particles/km² (calculated from Morris 1980), and in the subtropical North Pacific (the Hawaiian Islands region) it was 96,100 particles/km² (Day and Shaw 1987). The corresponding percentages of

Table 6.--Mean volume (cc) of the largest plastic fragment^a removed from the proventriculi and ventriculi of individual Hawaiian seabirds during 1986 and 1987.

Species	Proventriculus		Ventriculus	
	N	Mean volume	N	Mean volume
Black-footed albatross	51	2.056	16	0.089
Laysan albatross	47	0.649	63	0.073
Petrels (Bonin, Bulwer's, dark-rumped)	4	0.041	20	0.014
Shearwaters (wedge-tailed, Newell's, Christmas)	54	0.117	6	0.006
Sooty storm-petrel	30	0.017	--	--
Red-tailed tropicbird	14	0.051	--	--
Boobies (masked, red-footed)	2	0.054	--	--
Great frigatebird	5	0.329	--	--
Sooty tern	1	0.002	--	--

^aFragments were flat pieces of manufactured plastic, 1-3 mm thick, typically from broken plastic bottles or other containers.

Table 7.--Prevalence (number of stomach samples containing the color/the total number of samples examined \times 100) of different colors of plastic fragments^a removed from the proventriculi of Hawaiian seabirds during 1986 and 1987. Stomach samples were obtained by stomach pumping.

Species	N	Plastic color (%) ^b				
		White	Black	Blue	Green	Red
Black-footed albatross	82	93	27	43	41	13
Laysan albatross	78	95	32	49	50	27
Wedge-tailed shearwater	51	84	8	24	29	8
Sooty storm-petrel	30	87	13	10	27	40
Red-tailed tropicbird	14	57	43	21	7	0
Great frigatebird	5	80	40	0	0	0

^aFragments were flat pieces of manufactured plastic, 1-3 mm thick, typically from broken bottles or other containers.

^bColor groups were: white = white, yellow, tan and brown; black = gray and black; blue = purple and blue; green = green; red = pink and red.

seabird species that ingested plastic in these three regions were 41 (Day et al. 1985), 60 (Ryan 1987a), and 89 (this study), suggesting a positive relationship between plastic availability and the prevalence of ingestion. The same relationship may explain the lower prevalence of plastic in adult wedge-tailed shearwaters at Johnston Atoll compared with the same species at Midway. The beaches of Johnston Atoll have much less plastic refuse than those of the Northwestern Hawaiian Islands (S. I. Fefer, unpubl. observ.), suggesting a lower density of plastic in the surrounding waters used for feeding. Much of the plastic in these waters may be of Japanese origin as indicated by Pettit et al. (1981), who found that 108 of 109 identifiable plastic items in dead albatross carcasses at Midway were manufactured in Japan. Movement of floating plastic by ocean currents provides an explanation for the high prevalence of Japanese plastic in Hawaiian waters and the reduced amount of plastic on Johnston Atoll. The Kuroshio moves surface waters from near Japan southeast to the Hawaiian region, and may carry plastic with it (Fry et al. 1987). Johnston Atoll is affected primarily by the North Equatorial Current, which probably has less intense shipping and fishery activity and hence less plastic.

Day et al. (1985) found that subadult parakeet auklets, *Cyclorhynchus psittacula*, and tufted puffins, *Lunda cirrhata*, contained more plastic than adults, and Ryan (1988b) found the same association for blue petrel, *Halobaena caerulea*, chicks. The high prevalence in albatross chicks, and probably chicks of other species in the Hawaiian Islands, was likely due to the regurgitational chick-feeding process and the inability of very young chicks to regurgitate indigestibles. All Hawaiian seabirds except white terns feed their chicks by regurgitation, and plastic items ingested by the parents are probably passed to the chicks. Ryan (1988b) proposed that this "intergenerational" transfer of plastic reduces occurrence in the adult populations while increasing it in the chick populations. Plastic is usually expelled from the proventriculus of albatross chicks by regurgitation late in the chick-rearing period (Clarke et al. 1981), and the volume of plastic in a given chick, and perhaps the prevalence in the chick population, are reduced at that time. Other seabirds (giant petrels, cormorants, skuas, gulls, and terns) also regurgitate indigestible matter. The presence of plastic in the intestines of some birds indicates that plastic is also removed by defecation. Interspecific differences in physiology, the ratios of proventricular to ventricular volume, or the total volume of plastic ingested may have influenced the interspecific differences in the distribution of plastic through the gastrointestinal tract.

It was not determined if the color composition of fragments from the stomach samples reflected the color distribution of floating fragments at sea or some feeding specificity by the birds. Day et al. (1985) found that some Alaskan seabirds selectively consume plastic particles similar to prey items.

Black-footed albatrosses ingest about 10 times the volume of fish eggs ingested by Laysan albatrosses (Harrison et al. 1983), and might be expected to ingest more plastic pellets if these are mistaken for fish eggs. However, we found pellets more often in Laysan (52%) than black-

footed albatrosses (12%, $P < 0.001$). Black-footed albatrosses probably do not ingest single fish eggs, but instead consume large masses of eggs attached to floating objects, which may explain the higher (34%) prevalence of fibers in the proventriculi of black-footed albatrosses than Laysans (11%, $P < 0.001$).

The size of plastic particles relative to common food items may be important in explaining why albatrosses ingest much larger plastic pieces than other Hawaiian seabirds. Black-footed and Laysan albatrosses both ingest larger size classes of squid and other food items than most Hawaiian seabirds (Harrison et al. 1983). It is also possible that black-footed albatrosses have more frequent contact with large plastic items due to their habit of feeding on refuse dumped overboard by ships (Miller 1940).

ACKNOWLEDGMENTS

Funding for this project was provided by the National Marine Fisheries Service, and the U.S. Fish and Wildlife Service. We thank the U.S. Navy for permission to work on Midway and for billeting. Special thanks to Captains Robins and West, Barbers Point Naval Air Station, and Lt. Commanders D. Stevenson, E. Moormann, and G. Walsh at Midway. Logistical assistance on Midway was provided by Base Services, Inc. Field assistance on Midway was provided by C. Baggot and refuge staff. Refuge personnel conducted much of the field work on the remaining Northwestern Hawaiian Islands and Johnston Atoll, while volunteers of the Kilauea Point Natural History Association assisted on Kauai. C. S. Harrison and P. G. Ryan reviewed the manuscript and provided many helpful suggestions.

REFERENCES

- Bayer, P. D., and R. E. Olson.
1988. Plastic particles in 3 Oregon fulmars. *Oregon Birds* 14:155-156.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Miklas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. *Science* 178:749-750.
- Clarke, M. R., J. P. Croxall, and P. A. Prince.
1981. Cephalopod remains in regurgitations of the wandering albatross *Diomedea exulans* L. at South Georgia. *Br. Antarct. Surv. Bull.* 54:9-21.
- Connors, P. G., and K. G. Smith.
1982. Oceanic plastic particle pollution: Suspected effect on fat deposition in red phalaropes. *Mar. Pollut. Bull.* 13:18-20.
- Day, R. H., and D. G. Shaw.
1987. Patterns in the abundance of pelagic plastic and tar in the North Pacific Ocean, 1976-1985. *Mar. Pollut. Bull.* 18:311-316.

- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Dickerman, R. W., and R. G. Goelet.
1987. Northern gannet starvation after swallowing Styrofoam. *Mar. Pollut. Bull.* 18:293.
- Freeman, D. H.
1987. *Applied categorical data analysis*. Marcel Dekker, Inc., N.Y., 318 p.
- Fry, D. M., S. I. Fefer, and L. Sileo.
1987. Ingestion of plastic debris by Laysan albatrosses and wedge-tailed shearwaters in the Hawaiian Islands. *Mar. Pollut. Bull.* 18:339-343.
- Furness, R. W.
1985. Ingestion of plastic particles by seabirds at Gough Island, South Atlantic Ocean. *Environ. Pollut. (Ser. A)* 38:261-272.
- Harper, P. C., and J. A. Fowler.
1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptila* spp.) 1958-1977. *Notornis* 34:65-70.
- Harrison, C. S., T. S. Hida, and M. P. Seki.
1983. Hawaiian seabird feeding ecology. *Wildl. Monogr.* 85, 75 p.
- Harrison, C. S., M. B. Naughton, and S. I. Fefer.
1984. The status and conservation of seabirds in the Hawaiian Archipelago and Johnston Atoll. In J. P. Croxall, P. G. H. Evans, and R. W. Schreiber (editors), *Status and conservation of the world's seabirds*, p. 513-526. ICBP Tech. Publ. No. 2, Cambridge, U.K.
- Iudicello, S. K., and J. O'Hara.
1986. Use and disposal of nonbiodegradable plastics in the marine and Great Lakes environments. Report to U.S. Environmental Protection Agency by the Center for Environmental Education, 129 p.
- Kenyon, K. W., and E. Kridler.
1969. Laysan albatross swallow indigestible matter. *Auk* 86:339-343.
- Miller, L.
1940. Some tagging experiments with black-footed albatrosses. *Condor* 44:3-9.
- Morris, R. J.
1980. Plastic debris in the surface waters of the South Atlantic. *Mar. Pollut. Bull.* 11:164-166.

- Pettit, T. N., G. S. Grant, and G. C. Whittow.
1981. Ingestion of plastics by Laysan albatross. *Auk* 98:839-841.
- Ryan, P. G.
1987a. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.
1987b. The effects of ingested plastic on seabirds: Correlations between plastic load and body condition. *Environ. Pollut.* 46:119-125.
1988a. Effects of ingested plastic on seabird feeding: Evidence from chickens. *Mar. Pollut. Bull.* 19:125-128.
1988b. Intraspecific variation in plastic ingestion by seabirds and the flux of plastic through seabird populations. *Condor* 90:446-452.
- Ryan, P. G., A. D. Connell, and B. D. Gardner.
1988. Plastic ingestion and PCBs in seabirds: Is there a relationship? *Mar. Pollut. Bull.* 19:174-176.
- Ryan, P. G., and S. Jackson.
1986. Stomach pumping: Is killing seabirds necessary? *Auk* 103:427-428.
1987. The lifespan of ingested plastic particles in seabirds and their effect on digestive efficiency. *Mar. Pollut. Bull.* 18:217-219.
- Sileo, L., and S. I. Fefer.
1987. Paint chip poisoning of Laysan albatross at Midway Atoll. *J. Wildl. Dis.* 23:432-437.
- Sileo, L., P. R. Sievert, and M. D. Samuel.
1990. Causes of mortality of albatross chicks at Midway Atoll. *J. Wildl. Dis.* 26:329-338.
- Van Franeker, J. A.
1985. Plastic ingestion in the North Atlantic fulmar. *Mar. Pollut. Bull.* 16:367-369.
- Van Franeker, J. A., and P. J. Bell.
1988. Plastic ingestion by petrels breeding in Antarctica. *Mar. Pollut. Bull.* 19:672-674.
- Wilson, R. P.
1984. An improved stomach pump for penguins and other seabirds. *J. Field Ornithol.* 55:109-112.
-

THE INCIDENCE OF PLASTIC IN THE DIETS OF ANTARCTIC SEABIRDS

David G. Ainley, William R. Fraser, and Larry B. Spear
Point Reyes Bird Observatory
Stinson Beach, California 94970, U.S.A.

ABSTRACT

We investigated the diets of seabirds at sea in the Antarctic from 1976 to 1988. During the study period, on eight cruises in the Ross, southern Scotia, and Weddell Seas and Drake Passage, we collected or pumped the stomachs of 1,223 seabirds of 23 species. The stomach contents of species that feed below the sea surface contained little plastic, as expected; these birds live entirely on live prey. Among species that feed at the surface, most of which eat both live and dead organisms, incidence of plastic was highest among the smaller ones and those that are omnivores, or feed on zooplankton and micronekton. This includes the majority of Southern Ocean flighted birds. Incidence of plastic among them was a function of the degree to which their populations frequented waters outside of the Antarctic during the winter. Among those species that live south of the Antarctic Convergence year-round there was little evidence of plastic ingestion. Among those species that are summer visitors to the Antarctic, incidence of plastic in the diet decreased with increased latitude. These results indicate either that the Antarctic Convergence blocks plastic debris, which is commonly found at the sea surface in the north, from entering the Southern Ocean, or that other factors such as the northward movement of pack ice sweeps the sea clear of plastic. Results also suggest that floating plastic debris is not yet the problem in the Antarctic that it is in more northern waters.

INTRODUCTION

Much has been learned recently about the ingestion of plastic by seabirds, mainly through the efforts of Day (1980; Day et al. 1985) in the northern Pacific, and of Ryan (1987a, 1987b, 1988a, 1988b, 1988c; Ryan and Jackson 1987) in the southern Atlantic. At present, 69 seabird species are known to ingest plastic while feeding at sea; 37 of these species frequent oceans of the Southern Hemisphere (see reviews in Day et al. 1985; Ryan 1987b). The incidence of ingested plastic in seabirds has been rising steadily since the 1960's, with the earliest records from procellariiform birds (Harper and Fowler 1987). The large majority of species now known to

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

eat and retain plastic in their alimentary tracts are members of the order Procellariiformes.

Geographic variation in the incidence of plastic ingestion among seabirds is a function of proximity to areas of industrialization and human population centers. Among the 100 or so species investigated to date and for which sample sizes are greater than 30, in the following, about 80% of individuals carry plastic loads (see summaries in Day et al. 1985; Ryan 1987b; Sileo et al. 1990): Laysan and black-footed albatross, *Diomedea immutabilis* and *D. nigripes*, short-tailed shearwater, *Puffinus tenuirostris*, and parakeet auklet, *Cyphorhynchus psittacula*, which frequent polluted waters of the North Pacific Rim; northern fulmar, *Fulmarus glacialis*, in the polluted northeastern Atlantic and the North Sea; greater shearwater, *Puffinus gravis*, which ranges between southern Africa and the polluted northwestern Atlantic; white-faced storm-petrel, *Pelagodroma marina*, and cape petrel, *Daption capense*, in waters off southern Africa; and blue petrels, *Halobaena caerulea*, in waters off southern Africa and the southwestern Pacific. High incidences of plastic also have been detected in seabirds of various species sampled off the U.S. west coast (Balz and Morejohn 1976).

In this paper we present information on the incidence of plastic in the diets of Antarctic seabirds, and compare our findings with the background of information just reviewed. The only samples collected previously in the Antarctic were reported by Ryan (1987b) and van Franeker and Bell (1988), and included small samples of five species.

METHODS

Birds were collected at sea off the Antarctic Peninsula and in the Ross Sea (Fig. 1) during investigations of their diets and marine ecology (Ainley et al. 1984, 1988, in press; Fraser and Ainley 1986). One sample of 60 Adélie penguins, *Pygoscelis adeliae*, was obtained by pumping stomachs at Palmer Station, Anvers Island (lat. 64°S, long. 64°W), in the Drake Passage; other penguin samples, except those from the Ross Sea, were obtained at sea also by stomach pumping. A sample of 75 castings was obtained from blue-eyed shags, *Phalacrocorax atriceps*, also at Palmer Station. All other birds were shot and contents of the proventriculus and ventriculus were obtained by dissection, collecting areas in the Ross Sea, December-January 1977-80, are summarized in Ainley et al. (1984); those in the southern Scotia Sea (Weddel Confluence region), November 1983 and August 1988, are summarized in Ainley and Sullivan (1984); those in the Weddell Sea, March 1986, are in Sullivan and Ainley (1988); and those in the Drake Passage and adjacent areas, July-August 1985-87, are in Pietz and Strong (1987).

Unlike Day (1980), Ryan (1987a, 1987b, 1988a, 1988b, 1988c; Ryan and Jackson 1987), and others (e.g., Furness 1985) who actually were attempting to directly characterize plastic ingestion in seabirds, we collected information incidental to other work. Therefore, except for the most recent sampling in August 1988, we did not quantify the number, size, and color of the particles found in digestive tracts.




Figure 1.--The areas and time periods in which seabird stomach samples were collected.

RESULTS

We observed no large accumulations of plastic in any of the birds inspected. The greatest number of pieces was 22 in a blue petrel, but usually it was much smaller (the next largest quantity was 8, also in a blue petrel). The birds that contained plastic in August 1988 averaged (\pm SD) the following numbers per bird; blue petrel, 5.1 \pm 5.4 (n = 14); cape petrel, 1 (n = 4); and snow petrel, *Pagodroma nivea*, 1 (n = 4). As observed in other studies, most of the plastic consisted of small (3-6 mm diameter) fragments and "pellets." Styrofoam occurred rarely (1 of 74 pieces of plastic in 1988). One Antarctic petrel, *Thalassoica antarctica*, had eaten a piece of rubber from a (meteorological?) balloon; and a snow petrel had eaten threads of polypropylene rope.

No plastic was observed in any species that feed by diving beneath the sea surface (penguins, *Aptenodytes* and *Pygoscelis* spp.; the shag; and the diving petrel, *Pelecanoides urinatrix*), all of which feed on live micronekton (Croxall 1987, and papers therein; Table 1). Neither did we find plastic in any species that lack a well-developed gizzard or regularly regurgitate castings of indigestible material (albatrosses, *Diomedea* spp.; shags; skuas, *Catharacta* spp.; gulls, *Larus* spp.; and terns, *Sterna* spp.). Among those birds whose stomachs or gizzards did contain plastic (petrels, Procellariidae, only), incidence was higher in the smaller species. Such patterns are consistent with previously published information (Day et al. 1985; Ryan 1987b; but see Sileo et al. 1990, for the exception presented by the large albatrosses); these species tend to eat zooplankton or micronekton, and also scavenge dead organisms.

The frequency of occurrence of plastic decreased with increasing latitude in each of the four areas sampled, except to some extent in the Ross Sea (Table 2). The same pattern also emerged for only those species having a relatively high frequency of plastic ingestion. For example, in the Scotia Sea, the frequencies of plastic in blue petrels, cape petrels, Antarctic prions, and Wilson's storm-petrels combined, were 17, 6, 0, and 0% at lat. 56°-58°S (n = 60), 59°S (n = 51), 60°S (n = 50), and 61°-62°S (n = 17), respectively.

A pattern seemingly inconsistent with the latter finding is evident in a comparison between frequencies of occurrence of plastic in Scotia Sea and Weddell Sea samples (Table 1). Within a species, the frequency of plastic is much higher in birds from the more southerly Weddell Sea than in birds from the Scotia Sea. The Weddell Sea sample, which was from autumn, however, contained a much higher proportion of subadult nonbreeders, which would be expected to contain larger plastic loads (Ryan 1988c). Indeed the frequency of plastic in blue petrels in this sample was 88%, and in winter samples it was 90%, which were rates similar to those reported for this species off the coast of Africa. In spring, when there were many more adults in the sample, frequency was 21%.

Table 1.--The frequency of occurrence of plastic in the digestive tracts of seabirds collected in various sectors of the Southern Ocean.^a

Species	Ross Sea	Drake Passage	Scotia Sea	Weddell Sea	Total
<i>Aptenodytes forsteri</i>			0 (8)	0 (17)	0 (25)
<i>Pygoscelis adeliae</i>	0 (5)	0 (60)	0 (29)	0 (10)	0 (104)
<i>Pygoscelis papua</i>		0 (5)			0 (5)
<i>Phalacrocorax atriceps</i>		0 (1) ^b			0 (1)
<i>Diomedea palpebrata</i> ^c	0 (2)				0 (2)
<i>Macronectes giganteus</i> ^c	0 (2)	0 (5)	0 (4)	0 (2)	0 (13)
<i>Procellaria aequinoctialis</i> ^c			0 (10)		0 (10)
<i>Fulmarus glacialis</i> ^c	0 (13)	0 (4)	2 (49)	6 (18)	2 (84)
<i>Thalassoica antarctica</i> ^c	2 (40)	0 (25)	0 (66)	0 (53)	<1 (184)
<i>Pagodroma nivea</i> ^c	0 (108)	0 (77)	3 (139)	0 (39)	1 (363)
<i>Daption capense</i> ^c	50 (4)	5 (20)	11 (63)	31 (16)	14 (105)
<i>Pterodroma inexpectata</i> ^c	0 (4)				0 (4)
<i>Pterodroma brevirostris</i> ^c			0 (5)	9 (23)	7 (28)
<i>Halobaena caerulea</i> ^c			44 (45)	88 (17)	56 (62)
<i>Pachyptila vittata</i> ^c	67 (3)		4 (51)	20 (15)	10 (69)
<i>Pelecanoides urinatrix</i> ^c			0 (4)		0 (4)
<i>Oceanites oceanicus</i> ^c	37 (27)		4 (49)	33 (15)	19 (91)
<i>Fregetta tropica</i> ^c			0 (6)		0 (6)
<i>Catharacta maccormicki</i>	0 (25)				0 (25)
<i>Larus dominicanus</i>		0 (15)			0 (15)
<i>Sterna paradisaea</i>			0 (10)	0 (14)	0 (24)
<i>Sterna vittata</i>	0 (12)			0 (5)	0 (17)
<i>Chionis alba</i>	0 (2)				0 (2)

^aPercentage with sample sizes in parenthesis.

^bSamples of 30 pellets in 1977 and 45 in 1987 contained no plastic.

^cBirds of the order Procellariiformes; see Table 2.

DISCUSSION

Plastic was found in 36% of the 23 species examined, a percentage much lower than that of most other regional studies: 100% of 10 species in California (Balz and Morejohn 1976), 76% of 15 at Gough Island (Furness 1985), 71% of 14 in New Zealand (Imber in Day et al. 1985), 56% of 95 in a world survey (Day et al. 1985), 60% of 60 from the Southern Hemisphere (Ryan 1987b), and 98% of 22 in Hawaii (Sileo et al. 1990). In another study of seabirds in the equatorial Pacific, Ainley et al. (1990) found plastic in 59% of petrel species. Thus, an unusually low incidence of contamination is evident in the Antarctic as compared to other areas of the world ocean (see also van Franeker and Bell 1988).

Table 2.--Frequency of occurrence of plastic in procellariiform birds, by latitude and sampling sector in the Southern Ocean.^a

Latitude N	Ross Sea	Drake Passage	Scotia Sea	Weddell Sea
56°-58°			15 (183)	
59°			7 (119) ^b	
60°			1 (93)	
61°-62°		3 (37)	0 (97)	
63°-64°		0 (60) ^b		25 (60)
65°-66°		0 (34)		9 (137)
67°-69°	11 (35)			(b)
70°-75°	18 (63) ^b			
76°-78°	0 (46)			

^aPercentages, with sample size in parenthesis, are given; see Table 1 for procellariiform species.

^bApproximate northern edge of the pack ice.

The greatest frequency of occurrence found by us was in blue petrels, but the 56% rate was much lower than the 90% reported for this species by Ryan (1987b). A relatively high incidence was also evident in our Antarctic samples of the cape petrel (14%), Antarctic prion, *Pachyptila desolata* (10%), Kerguelen petrel, *Pterodroma brevirostris* (7%), and Wilson's storm-petrel, *Oceanites oceanicus* (19%). For these species, however, rates were much lower than those reported by Ryan (1987b), van Franeker and Bell (1988), Ainley et al. (1990), and Sileo et al. (1990) for other areas. Harper and Fowler (1987) reported a rate equivalent to ours for Antarctic prions found dead on beaches in New Zealand. In the present study, frequencies of plastic occurrence for the Antarctic fulmar, *Fulmarus glacialis*, white-chinned petrel, *Procellaria aequinoctialis*, and Antarctic petrel were negligible, as also noted by van Franeker and Bell (1988) for the Indian Ocean sector of Antarctica. Ryan (1987b) found a similarly low rate for this sample of Antarctic petrels from the Antarctic, but higher rates of 11% in fulmars found dead on beaches in southern Africa and of 57% in white-chinned petrels from southern African waters. Ainley et al. (1990) found a rate of 67% for the latter species in the Peru Current. Not the present, nor Ryan's, nor van Franeker and Bell's studies detected much plastic in snow petrels, which are restricted to the Antarctic. Thus, a low incidence of plastic contamination is again indicated for Antarctic waters.

Our findings also indicate that the greater the distance south from the Antarctic Convergence the less likely birds are to have eaten plastic. More southerly individuals either have weaker ties to waters outside the Antarctic (where densities of plastic are greater, Morris 1980; Pruter 1987), or they have not frequented northern waters recently. If bird

residency time in pollution-free waters is a factor, such a pattern might also support the idea that over time there is a gradual attrition of plastic from digestive tracts (Day et al. 1985; Ryan and Jackson 1987; Ryan 1988a; van Franeker and Bell 1988; Ainley et al. 1990). Again, however, the pattern indicates that the density of plastic is very low in Antarctic waters. In other words, once south of the Antarctic Convergence, northern seabirds lose plastic from their digestive tracts faster than they gain it.

Both the patterns described above and other patterns indicate that at present there is little plastic floating on the surface of ocean waters south of the Antarctic Convergence. Antarctic surface waters flow northward away from the continent, and then sink beneath subantarctic waters at the Antarctic Convergence (Deacon 1964). Where the Antarctic Convergence is particularly well developed, flotsam (e.g., kelp fragments) is much in evidence along its northern edge (D. G. Ainley, pers. observ.; S. S. Jacobs, Lamont Doherty Geological Observatory, pers. commun.). Thus, the convergence may act to some degree as a barrier to flowing debris and pollutants from the north (though not an absolute barrier, because eddies are able to transport some northern waters across the convergence; Jacobs, pers. commun.). Because there is little human activity in the Antarctic, there is at present a relatively low rate of disposal of plastics. This helps to maintain the apparent low densities of floating plastic there, but few efforts to directly sample the abundance of floating plastic in Antarctic waters have been reported (Gregory et al. 1984; Pruter 1987). One additional factor that may help to sweep Antarctic waters clear of plastic is the seasonal, northward advance of the pack ice (Jacobs, pers. commun.). In that each of our samples was collected with respect to the pack ice edge (because we were comparing the diets of birds in and out of the ice), and the decreasing plastic loads we detected were not a function of absolute latitude south of the convergence (i.e., position of the pack ice edge differed for each sample), our results indicate that northward movement of ice would indeed help to clear any plastic from the sea surface in the Antarctic. In fact, a great deal of organic detritus (e.g., dead diatoms) is scoured from the water column by the freezing process, transport north, and released by melting at the ice pack edge, where a large amount of detrital material can be found (C. W. Sullivan, Department of Biological Sciences, University of Southern California, pers. commun.).

Assuming that most of the plastic found in the birds inspected in this study was ingested near to or north of the Antarctic Convergence, the frequency of occurrence of plastic in seabirds provides an index of how strongly certain petrel species are tied to Antarctic seas. Considering the dearth of ecological studies in Antarctic waters during the spring and winter, such an index is useful in characterizing the avifauna. Those seabird species with low frequencies of occurrence of plastic should exhibit the weakest tendencies to exit the Antarctic during the winter. These data, therefore, suggest that Antarctic and snow petrels, and a significant proportion of Antarctic fulmars, do not leave the Antarctic during winter, as do so many other "Antarctic" seabirds. In fact, the only plastic found in snow petrels was in individuals that had moved north with the pack ice during winter, and were thus close to the Antarctic Convergence. Along these same lines of reasoning, one might expect a large

amount of plastic to accumulate along the Antarctic Convergence, and the higher incidence of plastic in certain species, especially the blue and Kerguelen petrels, may indicate that these species frequent the convergence area more than most other species or that in general they frequent water mass convergences (cf. Bourne and Clark 1984). Census results support these distributional patterns (Ainley and Fraser, unpubl. data).

ACKNOWLEDGMENTS

Logistical support was provided through the U.S. Antarctic Program. We thank support personnel of Holmes and Narver, Inc., ITT-Antarctic Services, Inc., and the officers and crew of U.S. Coast Guard cutters *Burton Island*, *Glacier*, and *Northwind*, and of the RV *Melville* and MV *Polar Duke*. Assistance in the field was provided by G. J. Divoky, R. Ferris, E. F. O'Connor, P. Pietz, R. Pitman, C. Strong, and G. Wallace. R. H. Day, S. S. Jacobs, and P. G. Ryan provided many helpful comments on the manuscript. Our work was financed by grants from the U.S. National Science Foundation, mainly grants DPP 8304815 and 8419894 in support of Antarctic Marine Ecosystem Research in the Ice Edge Zone (AMERIEZ). This is Publication No. 393 from the Point Reyes Bird Observatory.

REFERENCES

- Ainley, D. G., W. R. Fraser, and K. A. Daly.
1988. Effects of pack ice on the composition of micronektonic communities in the Antarctic. In D. Sahrhage (editor), *Southern Ocean resources variability*, p. 140-146. Springer Verlag, Heidelberg, Berlin.
- Ainley, D. G., W. R. Fraser, W. O. Smith, T. L. Hopkins, and J. J. Torres.
In press. The structure of upper level pelagic food webs in the Antarctic: Effect of phytoplankton distribution. *J. Mar. Systems*.
- Ainley, D. G., E. F. O'Connor, and R. J. Boekelheide.
1984. The marine ecology of seabirds in the Ross Sea, Antarctica. *Ornithol. Monogr. No. 32*. Am. Ornithol. Union, Wash., D.C.
- Ainley, D. G., L. B. Spear, and C. A. Ribic.
1990. The incidence of plastic in the diets of pelagic seabirds in the eastern equatorial Pacific region. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Ainley, D. G., and C. W. Sullivan.
1984. AMERIEZ 1983: A summary of activities on board R/V *Melville* and USCGC *Westwind*. *Antarct. J. U. S.* 19(5):100-103.
- Balz, D. M., and G. V. Morejohn.
1976. Evidence from seabirds of plastic particle pollution off central California. *West. Birds* 7:111-112.

- Bourne, W. R. P., and G. C. Clark.
1984. The occurrence of birds and garbage at the Humboldt Front off Valparaiso, Chile. *Mar. Pollut. Bull.* 15:343-344.
- Croxall, J. P. (editor).
1987. *Seabirds: Feeding ecology and role in marine ecosystems.* Cambridge Univ. Press, Cambridge, 408 p.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Deacon, G. E. R.
1964. A discussion of the physical and biological changes across the Antarctic Convergence. Introduction. *Proc. Royal Soc. (Ser. A)* 281:1-6.
- Fraser, W. R., and D. G. Ainley.
1986. Ice edges and seabird occurrence in Antarctica. *BioScience* 36:258-263.
- Furness, R. W.
1985. Ingestion of plastic particles by seabirds at Gough Island, South Atlantic Ocean. *Environ. Pollut. (Ser. A)* 38:261-272.
- Gregory, M. R., R. M. Kirk, and M. C. G. Mabin.
1984. Pelagic tar, oil, plastics and other litter in the surface waters of the New Zealand sector of the Southern Ocean, and on Ross Dependency shores. *N.Z. Antarct. Res.* 6:12-28.
- Harper, P. C., and J. A. Fowler.
1987. Plastic pellets in New Zealand storm-killed prions (*Pachyptilla* spp.). *Notornis* 34:65-70.
- Harrison, C. S., T. S. Hida, and M. P. Seki.
1983. Hawaiian seabird feeding ecology. *Wildl. Monogr.* 85:1-71.
- Morris, R. J.
1980. Plastic debris in the surface waters of the South Atlantic. *Mar. Pollut. Bull.* 11:164-166.
- Pietz, P. J., and C. S. Strong.
1987. Ornithological observations west of Antarctic Peninsula. *Antarct. J. U. S.* 21(5):203-204.
-

- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.
- Ryan, P. G.
1987a. The effects of ingested plastic on seabirds: Correlations between plastic load and body condition. *Environ. Pollut.* 46: 119-125.
1987b. The incidence and characteristics of plastic particles ingested by seabirds. *Mar. Environ. Res.* 23:175-206.
1988a. Effects of ingested plastic on seabird feeding: Evidence from chickens. *Mar. Pollut. Bull.* 19:125-128.
1988b. The characteristics and distribution of plastic particles at the sea-surface off the southwestern Cape Province, South Africa. *Mar. Environ. Res.* 25:249-273.
1988c. Intraspecific variation in plastic ingestion by seabirds and the flux of plastic through seabird populations. *Condor* 90:446-452.
- Ryan, P. G., and S. Jackson.
1987. The lifespan of ingested plastic particles in seabirds and their effect on digestive efficiency. *Mar. Pollut. Bull.* 18: 217-219.
- Sileo, L., P. R. Sievert, M. D. Samuel, and S. I. Fefer.
1990. Prevalence and characteristics of plastic ingested by Hawaiian seabirds. In R. S. Shomura and M. L. Godfrey (editors), *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- Sullivan, C. W., and D. G. Ainley.
1988. AMERIEZ 1986: A summary of activities on board R/V *Melville* and USCGC *Glacier*. *Antarct. J. U. S.* 22(5):167-169.
- Van Franeker, J. A., and P. J. Bell.
1988. Plastic ingestion by petrels breeding in Antarctica. *Mar. Pollut. Bull.* 19:672-674.

PLASTIC DEBRIS INCORPORATED INTO DOUBLE-CRESTED
CORMORANT NESTS IN THE GULF OF MAINE

Richard H. Podolsky
Research and Academics
Island Institute
Rockland, Maine 04841, U.S.A.

and

Stephen W. Kress
Ornithological Research Unit
National Audubon Society Research Department
Ithaca, New York 14850, U.S.A.

ABSTRACT

The incorporation of plastic debris into double-crested cormorant, *Phalacrocorax auritus*, nests is reported on three islands in the Gulf of Maine. Of the 497 nests examined during 1987 and 1988, 188 nests (37%) contained plastic debris. Sections of lobster trap line, plastic bags, and pieces of fishing net dominated this debris. The significance of this is discussed and future monitoring of plastics in seabird nests is recommended.

INGESTION OF PLASTICS BY TELEOST FISHES

Donald E. Hoss and Lawrence R. Settle
Southeast Fisheries Science Center Beaufort Laboratory
National Marine Fisheries Service, NOAA
Beaufort, North Carolina 28516-9722, U.S.A.

ABSTRACT

Ingestion of plastic debris by many types of animals such as turtles and seabirds is well documented and considered to be a serious threat to their survival. Marine fishes also ingest plastic debris but the amount ingested and the effect of the ingested debris are not well documented. If large amounts of inert plastic debris were ingested, it might affect the fishes' well-being by blocking the digestive tract and reducing the feeding drive. Also, certain types of debris could cause injury to the digestive tract and, depending on its chemical composition, might even have a toxic effect.

In this paper we review the literature to determine what is known about ingestion of plastics by marine fishes and report on our studies on ingestion of plastic particles by larvae and juveniles. There is at present no comprehensive list of fishes known to have ingested plastic. However, observations made incidental to other studies indicate that many species do at least occasionally ingest plastic. Plastics have been found in larvae, juveniles, and adults of both pelagic and demersal species. Currently, there is no clear evidence that juvenile and adult fish have been affected by ingesting plastic. Studies in the field on larval fish have suggested that swallowed plastic spheres could cause intestinal blockage and that polychlorinated biphenyls associated with the surface of the spherules could have toxic effects.

Laboratory experiments to determine the effects of plastic ingestion on larval and juvenile fish have been equivocal. In some cases the fish were observed to take particles, but then reject them.

We have found in our laboratory studies on larvae that five of six species tested--Atlantic menhaden, *Brevoortia tyrannus*, pinfish, *Lagodon rhomboides*, spot, *Leiostomus xanthurus*, striped mullet, *Mugil cephalus*, and two species of flounder, *Paralichthys* spp.--will feed on polystyrene microspheres. However, only spot and mullet were found to have particles in their gut. Particles passed from the gut after a period of time and larvae subsequently fed on brine shrimp larvae.

INTRODUCTION

Plastic debris is a common contaminant of marine waters and is potentially available for ingestion by marine life. Since the report of Carpenter and Smith (1972) on contamination of the Sargasso Sea surface by plastic particles, numerous surveys have reported on finding various types of plastics in waters from around the world (Carpenter et al. 1972; Kartar et al. 1973; Venrick et al. 1973; Colton et al. 1974; Hays and Cormons 1974; Morris and Hamilton 1974; Wong et al. 1974; Gregory 1977; Shaw 1977; Shaw and Mapes 1979; Shiber 1979, 1987; Morris 1980; Dahlberg and Day 1985; Day et al. 1986; Ignell and Dahlberg 1986). A more extensive discussion of the worldwide distribution of plastics in the sea is given by Pruter (1987).

Ingestion of plastic debris by many types of animals (e.g., marine turtles and birds) is, in fact, well documented and in many cases considered to be a serious hazard (Balazs 1985; Day et al. 1985; Azzarello and Van Vleet 1987; Fry et al. 1987; Gramentz 1988). For marine fishes, the ingestion of plastic debris and its subsequent effect is not well documented, but it is assumed that they, like other marine animals, will be unable to distinguish between normal prey and small pieces of plastics. Fish may swallow pieces mistaken for prey or ingest pieces incidental to normal feeding. Once ingested, this debris may block the digestive tract, lessen feeding, and cause ulceration or other physical injury to the stomach lining. It has been suggested that ingested plastics may also release toxic chemicals (Day et al. 1985). Animals weakened by the adverse effects of ingestion may then be more susceptible to disease and predators (Laist 1987).

The objectives of this paper are twofold:

1. to review what is known about the ingestion of plastics by marine fishes, and
2. to present recent field and laboratory data on plastic ingestion in larval and juvenile fishes.

REVIEW OF INGESTION

Larvae and Juveniles

The best documentation for ingestion of plastic by marine fishes is, somewhat surprisingly, for larval and juvenile stages. Carpenter et al. (1972) were the first to report larval fishes feeding on plastic. They reported that of 14 species of fishes collected by oblique plankton tows, 8 species contained plastic in their guts (Table 1). These authors found bacteria and polychlorinated biphenyls (PCB's) present on surfaces of the plastic particles. They speculate that a main effect of ingesting the particles may be intestinal blockage in some of the smaller fish.

Kartar et al. (1973), working in the Severn Estuary, United Kingdom, in 1972-73, found as many as 30 polystyrene particles in the stomachs of

Table 1.--Larval and juvenile fishes collected in the field with plastics in their gut.

Species	Mean size (mm)	Source
Clupeidae		
<i>Brevoortia patronus</i> , gulf menhaden	7.6	Govoni (pers. commun.)
<i>Clupea harengus</i> , Atlantic herring	42	Carpenter et al. 1972
Gadidae		
<i>Ciliata mustela</i> , five-beard rockling	--	Kartar et al. 1976
<i>Pollachius virens</i> , pollock	30	Carpenter et al. 1972
Atherinidae		
<i>Menidia menidia</i> , Atlantic silverside	16	Carpenter et al. 1972
Sciaenidae		
<i>Micropogonias undulatus</i> , Atlantic croaker	6.3	Govoni (pers. commun.)
Labridae		
<i>Tautoglabrus adspersus</i> , tautog	91	Carpenter et al. 1972
Gobiidae		
<i>Govius minutus</i> , sand goby	--	Kartar et al. 1976
Cottidae		
<i>Myoxocephalus aenus</i> , grubby	5.8	Carpenter et al. 1972
Cyclopteridae		
<i>Liparis liparis</i> , striped seasnail	--	Kartar et al. 1976
Pleuronectidae		
<i>Platichthys flesus</i> , flounder	20-50	Kartar et al. 1973
<i>Pseudopleuronectes americanus</i> , winter flounder	4.6	Carpenter et al. 1972

0+ and 1+ year class flounder, *Platichthys flesus* (Table 1). In more recent work in the same estuary, Kartar et al. (1976) found only a few particles in the sediment and none in four common species of fish which previously contained plastics. Flounder contained particles, but the numbers found per fish had declined between 1973 and 1975. They conclude that this type of plastic pollution has almost ceased in this particular estuary.

The gut contents of over 3,000 larval gulf menhaden, *Brevoortia patronus*, spot, *Leiostomus xanthurus*, and Atlantic croaker, *Micropogonias undulatus*, from the northern Gulf of Mexico were examined at the Beaufort

Laboratory, National Marine Fisheries Service (NMFS), between 1979 and 1982. Inert material, some of which was plastic, was found in only 20 of the fish (Govoni pers. commun.). Although this research was not designed to look specifically for plastic, it is certain that particles would have been observed had they been present in the gut in amounts found by Carpenter et al. (1972) and Kartar et al. (1973).

Colton et al. (1974) examined over 500 larvae from 22 species collected in water containing high concentrations of plastic spheres and found no plastic particles in the gut contents. They followed up their field work with laboratory experiments to determine if fish held in tanks would feed on these plastic particles and, if so, to measure any resulting effects of ingestion. Five species were tested over a 2-week period (Table 2). Samples were taken at regular intervals to determine if they had fed on the plastic particles. No particles were found in the guts of juveniles or larvae. Tomcod, *Microgadus tomcod*, and striped killifish, *Fundulus majalis*, juveniles were observed to feed on the particles, but they either rejected the particle or it passed through the gut with no harmful effect. These authors concluded that at present levels of abundance, the ingestion of plastics by larvae and juveniles would be minor compared to other pollution problems.

In the laboratory, Hjelmeland et al. (1988) demonstrated that larval Atlantic herring, *Clupea harengus*, would ingest polystyrene spheres (Table 2). The spheres, which had no nutritional value and were not degradable by digestive enzymes, nevertheless induced digestive secretion. However, the response was significantly lower than that obtained when the larvae were fed living prey.

Adults

To our knowledge, there has been no study specifically directed at ingestion of plastics or the effects of ingestion of plastics on adult fish. Most available information has been collected incidental to other studies. This is in spite of the fact that ingestion of plastics is continually cited as a potential hazard to fish (Laist 1986; U.S. Congress 1986).

There are several feeding studies that report finding plastics in the guts of fish incidental to the main objective of the study. A series of papers by Manooch (1973) and various coauthors (Manooch and Hogarth 1983; Manooch and Mason 1983; Manooch et al. 1984, 1985) are a good example. These authors found plastics of various types in five species of pelagic fishes and one anadromous fish (Table 3).

It is assumed that these plastic items were eaten accidentally or that they were mistaken for natural prey. Tuna, *Thunnus* spp., and dolphin, *Coryphaena hippurus*, seem to have the most diverse collection of plastics in their guts (Fig. 1), and this is probably due to both their feeding habits and their association with drift lines where plastic and other debris are known to collect (Manooch and Mason 1983; Manooch et al. 1984). These authors suggested that gut contents of dolphins could serve as indicators of surface water quality.

Table 2.--Results of laboratory experiment using plastic microspheres.

Species	Life stage	Results
Clupeidae <i>Clupea harengus</i> , Atlantic herring	Larval	Ingested pellets. ^a
Gadidae <i>Melanogrammus aeglefinus</i> , haddock	Larval	Ingestion negative no plastic in gut. ^b
<i>Microgadus tomcod</i> , tomcod	Juvenile	Ingested plastic but rejected or passed it. ^b
Cyprinodontidae <i>Fundulus majalis</i> , striped killifish	Juvenile	Ingested plastic but rejected or passed it. ^b
Gasterosteidae <i>Gasterosteus aculeatus</i> , threespine stickleback	Juvenile	Ingestion negative no plastic in gut. ^b
Pleuronectidae <i>Pseudopleuronectes americanus</i> , winter flounder	Larval and juvenile	Ingestion negative no plastic in gut. ^b

^aHjelmeland et al. 1988.

^bColton et al. 1974.

There is some observational evidence (Manooch pers. commun.) that plastics may remain in the guts of fish for long periods of time and be encysted in the stomach or gut lining. The long-term effect of this is not known but could hardly be beneficial to the fish.

Plastic cups were reported from the stomachs of cod, *Gadus morhua*, whiting, *Micromesistius poutassou*, and pollock, *Pollachius virens*, off the coast of the United Kingdom (Anonymous 1975). One pollock was found to contain four cups. Apparently the source of the cups was from the cross-channel ferries. The author concludes that the fish will eventually die since the cups are indigestible, but no evidence is presented for this statement.

CURRENT NATIONAL MARINE FISHERIES SERVICE RESEARCH

Previous studies have shown a high degree of patchiness in plastic distribution in the sea. This patchiness is attributable to currents, winds, and differential inputs (Shaw and Mapes 1979). In recent years scientists have focused increasingly on oceanographic fronts for numerous

Table 3.--Plastic found in adult marine fishes.

Species	Type of plastic	Source
Gadidae		
<i>Gadus morhua</i> , Atlantic cod	Plastic cups	Anonymous 1975.
<i>Micromesistius poutassou</i> , blue (pout) whiting	Plastic cups	Anonymous 1975.
<i>Pollachius virens</i> , pollock	Plastic cups	Anonymous 1975.
Percichthyidae		
<i>Morone americana</i> , white perch	Plastic pellets	Carpenter et al. 1972.
<i>Morone saxatilis</i> , striped bass	Plastic cigar holder	Manooch 1973.
Coryphaenidae		
<i>Coryphaena hippurus</i> , dolphin	Nylon rope, bottle, packaging, colored fragments	Manooch et al. 1984.
Scombridae		
<i>Acanthocybium solanderi</i> , wahoo	Fragment of black plastic sheeting	Manooch and Hogarth 1983.
<i>Euthynnus alletteratus</i> , little tunny	Packaging	Manooch et al. 1985.
<i>Thunnus albacares</i> , blackfin tuna	Plastic bag, colored fragments	Manooch and Mason 1983.
<i>Thunnus atlanticus</i> , yellowfin tuna	Colored fragments	Manooch and Mason 1983.
Triglidae		
<i>Prionotus evolans</i> , striped searobin	Plastic pellets	Carpenter et al. 1972.

reasons; among them are the observations that fishes (as well as sea turtles, marine mammals, and seabirds) are often aggregated about these zones along with the flotsam and other debris.

Both adult and larval fishes, including species of economic importance, have been observed in aggregations along frontal zones, but there has been little work describing the possible effects of associated debris.

1977; Shiber 1979; Day 1980), their distribution and abundance in the Gulf of Mexico is not well documented. We examined samples from three sites in the northern Gulf of Mexico (Cape San Blas, Florida, the plume of the Mississippi River, and Galveston, Texas) collected on a cruise in 1981. At each of these sites, sample tows were made with a multiple opening and closing net and environmental sensing system (MOCNESS) (Wiebe et al. 1976) at the surface, mid-depth, and bottom of the water column. Water samples from these stations were examined for the presence of small plastic particles such as those found by Carpenter et al. (1972) and Colton et al. (1974) (Fig. 2).

Of the 51 samples examined from the December collection, 27 were from the surface and the remaining 24 were from the middle of the water column. The greatest number of particles were found in the upper 7 m of the water column in the vicinity of Southwest Pass (Tester et al. 1987) (Table 4). This may be a reflection of both the high utilization of this area by shipping and industry and the outflow of the Mississippi River.



Figure 2.--Plastic material removed from samples collected at three sites in the northern Gulf of Mexico. Scale at bottom in millimeter.

Feeding Experiments

During 1988 and 1989, we conducted a series of feeding experiments (Settle et al. in prep.) to determine 1) if early life stages of marine fishes would ingest plastic particles in the laboratory, and 2) what effects ingestion might have. A similar, but inconclusive, investigation was attempted by Colton et al. (1974). We used polystyrene microspheres sorted to appropriate food particle size (100-500 μm). All plastic particles were "aged" in algae-rich seawater for at least 2 weeks. Six species of fish were used: Atlantic menhaden, *Brevoortia tyrannus*, pinfish, *Lagodon rhomboides*, spot, *Leiostomus xanthurus*, striped mullet, *Mugil cephalus*, southern flounder, *Paralichthys lethostigma*, and flounder, *Paralichthys* spp. Menhaden were laboratory spawned; all others were collected from the Newport River estuary, North Carolina. Fish were maintained in 5-L tanks and starved for 48 h prior to the introduction of plastic particles. Particle concentrations ranged from 200 to 1,000 L^{-1} .

All species except *Paralichthys* spp. were observed ingesting plastics, but rejection was also commonly observed (Table 5). Experiments lasted from 10 min to 19 h. At the end of the experimental period, fish were killed and their guts examined. Four of the six species had plastic particles in their alimentary tract. Thus, even though some plastics were rejected, some were fully ingested as well. Mullet and spot ingested the greatest quantity of particles, with some containing over 30 particles (maximum 45) (Fig. 3).

These results showed conclusively that these species would ingest aged plastic particles when deprived of food for 48 h, and in some cases retain particles in the gut for several hours.

Based on these results, a second series of experiments were conducted on mullet (21-25 mm SL) and spot (16-23 mm) to investigate if plastic ingestion would cause mortality. As in the previous work, the fish were starved for 2 days prior to the start of the experiment. The fish were initially fed plastic spheres (1,000 L^{-1}), with brine shrimp, *Artemia* spp., added after 10 min. These experiments were conducted for 10 days during which brine shrimp were added on a daily basis. Plastic spheres were left in the tank throughout the experiment.

Both spot and mullet were observed to ingest plastic particles when they were first added. They also were observed to reject some of the particles. Spot took plastic from the water column and off the bottom while mullet fed only from the water column. When brine shrimp were present, both species appeared to select them over the plastic and usually rejected plastic if ingested. There was no experimental mortality observed during the 10-day period and the fish were observed defecating. Therefore, it does not appear that the plastic blocked the gut.

At the end of the experiment the fish were sacrificed, measured, and examined for plastic in their guts. Six of twenty-four spot contained plastic. It is likely that spot ingested particles throughout the experiment, either those resuspended in the water each day or those on the

Table 4.--Small plastic particles in the Gulf of Mexico. Samples were taken from the surface to near bottom. Stations A1, B1, and D1 were only in 18.3 m (10 fathoms) of water, and A2, B2, and D1 were in 91.4 m (50 fathoms) of water. Plastics were collected only at the depths indicated.

Region	Station	Sample depth (m)	Particles per 100 m ³
Mississippi River	A1	1	26
		2	67
		5	31
		5	19
		6	5
		7	60
	A2	1	2
		2	1
Cape San Blas, Florida	B1	1	5
		3	1
		8	2
	B2	1	1
		30	1
		31	2
Galveston, Texas	D1	1	4
		5	9
	D2	1	1

bottom. Particles were well distributed throughout the alimentary tract, giving the impression that they were being effectively passed (Fig. 4). None of 20 mullet contained pellets at the end of the 10 days although they were observed to feed on them during the course of the experiment.

DISCUSSION AND CONCLUSIONS

There is now ample evidence to state that marine fish of many species will eat plastic debris. Larval and juvenile fishes have been collected in the field with plastic fragments and raw plastic pellets in their guts. Adult fishes have been found with a wide variety of material in their guts ranging from unidentified fragments to whole cups and bottles. There is almost no evidence, however, to determine the magnitude of the problem or to determine if ingestion is an important cause of mortality in fish.

Table 5.--Results of aged polystyrene microsphere feeding experiments (Settle et al. in prep). (+ indicates plastics were ingested, - indicates plastics were not ingested.)

Species	Size range (mm)	Particle size (m)	Ingestion	Percent with plastic in gut
Clupeidae				
<i>Brevoortia tyrannus</i> , Atlantic menhaden	9-29	100-500	+	0
Sparidae				
<i>Lagodon rhomboides</i> , pinfish	11-14	350-500	+	15
Sciaenidae				
<i>Leiostomus xanthurus</i> , spot	19-25	350-500	+	15
Mugilidae				
<i>Mugil cephalus</i> , striped mullet	18-25	210-350	+	75
Bothidae				
<i>Paralichthys lethostigma</i> , southern flounder	13-15	210-250	+	6
<i>Paralichthys</i> spp., flounder	10-15	350-500	-	0

It has been suggested that ingestion of plastic production pellets by larval and juvenile fishes may cause blockage of the digestive tract and prevent normal feeding. There is no experimental evidence that we know of to support this. In those laboratory experiments where larvae have fed on pellets (Colton et al. 1974; Hjelmeland et al. 1988; Settle et al. in prep.), the pellets have either been rejected or passed through the gut. In our experiments the larvae subsequently fed on brine shrimp nauplii and appeared healthy. Had the larvae been fed angular particles or particles containing toxic chemicals, the results may have been different. In the sea, dead larvae would seldom if ever be collected because of rapid decomposition.

Food habit studies confirm that large fish also eat plastic material, but the frequency and quantity of material eaten is not well documented. Ingestion of large pieces of plastic by fish may cause a health problem. Many predatory fish have large mouths and can swallow large pieces of plastic. They cannot digest the plastic, however, and it may prove to be too large to pass from the stomach into the gut and out the anus. If the fish cannot regurgitate the piece, it may block the intestine or cause ulceration.




Figure 3.--Spot, *Leiostomus xanthurus* (17 mm standard length), with ingested polystyrene microspheres (350-500 μm) in the gut.

We conclude that the overall ingestion of inert plastic by larval and juvenile fish is probably not a significant mortality factor at this time in the ocean environment. Monitoring of larval fishes from different areas to determine if the frequency of occurrence of plastic in the guts is changing should be continued and incorporated into ongoing ichthyoplankton studies.

We also recommend that studies be conducted to determine if larger predatory fishes can swallow and subsequently pass large, irregular pieces of plastic. Additional mortality caused by plastic ingestion might be detrimental to populations of certain species of sport fish already under intense fishing pressure.




Figure 4.--Spot, *Leiostomus xanthurus* (17 mm standard length), partially dissected to show polystyrene microspheres (350-500 μm) distributed throughout the alimentary tract.

ACKNOWLEDGMENTS

We thank Patricia Tester and Xiaoyen Zheng for assistance in the laboratory and Curtis Lewis for the photography. Charles Manooch III, provided critical review of the manuscript. Beaufort Laboratory research was funded in part by funds provided by the Marine Entanglement Research Program, NMFS.

REFERENCES

- Anonymous.
1975. Plastic cups found in fish. *Mar. Pollut. Bull.* 6:148.
- Azzarello, M. Y., and E. S. Van Vleet.
1987. Marine birds and plastic pollution. *Mar. Ecol. Prog. Ser.* 37:295-303.

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Carpenter, E. J., S. J. Anderson, G. R. Harvey, H. P. Milkas, and B. B. Peck.
1972. Polystyrene spherules in coastal waters. Science (Wash., D.C.) 178:749-750.
- Carpenter, E. J., and K. L. Smith.
1972. Plastics on the Sargasso Sea surface. Science (Wash., D.C.) 174:1240-1241.
- Colton, J. B., Jr., F. D. Knapp, and B. R. Burns.
1974. Plastic particles in surface waters of the northwestern Atlantic. Science (Wash., D.C.) 185:491-497.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Day, R. H.
1980. The occurrence and characteristics of plastic pollution in Alaska's marine birds. M.S. Thesis, Univ. Alaska, Fairbanks, 111 p.
- Day, R. H., D. M. Clausen, and S. E. Ignell.
1986. Distribution and density of plastic particulates in the North Pacific Ocean in 1986. Document submitted to the International North Pacific Fisheries Commission, Anchorage, Alaska, November 1986. Northwest and Alaska Fisheries Center, Auke Bay Laboratory, NMFS, Auke Bay, Alaska.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Fry, D. M., S. I. Fefer, and L. Sileo.
1987. Ingestion of plastic debris by Laysan albatrosses and wedge-tailed shearwaters in the Hawaiian Islands. Mar. Pollut. Bull. 18:339-343.
- Gramentz, D.
1988. Involvement of loggerhead turtles with the plastic, metal, and hydrocarbon pollution in the central Mediterranean. Mar. Pollut. Bull. 19:11-13.

- Gregory, M. R.
1977. Plastic pellets on New Zealand beaches. *Mar. Pollut. Bull.* 8:82-84.
- Hays, H., and G. Cormons.
1974. Plastic particles found in tern pellets, on coastal beaches and at factory sites. *Mar. Pollut. Bull.* 5:44-46.
- Hjelmeland, K., B. H. Pedersen, and E. M. Nilssen.
1988. Trypsin content in intestines of herring larvae, *Clupea harengus*, ingesting inert polystyrene spheres or live crustacea prey. *Mar. Biol. (Berl.)* 98:331-335.
- Ignell, S. E., and M. L. Dahlberg.
1986. Results of 1986 cooperative research on the distribution of marine debris in the North Pacific Ocean. (Document submitted to the International North Pacific Fisheries Commission, Anchorage, Alaska, November 1986. Northwest and Alaska Fisheries Center, Auke Bay Laboratory, NMFS, Auke Bay, Alaska.)
- Kartar, S., F. Abou-Seedo, and M. Sainsbury.
1976. Polystyrene spherules in the Severn Estuary - A progress report. *Mar. Pollut. Bull.* 7:52.
- Kartar, S., R. A. Milne, and M. Sainsbury.
1973. Polystyrene waste in the Severn Estuary. *Mar. Pollut. Bull.* 4:144.
- Laist, D. W.
1986. An overview of impacts associated with lost and discarded fishing gear, packaging material, and other persistent synthetic materials on living marine resources. *TCS Bull.* 9(4):5-13.

1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. *Mar. Pollut. Bull.* 18:319-326.
- Manooch, C. S. III.
1973. Food habits of yearling and adult striped bass, *Morone saxatilis* (Walbaum), from Albemarle Sound, North Carolina. *Chesapeake Sci.* 14:73-86.
- Manooch, C. S. III, and W. T. Hogarth.
1983. Stomach contents and giant trematodes from wahoo, *Acanthocybium solanderi*, collected along the South Atlantic and gulf coasts of the United States. *Bull. Mar. Sci.* 33:227-238.
- Manooch, C. S. III, and D. L. Mason.
1983. Comparative food studies of yellowfin tuna, *Thunnus albacares*, and blackfin tuna, *Thunnus atlanticus* (Pisces: Scombridae) from the southeastern and gulf coasts of the United States. *Brimleyana* 9:33-52.

- Manooch, C. S. III, D. L. Mason, and R. S. Nelson.
1984. Food and gastrointestinal parasites of dolphin *Coryphaena hippurus* collected along the southeastern and gulf coasts of the United States. Bull. Jpn. Soc. Sci. Fish. 50:1511-1525.
1985. Foods of little tunny, *Euthynnus alletteratus*, collected along the southeastern and gulf coasts of the United States. Bull. Jpn. Soc. Sci. Fish. 51:1207-1218.
- Morris, A. W., and E. S. Hamilton.
1974. Polystyrene spherules in the Bristol Channel. Mar. Pollut. Bull. 5:26-27.
- Morris, R. J.
1980. Plastic debris in the surface waters of the South Atlantic. Mar. Pollut. Bull. 11:164-166.
- Pruter, A. T.
1987. Sources, quantities and distribution of persistent plastics in the marine environment. Mar. Pollut. Bull. 18:305-310.
- Settle, L. R., et al.
In prep.
- Shaw, D. G.
1977. Pelagic tar and plastic in the Gulf of Alaska and Bering Sea: 1975. Sci. Total Environ. 8:13-20.
- Shaw, D. G., and G. A. Mapes.
1979. Surface circulation and the distribution of pelagic tar and plastics. Mar. Pollut. Bull. 10:160-162.
- Shiber, J. G.
1979. Plastic pellets on the coast of Lebanon. Mar. Pollut. Bull. 10:28-30.
1987. Plastic pellets and tar on Spain's Mediterranean beaches. Mar. Pollut. Bull. 18:84-86.
- Tester, P. A., L. R. Settle, and D. E. Hoss.
1987. Association of marine debris with upwelling and frontal zones: A progress report to NMFS Marine Entanglement Research Program.
- U.S. Congress. Senate.
1986. The Plastic Waste Reduction Act of 1986. Congressional Record, Senate 25 June 1986, p. 58397.
- Venrick, E. L., T. W. Backman, W. C. Bartman, C. J. Platt, M. S. Thornhill, and R. E. Yates.
1973. Man-made objects on the surface of the central North Pacific Ocean. Nature 241:271.
-

- Wiebe, P. H., K. H. Burt, S. H. Boyd, and A. W. Morton.
1976. A multiple opening/closing net and environmental sensing system for sampling zooplankton. *J. Mar. Res.* 34:313-326.
- Wong, C. S., D. R. Green, and W. J. Cretney.
1974. Quantitative tar and plastic waste distributions in the Pacific Ocean. *Nature* 247:30-32.

**SYNTHETIC MATERIALS FOUND IN THE STOMACHS OF LONGNOSE
LANCETFISH COLLECTED FROM SURUGA BAY, CENTRAL JAPAN**

Tadashi Kubota
Department of Fisheries, Faculty of Marine
Science and Technology, Tokai University
Shimizu-shi, Shizuoka, 424 Japan

ABSTRACT

Stomach contents of a total of 372 longnose lancetfish, *Alepisaurus ferox* Lowe, 296 stranded on the beach of Miho Key in Suruga Bay between 1964 and 1983 and 76 fished by gillnets in waters near the key between 1969 and 1975, were examined. In addition to food organisms, many synthetic items such as pieces of polyethylene and vinyl were found in the stomachs. This paper examines the presence of these synthetic materials in the stomachs of longnose lancetfish. Major results of this study were as follows:

- Synthetic materials found in the stomachs were mostly soft polyethylene and vinyl pieces of various sizes and colors. Intact plastic soft drink bottles were also found.
- The feeding ratio of synthetic materials in the stomach of lancetfish was 62.2% for stranded specimens and 63.2% for gillnet specimens.
- Average number of pieces of synthetic materials in the stomach was 3.1 for stranded specimens and 2.2 for gillnet specimens.
- The feeding ratio and number of synthetic pieces in the stomachs of longnose lancetfish have increased sharply during the past several years, suggesting that there have been increases in the amounts of synthetic materials in Suruga Bay and neighboring waters. There are concerns that the neglected synthetic materials may impact large marine organisms adversely.

INTRODUCTION

Longnose lancetfish, *Alepisaurus ferox* Lowe (Alepisauridae), is widely distributed in the Pacific, Atlantic, and Indian Oceans. It has a large mouth, large eyes, and very sharp bladelike teeth. Its body tissue is soft and watery. It is well known as voracious fish (Fig. 1).

In Suruga Bay and Sagami Bay, located at the center of Honshu in Japan, longnose lancetfish are often stranded alive on the shore by waves. Strandings are especially frequent between December and May on the shores of Kambara, Numazu, Miho, and Ohsesaki, which are located at the inmost part of Suruga Bay (Kubota and Uyeno 1970).

Since 1964, the author has been collecting lancetfish caught with gillnets and boat seines and those stranded on the shore of Miho Key to study their morphology and food habits (Fig. 2; Kubota and Uyeno 1970, 1978; Kubota 1971, 1973, 1977; Kubota and Mori 1975; Okutani and Kubota 1976).

The stomachs of the fish examined contained many pieces of synthetic materials such as polyethylene and vinyl in addition to ordinary food items (e.g., fishes, cephalopods, shrimps, salps, *Pyrosoma*).

It was pointed out that synthetic materials found in the stomachs of lancetfish are from pollution of the ocean and that they served as an index for effects on large nekton such as fish (Kubota 1977). No previous study has examined the effects of synthetic materials on marine nekton.

The objectives of this study were to determine the amounts of synthetic material ingested by longnose lancetfish and to determine how it had changed with the time.

MATERIALS AND METHODS

In this study, 372 fish were examined. Of these, 296 were found stranded on the shore of Miho Key and the remaining 76 were caught in gillnets in the area near Miho Key between December and May. The lengths of the fish range from 50 to 125 cm. Immediately after collection, measurements of meristic characters were made in the laboratory. Food items found in stomachs were removed for identification. The amount and size of nonfood items were also recorded. Nonfood items included leaves, pieces of wood, straw, fragments of orange, fragments of vegetable, rubber, vinyl pieces, polyethylene pieces, and intact plastic soft drink bottles.

RESULTS

Synthetic materials eaten by lancetfish were mostly soft polyethylene and vinyl pieces. Both size and color of these items varied (Fig. 3). Besides these materials, intact plastic soft drink bottles (38 mm in diameter and 74 mm in height) were found in the stomachs of 11 lancetfish stranded on the shore between 1971 and 1973. Of these, four lancetfish had two bottles each in their stomachs in addition to food items.




Figure 1.--A longnose lancetfish collected from Suruga Bay. Date collected: 27 April 1967, body length: 887 mm. Scale in figure indicates 300 mm.




Figure 2.--Suruga Bay, central Japan. The shaded portion of Miho Key is the beach where longnose lancetfish have been stranded and the shaded area off Miho Key indicates a gillnet fishing ground.




Figure 3.--Synthetic materials from stomachs of longnose lancetfish stranded on the beach of Miho Key. Date collected: May 1971. Scale in figure indicates 100 mm.

The feeding ratio of synthetic materials to food was examined for each year. The feeding ratio for 184 of the 296 specimens that were stranded on the shore was 62.2%, whereas the feeding ratio for 48 specimens among 76 caught with gillnets was 63.2%. The average amount of synthetic material per specimen for each year was also studied. The average amount of synthetic materials per specimen was 3.1 pieces for specimens stranded on the shore and 2.2 pieces for those caught in gillnets.

The results from stranded specimens cover a long time period. Therefore, the study period was divided into two parts, 1964-75 and 1978-83. The feeding ratio of synthetic materials in stomachs of lancetfish was 58.0% in the period 1964-75, and it increased to 72.0% in 1978-83. The average amount of synthetic material per specimen increased from 2.2 pieces in 1964-75 to 4.5 pieces (more than double) in 1978-83 (Tables 1 and 2).

For the samples that had synthetic materials in their stomachs, the frequency of the amount of synthetic materials was studied to see how many pieces were eaten per specimen.

Of those fish on the shore, 112 specimens did not have synthetic materials in their stomachs at all. Thirty-six samples had one piece. Of 184 fish, 135 (73.4%) ate 1 to 6 pieces of synthetic material. One lancetfish ate 17 pieces.

Table 1.--Number of synthetic pieces found in the stomachs of longnose lancetfish stranded on the beach of Miho Key.

Year	Number of lancetfish	Number of lancetfish with pieces of synthetic materials	Total number of pieces of synthetic materials	Average number of pieces of synthetic materials
1964	2	1	5	2.5
1965	2	1	7	3.5
1966	2	1	15	7.5
1967	9	2	10	1.1
1968	19	12	33	1.7
1969	19	11	29	1.5
1970	18	9	38	2.1
1971	56	30	99	1.8
1972	21	15	70	3.3
1973	24	17	72	3.0
1974	2	1	3	1.5
1975	2	1	1	0.5
1976	--	--	--	--
1977	--	--	--	--
1978	57	39	281	4.9
1979	37	24	145	3.9
1980	16	12	76	4.8
1981	--	--	--	--
1982	--	--	--	--
1983	10	8	38	3.8
Total	296	184	922	

For the fish caught with gillnets, 28 did not have any synthetic material pieces and 12 fish ate only 1 piece. One specimen had 15 pieces in its stomach. All the others contained 10 or fewer pieces (Tables 3 and 4).

These results show that synthetic material in the stomachs of longnose lancetfish is increasing. This is from the increase in synthetic materials being discarded by people into rivers and the ocean.

DISCUSSION

The longnose lancetfish is a voracious feeder. It has nonselective food habits and will catch anything in the ocean it can swallow. In most cases, the stomach contents can be identified to the species level. The feeding habits of lancetfish are the same in other areas (Haedrich 1964; Haedrich and Nielsen 1966; Fourmanoir 1969; Rancurel 1970; Fujita and Hattori 1976). Therefore, it is possible to identify the organisms in the

Table 2.--Number of synthetic pieces found in the stomachs of longnose lancetfish caught by gillnet.

Year	Number of lancetfish	Number of lancetfish with pieces of synthetic materials	Total number of pieces of synthetic materials	Average number of pieces of synthetic materials
1969	1	1	1	1.0
1970	14	6	12	0.9
1971	33	23	74	2.2
1972	21	12	63	3.0
1973	5	4	10	2.0
1974	1	1	3	3.0
1975	1	1	5	5.0
Total	76	48	168	

Table 3.--Number and frequency of synthetic pieces found in each stomach of longnose lancetfish stranded on the beach of Miho Key.

Number of pieces of synthetic materials found in each stomach	Number of lancetfish
0	112
1	36
2	29
3	25
4	16
5	16
6	13
7	6
8	9
9	6
10	5
11	4
12	4
13	3
14	4
15	3
16	4
17	1

Table 4.--Number and frequency of synthetic pieces found in each stomach of longnose lancetfish caught by gillnet.

Number of pieces of synthetic materials found in each stomach	Number of lancetfish
0	28
1	12
2	9
3	8
4	7
5	4
6	3
7	2
8	0
9	1
10	1
15	1

habitat where lancetfish live using stomach analysis. The distribution of nonfood items such as synthetic materials can also be determined.

Results showing that polyethylene and vinyl pieces found in the stomachs of lancetfish have increased over time imply that fairly large quantities of synthetic materials are present in the waters near Miho Key. In the last few years, the author has observed water surfaces of the area from the innermost part to the central part of Suruga Bay from on board a research vessel, and has seen large floating vinyl pieces. These items were not seen at all in the sea 8 years ago. This study documents the notion that quantities of discarded synthetic materials have increased in recent years. Synthetic materials mass-produced to meet the consumer demands will continue to contaminate the seas around Japan because they are discarded from houses and factories as waste and enter the sea through rivers.

Because of their feeding habits, lancetfish can serve as a biological monitor of synthetic pollution in the ocean.

REFERENCES

- Fourmanoir, P.
1969. Contenus stomachaus d'*Alepisaurus* (Poissons) dans le Sud-Ouest Pacifique. Cah. ORSTOM Ser. Oceanogr. 7:51-60.

- Fujita, K., and J. Hattori.
1976. Stomach content analysis of longnose lancetfish, *Alepisaurus ferox* in eastern Indian Ocean and the Coral Sea. Jpn. J. Ichthyol. 23:133-142.
- Haedrich, R. L.
1964. Food habits and young stages of North Atlantic *Alepisaurus* (Pisces, Iniomi). Breviora 201:1-15.
- Haedrich, R. L., and J. G. Nielsen.
1966. Fishes eaten by *Alepisaurus* (Pisces, Iniomi) in the southeastern Pacific Ocean. Deep-Sea Res. 13:909-919.
- Kubota, T.
1971. Food of anglerfish, *Lophius litulon*, obtained from stomachs of lancetfish, *Alepisaurus ferox*, in Suruga Bay. Bull. Plankton Soc. Jpn. 18:28-31. [In Jpn.]

1973. Four links of food chain from the lancetfish, *Alepisaurus ferox*, to zooplankton in Suruga Bay, Japan. J. Fac. Mar. Sci. Technol., Tokai Univ. 7:231-243.

1977. Food of lancetfish, *Alepisaurus ferox* Lowe, fished by gill-net in Suruga Bay, Japan. J. Fac. Mar. Sci. Technol., Tokai Univ. 10:137-146.
- Kubota, T., and T. Mori.
1975. Morphology of "barrel" of *Phronima* eaten by lancetfish, *Alepisaurus ferox* Lowe, in Suruga Bay, Japan. Annu. Rep. Mar. Sci. Mus., Tokai Univ. 2/3:61-65. [In Jpn.]
- Kubota, T., and T. Uyeno.
1970. Food habits of lancetfish, *Alepisaurus ferox* (order Myctophiformes) in Suruga Bay, Japan. Jpn. J. Ichthyol. 17:22-28.

1978. On some meristic characters of lancetfish, *Alepisaurus*, collected from Suruga Bay, Japan. J. Fac. Mar. Sci. Technol., Tokai Univ. 11:63-69.
- Okutani, T., and T. Kubota.
1976. Cephalopods eaten by lancetfish, *Alepisaurus ferox* Lowe, in Suruga Bay, Japan. Bull. Tokai Reg. Fish. Res. Lab. 84:1-9.
- Rancurel, P.
1970. Les contenus stomacaux d'*Alepisaurus ferox* dans les Sud-Ouest Pacifique (Cephalopodes). Cah. ORSTOM Ser. Oceanogr. 8:3-87.

**ECOLOGICAL ASPECTS OF MARINE TURTLES IMPACTED
BY OCEAN DEBRIS: A 1989 PERSPECTIVE**

George H. Balazs and Barry K. Choy
Southwest Fisheries Science Center Honolulu Laboratory
National Marine Fisheries Service, NOAA
Honolulu, Hawaii 96822, U.S.A.

ABSTRACT

Authenticated reports of debris entanglement and ingestion by marine turtles have continued to accumulate since a comprehensive, worldwide list of such events was first assembled in 1984. Although fragmentary, available evidence indicates that ingestion of man-made debris floating on the high seas has the greatest potential for adversely impacting sea turtle populations. A major problem in gathering detailed information on this phenomenon is the inability of researchers to locate and study pelagic habitats used by juvenile turtles of all species. Consequently, those cases of debris ingestion that do become known should be considered as the tip of the iceberg. Due to the insights of the late Archie Carr, pelagic habitats used as foraging sites by sea turtles are not believed to be frontal systems (convergences, rips, drift lines) where buoyant food and debris are drawn together by advection. International concern for the impact of buoyant wastes in the ocean is heightened by the fact that many sea turtle populations are endangered and have experienced serious declines from overfishing and other adverse factors.

STUDIES ON THE INGESTION OF PLASTIC AND LATEX BY SEA TURTLES

Peter L. Lutz*
Rosenstiel School of Marine and Atmospheric Science
University of Miami
Miami, Florida 33149, U.S.A.

*Present address: Department of Biological Sciences, Florida Atlantic University, Boca Raton, Florida 33431.

ABSTRACT

Small pieces of latex and plastic sheeting were offered to sea turtles on different occasions and the turtles' feeding behavior was noted, as well as the time taken for the turtles to pass ingested materials. The physiological and clinical status of turtles that had consumed plastic sheeting was also monitored. We observed that green sea and loggerhead turtles actively seek out and consume the offered material. Some color preference was shown, clear plastic having the lowest acceptance rate. The amount consumed was influenced by appetite. At the low feeding levels allowed in these experiments, we detected no effects of plastic ingestion on gut function, metabolic rate, blood chemistry, liver function, or salt balance. However, blood glucose declined for 9 days following ingestion, indicating a possible interference in energy metabolism or gut function. The sojourn of the ingested latex material in the gut ranged from a few days to 4 months. Moreover, some of the turtles passed multiple pieces all bound together, although they had ingested the individual pieces at different times. Since the gut clearance time for food is in the order of days, it appears that some of the latex pieces were being held up in the intestine. Latex pieces that had been retained for the longest time in the gut showed evidence of deterioration.

INTRODUCTION

As man's use of nonbiodegradable products increases, so does the amount of such material dumped into the ocean. Offshore garbage dumping by ships at sea was legal until recently and the ocean is considered by some (e.g., Osterburg 1986) as "nature's trash basket." However, one consequence of this practice is that contact by marine animals with nonbiodegradable refuse such as plastic bags and Styrofoam products also increases. Hopefully, the ratification of the MARPOL V agreement will help

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

to alleviate the problem, but recent incidents of entanglement and ingestion in marine mammals and seabirds (Cawthorn 1985) suggest that harmful contact with refuse may occur much more frequently than previously thought.

It is becoming increasingly recognized that the ocean dumping of plastic waste presents a particularly serious hazard for sea turtles. Sea turtles consume a wide variety of debris and, in the man-made category, plastic bags and sheets appear to be the most prevalent material ingested (Balazs 1985). In some instances, the level of contamination can be very high. For example, plastic bags were found in 23% of a sample of green sea turtles in Peruvian waters (Hays de Brown and Brown 1982), and in one analysis 44% of adult nonbreeding leatherbacks were found to have plastic in their stomachs (Mrosovsky 1981). It has been suggested that one cause for ingestion is that turtles mistake the plastic for their natural jellyfish prey (Fritts 1982). More recently concern has been expressed over spent balloon material in the ocean, the result of increasing popularity of massive balloon launches.

Is the ingestion of plastic and latex by sea turtles any cause for concern? Clearly, if sufficient material is swallowed to cause a complete stoppage of the gut, death will result from starvation. However, there are only a few such documented cases (Balazs 1985; Cawthorn 1985), and most of the evidence for turtles swallowing plastic comes from butchered turtles (Balazs 1985). In domestic vertebrates, persistent partial blockage of the intestine can interfere with gut function (Fraser 1986). In the sea turtle, a coating of the gut wall by plastic could cause a reduction in absorption efficiency and also cause mechanical damage to the gut lining. Sublethal ingestion, therefore, where complete intestinal blockage does not occur, may be quite common and could adversely affect behavior, growth, reproduction, and general homeostatic physiological functioning and lead to other potentially lethal situations.

There is, unfortunately, no information on whether the ingestion of such material is accidental or deliberate, or information on the effects of sublethal ingestion of plastics by sea turtles. Given the critical position of most sea turtle populations and the huge magnitude of ocean dumping (van Dolah et al. 1980; Horsman 1982), it is clearly important to determine if the swallowing of such inert material by sea turtles is harmful and to establish the seriousness of any harm.

The purpose of this study was to document the mode of plastic and latex ingestion in sea turtles and to give a first estimate of how serious the resultant harm might be.

MATERIALS AND METHODS

This is the first study of its kind, and since there were no previous data to use as a guide, and as we did not wish to cause any lasting harm to the experimental sea turtles, we were particularly careful and cautious in designing our experimental protocol.

Animals

Green sea and loggerhead turtles were kept in tanks of approximately 3,785 L (1,000 gal) capacity. Each tank was supplied with running, filtered seawater. The turtles were fed a specially formulated feed for sea turtles (Purina sea turtle chow) each day during the experiments unless otherwise noted.

Ingestion

In the initial experiment, green and loggerhead yearlings (ca. 1 kg weight) and juvenile (10 to 18 kg) turtles were allowed to consume a small single piece of plastic sheeting (1 to 10 cm²) and were observed for about 2 weeks during which time various behavioral (yearling and juvenile) and physiological (juvenile) measurements were taken. The animals were fed turtle chow daily during this experiment. Since a preliminary examination of the data showed no adverse effects, a second set of experiments was undertaken at an increased (but still modest) level of plastic ingestion. Seven loggerheads weighing 13 to 18 kg were used in this section (four experimental, three control) and were fed five to seven small pieces of plastic. They were also fed daily and observed for 2 weeks. In these experiments the initial measurements before feeding plastic served as individual controls. In order to understand the effects of simple food limitation per se, a third set of turtles was starved for 2 weeks and the various physiological parameters were monitored. This set also served as a control for those turtles in the previous experiments that occasionally refused food for a few days.

An additional study on latex was undertaken in order to determine whether the ingestion of balloon material was accidental or deliberate and if the latex material was altered on passage through the gut. Five turtles were isolated in separate tanks and were offered small (ca. 1 cm²) pieces of colored latex and clear plastic sheeting under different conditions. The turtles' feeding behavior was noted, as well as the time taken for the turtles to pass ingested materials. The passed material was collected for examination.

Gut Function

Food consumption was measured as the number of pellets consumed each day. The pellets weighed on average 0.918 ± 0.085 g. Feces were collected in plastic bags attached to the turtles and stored frozen at -20°C. It was noted that defecation usually started 1 to 2 h after feeding. Samples of food and feces were dried at 67°C for 48 h and their calorific value measured using a Parr 1241 Adiabatic Calorimeter.

The ash content of food and feces was estimated by weighing samples before and after being heated in a muffle furnace at 600°C for 24 h. Ash was used as a digestibility marker (Conover 1966). Although this method has been criticized because of its unproven assumption that ash-forming materials are neither added nor absorbed as food passes through the gut (Bjorndal 1985; Newman et al. 1985), it is used fairly commonly in studies

of digestibility in marine organisms, and gives values in reasonable agreement with the acid insoluble method in sea turtles (Vargo et al. 1986). It also has value as a comparative estimate.

Gut passage time was determined from the first appearance in the feces of the plastic sheets and of small plastic markers (Teflon disks, 2-3 mm diameter) that had been included in the food.

Occult blood in the feces was tested for using the benzidine reaction (Henry 1974).

Dive Time

Dive time was recorded on a stopwatch while observing the turtles' diving behavior in the tank. Surface time was not measured since the interval was, almost without exception, less than 3 sec (usually one breath).

Oxygen Consumption

A closed circuit method was used for oxygen consumption measurements. The turtle was placed in a sealed humidified air chamber connected to an Applied Electrochemistry oxygen analyzer. Chamber air was pumped through the analyzer and returned to the chamber. Carbon dioxide and water vapor were removed from the analyzer input line by chemical scrubbers (Ascarite and Dririte). The experiments were run for approximately 1 h, and the minimal chamber partial oxygen pressures (PO_2) were always >100 torr.

Blood Chemistry

Blood was taken from the dorsal cervical sinus as previously described (Bentley and Dunbar-Cooper 1980).

Blood gases (PO_2 , PCO_2) and pH were determined immediately on whole blood using a Radiometer BMS Mk 2 blood-gas analyzer set to the experimental temperature (22°C). Plasma bicarbonate was calculated from the pH and PCO_2 data using the temperature- and pH-dependent CO_2 solubility and dissociation constants of Severinghaus (1965).

The blood was then centrifuged and the plasma divided into two parts. One part was deproteinized with 8% chilled perchloric acid and served for plasma lactate and urea measurements using the Sigma kit No. 826-uv for lactate and the Sigma kit No. 640 for urea. The untreated plasma was analyzed for osmotic pressure using a Wescor 6100 osmometer and saved frozen for measurement of ions and metabolites. Plasma chloride was measured by an Aminco chloride titrator and plasma cations by atomic absorption spectrophotometry (Perkin Elmer PE 403). Column chromatography was used to estimate plasma cortisol, and glutamic pyruvate transaminase levels were measured by spectrophotometry using Sigma kit No. 505. The hematocrit and the percentage volume of white blood cells were read after centrifugation.

RESULTS

Feeding and Digestion

Ingestion

During normal feeding, green sea turtles were each offered, on different occasions, five pieces of pink, blue, and yellow latex, and clear plastic (Fig. 1). Each turtle had its own preference: No. 1, blue; No. 2, pink; and No. 3, yellow; Nos. 4 and 5 refused all. Surprisingly, none of the turtles accepted the clear plastic. On offering yellow material to turtles that had been fasted for 3 days, there was a substantial increase in the amount of ingestion. Turtles No. 1, No. 2, and No. 3 consumed all of the material offered, but turtle No. 5 continued to hold itself aloof from this experiment (Fig. 2). In two additional sets of experiments on fasted turtles, turtle No. 1 ingested clear plastic but the others continued to ignore it (Fig. 3).

Gut Passage Time

The ingested material started appearing in the tank water after a few days and then declined over the next few weeks (Fig. 4). This time course corresponded with normal gut passage time as measured by the Teflon markers (11.3 days, range 10 to 13 days, $n = 3$). Quite unexpectedly, latex material continued to appear in the tank for up to as long as 4 months, peaking at about 8 weeks. Some of the turtles passed multiple pieces all bound together, although they had ingested the individual pieces at different times. The latex pieces that had been held for the longest time in the gut showed evidence of deterioration.

Food Consumption

In the loggerheads, daily food consumption did not vary much on an individual basis and when changes occurred they were fairly smooth (Fig. 5A). There was no noticeable pattern after feeding plastic. The average daily rate of consumption (grams of food per kilogram body weight per day) for individual loggerheads was 5.07 ± 1.97 , $n = 7$; 5.9 ± 3.08 , $n = 7$; 9.2 ± 1.59 , $n = 11$; 9.3 ± 2.06 , $n = 8$. In the green sea turtles, the average rates were similar, i.e., 6.7 ± 3.8 , $n = 8$; 10.9 ± 1.93 , $n = 8$; 11.82 ± 2.8 , $n = 9$. However, in one of the green sea turtle consumption gradually diminished to zero on day 4 and then recovered (Fig. 5B). The consumption patterns for the other two turtles were similar to those observed in the loggerheads.

Energy Adsorption

The calorific value of the feces showed no consistent change with time in either the green sea turtles or the loggerheads (Fig. 6). Interestingly, the green sea turtle feces had a higher calorific content than the loggerhead (loggerhead feces $3,328 \pm 145$ cal/g, $n = 10$; green $4,126 \pm 324$ cal/g, $n = 9$). These differences were statistically significant ($P < 0.01$). It can be calculated that an amount of loggerhead food containing 1 g of ash




Figure 1.--Voluntary ingestion of latex pieces in green sea turtles.




Figure 2.--Effect of 3 days fasting on latex ingestion in green sea turtles.




Figure 3.--Effect of 3 days fasting on the ingestion of clear plastic in green sea turtles.




Figure 4.--Gut passage time for ingested pieces of latex in the green sea turtle.




Figure 5.--The effect of plastic ingestion (†) on food consumption in four individual loggerhead (A) and three green sea turtles (B).




Figure 6.--The calorific value of green and loggerhead turtle feces after being fed plastic (†).

would have a gross energy content of 51,381 cal, while feces with the same amount of ash would have 12,743 cal. Assuming constancy of ash, this indicates a digestible energy adsorption efficiency of 75.2%.

Stool Culture

In the two control loggerheads, the fecal flora was respectively 99% g positive, 1% g negative; and 98% g positive, 2% g negative. In three turtles that had been fed plastic bags, the fecal floral composition was as follows: 100% g positive; 85% g positive, 15% g negative; 100% g positive. The gut bacterial composition was, therefore, substantially gram positive in nature and this feature was not altered by plastic ingestion.

Occult Blood

No occult blood was observed in any of the fecal samples examined either in the control or the experimental animals. The plastic ingestion, therefore, had not caused intestinal bleeding.

Respiration

Oxygen Consumption

Plastic ingestion had no apparent effect on the oxygen consumption of either the green sea or the loggerhead turtles, and on an individual basis they were remarkably constant over the 2 weeks of monitoring. Metabolic rates for the green sea turtles ranged from 47.9 to 73.8 ml/kg/h, and the loggerhead values showed an almost identical range of from 38.1 to 70.2 ml/kg/h. Similar oxygen consumptions have been obtained for green sea turtles (70.8 ml/kg/h at 25°C, Kraus and Jackson 1985) and loggerheads (62.0 ml/kg/h, Lutz and Bentley 1985) measured in air.

Blood Chemistry and Acid Base Balance

Oxygen

Venous oxygen levels remained relatively constant in both the experimental turtles and in the starved group. There was no significant difference between groups. Since venous oxygen levels are determined by the difference between oxygen supply and tissue use, and since oxygen consumption did not change, it seems likely that the mechanisms for oxygen transport have not been affected by plastic ingestion. The mean venous value for all of the data ($PO_2 = 56.69 \pm 1.59$, $n = 38$) is very similar to that found in an earlier study on the same animals (Lutz and Dunbar-Cooper 1987).

Carbon Dioxide

Venous carbon dioxide remained similarly constant over the course of the experiment, and no statistical difference was found between the control and the experimental groups. The mean value for all of the data is $PO_2v = 24.79 \pm 0.976$, $n = 38$.

Blood pH

For the group of experimental turtles fed plastic, venous blood pH appeared to decline on the first day after feeding plastic ($P \leq 0.5$) and continued to fall in two turtles on day 2 and in one until day 3 (Fig. 7A). No such trend was noted for the starved controls (Fig. 7B). However, the range in pH shifts was very narrow, and for the whole set the average pH was 7.550 ± 0.008 , $n = 38$, close to the predicted normal venous pH for the prevailing body temperature (25°C , $\text{pH} = 7.442$, Lutz et al. 1988).

Bicarbonate

There was no change in venous bicarbonate on the day following plastic ingestion. The overall bicarbonate concentration was 22.6 ± 0.971 mM, $n = 38$.

Glucose

In the loggerheads fed plastic, blood glucose levels declined for 10 days (Fig. 8), but recovered to initial values by day 14, about the time plastic was expelled from the gut (see below). A least squares linear regression of the relationship between blood glucose (G) and days after plastic ingestion (T) produced the following equation illustrated in Figure 8.

$$G \text{ (mM)} = 6.683 - 0.445 T \quad r = 0.866, n = 12$$

The average rate of decline in blood glucose was therefore 0.45 mM/day. Interestingly, starvation by itself caused a marked fall in blood glucose levels (Fig. 9). In both the loggerhead and green sea turtles, blood glucose levels declined sharply on the second day of starvation at much greater rates than the fed loggerheads who had consumed plastic viz., 2.52 mM/day in the green and 2.42 mM/day in the loggerhead.

Glutamic Transaminase

The initial concentration of loggerhead glutamic transaminase plasma (GTP) was 1.67 ± 0.608 , $n = 7$, international units/ml. The GTP values varied somewhat in both the control turtles and the plastic-fed turtles for the first 3 days of the experiment (Fig. 10), but after the fourth day there was a marked decline in values in both groups, possibly related to the fall in plasma glucose.

Cortisol

In all samples tested, the blood cortisol levels were extremely low (≤ 1.0 $\mu\text{g/dl}$), indicating that the turtles were not stressed by the experimental protocol. Blood cortisol levels have been seen to increase in stressed loggerheads from similar low initial levels (1-3 $\mu\text{g/dl}$ to as high as 37 $\mu\text{g/dl}$ (D. Owens pers. commun.).




Figure 7.--The effect of plastic ingestion (A, ↑) and starvation (B, ↑) on the loggerhead turtle venous pH.

Hematocrit

The hematocrit values did not change over the course of the experiments in either the loggerhead or the green sea turtle. The loggerhead mean value (28.6%) is less than that found for loggerheads sampled in the wild (35.5%, Lutz and Dunbar-Cooper 1987) and less than that found for the green sea turtle (33.3%); the latter difference is significant ($P < 0.01$).

White Blood Cells

No change was seen in white blood cell volume following plastic ingestion. In the loggerheads, the white blood cells initially made up about 0.2% of the whole blood, and with one exception the values were reasonably constant, ranging between 0.2 and 0.4% for 10 days after plastic ingestion.




Figure 8.--The effect of plastic ingestion (↑) on blood glucose levels in the loggerhead turtle.

DISCUSSION

We have been able to demonstrate that both green sea and loggerhead turtles do not discriminate against plastic sheeting when they engulf food intermingled with plastic. The experiments with latex ingestion in loggerheads demonstrated that if their appetite is sufficient, they will actively swim towards and ingest latex materials, that all colors are acceptable, and that the amount ingested will depend on their nutritional state. Indeed, it was our impression that hungry sea turtles will swallow almost any material of a suitable size and consistency and will continue to do so until satiation.

No clear evidence of ill effects from plastic ingestion was found in this set of experiments though it should be noted that the turtles were only allowed to consume very small amounts. In fact, the constancy of many of the physiological parameters over the 2 weeks of monitoring is evidence that the experimental setup was not, by itself, a perturbing influence.

Further evidence of a lack of stress is seen in the low blood cortisol levels. The values are similar to those reported for resting blood cortisol levels for vertebrates in general which are around 1-5 $\mu\text{g}/100\text{ ml}$ (rainbow trout, 3.8 $\mu\text{g}/100\text{ ml}$, Donaldson 1981; loggerhead, 1-3 $\mu\text{g}/100\text{ ml}$, Owens pers. commun.; dog, 1-5 $\mu\text{g}/100\text{ ml}$, Fraser 1986). For many animals, stress produces a surge in blood corticosteroids, often within hours of the stress, that will persist during the stress and sometimes for days afterwards (Fraser 1986). Compared to resting values, the expected increases in blood cortisol concentrations under stressful conditions can be substantial (16 $\mu\text{g}/100\text{ ml}$ in the stressed rainbow trout, Donaldson 1981).




Figure 9.--The effect of starvation (↑) on blood glucose concentrations in the loggerhead (A) and green sea turtles (B).

There was no evidence of plastic ingestion affecting feeding and the handling of food. The rate of food consumption did not change after eating plastic in either the loggerhead or the green sea turtles, and the average daily consumption was similar for both species (9.79 g/kg/day, green; 7.37 g/kg/day, loggerhead). Wood and Wood (1981) found a similar food intake for green sea turtles fed pellets (8 to 12 g/kg/day). The food consumption rates found in this study are equivalent to a calorific intake of 44.2 kcal/kg/day for the green and 33.3 kcal/kg/day for the loggerhead.

The efficiency of food adsorption and the calorific value of the feces were unaltered, and the bacterial composition of the gut was not changed. There was no evidence of blood in the feces, pointing to an absence of mechanical damage as plastic passed through the gut.

No effect of plastic ingestion was detected with respect to any of the measured parameters that are directly associated with respiratory




Figure 10.--The effect of plastic ingestion (†, A) and starvation (†, B) on glutamic transaminase levels in loggerhead plasma.

physiology, viz., metabolic rate, blood oxygen and carbon dioxide levels, blood acid base status.

The hematocrit was remarkably constant, an indicator of health, and no marked changes were seen in the proportion of white blood cells. A very substantial increase in white blood cell numbers (400%) was one of the most notable features of sea turtles affected by oil pollution (Vargo et al. 1986). No evidence of liver malfunctioning was seen in the lack of increase in plasma glutamic pyruvic transaminase (Fraser 1986).

The rates of change in blood glucose are a possible exception to this pattern. The key observation was that blood glucose declined rapidly in loggerheads that were starved and also fell, although at a lesser rate, in turtles that had been fed plastic sheets. The implication is, therefore,

that blood glucose levels in sea turtles are especially sensitive to nutrient uptake from the gut and that this process had been interfered with in those animals that had consumed plastic. Interestingly, the blood glucose concentrations for the control fed loggerheads in this study ($5.23 \text{ mM} \pm 1.279 \text{ mM}$, $n = 10$) were much higher than those recorded in the wild from loggerheads sampled in the Port Canaveral ship channel (ca. 1 mM , Lutz and Dunbar-Cooper 1987) evidence perhaps that the Canaveral turtles had not been feeding. Blood glucose levels, therefore, may serve as a sensitive index of nutritional status for turtles both in the laboratory and in the wild.

The study did point to some interesting differences in the physiology of green sea turtles and loggerheads. On average the green sea turtles had a higher hematocrit than the loggerheads (33.3%, green; 28.6%, loggerhead) and a higher proportion of white blood cells (0.2%, loggerhead; 1.02%, green). In the green sea turtles, the average daily food consumption of the pelleted food was about 32% higher. On the other hand, this was offset somewhat by green sea turtles having a higher feces energy content (24% higher in the green) and, therefore, a lower efficiency in extracting energy from the food.

In summary, when hungry, sea turtles will actively consume plastic and latex material. Except for a possible interference in energy metabolism (declining blood glucose levels), at the levels allowed in this study ingestion produced no measurable changes in the physiological parameters that were measured. However, the observation that pieces of latex can gather up in the gut and remain there for considerable periods of time should be viewed with some concern and certainly needs more detailed investigation.

ACKNOWLEDGMENTS

This work was supported by the U.S. Department of Commerce RFP No. FSN-85-0178 and by a grant from the Toy Manufacturers Association of America.

REFERENCES

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Bentley, T. B., and A. Dunbar-Cooper.
1980. A blood sampling technique for sea turtles. Contract No. NA-80-GE-A-00082 for the National Marine Fisheries Service.
- Bjorndal, K.
1985. Use of ash as an indigestible dietary marker. Bull. Mar. Sci. 36:224-230.

- Cawthorn, M. W.
1985. Entanglement in, and ingestion of, plastic litter by marine mammals, sharks, and turtles in New Zealand waters. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 336-343. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Conover, R. J.
1966. Assimilation of organic matter by zooplankton. *Limnol. Oceanogr.* 11:338-345.
- Donaldson, E. M.
1981. The pituitary - inter renal axis as an indicator of stress in fish. In A. D. Pickering (editor), *Stress and fish*, p. 12-47. Acad. Press, Lond.
- Fraser, C. M. (editor).
1986. The Merck veterinary manual. Sixth ed. Merck and Co., Rahway, N.J.
- Fritts, T. H.
1982. Plastic bags in the intestinal tracts of leatherback marine turtles *Eromochelys-Coracea*. *Herpetol. Rev.* 13(3):72-73.
- Hays de Brown, C., and W. M. Brown.
1982. The status of sea turtles in the southeastern Pacific: Emphasis on Peru. In K. A. Bjorndal (editor), *Biology and conservation of sea turtles*. Proceedings of the World Conference on Sea Turtle Conservation, November 1979, Washington, D.C., p. 235-240. Smithsonian. Inst. Press, Wash., D.C.
- Henry, R. J.
1974. *Clinical chemistry: Principle and techniques*. Second ed. Harper and Row, N.Y.
- Horsman, P. V.
1982. The amount of garbage pollution from merchant ships. *Mar. Pollut. Bull.* 13:167-169.
- Kraus, D. R., and D. C. Jackson.
1985. Temperature effects on ventilation and acid-base balance of the green turtle. *Am. J. Physiol.* 239:254-258.
- Lutz, P. L., and T. Bentley.
1985. Adaptations for diving in the sea turtle. *Copeia* 1985:671-697.
- Lutz, P. L., and A. Dunbar-Cooper.
1987. Variations in the blood chemistry of the loggerhead sea turtle *Caretta caretta*. *Fish. Bull., U.S.* 85:37-44.

- Lutz, P. L., A. Dunbar-Cooper, and M. Bergey.
1988. Effects of temperature on gas exchange and acid base balance in the sea turtle *Caretta caretta* at rest and during routine activity. Submitted.
- Mrosovsky, N.
1981. Plastic jellyfish. Mar. Turtle Newsl. 17:5-7.
- Newman, M. W., P. L. Lutz, and S. C. Snedaker.
1985. Temperature effects on feed ingestion and assimilation efficiency of nutrient by the Malaysian prawn, *Macrobrachium rosenbergii*. J. World Maricult. Soc. 13:95-103.
- Osterburg, C.
1986. Waste disposal - where should it be? Land or sea? Sixth International Ocean Disposal Symposium 181.
- Severinghaus, J. W.
1965. Blood gas concentration. In Handbook of physiology respiration, p. 1475-1487. Am. Physiol. Soc., Wash., D.C.
- Van Dolah, R. F., V. G. Burrell, Jr., and S. B. West.
1980. The distribution of pelagic tars and plastics in the South Atlantic Bight. Mar. Pollut. Bull. 11:352-356.
- Vargo, S., P. L. Lutz, D. Odell, E. van Vleet, and G. Bossart.
1986. Effects of oil on sea turtles. Final report to M.M.S., U.S. Department of the Interior. Rep. No. 14-12-0001-30063.
- Wood, F. E., and J. R. Wood.
1981. Growth and digestibility for the green turtle (*Chelonia mydas*) fed diets containing varying protein levels. Aquaculture 25:269-274.

EFFECTS OF ANTHROPOGENIC DEBRIS ON SEA TURTLES IN THE NORTHWESTERN GULF OF MEXICO

Pamela Plotkin and Anthony F. Amos
The University of Texas at Austin
Marine Science Institute
Port Aransas, Texas 78373, U.S.A.

ABSTRACT

Reports of sea turtles ingesting and becoming entangled in marine debris and the adverse effects associated with these encounters exist worldwide, but the magnitude of this problem has yet to be determined. Data collected from sea turtles stranded on the south Texas coast from 1986 through 1988 indicate that they are significantly affected by ingestion of and, to a lesser extent, by entanglement in marine debris. All five species of sea turtles found in the Gulf of Mexico, both male and female, posthatchling through adult, had eaten or were ensnared by debris. Plastics discarded at sea were involved in the majority of these incidents. The offshore oil industry, cargo ships, research vessels, commercial and recreational fishing boats, and other seagoing vessels are primarily responsible for the trash discarded at sea which threatens sea turtles in the Gulf of Mexico.

INTRODUCTION

Because of their widespread intentional exploitation by man in the past, sea turtle populations in the United States have declined and all species are currently considered either threatened with or in danger of extinction. The greatest threat to their survival today is man's incidental exploitation. Every year thousands of sea turtles are incidentally caught and drowned in the net trawls of shrimp fishermen, beach front development encroaches on valuable sea turtle nesting beaches and threatens their reproductive efforts, newly hatched sea turtles are run over by cars or die from heat and exhaustion after they are enticed to crawl from their nests towards the bright lights of a condominium instead of towards the comparatively dimly lit sea, and an unknown number of sea turtles die when they become entangled in or ingest nonbiodegradable anthropogenic marine debris.

Balazs (1985) was the first to examine the widespread effects and impacts of marine debris on sea turtles. He compiled reports from the literature and through personal communication on the incidences of

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

entanglement in and ingestion of marine debris by sea turtles worldwide. Collectively, these reports painted a rather grim picture for the recovery of sea turtle populations. But precisely how much of a threat marine debris poses to sea turtles has not yet been determined. Because sea turtles spend most of their lives at sea and are generally inaccessible to researchers, it has been difficult to assess the magnitude of this problem on any population. The objective of the present study was to determine the extent of entanglement and ingestion for sea turtles found stranded on the south Texas coast.

METHODS

Data were collected from sea turtles found stranded on Mustang Island, North Padre Island, and South Padre Island, Texas, from 1986 through 1988.

Entanglement

Stranding forms submitted to the Texas Sea Turtle Stranding and Salvage Network coordinator were used to obtain information on entangled sea turtles. Information culled from these forms included species stranded, date stranded, condition of the turtle (i.e., alive or dead), size of the turtle (curved carapace length (CCL)), type of entanglement, and fate of the turtle.

Ingestion

Stranded turtles were necropsied following Wolke and George (1981). Prior to necropsy, the species was identified and CCL and width measurements were recorded. During necropsy, the sex of the turtle was determined by visual examination of the gonads. The esophagus, stomach, and intestinal tract were removed from the body cavity and all organs were examined for abnormalities: lesions, ruptures, and parasites. The contents of the digestive tracts were emptied onto a fine-meshed sieve and rinsed with water. Anthropogenic debris was separated from the other food items, catalogued, and saved for later analysis. The remaining food items were preserved in 10% buffered formalin.

RESULTS

Entanglement

Sea turtles became entangled when their head, limbs, or entire bodies accidentally were ensnared in debris or active fishing gear. During the 3-year study, 30 (7.5%) of the 400 sea turtles reported stranded were entangled (Table 1). All of the sea turtle species found in the northwestern Gulf of Mexico had been ensnared. These included 13 Kemp's ridleys, *Lepidochelys kempi*, 7 loggerheads, *Caretta caretta*, 6 hawksbills, *Eretmochelys imbricata*, 3 green turtles, *Chelonia mydas*, and 1 leatherback, *Dermochelys coriacea*. Commercial and recreational fishermen and their lost or discarded gear were responsible for the majority of these incidents. Sea turtles were found entangled in fishing line or hook (9), shrimp trawl (7), net or rope (5), plastic woven produce sacks (4), tar (3), trotline

Table 1.--Incidence of entanglement in sea turtles found stranded on the south Texas coast from 1986 through 1988.

Year	Number of turtles entangled (%)	Total number of turtles stranded
1986	14 (7.8)	179
1987	11 (10.1)	109
1988	5 (4.5)	112
All years	30 (7.5)	400

(1), and crab pot (1). Injuries resulting from their entanglement were responsible for the deaths of seven of these turtles. The remaining 23 turtles were rehabilitated at the University of Texas Marine Science Institute and, with the exception of 1 permanently injured (blind) turtle, were released back into the Gulf of Mexico.

Ingestion

Marine debris was found in the stomachs or intestinal tracts of 60 (54.1%) of the 111 turtles necropsied (Table 2). It was present in 52.3% of the loggerheads, 46.7% of the green turtles, and 87.5% of the hawksbills (Table 3). (No leatherbacks were necropsied during the study.) Shaver (pers. commun.) examined the gut contents of Kemp's ridleys stranded within the same study area and found debris in 29.8% of those turtles (Table 3). Plastic materials were most frequently eaten (Table 4). Most of this material (ca. 60%) was buoyant in nature, but some was not, indicating that sea turtles not only feed on debris floating on the surface of the water, but also feed on debris that is suspended in the water column or is on the bottom.

The incidence of debris ingestion was highest in those turtles stranded during December and lowest in turtles stranded during August (Fig. 1). However, seasonal trends should not be interpreted from these data because recent work by Lutz (pers. commun.) has revealed that sea turtles have the ability to retain plastic in their digestive tracts for prolonged periods of time.

Our ingestion data support Carr (1987), who warned that the young, advanced pelagic stage sea turtles were most vulnerable because they spend the first few years of their life in the open ocean, dependent upon drift lines (areas of high debris concentrations) for their food supply and shelter. Information on the size (carapace length) at which sea turtles become sexually mature (adult) is based upon data collected from females at their nesting beaches. The size at sexual maturity differs among the sea turtle species, varies geographically within a species, and is unknown for male sea turtles. For the purposes of this study, we defined posthatchling

Table 2.--Incidence of debris ingestion in sea turtles found stranded on the south Texas coast from 1986 through 1988.

Year	Number of turtles with debris (%)	Total number of turtles necropsied
1986	10 (40.0)	25
1987	32 (59.3)	54
1988	18 (56.3)	32
All years	60 (54.1)	111

Table 3.--Incidence of debris ingestion by the different sea turtle species found stranded on the south Texas coast from 1986 through 1988.

Species	Number of turtles with debris (%)	Total number of turtles necropsied
Loggerhead, <i>Caretta caretta</i>	46 (52.3)	88
Green, <i>Chelonia mydas</i>	7 (46.7)	15
Hawksbill, <i>Eretmochelys imbricata</i>	7 (87.5)	8
Kemp's ridley, <i>Lepidochelys kempii</i> ^a	31 (29.8)	104

^aD. J. Shaver pers. commun.

Table 4.--Types of debris (and their occurrence) collected from the intestinal tracts of sea turtles found stranded on the south Texas coast from 1986 through 1988.

Type of debris	Number of turtles that had ingested that type	Percent (N = 111)
Plastic bag, pieces	39	35.1
Styrofoam	17	15.3
Plastic, hard pieces	15	13.5
Plastic, line or rope	10	9.0
Plastic beads or pellets	8	7.2
Balloons	7	6.3
Tar	7	6.3
Glass	2	1.8
Paper or cardboard	2	1.8
Aluminum	2	1.8
Stainless steel hook	1	0.9
Latex or rubber	1	0.9
Heat-sealed drink tab	1	0.9




Figure 1.--Percent occurrence (by month) of anthropogenic debris found in the digestive tracts of sea turtles stranded on the south Texas coast.

to 40-cm CCL as advanced pelagic stage turtles, 40-80 cm CCL as subadult turtles, and ≥ 80 -cm CCL or greater as adult turtles. We found debris in 70.8% of the advanced pelagic stage turtles, 55.4% of the subadult turtles, and 31.8% of the adult turtles (Fig. 2).

Debris ingestion resulted in the deaths of four of the turtles necropsied during this study (a noticeable obstruction or blockage in the digestive tract was observed), but could not be implicated in the deaths of the remaining 56 turtles. It was difficult to determine if the debris eaten had caused a turtle's death. For most cases observed, only small quantities of debris were present, and they were usually well mixed in the digestive tracts with the other food items and probably did not contribute to death.

DISCUSSION

A number of the turtles that washed ashore during the study were already missing a limb. Many of these losses were suspected to be the result of a prior entanglement, but because there was no proof, these turtles were not counted as having been entangled. Therefore, we feel that our entanglement numbers may be too small. The reasons why sea turtles become entangled remains unclear. Their natural curiosity towards objects




Figure 2.--Occurrence of anthropogenic debris found in the digestive tracts of sea turtles stranded on the south Texas coast from 1986 through 1988 (by carapace length (cm)).

adrift in the water is most often cited as the reason for their propensity for probing near and becoming ensnared in debris. It is likely that sea turtles are attracted to these floating objects because they are seeking food or shelter.

An unusual relationship was found between hawksbills and plastic woven produce sacks (onion sacks). The four incidents of entanglement in those sacks reported here all involved advanced pelagic stage hawksbills (their CCL ranged from 19.4 to 28.5 cm) had their head or limbs caught in the plastic fibers of a produce sack. In addition to our four reports, we know of two other hawksbills that were found entangled in the exact same manner. In 1988, one was found stranded on Galveston Island, Texas (M. Duronslet pers. commun.), and the other was found in April 1989 on the beach at Rancho Nuevo, Mexico (R. Byles pers. commun.). What affinity, if any, hawksbills have for onion sacks is unknown. More behavioral studies of all of the sea turtle species are necessary before we can explain how and why they become involved in these situations.

Debris was eaten by more than half of the turtles necropsied during this study, and while this ingestion did not appear to result in the deaths

of the majority of these turtles, its presence in the digestive tracts of so many is indicative of the pervasiveness of anthropogenic debris in the northwestern Gulf of Mexico. It has been suggested that sea turtles eat debris because it resembles their natural prey or perhaps because epizoic or epiphytic growth on the debris has attracted the turtle. Before man began discarding his nonbiodegradable wastes into the oceans, sea turtles did not have to differentiate between what was edible and what was not, because essentially everything was edible. In the Gulf of Mexico, the offshore oil industry, cargo ships, research vessels, commercial and recreational fishing boats, and other seagoing vessels are primarily responsible for the trash discarded at sea which eventually is consumed by many sea turtles. Prevailing currents and winds drive virtually all of the trash that is dumped into the Gulf of Mexico (and to a lesser extent the Caribbean) to the northwestern Gulf of Mexico and onto the Texas coast.

Annex V of MARPOL (implemented domestically by the Act to Prevent Pollution from Ships) came into effect on 31 December 1988. This annex prohibits the dumping of plastics at sea and regulates how far from shore other anthropogenic debris may be discarded. The passage of this law probably will not deter the many who have grown accustomed to dumping their trash overboard. This law needs to be enforced at sea and at the ports, and those who are guilty should be fined as one means of controlling the oceanic debris problem. Most importantly, people need to be educated and convinced to save their refuse until they can properly dispose of it on land.

Certain bodies of water such as the Mediterranean Sea were given special designation under Annex V of MARPOL. These areas have been afforded extra protection because of their unique oceanographic or ecological conditions, and it is now illegal to discard any type of debris in these waters. The Gulf of Mexico was considered a candidate for this special protection, but was not designated as such when Annex V was passed. The semienclosed nature of the Gulf of Mexico, the prevalence of marine debris in these waters and on adjacent beaches, and the importance of this area as a habitat for sea turtles (in particular the critically endangered Kemp's ridley sea turtle) should be enough justification for its designation as a special area. The likelihood that a sea turtle inhabiting the Gulf of Mexico will come into contact with anthropogenic debris is quite substantial.

ACKNOWLEDGMENTS

We would like to thank the University of Texas Marine Science Institute; Texas A&M University Sea Grant; the Galveston Laboratory, National Marine Fisheries Service, NOAA; Sea Turtles Inc.; and Sigma Xi, the scientific research society, for their financial support of this study. This work would not have been possible without the many who have helped in reporting and retrieving stranded sea turtles: Padre Island National Seashore, Texas Parks and Wildlife Department, Nueces County Parks Department, Port Aransas Police Department, the Pan American University Coastal Studies Laboratory, Donna Shaver, Robert Whistler, Jenny Bjork, Ed Hegen, Page Campbell, Rosemary Breedlove, and Don Hockaday. Special thanks are due to Richard Byles for his comments and review of this paper.

REFERENCES

- Balazs, G.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Carr, A.
1987. Impact of non-degradable marine debris on the ecology and survival outlook of sea turtles. Mar. Pollut. Bull. 18:352-356.
- Wolke, R. E., and A. George.
1981. Sea turtle necropsy manual. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SEFC-24.

ON THE SYNTHETIC MATERIALS FOUND IN THE DIGESTIVE SYSTEMS OF, AND
DISCHARGED BY, SEA TURTLES COLLECTED IN WATERS ADJACENT TO JAPAN

Itaru Uchida
Himeji City Aquarium
Nobusue Tegarayama
Himeji, Hyogo, Japan

ABSTRACT

It has been recognized that five species of sea turtles are common in waters adjacent to Japan: *Caretta caretta*, *Chelonia mydas*, *Eretmochelys imbricata*, *Lepidochelys olivacea*, and *Dermochelys coriacea*. Stranded, live turtles collected along the coast of central and southern Japan were examined, and most were found to have ingested synthetic materials into their systems during the research period. The main types of plastics found were transparent bags or sheeting, monofilament fishing line, and rope parts. Large plastic sheets were ingested particularly by the leatherback turtle, *D. coriacea*. It is concluded that most of the sea turtles found in waters adjacent to Japan have a high frequency of plastic ingestion.

PLASTIC INGESTION IN A PYGMY SPERM WHALE, *KOGIA BREVICEPS*

Raymond J. Tarpley
The Texas Veterinary Medical Center
Texas A&M University
College Station, Texas 77843, U.S.A.

ABSTRACT

An adult female pygmy sperm whale, *Kogia breviceps*, (2.9 m body length) and a young male (1.8 m body length) thought to be her calf stranded alive on Galveston Island on 1 January 1984. Both animals were transported to a holding tank for observation and treatment. The female was extremely weak and died on the third day of captivity. Severe multiple mucosal ulcerations were found throughout all stomach chambers during necropsy. In contrast, the calf initially appeared to be thriving. He was able to swim unassisted and eventually began to make shallow dives. Force feeding was begun, and on the eighth and ninth days he voluntarily accepted squid placed in front of him. However, on the tenth day he suddenly weakened, lost interest in feeding, and died. On necropsy, the first two stomach compartments (forestomach and fundic chamber) were found to be completely occluded by a plastic garbage can liner, a bread wrapper, a corn chip bag, and two other pieces of plastic sheeting. The small third stomach chamber (connecting channel) prevented passage of the debris farther along the gastrointestinal tract. A severe inflammation within the abdominal cavity was also found which was diagnosed as the immediate cause of death and thought to be secondary to the gastric obstruction. The primary food item for this species is squid, and it is feasible that suspension of this debris in the water column may have been mistaken as prey by the inexperienced calf.

INGESTION OF PLASTIC DEBRIS BY STRANDED
MARINE MAMMALS FROM FLORIDA

Nélio B. Barros, Daniel K. Odell
Rosenstiel School of Marine and Atmospheric Science
University of Miami
Miami, Florida 33149, U.S.A.

and

Geoffrey W. Patton
Mote Marine Laboratory
1600 City Island Park
Sarasota, Florida 33577, U.S.A.

ABSTRACT

Ocean pollution in the form of plastic debris has been recently recognized as a major threat to marine wildlife. Injuries and fatalities caused by entanglement and ingestion of floating and submerged debris have been documented for an increasing number of marine vertebrates (mammals, birds, turtles, fish). Ingested plastics may cause a false sensation of fullness, decreasing feeding bouts, or may obstruct normal passage of food through the digestive tract.

Ingestion of plastic material is reported here for five species of marine mammals stranded along the Florida coast. These include four cetacean species (bottlenose dolphin, *Tursiops truncatus*; false killer whale, *Pseudorca crassidens*; pygmy sperm whale, *Kogia breviceps*; dwarf sperm whale, *K. simus*) and the sirenian, *Trichechus manatus* (West Indian manatee), from both coastal (*Tursiops truncatus*, *T. manatus*) and pelagic (*P. crassidens*, *K. breviceps*, *K. simus*) habitats. Debris included plastic jugs, disposable surgeon gloves, plastic bags, and monofilament lines. Plastic debris was usually found with food items (vegetation, and fish and cephalopod remains). On at least one occasion (an emaciated *T. truncatus*), ingestion of such material is believed to have contributed to death. With the increasing littering of the Florida coastline, plastic impact on marine mammal populations may be much more widespread than previously reported.

SURVEY OF MARINE DEBRIS INGESTION BY ODONTOCETE CETACEANS

William A. Walker
Section of Mammalogy
Natural History Museum of Los Angeles County
Los Angeles, California 90007, U.S.A.

and

James M. Coe
Alaska Fisheries Science Center
National Marine Fisheries Service, NOAA
Seattle, Washington 98115, U.S.A.

ABSTRACT

Odontocete cetaceans are affected to an unknown degree by the ingestion of oceanic debris. Published accounts discuss primarily the sperm whale, *Physeter macrocephalus*.

The pathologic effects of foreign body ingestion on captive cetaceans are well documented, and provide background information on the potential effects of debris ingestion on wild, free-ranging animals. A survey of major institutions reveals 40 incidences of debris ingestion in 16 species of stranded odontocete cetaceans. Plastic debris was prevalent, with a total occurrence of 80.0%. Evidence indicates ingestion of debris may be secondary to the stranding syndrome. A survey of prior food habits analyses on 10 species of odontocete cetaceans was conducted. All species combined, a total of 1,790 stomachs were examined. Marine debris was encountered only in Baird's beaked whale, *Berardius bairdii*, taken at two localities in the coastal waters of Japan. In *B. bairdii* taken off the Pacific coast of central Japan, debris incidence in 86 stomachs was 26.7%. Plastic debris made up 39.1% of the foreign material ingested. Off northern Hokkaido, in the southern Okhotsk Sea, incidence of debris in 20 stomachs was 15.0%. Food habits data indicate that the lower frequency of debris ingestion is related to differences in feeding strategy in the northern region.

In the wild state odontocete cetaceans are probably discriminating feeders. Evidence indicates that the high occurrence of debris in *Physeter macrocephalus* and *B. bairdii* is due primarily to incidental ingestion along with benthic prey.

INTRODUCTION

The quantity of increasingly diverse marine litter discarded into the world's oceans is reaching enormous proportions. Billions of pounds of debris are dumped into the sea each year (Carpenter and Smith 1972; Venrick et al. 1973; Wong et al. 1974; Morris 1980a, 1980b; Van Dolah et al. 1980; Eldridge 1982; O'Hara et al. 1988). Recent studies reveal this to be more than an aesthetic problem. Debris, particularly nonbiodegradable plastics, is accumulating in the marine environment and causing significant mortality in some marine animals (Wallace 1985).

In 1984, the first Workshop on the Fate and Impact of Marine Debris was held in Hawaii. Papers presented confined their data on ingestion of marine debris largely to marine birds (Day et al. 1985) and turtles (Balazs 1985; Cawthorn 1985). The potential problem of debris ingestion by marine mammals was not addressed, with only anecdotal accounts appearing in the proceedings (Cawthorn 1985; Mate 1985).

Ingestion of debris by cetaceans does occur. Early accounts of an impressive array of nonfood items ingested by the sperm whale, *Physeter macrocephalus*, are well known (Turner 1903; Millais 1906; Hollis 1939; Pike 1950; Sleptsov 1952; Clarke 1956; Berzin 1959, 1971; Caldwell et al. 1966). Berzin (1971) discusses accounts of "several vinyl chloride bags" found in North Pacific sperm whale stomachs as early as 1961. More recent accounts of debris ingestion involve stranded cetaceans (Wehle and Coleman 1983; Cawthorn 1985; Mate 1985; Cowan et al. 1986).

It is evident that there is a need for research on the occurrence and potential impact of cetacean debris ingestion. This study, though confined to the toothed cetaceans (odontoceti), begins to address this subject. It is divided into three major areas of inquiry. 1) Investigate the effects of foreign body ingestion by captive cetaceans maintained by marine aquariums in order to assess the potential effects of debris ingestion in the wild. 2) Conduct a survey on incidences of debris ingestion in stranded cetaceans. 3) Survey incidence of marine debris in food habits analyses conducted on free-ranging cetacean populations.

METHODS OF DATA COLLECTION

Information on the pathologic effects of captive cetacean foreign body ingestion was derived from the literature and the senior author's personal experience as biologist and curator at Marineland of the Pacific during the period 1968-74.

Records of debris ingestion in stranded odontocete cetaceans were solicited from institutions and persons known to include stomach content examination as part of a coordinated stranded animal recovery program. Due to variation in recordkeeping and necropsy techniques imposed by a 24-year time frame (1963-86), frequency of occurrence data on debris ingestion could not be reliably derived. Accounts from the literature were not included unless sufficient information on time frame, locality, and nature of ingested debris was available.

Data on evidence of debris ingestion in free-ranging odontocete cetaceans were obtained from food habits studies conducted on animals collected at sea as a result of incidental fisheries interactions or directly taken for research or commercial purposes. In each case, personnel directly involved in the preliminary stomach content sorting procedures were interviewed. Unusual items encountered during this stage tend to be remembered (though not necessarily recorded). An "I don't recall" response during the interview resulted in elimination of the study from the data base.

In some instances determinations as to whether ingested objects constitute debris directly ingested or introduced secondarily through prey species presented a problem. Tiny bits of plastic and isolated fishhooks are particularly suspect. Fish are well known to be attracted to and ingest man-made objects. Recently ingestion of plastic particles by oceanic squid has been documented (Araya 1983; Machida 1983). For purposes of this study, animals containing only isolated fishhooks were noted but not included in the frequency data presented.

As this study progressed, it became apparent that there are probably more isolated incidences of debris ingestion in odontocete cetaceans than we were able to locate in the time allowed. We welcome any oversights brought to our attention so the records may be included in a future revision of this report.

RESULTS

Captive Cetacean Foreign Body Ingestion

Mortality in marine parks and zoos due to foreign body ingestion is well documented in the literature. Brown et al. (1960) described causes of mortality in a major oceanarium, Marineland of the Pacific, during the first 5 years of operation. Three of those years were summarized as follows: "The losses occurring during the years 1955 to 1958 were, with few exceptions, caused by animals swallowing indigestible foreign material and resulting in gastric and enteric impactions." Nakajima et al. (1965) reported that 18 of 92 (19.6%) dolphin casualties at Enoshima Aquarium in Japan between 1958 and 1965 had foreign material in their stomachs. Caldwell et al. (1965) described a simultaneous mortality of three trained bottlenose dolphins at Gulfarium in Florida. They died from ingesting plastic strips from the tank enclosure. "Balls of plastic up to four inches in diameter were found in the first stomach of these animals."

During the senior author's tenure at Marineland of the Pacific, numerous cetaceans died as a result of ingestion of foreign material. One trained Pacific bottlenose dolphin was particularly noteworthy in that it had ingested a piece of a polyethylene plastic bag ca. 0.19 m (2 ft square). Necropsy findings revealed that while the major portion of the bag remained in the forestomach, a small section extended through the sphincter and into the gastric stomach. Tissue pathology was extensive. Approximately one-half of the forestomach lining and submucosa had eroded away, with necrotic tissue and inflammation extending deep into the

musculature of the stomach wall. The stomach wall and serosa adjacent to this lesion were edematous and thickened five to six times beyond normal. It was surmised that the effect of the plastic protruding through the sphincter into the gastric stomach caused an excess of digestive fluids to be released into the forestomach, severely injuring those portions of stomach lining insulated by plastic.

Captive cetaceans have been known to ingest a wide variety of foreign material. Objects such as cotton gloves, tin cans, plastic bags, bottles, pens, coins, flashbulbs, plastic combs, nails, steel wool cleaning pads, plastic toys, and women's jewelry are some of the articles reported (Brown et al. 1960; Amemiya 1962; Caldwell et al. 1965; Nakajima et al. 1965; Ridgway 1965, 1972; Brown et al. 1966).

The reasons for the high incidence of foreign body ingestion in captive cetaceans are not clear. The captive environment, due to its obvious spatial limitations, is at best an abnormal one. The social behavior of these animals has been severely altered (Caldwell et al. 1968). Ridgway (1972) suggested that since captive animals are taught to consume dead fish, they may consider any object entering the pool as edible. Excitement of training, performing, play behavior, and competition for food may also be contributing factors (Nakajima et al. 1965).

What is clear from the accounts on captive cetacean ingestion of foreign objects is that it has the potential for being a direct cause of mortality, or at least debilitating to a degree which could predispose animals to disease or predation in the wild state.

Stranded Animal Debris Ingestion

Case descriptions of stranded animal debris ingestion by species are presented in Table 1. A total of 43 accounts were available, spanning a period of time from 1963 to 1986. Sixteen species of odontocete cetaceans were involved.

Reports on debris ingestion came primarily from the east and west coasts of North America (37 and 58%, respectively). Only one record each was obtained from the Gulf of Mexico (Texas) and Hawaii. The differences in frequency probably reflect regional variation in stranded cetacean recovery and detailed necropsy techniques rather than true geographic differences in abundance of marine debris.

The kinds of debris ingested varied considerably. Plastic bags and plastic sheeting were the most prevalent items (62.5%). Other miscellaneous plastic articles such as drinking straws, bottle caps, discarded fishing net, synthetic rope, and a small container occurred in 17.5% of the cases. The total occurrence for all plastic debris was 80.0%. Nonplastic debris such as rubber balloons, asphalt, cellophane, cloth, paper, and metal articles (excluding fishhooks) was encountered in 37.5% of the cases reported. Fragments of marine plants, which are also abnormally ingested, were encountered in 32.6% of the stomachs examined.

Table 1.--Records of the ingestion of marine debris by stranded odontocete cetaceans.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
--	6/79	Florence, OR	--	--	Physeter macrocephalus, sperm whale About 1 L of tightly packed trawl net in stomach. One of 38 stomachs examined from a mass stranding of 41 animals	Mate 1985.
Sp-1	6/7/79	Purgatory Bay, Bonavista Cove, Newfoundland, Canada	1,030	F	Small length of nylon rope and unidentified debris	U.S. National Museum. ^a
MME01362	7/1/85	Seaside, NJ	510	--	Mylar balloon	U.S. National Museum. ^a
USNM504132	12/4/74	Corolla, NC	178	F	<i>Kogia simus</i> , dwarf sperm whale Plastic Wonderbread bread wrapper	U.S. National Museum. ^a
CMNH0216	4/27/76	Sullivan's Island, SC	318	F	<i>Kogia breviceps</i> , pygmy sperm whale Two small pieces of thin plastic	Charleston Museum. ^b
MME00549	1/1/84	Galveston, TX	182	M	Pounds of plastic bags clogging its stomach chambers	U.S. National Museum; ^a Wehle and Coleman 1983.
MME01263	5/17/85	Brevard Co., FL	320	M	Plastic bag	U.S. National Museum. ^a

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
JRH-087	11/20/80	San Diego, CA	526	M	Piece of asphalt	Southwest Fisheries Science Center. ^c
USNM550111	1/7/81	Assawaman, VA	580	F	Large plastic bag and plastic wrappers	U.S. National Museum. ^a
USNM550734	1/27/86	Seaford, VA	512	F	Plastic straw and a horse chestnut	U.S. National Museum. ^a
USNM550018	11/22/80	Hatteras Island, NC	311	M	Large piece of clear plastic bag	U.S. National Museum. ^a
USNM550362	12/28/83	Cape May, NJ	371	F	Stomach filled with plastic bag	U.S. National Museum. ^a
USNM550754	2/14/86	East Hampton, NY	420	M	One plastic bottle cap	U.S. National Museum. ^a
USNM550310	5/18/83	Corolla, NC	275	M	Small plastic container	U.S. National Museum. ^a
USNM504462	6/28/76	Mauai, HI	215	F	Plastic bag in stomach. This animal was one of nine mass stranded on 6/28/76.	U.S. National Museum. ^a

Ziphius cavirostris, Cuvier's beaked whale*Mesoplodon europaeus*, Gervais' beaked whale*Mesoplodon densirostris*, Blainville's beaked whale*Globicephala macrorhynchus*, short-finned pilot whale*Steno bredanensis*, rough-toothed dolphin

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
USNM504486	10/12/76	Sandbridge, VA	206	M	Two pieces of heavy black plastic. This animal and one below are part of mass stranding of 13 on 10/12/76	U.S. National Museum. ^a
USNM504494	10/12/76	Sandbridge, VA	233	M	One large fishhook loose in stomach	U.S. National Museum. ^a
--	8/29/63	Santa Monica, CA	165	M	Piece of paper wadded into a 5.1 cm (2-in) ball along with seaweed, squid beaks, and roundworms.	Caldwell et al. 1965.
WAW-130	8/15/71	Santa Monica, CA	167	M	Stomach contained numerous small plastic bags, pieces of cardboard, and waxed paper. Numerous kelp fronds (<i>Macrocystis pyrifera</i>) also present. This animal had been observed inside a yacht harbor for 10 days prior to stranding. Necropsy diagnosed parasitic central nervous system pathology and hepatitis as cause of death.	W. Walker, unpubl. data; Cowan et al. 1986.
WAW-174	9/21/72	Long Beach, CA	176	F	Forestomach half full of four plastic bags, two plastic bottle caps, and numerous small sticks, twigs, leaves, and kelp fronds	W. Walker, unpubl. data. Pathology data from Cowan et al. 1986

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
WAW-192	7/18/83	Santa Monica, CA	188	F	<p>(<i>Macrocystis</i> and <i>Eggregia</i>). One No. 4 fishhook snagged in stomach lining. Necropsy diagnosed central nervous system pathology due to parasitism by the trematode, <i>Nasitrema</i> sp.</p> <p>Three small plastic bags, one plastic drinking straw, one Calif. Dep. Fish Game mackerel fish tag No. M-11283. Tag is 5.1 cm (2 in) long, yellow, spaghetti type. Necropsy diagnosed central nervous system pathology due to parasitism as cause of stranding.</p>	W. Walker unpubl. data. Pathology data from Cowan et. al. 1986.
WAW-148	2/12/72	Malibu, CA	173	F	<p><i>Delphinus delphis</i>, common dolphin</p> <p>One 15 * 15 cm plastic bag, several kelp fronds (<i>Macrocystis pyrifera</i>). Cause of stranding diagnosed as parasitism of central nervous system.</p>	W. Walker unpubl. data. Pathology data from Cowan et al. 1986.
WAW-172	9/8/72	Will Rogers State Beach, Los Angeles County, CA	190	M	<p>Two 20-cm² pieces of cellophane; one small piece of black plastic (approximately 3 cm²), and portions of marine plants (<i>Macrocystis</i>, <i>Eggregia</i>, and</p>	W. Walker unpubl. data. Pathology data from Cowan et al. 1986.

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
LACM72286	11/24/80	Hermosa Beach, CA	197	F	<i>Phyllospadix</i> . Cause of strangling diagnosed as parasitism of central nervous system.	Los Angeles County Museum of Natural History.
JEH331	4/25/86	Will Rogers State Beach, Los Angeles County, CA	193	F	One rusted fishhook embedded in stomach wall. Necropsy revealed an apparently unrelated massive tumor or abscess in abdomen adjacent to left kidney.	Los Angeles County Museum of Natural History.
WFP-559	2/5/77	La Jolla, CA	302	M	Stomach contained one partial red balloon (3 x 13 cm), one piece of clear plastic (8 x 13 cm), and kelp fronds (<i>Macrocystis pyrifera</i>). <i>Tursiops truncatus</i> , bottlenose dolphin (All southern California coastal population)	Los Angeles County Museum of Natural History.
WAW-141	12/26/71	Huntington Beach, CA	207	F	One rusted metal bottle cap beach sand, fragments of kelp fronds Three approximately 20-cm ² pieces of heavy clear plastic approximately 3 mil thick; several littorinid snail shells	W. Walker unpubl. data.
WFP-535	8/9/76	La Jolla, CA Diego County, CA	313	M	Two cellophane cigarette wrappers one rusted fishhook, kelp fronds (<i>Egretta</i> sp.)	W. Walker unpubl. data.

Table 1. --Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
WFP-537	8/31/76	Encinitas, CA	--	F	One blue vinyl plastic strip (3 x 30 cm); kelp frond fragments, one gastropod operculum.	W. Walker unpubl. data.
WFP-565	1/27/77	Del Mar, CA	267	M	One black rubber "bungee cord" with metal hooks at ends (2 x 1 x 40 cm), sand, mollusc shell fragments	W. Walker unpubl. data.
WAW-553	9/2/78	Huntington Beach, CA.	231	F	One partial shoeleace, beach sand, mollusc shell fragments	W. Walker unpubl. data.
JRH-057	5/13/80	La Jolla, CA	251	F	Two plastic bags, one 20 x 20 cm, other partial 40 x 15 cm; kelp fronds (<i>Macrocystis pyrifera</i>), beach sand, gravel, shell fragments	W. Walker unpubl. data. data.
LJH-006	11/14/81	San Diego, CA	236	F	Metal spring (2.0 x 20 cm)	Southwest Fisheries Science Center. ^d
HJB-036	9/3/86	Solana Beach, CA	U	M	Two fishhooks ca. 2.5 cm (1 in) long	Southwest Fisheries Science Center. ^c
SEAN7595	5/6/82	Martha's Vineyard, MA	230	M	Plastic bag in throat	New England Aquarium and U.S. National Museum of Natural History. ^a

Grampus griseus, Risso's dolphin

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
LACM47145	12/8/84	Manhattan Beach, CA	225	--	Blue balloon, partial (20 x 2.5 cm)	Los Angeles County Museum of Natural History. ^d
					<i>Stenella coeruleoalba</i> , striped dolphin	
DAP-001	3/22/83	Cape Point, NC	220	M	Plastic bag in stomach	U.S. National Museum ^a of Natural History, Smithsonian Institution, Wash., D.C.
					<i>Lissodelphis borealis</i> , northern right whale dolphin	
WAW-194	8/2/73	Will Rogers State Beach, Los Angeles County, CA.	211	F	One partial plastic bag in mouth, remainder 25 x 30 cm in forestomach; fronds of marine plants <i>Macrocystis</i> , <i>Cystoseira</i> , and <i>Egregia</i> ; one honey bee (hymenoptera); three white bird feathers. Necropsy diagnosed parasitism of central nervous system as cause of stranding	W. Walker unpubl. data.
					<i>Cystoseira</i> , and <i>Egregia</i> ; one honey bee (hymenoptera); three white bird feathers. Necropsy diagnosed parasitism of central nervous system as cause of stranding	

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
WAW-209	10/4/73	Santa Monica, CA	225	F	Several small bits of blue vinyl plastic (ca. 2 cm ²); one rusted metal bottle cap; 10-12 pieces of kelp fronds (<i>Macrocystis pyrifera</i>). Cause of stranding undetermined.	W. Walker unpubl. data.
<i>Phocoena phocoena</i> , harbor porpoise						
USNM504220	3/1/75	Corolla, NC	--	--	Piece of cloth and plastic in stomach	U.S. National Museum. ^a
<i>Phocoenoides dalli</i> , Dall's porpoise						
LACM54739	7/2/73	Venice Beach, CA	222	M	Stomach jammed with debris as follows: 13 pieces of clear plastic sheets ranging in size from 4 x 9 cm to 35 x 41 cm; 1 piece black plastic 5 x 16 cm; 3 heavy, clear plastic bags 20 x 39 cm; 2 sandwich bags both 14 x 20 cm; 2 plastic bread bags both 23 x 47 cm; 1 plastic drinking straw; 2 pieces of crumpled cardboard approximately 10 x 13 cm; kelp fronds (<i>Macrocystis pyrifera</i>). Necropsy not performed due to autolysed condition of tissues. Cause of stranding undetermined.	Los Angeles County Museum of Natural History ^d and Walker unpubl. data.

Table 1.--Continued.

Specimen No.	Date	Location	Length (cm)	Sex	Description	Information source
WAW-197	8/10/73	Santa Barbara Yacht Harbor, CA	190	F	One blue plastic bottle cap; one 20 x 20 cm plastic bag. The remainder of the fore stomach is jammed with kelp, <i>Macrocystis pyrifera</i> . Necropsy diagnosed parasitic central nervous system disorder as cause of stranding. Animal had been observed in the harbor 3 days prior to stranding.	W. Walker unpubl. data; Cowan et al. 1986.
SBMNH 78-47	11/8/78	Carpenteria State Beach, Santa Barbara County, CA	204	M	Pieces of plastic bags and kelp in stomach	Santa Barbara Museum of Natural History, Santa Barbara, CA.

^aCharleston Museum of Natural History, Charleston, S.C.

^bU.S. National Museum of Natural History, Smithsonian Institution Washington, D.C.

^cSouthwest Fisheries Science Center, National Marine Fisheries Service, La Jolla, Calif.

^dLos Angeles County Museum of Natural History, Los Angeles, Calif.

In one case (WAW-192), a small spaghetti-type mackerel fish tag was found in the stomach. This item was undoubtedly introduced secondarily through ingested prey (*Scomber japonicus*).

Autopsy data were available in eight (18.6%) of the cases (WAW-130, 174, 192, 148, 172, 194, 197, and LACM 72286). Chronic pre-existing disease was present in all instances. In seven of these cases, brain parasitism by the trematode *Nasitrema* sp. was diagnosed as the primary cause of stranding. All these cases occurred in southern California. In two isolated instances (WAW-130 and 197), the animals were observed for up to 10 days inside a boat marina breakwater prior to stranding.

In the southern California area, brain parasitism due to *Nasitrema* sp. has proven to be a common pathologic factor in individual strandings of small cetaceans (Ridgway 1965; Ridgway and Dailey 1972; Dailey and Walker 1978; Cowan et al. 1986). Cowan et al. (1986) found 91% parasitized brains in 44 brains examined in 4 species of stranded cetaceans. No marine debris-related gastrointestinal pathology was evident in any of the 23 southern California strandings summarized in Table 1.

Naturally occurring disease factors may predispose these animals to ingest abnormal objects. The high incidence of pre-existing brain parasitism and the absence of debris-induced gastrointestinal pathology suggest that the significance of marine debris in stranded cetaceans should remain questionable unless accompanied by related pathologic changes and a complete necropsy and tissue analysis of all major organ systems.

Marine Debris Encountered in Food Habits Analyses of Free-Ranging Animals

Data on 10 species of odontocete cetaceans were available (Table 2). All species combined, a total of 1,790 stomachs had been examined. The geographic regions covered in the sample are diverse. Localities in the North Pacific and Bering Sea represented 81.5% of the sample. The Okhotsk Sea represented 5.9% and the remaining 12.6% were collected off the coast of Uruguay in the South Atlantic. Of the 10 species of cetaceans reported, only the Baird's beaked whale, *Berardius bairdii*, taken at 2 different localities in the coastal waters of Japan had ingested debris. In 86 *B. bairdii* taken off Chiba Prefecture, central Japan, the frequency of debris ingestion was 26.7%. A lower incidence of debris was evident in 20 *Berardius* examined from the southern Okhotsk Sea, where frequency of ingestion was 15.0%. Overall frequency for both areas sampled was 24.5%.

The nature of debris material ingested by *B. bairdii* from both regions was diverse (see Tables 3 and 4 for detailed accounts). Occurrence of plastic bags and sheeting was 30.8%. Other plastic articles including discarded fishing gear had a frequency of 11.5%. All plastic material combined was 42.3%. Miscellaneous nonplastic material such as vegetable refuse, wood boards, concrete fragments, pieces of glass, cigarette filters, cellophane, rubber material, a roof tile fragment, bottle caps, rusty hinge, aluminum can pull tabs, and a metal butane lighter top had an occurrence of 76.0%. Observations made during collection of stomach

content samples revealed no debris-associated lesions or evidence of impaction.

Examination of the data in Tables 3 and 4 reveals that the major portion of debris found in *B. bairdii* stomachs were negatively buoyant items probably ingested at or near the ocean bottom.

Differences in the frequency of debris ingestion between the Pacific coast of central Japan (26.7%) and the southern Okhotsk Sea (15.0%) are probably due to regional differences in feeding strategy. Off the Pacific coast of central Japan, *Berardius* are known to feed primarily on benthic prey. In this region they are documented to be feeding on bottom-dwelling morid and macrourid fishes (81.7%). Cephalopods represented only 18.0% of the consumed prey. In addition, stones and gravel were encountered in all stomachs examined and were undoubtedly consumed incidentally during bottom feeding (Walker and Mead 1988). In the southern Okhotsk Sea, *Berardius* feeding strategy changes considerably. In this region, cephalopod prey are dominant (87.6%) in the food habits sample. The predominantly benthic morid and macrourid fishes represent only 8.2% of the prey. Stones and gravel were encountered in only 10.0% of the stomachs examined (W. A. Walker unpubl. data).

The other nine species of cetaceans summarized in Table 2 are known to feed primarily in the epipelagic and mesopelagic zones in the upper water column. Debris occurrence in all 1,684 stomachs examined was 0. The stomach sample for these nine species is small compared to conservative, stock-level, population estimates. As a result, the absence of marine debris in the species summarized in Table 2 is inconclusive. However, some inference can be made from the Dall's porpoise, *Phocoenoides dalli*, samples from the northern North Pacific and Bering Sea. Both these regions are documented to have a high density of marine debris (Venrick et al. 1973; Feder et al. 1978; Shaw and Mapes 1979; Dahlberg and Day 1985). Eight hundred fifteen *Phocoenoides* stomachs were examined from these regions and no debris items were encountered.

DISCUSSION

Accounts of mortality and pathology caused by foreign body ingestion in captive cetaceans leave little doubt as to the potential effects of marine debris in wild, free-ranging cetaceans.

Most of the available records of debris ingestion are from stranded odontocete cetaceans. However, debris ingestion in singly stranded animals may be, in a large percentage of cases, part of the stranding syndrome. Pre-existing disease factors related to parasitism occurred in almost all cases accompanied by complete necropsy observations.

Debris ingestion data on free-ranging animals derived from previously conducted food habits studies were, with the exception of Baird's beaked whale, negative. Of the 10 species summarized in Table 2, *B. bairdii* is the only species of cetacean known to demonstrate regionally varying degrees of deepwater bottom-feeding strategy. All the remaining nine

Table 2.--Summary of ingested marine debris encountered in food habits analyses of free-ranging small cetaceans.

Species	Date	General location	Collection method	Sample size	Debris occurrence	Information source
<i>Berardius bairdii</i>	1988-89	Taken in the southern Okhotsk Sea off northern Hokkaido, Japan	Shore-based harpoon fishery	20	15.0%	W. Walker unpubl. data. For detailed summary see Table 4.
<i>Berardius bairdii</i>	1985-87	Taken off the Pacific coast of central Japan	Shore-based harpoon fishery	86	26.7%	W. Walker and J. G. Mead unpubl. data. Food habits data in Walker and Mead 1988. For detailed summary see Table 3.
<i>Steno bredanensis</i>	1977-87	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna fishery	16	0	W. Walker and J. G. Mead unpubl. data.
<i>Lagenorhynchus obliquidens</i>	1979-72	Collected off the coasts of Washington and California	Collected at sea for research purposes	44	0	C. Fiscus, food habits data presented in Stroud et al. 1981.
<i>Delphinus delphis</i>	1977-80	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna purse seine fishery	32	0	Southwest Fisheries Science Center, La Jolla, CA, unpubl. data.
<i>Tursiops truncatus</i>	1972-87	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna purse seine fishery	35	0	W. Walker unpubl. data; food habits data presented (in part) in Walker 1981.
<i>Stenella attenuata</i>	1968-77	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna purse seine fishery	231	0	Southwest Fisheries Science Center, La Jolla, CA, food habits data (in part) published in Perrin et al. 1973.

Table 2.---Continued.

Species	Date	General location	Collection method	Sample size	Debris occurrence	Information source
<i>Stenella coeruleoalba</i>	1977-80	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna purse seine fishery	104	0	Southwest Fisheries Science Center, La Jolla, CA, unpubl. data.
<i>Stenella longirostris</i>	1968-80	Offshore waters of the eastern tropical Pacific	Incidental take in the yellowfin tuna purse seine fishery	78	0	Southwest Fisheries Science Center, La Jolla, CA, published (in part) in Perrin et al. 1973.
<i>Phocoenoides dalli</i>	1988	Okhotsk Sea, Japan ca. lat. 44#10'N, long. 144#30'E	Shore-based hand-harpoon fishery	86	0	W. Walker unpubl. data.
<i>Phocoenoides dalli</i>	1979-86	Northern North Pacific and Bering Sea	Incidental take of Japanese high seas salmon gillnet fishery	815	0	T. Crawford and L. Tsunoda, Northwest and Alaska Fisheries Center, National Marine Mammal Laboratory, Seattle, Wash. Food habits data presented (in part) in Crawford 1981.
<i>Phocoenoides dalli</i>	1958-72	Collected off the coasts of Washington and California	Collected at sea for research purposes	17	0	C. Fiscus; food habits data in Stroud et al. 1981.
<i>Pontoporia blainvilli</i>	1969-75	Off the coast of Uruguay, South America (ca. lat. 34#30'S)	Incidental take in local shark gillnet fisheries	226	0	R. Brownell, Jr. and W. Walker. Prey species accounts on 11 animals in Fitch and Brownell 1971.

Table 3.--Summary of ingested marine debris in 86 *Berardius bairdii* taken at Wadaura, Chiba Prefecture, Japan, 1985-87.

Specimen No.	Date	Length (m)	Sex	Age (year)	Description of debris
85-008	7/23/85	10.10	M	23	Vegetable refuse--approximately 1 dozen coffee beans.
85-015	7/28/85	9.85	M	73	Three small glass fragments (two clear, one brown) approximately 1.5 x 2 x 0.5 cm, edges worn; two cigarette filters.
85-017	7/29/85	10.05	F	23	One cigarette filter and a piece of tree bark.
85-018	7/30/85	10.43	M	31	One No. 2 size rusted fishhook.
85-021	8/1/86	10.00	M	51	One piece of wadded-up longline approximately 15 cm diameter with 15-20 rusted No. 2 size hooks. The main lines and branch lines are made up of braided No. 7 nylon net twine with hooks set approximately 120 cm apart. Condition of this object suggests recent ingestion of discarded fishing gear.
85-022	8/1/85	9.90	M	21	One piece of black plastic (25 x 15 cm) approximately 1.5 mil thick and vegetable refuse--two corn kernels, <i>Zea maise</i> .
85-023	8/2/85	10.70	F	54	One fishhook--only rusted shank and small portion of leader remain.
85-024	8/2/85	9.90	M	84	One piece of black vinyl plastic (130 x 135 x 0.3 cm).
85-026	8/3/85	9.65	M	8	One 45 x 3 cm mahogany stick with staples.
85-033	8/6/85	9.62	M	70	One 20 x 15 cm thin plastic sheet (food wrapper?); cellophane package material (8 x 6 cm). Vegetable refuse--one undigested potato (5 x 6 x 3 cm) two pieces of tree bark (3-4 x 5-6 cm).

Table 3.--Continued.

Specimen No.	Date	Length (m)	Sex	Age (year)	Description of debris
85-031	8/5/85	9.50	M	60	One 8 × 10 × 0.5 cm irregular-shaped piece of clear plastic (PVC?).
86-004	7/27/86	10.75	F	22	One 10 × 15 × 1.0 cm piece of pine board.
86-011	7/29/86	10.40	F	53	One 12 × 2 × 1 cm piece of wood; one 20 × 30 cm black plastic sheet approximately 3 mil thick.
86-012	7/30/86	9.70	M	17	Two fragments of clear plastic 4 × 3 cm and 6 × 2 cm, both approximately 3 mil thick.
86-020	8/7/86	9.10	F	8	One 5 × 6 × 2 cm fragment of blue glazed roofing tile, two cigarette filters.
86-026	8/9/86	10.20	M	26	One 10 × 10 cm piece of concrete; one bottle cap; one 6 × 6 × 2 cm piece of tree bark.
86-028	8/10/86	9.70	M	38	One fragment of 20 × 15 × 0.3 cm rubber mat and 3-4 bird feathers.
87-013	7/29/87	10.32	M	56	Vegetable refuse--25-30 soybeans; one badly rusted metal hinge.
87-014	7/29/87	10.80	F	40	One 3 × 4 × 3 concrete fragment; two pieces of cellophane (both approximately 8 × 15 cm).
87-015	8/1/87	10.60	F	19	Vegetable refuse--10-15 soybeans, 8 corn kernels, <i>Zea maise</i> ; 1 aluminum pull tab.
87-016	8/1/87	10.20	F	17	One corroded fishhook; one metal portion (top) of a butane cigarette lighter.
87-017	8/2/87	10.35	M	68	Four clear glass fragments 2 to 4 cm ² and approximately 0.3 cm thick, all worn smooth on edges.

Table 3.--Continued.

Specimen No.	Date	Length (m)	Sex	Age (year)	Description of debris
87-021	8/10/87	10.40	F	51	One piece of wood with nail (not protruding) 17.5 × 4.5 × 7.0 cm (weight 130 g).
87-023	8/13/87	10.30	F	44	One large concrete fragment 9.5 × 6.5 × 3.0, weight 230 g; one piece of 3 mil thick black plastic 18 × 23 cm.
87-024	8/14/87	10.58	M	41	One fragment of brown glass, 4 × 2 × 0.3 cm, edges smooth, one piece of brown plastic sheet 30-35 cm ² .

species summarized are known to feed primarily in the epipelagic and mesopelagic zones.

Experimental evidence suggests that odontocete cetaceans are probably very discriminating feeders. Mistaken ingestion of oceanic debris due to its resemblance to preferred prey species is unlikely because of odontocete cetacean echolocation capabilities. In captive experiments, the bottlenose dolphin, *Tursiops truncatus*, has been shown to be capable of making fine discriminations in size, shape, texture, and composition of objects (Kellogg 1958, 1959a, 1959b; Norris et al. 1961; Evans and Powell 1967; Norris 1969). Kellogg (1959a) demonstrated the use of echolocation by *T. truncatus* to locate preferred food fish but avoid inedible objects. Selection between a water-filled 2-cm gelatin capsule and an equal-sized piece of cut fish has been demonstrated (Norris et al. 1961). Evans and Powell (1967) determined that *T. truncatus* could discriminate between identical-sized sheets of different metals, or even between sheets of the same metal but of different thicknesses. Monofilament about 1 mm in diameter was determined to be at the threshold of detection for a captive harbor porpoise, *Phocoena phocoena* (Busnel et al. 1965).

The two species of free-ranging odontocete cetaceans documented to ingest marine debris are the sperm whale *Physeter macrocephalus*, and the Baird's beaked *B. bairdii*. Both these cetaceans are known to spend some time feeding at or near the bottom, particularly in coastal waters. This is verified by the behavior of preferred prey species and by the common occurrence of stones and gravel in examined stomachs (Betesheva and Akimushkin 1955; Nemoto and Nasu 1963; Tomilin 1967; Berzin 1971; Walker and Mead 1988). Ingestion of debris in these two species of cetaceans is very likely to be incidental, the debris being consumed along with bottom-dwelling prey.

Table 4.--Summary of ingested marine debris in 20 *Berardius bairdii* taken at Abashiri, Hokkaido, Japan, 1988-89.

Specimen No.	Date	Length (m)	Sex	Description of debris
Ab-88-03	8/28/88	10.70	M	One No. 2 size fishhook with short, approximately 15 cm, portion of leader attached.
Ab-88-08	9/7/88	10.10	M	Two No. 2 size fishhooks.
Ab-88-18	9/24/88	10.20	F	One badly rusted, partial No. 2 size fishhook.
89-HK-101	9/1/89	9.40	M	One cotton sleeve from rubber glove, black rubber fragments still attached to anterior edges.
89-HK-102	9/2/89	10.78	F	One No. 2 fishhook and three ca. 30 cm ² portions of clear plastic sheeting. All appear to be portions of the same material and are ca. 3 cm thick. Vegetable refuse--three pieces of citrus fruit (orange?) peels.
89-HK-104	9/9/88	10.20	M	One ca. 10-cm diameter wad of thin, clear plastic sheeting ca. 1.5 mil thick.

Records presented in this report of marine debris ingested by *Berardius* off central Japan all came from off the Boso Peninsula, Chiba Prefecture, an area extending from the northern edge of the entrance to Tokyo Bay north to Choshi off the Pacific coast of Japan (lat. 34°39'-35°57'N). The animals were taken primarily along the 1,000-m depth contour line. Due to the proximity to Tokyo Bay and Choshi (a major fishing port), the entire area is subject to extremely heavy merchant and commercial fishing vessel traffic. The *Berardius* stomach samples from off northern Hokkaido were from two general areas in the southern Okhotsk Sea: 1) The immediate vicinity of the major commercial fishing port of Abashiri (ca. lat. 44°30'N, long. 144°30'E) and 2) Nemuro Strait between the Shiretoko Peninsula (Japan) and Kunashir Island (U.S.S.R.) (ca. lat. 44°15'N, long. 145°30'E). Both these areas are also subject to heavy commercial fishing activity.

Debris encountered in *Berardius* stomachs from both the Pacific coast of Japan and the southern Okhotsk Sea consisted almost exclusively of negatively bouyant material. The varied nature of the debris (e.g., broken glass, vegetable refuse, aluminum pull tabs, bottle caps) and the deep

offshore location strongly suggest shipboard refuse as the primary source of the ingested debris.

It should be noted that the absence of debris-related pathology in the Baird's beaked whale sample does not rule out the occurrence of debris-related mortality in the study areas. To the contrary, the high incidence of ingested debris, the varied nature of the debris material, and records of debris-related mortality in captive cetaceans permit some speculation on the potential for incidental debris ingestion as a mortality factor in this species. *Berardius* is a large animal (up to 12.8 m in length). Death or debilitation through gastrointestinal impaction would probably require a relatively large volume of indigestible material. However, some of the debris material summarized in Tables 3 and 4 has considerable pathologic potential. Ingestion of relatively small items such as sharp metallic objects or freshly broken glass poses a real identifiable hazard and is a well-documented factor in human as well as veterinary disease. Complications through mechanical trauma to the gastrointestinal tract by such objects range from laceration and hemorrhage to perforation of the gut wall and acute bacterial peritonitis. These conditions are eminently life threatening. In the wild state, the elapsed time from onset to death would probably be short and accompanied by marked behavioral alterations. As a result, the probability of encountering these acute states in the fisheries sample would be very low. In the commercial fisheries sample we should expect only chronic, prolonged disease conditions to be manifest. Some bias toward the taking of chronically ill whales in commercial harpoon fisheries is suspected. Walker (1988) suggested, on the basis of a parasite survey, that the prolonged chase procedure in the Japanese *Berardius* fishery may involve selection toward the taking of some physically infirm individuals.

Wallace (1985) raised the question of whether sinking truly constituted debris removal. Results of this study indicate that negatively buoyant debris are not neutral, but still may pose a potential hazard to large predators feeding on benthic prey.

RECOMMENDATIONS

The worldwide trend toward curtailment of the commercial take of marine mammals restricts opportunities for future food habits sampling. In the future, food habits studies are likely to become more reliant on samples obtained from incidental fisheries' interactions and strandings. The following recommendations should be incorporated into future research:

- Develop and incorporate a consistent format for recording the presence as well as the absence of marine debris in stomach content analyses conducted on both free-ranging and stranded marine mammals. The recording of negative findings should also be emphasized, as they are crucial in establishing reliable information on the frequency of debris occurrence in future studies.
- Whenever possible, occurrence of ingested debris should be

of the debris should also be kept (e.g., size, shape, consistency). In the case of stranded animals, the necropsy should include examination and evaluation of all major organ systems. Data presented in this report suggest that ingestion of debris may be secondary to other naturally occurring disease factors.

- The filter-feeding strategy of the baleen whales, Mysticeti, may make them particularly vulnerable to incidental debris ingestion in both the benthic zone and zones of the upper water column. Researchers should be encouraged to take advantage of the few remaining commercial fisheries to record evidence of ingested debris. The stomachs and oral cavities of stranded baleen whales should be examined whenever possible. Individuals involved in past food habit studies of baleen whale species should be interviewed.
- Research on marine mammal food habits as well as debris ingestion should continue in order to establish preferred prey species and feeding behavior, and increase the data base on debris ingestion.
- The data presented in this report are limited to gross observations on the acute effects of ingested debris (e.g., gastrointestinal impaction, ulceration). Research into the potentially insidious effects of absorption of hydrocarbon contaminants such as plasticizers should be conducted.

ACKNOWLEDGMENTS

The institutions which provided data are acknowledged in Tables 1 and 2. I thank the following people for contributing additional data or assistance in compiling this report: R. L. Brownell, Jr., T. W. Crawford, M. E. Dahlheim, C. H. Fiscus, J. W. Gilpatrick, Jr., J. E. Heyning, L. L. Jones, T. Kasuya, S. K. Lafferty, J. G. Mead, Y. Mori, S. F. Noel, W. F. Perrin, Y. Shimomichi, H. Shogi, J. Taguchi, L. M. Tsunoda, and C. D. Woodhouse. The Baird's beaked whale research was partially funded by The Institute for Cetacean Research, Tokyo, Japan. This report was funded by the Marine Entanglement Research Program, National Marine Fisheries Service (Contract No. 40ABNF6-3361).

REFERENCES

- Amemiya, I.
1962. The dolphin that swallowed a football. Intern. Zoo Yearbook 4:34.
- Araya, H.
1983. Fishery biology and stock assessment of *Ommastrephes bartrami* in the North Pacific Ocean. Mem. Natl. Mus. Victoria 44:269-283.

- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Berzin, A. A.
1959. O pitanii kashalota v Beringovom more (Feeding of the sperm whale in the Bering Sea). Izvestiya TINRO 47.

1971. The sperm whale. In A. V. Yablokov (editor), Pishchevaya Promyshlennost, Moscow. (Translation by Israel Program for Scientific Translations, 1972, 394 p.)
- Betesheva, E. I., and I. I. Akimushkin.
1955. Food of the sperm whale (*Physeter catadon*) in the Kurile Islands region. Tr. Inst. Okeanol. 18:86-94. [In Russ.]
- Brown D. H., D. K. Caldwell, and M. C. Caldwell.
1966. Observations on the behavior of wild and captive false killer whales, with notes on associated behavior of other genera of captive delphinids. Los Ang. Cty. Mus. Contrib. Sci. 95:1-32.
- Brown, D. H., R. W. McIntyre, C. A. Delli Quadri, and R. J. Schroeder.
1960. Health problems of captive dolphins and seals. J. Am. Vet. Med. Assoc. 137:534.
- Busnel, Renee-Guy, A. Dziedzic, and S. Andersen.
1965. Seuils de perception du systeme sonar du Marsoiun *Phocaena phocaena* L., en fonction du diametre d'un obstacle filiforme. C. R. Acad. Sci. Paris 260:295-297.
- Caldwell, D. K., M. C. Caldwell, and D. W. Rice.
1966. Behavior of the sperm whale, *Physeter catadon* L. In K. S. Norris (editor), Whales, dolphins, and porpoises, p. 678-716. Univ. Calif. Press, 789 p.
- Caldwell, M. C., D. K. Caldwell, and J. B. Siebenaler.
1965. Observations on captive and wild Atlantic bottlenosed dolphins, *Tursiops truncatus*, in the northeastern Gulf of Mexico. Los Ang. Cty. Mus. Contrib. Sci. 91:1-10.
- Caldwell, M. C., D. K. Caldwell, and B. C. Townsend, Jr.
1968. Social behavior as a husbandry factor in captive cetaceans. Proceedings of the Second Symposium on Diseases and Husbandry of Aquatic Mammals, St. Augustine, Florida, p. 1-9. Marineland Research Laboratory.
- Carpenter, E. J., and K. L. Smith, Jr.
1972. Plastics on the Sargasso Sea surface. Science (Wash. D.C.) 175:1240-1241.
-

Cawthorn M. W.

1985. Entanglement in, and ingestion of, plastic litter by marine mammals, sharks, and turtles in New Zealand waters. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 336-343. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Clarke, R.

1956. Sperm whales of the Azores. Discovery Rep. 28:98-237.

Cowan, D. R., W. A. Walker, and R. L. Brownell, Jr.

1986. Pathology of small cetaceans stranded along southern California beaches. In M. M. Bryden and R. Harrison (editors), Research on dolphins, p. 323-367. Oxford Univ. Press, Oxford, 478 p.

Crawford, T. W.

1981. Vertebrate prey of *Phocoenoides dalli*, (Dall's porpoise), associated with the Japanese high seas salmon fishery in the North Pacific Ocean. M.S. Thesis. Univ. Wash. College of Fisheries, 72 p.

Dahlberg, M. L., and R. H. Day.

1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Dailey, M. D., and W. A. Walker.

1978. Parasitism as a factor (?) in single strandings of southern California cetaceans. J. Parasitol. 64:593-596.

Day, R. H., D. H. S. Wehle, and F. C. Coleman.

1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Eldridge, I. M.

1982. Sea-lane litter survey. Environ. Conserv. 164.

Evans, W. E., and B. A. Powell.

1967. Discrimination of metallic plates by an echolocating delphinid. Proc. Symp. Bionic Models of Animal Sonar System, Frascati, Italy, 1966, p. 363-398. Labor. d'Acoustique Animale, Jouy-en-Josas, France.

Feder, H. M., S. C. Jewett, and J. R. Hilsinger.

1978. Man-made debris on the Bering Sea floor. Mar. Pollut. Bull. 9:52-53.

- Fitch, J. E., and R. L. Brownell, Jr.
1971. Food habits of the franciscana, *Pontoporia blainvillei*, (Cetacea: Platanistidae) from South America. *Bull. Mar. Sci.* 21:626-636.
- Hollis, H.
1939. Biological report of the United States Bureau of Fisheries. *Norsk Hvalf. Tid.* 1.
- Kellogg, W. N.
1958. Echo ranging in the porpoise. *Science (Wash. D.C.)* 128:982-988.
1959a. Auditory perception of submerged objects by porpoises. *J. Acoust. Soc. Am.* 31:1-6.
1959b. Size-discrimination by reflected sound in a bottlenose dolphin. *J. Comp. Physiol. Psychol.* 52:509-514.
- Machida, S.
1983. A brief review of the squid fishery by *Hoyo Maru No. 67* in southeast Australian waters in 1979/80. *Mem. Natl. Mus. Victoria* 44:291-295.
- Mate, B. R.
1985. Incidents of marine mammal encounters with debris and active fishing gear. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii*, p. 453-457. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Millais, J. S.
1906. *The mammals of Great Britain and Ireland.* London 3:215-348.
- Morris, R. J.
1980a. Floating plastic debris in the Mediterranean. *Mar. Pollut. Bull.* 11:125.
1980b. Plastic debris in the surface waters of the South Atlantic. *Mar. Pollut. Bull.* 11:164-166.
- Nakajima, M., K. Sawaura, A. Fujimoto, and T. Oda.
1965. Foreign bodies in the stomachs of the captive dolphins. *Enoshima Marineland Rep.* 2:27.
- Nemoto, T., and K. Nasu.
1963. Stones and other aliens in the stomachs of the sperm whales in the Bering Sea. *Sci. Rep. Whales Res. Inst.* 17:83-91.
- Norris, K. S.
1969. The echolocation of marine mammals. In H. T. Andersen (editor), *The biology of marine mammals*, p. 391-421. Acad. Press, Lond. and N.Y., 511 p.
-

- Nemoto, T., and K. Nasu.
1963. Stones and other aliens in the stomachs of the sperm whales in the Bering Sea. *Sci. Rep. Whales Res. Inst.* 17:83-91.
- Norris, K. S.
1969. The echolocation of marine mammals. In H. T. Andersen (editor), *The biology of marine mammals*, p. 391-421. Acad. Press, Lond. and N.Y., 511 p.
- Norris, K. S., J. H. Prescott, P. V. Asa-Dorian, and P. Perkins.
1961. An experimental demonstration of echolocation behavior in the porpoise, *Tursiops truncatus* (Montagu). *Biol. Bull. (Woods Hole)* 120:
- O'Hara, K. J., S. Iudicello, and R. Bierce.
1988. A citizen's guide to plastics in the ocean: More than a litter problem. Center for Environmental Education, Wash., D.C., 131 p.
- Perrin, W. F., R. R. Warner, C. H. Fiscus, and D. B. Holts.
1973. Stomach contents of porpoise, *Stenella* spp., and yellowfin tuna, *Thunnus albacares*, in mixed-species aggregations. *Fish. Bull., U.S.* 71:1077-1096.
- Pike, G.
1950. Stomach contents of whales caught off the coast of British Columbia. *Progr. Rep. Pacific Coast Stat. Fish. Res. Board Can.* 83.
- Ridgway, S. H.
1965. Medical care of marine mammals. *J. Vet. Med. Assoc.* 147:1077.
1972. Homeostasis in the aquatic environment. In S. H. Ridgway (editor), *Mammals of the sea: Biology and medicine*, p. 590-731. Charles C. Thomas, Springfield, Illinois, 812 p.
- Ridgway, S. H., and M. D. Dailey.
1972. Cerebral and cerebellar involvement of trematode parasites in dolphins and their possible role in stranding. *J. Wildl. Dis.* 8:33-43.
- Shaw, D. G., and G. A. Mapes.
1979. Surface circulation and the distribution of pelagic tar and plastic. *Mar. Pollut. Bull.* 10:160-162.
- Sleptsov, M. M.
1952. Kitoobraznye dal'nevostochnykh morei (Whales of the Far East). *Izvestiya TINRO* 38.
- Stroud, R. K., C. H. Fiscus, and H. K. Kajimura.
1981. Food of the Pacific white-sided dolphin, *Lagenorhynchus obliquidens*, and northern fur seal, *Callorhinus ursinus*, off California and Washington. *Fish. Bull., U.S.* 78:951-959.


- Tomilin, A. G.
1967. Mammals of the U.S.S.R. and adjacent countries (1957).
Jerusalem, Israel Program for Scientific Translations 9, 717 p.
- Turner, W.
1903. The occurrence of the sperm whale or cachalot in the Shetland Seas, with notes on the tympanopetrous bones of *Physeter*, *Kogia* and other odontoceti. Proc. Roy. Soc. Edinburgh 24:632-644.
- Van Dolah, R. F., V. G. Burrell, Jr., and S. B. West.
1980. The distribution of pelagic tars and plastics in the South Atlantic Bight. Mar. Pollut. Bull. 11:352-356.
- Venrick, E. L., T. W. Backman, W. C. Bartram, C. J. Platt, M. S. Thornhill, and R. E. Yates.
1973. Man-made objects on the surface of the North Pacific Ocean. Nature (Lond.) 241:271.
- Walker, W. A.
1981. Geographical variation in morphology and biology of bottlenose dolphins (*Tursiops*) in the eastern North Pacific. Southwest Fish. Cent., Natl. Mar. Fish. Serv., NOAA, La Jolla, Calif. Southwest Fish. Cent. Admin. Rep. LJ-81-03C, 24 p.

1988. Preliminary report on the parasites and pathology of Baird's beaked whales taken off the Pacific coast of central Japan 1985-1987. Int. Whaling Comm., Small Cetacean Subcommittee Doc. No. SC/40/SM15, 13 p.
- Walker, W. A., and J. G. Mead.
1988. Preliminary report on the food habits of Baird's beaked whales taken off the Pacific coast of central Japan 1985-1987. Int. Whaling Comm., Small Cetacean Subcommittee Doc. No. SC/40/SM16, 8 p.
- Wallace, N.
1985. Debris entanglement in the marine environment: A review. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 259-277. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Wehle, D. H. S., and F. C. Coleman.
1983. Plastics at sea. Nat. Hist. 92(2):20-26.
- Wong, C. S., D. R. Green, and W. J. Cretney.
1974. Quantitative tar and plastic waste distributions in the Pacific Ocean. Nature (Lond.) 246:30-32.
-

RECENT TECHNICAL MEMORANDUMS

Copies of this and other NOAA Technical Memorandums are available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22167. Paper copies vary in price. Microfiche copies cost \$4.50. Recent issues of NOAA Technical Memorandums from the NMFS Southwest Fisheries Center are listed below:


- NOAA-TM-NMFS-SWFSC-144 Atlas of eastern tropical Pacific oceanographic variability and cetacean sightings, 1986-1989.
P.C. FIELDER, L.J. LIERHEIMER, S.B. REILLY, S.N. SEXTON, R.S. HOLT
and D.P. DEMASTER
(July 1990)
- 145 Trends in landings, species composition, length-frequency distributions, and sex ratios of 11 rockfish species (Genus *Sebastes*) from central and northern California ports (1978-88).
D.E. PEARSON and S. RALSTON
(July 1990)
- 146 Field manual for phocid necropsies (specifically *Monachus schauinslandi*).
J. M. WINCHELL
(July 1990)
- 147 Survey of the abundance and distribution of pelagic young-of-the-year rockfishes, *Sebastes*, off central California.
T. W. ECHEVERRIA, W. H. LENARZ and C. A. REILLY
(September 1990)
- 148 United States Agency for International Development and National Marine Fisheries Service workshop on tropical fish stock assessment, 5-26 July 1989, Honolulu, Hawaii.
J. J. POLOVINA and R. S. SHOMURA
(September 1990)
- 149 Summary of the 1988 U.S. tuna/porpoise observer data.
A. R. JACKSON
(September 1990)
- 150 Population monitoring of the Hawaiian monk seal, *Monachus schauinslandi*, and captive maintenance project at Kure Atoll, 1988.
J. R. HENDERSON and M. R. FINNEGAN
(September 1990)
- 151 The Hawaiian monk seal on Laysan Island, 1988.
T. C. JOHANOS, B. L. BECKER, M. A. BROWN,
B. K. CHOY, L. M. HURUKI, R. E. BRAINARD and
R. L. WESTLAKE
(September 1990)
- 152 A personal computer based system for analog-to-digital and serial communication data acquisition.
R. C. HOLLAND
(November 1990)
- 153 The nearshore physical oceanography off the central California coast during May-June, 1989: A summary of CTD from juvenile rockfish surveys.
F. B. SCHWING, S. RALSTON, D. M. HUSBY
and W. H. LENARZ
(December 1990)



file copy


NOAA Technical Memorandum NMFS

DECEMBER 1990




**PROCEEDINGS OF THE SECOND INTERNATIONAL
CONFERENCE ON MARINE DEBRIS
2-7 APRIL 1989, HONOLULU, HAWAII**

VOLUME II



Richard S. Shomura
Mary Lynne Godfrey
(Editors)



NOAA-TM-NMFS-SWFSC-154

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Southwest Fisheries Science Center
University of Hawaii Sea Grant College Program

NOAA Technical Memorandum NMFS

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.

NOAA Technical Memorandum NMFS

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information; and have not received complete formal review, editorial control, or detailed editing.

DECEMBER 1990

**PROCEEDINGS OF THE SECOND INTERNATIONAL
CONFERENCE ON MARINE DEBRIS
2-7 APRIL 1989, HONOLULU, HAWAII**

Richard S. Shomura and Mary Lynne Godfrey (Editors)
University of Hawaii
School of Ocean and Earth Science and Technology
Hawaii Institute of Marine Biology
1000 Pope Road
Honolulu, Hawaii 96822

Canada Department of Fisheries and Oceans
Council for Solid Waste Solutions
Intergovernmental Oceanographic Commission (UNESCO)
National Coastal Resources Research and Development Institute (U.S.A.)
Pacific Rim Fishing Industries
Sea Grant College Program, University of Hawaii
SUNY, Stony Brook, Marine Sciences Research Center's
Waste Management Institute
School of Ocean and Earth Science and Technology, University of Hawaii
U.S. Environmental Protection Agency
U.S. Fish and Wildlife Service
U.S. Marine Mammal Commission
U.S. Minerals Management Service
U.S. National Oceanic and Atmospheric Administration
U.S. Navy

NOAA-TM-NMFS-SWFSC-154

U.S. DEPARTMENT OF COMMERCE
Robert A. Mosbacher, Secretary
National Oceanic and Atmospheric Administration
John A. Knauss, Under Secretary for Oceans and Atmosphere
National Marine Fisheries Service
William W. Fox, Jr., Assistant Administrator for Fisheries

TABLE OF CONTENTS


	Page
SESSION IV. ECONOMIC IMPACTS ON VESSELS AND SHORELINES	775
N. F. Meade, K. M. Drazek, and V. R. Leeworthy	777
An economic perspective on the problem of marine debris	
S. Takehama	792
Estimation of damages to fishing vessels caused by marine debris, based on insurance statistics	
R. E. Weisbrod	810
New York State marine debris program [Abstract]	
K. D. Wagner	811
Medical wastes and the beach washups of 1988: Issues and impacts	
SESSION V. SOLUTIONS THROUGH TECHNOLOGY	825
G. Scott	827
The philosophy and practice of degradable plastics	
A. L. Andrady	848
Environmental degradation of plastics under land and marine exposure conditions	
M. Xanthos, P. G. Kelleher, A. Patel, and W. G. Gordon	870
Recycling of marine plastics debris through melt reprocessing: A case for lost or abandoned fishing gear	
C. S. Alig, L. Koss, T. Scarano, and F. Chitty	879
Control of plastic wastes aboard naval ships at sea	
L. A. Martinez	895
Shipboard waste disposal: Taking out the trash under the new rules	
T. J. Chang	915
Low technology (burn barrel) disposal of shipboard-generated (MARPOL V) wastes	
F. Recht and S. Lasseigne	921
Providing refuse reception facilities and more: The port's role in the marine debris solution	

M. Aizawa and A. Satou	935
Disposition and recycling of plastic products including used nets	
SESSION VI. SOLUTIONS THROUGH LAW AND POLICY	945
D. P. Redford	947
Status of the U.S. Environmental Protection Agency marine debris activities and programs	
D. T. Edwards and E. Rymarz	956
International regulations for the prevention and control of pollution by debris from ships	
M. J. Bean	989
Redressing the problem of persistent marine debris through law and public policy: Opportunities and pitfalls	
R. E. Seebald	998
Implementation and enforcement of Annex V of MARPOL 73/78 in the United States [Abstract]	
C. A. Crampton	999
The Coast Guard's Annex V compliance report: A case study	
J. Sawaki and I. Yamaguchi	1008
Control regulations for discharge of onboard wastes from ships	
R. Rose	1020
Marine plastic debris: What Washington State has done	
G. Peet	1034
The protection of specific sea areas against marine debris	
L. M. Paul	1045
Using the protective principle to unilaterally enforce transnational marine pollution standards	
SESSION VII. SOLUTIONS THROUGH EDUCATION	1075
R. G. Bruner	1077
The plastics industry and marine debris: Solutions through education	
T. R. Dixon	1090
MARPOL 73/78 information, education, and training: Meeting the challenge	




P. Debenham	1100
Education and awareness: Keys to solving the marine debris problem	
B. Wallace	1115
Shipping industry marine debris education plan	
N. Yagi and Y. Otsuka	1123
Cleanup program in Japan	
L. Koss, F. Chitty, and W. A. Bailey	1132
U.S. Navy's plastics waste educational efforts	
B. Wallace	1140
How much do commercial and recreational fishermen know about marine debris and entanglement? Phase 1	
J. Neilson	1149
The Oregon experience--four years later	
A. Farias	1154
The Texas Adopt-A-Beach Program: A public/private approach to clean beaches	
J. Humphreys and P. Mullin	1165
Marine debris demonstration and education project at Squalicum Harbor, Bellingham, Washington, U.S.A.	
S. Friday	1174
Marine debris: North Carolina's solutions through education	
S. Laska	1179
Designing effective educational programs: The attitudinal basis of marine littering	
POSTER PRESENTATION	1191
G. J. Vauk and E. Schrey	1193
Animals as litter victims at the German North Sea coast	
J. N. Kushima and R. P. Clarke	1194
Marine debris and epipelagic fishes	
VIDEO PRESENTATION	1195
A. F. Amos	1197
Portrait of a barrier island beach	

Offshore Operators Committee	1198
U.S. oil industry efforts in addressing beach debris problem	
Government of Japan	
Plastic pollution: The sea is crying	
Government of Japan	
Cleanup operation of the sea floor	
Government of Japan	
Japanese research activities on marine debris	
WORKING GROUP REPORTS	1199
Report of the working group on methods to	1201
assess the amount and types of marine debris	
(C. A. Ribic, Chair)	
Report of the working group on entanglement	1207
of marine life	
(W. R. P. Bourne, Chair)	
Report of the working group on ghost fishing	1216
(P. A. Breen, Chair)	
Report of the working group on ingestion	1226
(L. Sileo, Chair)	
Report of the working group on economic	1235
aspects of marine debris	
(K. E. McConnell, Chair)	
Report of the working group on technology	1240
(W. G. Gordon, Chair)	
Report of the working group on law and policy	1249
(D. Cottingham, Chair)	
Report of the working group on marine debris education	1256
(K. J. O'Hara, Chair)	
APPENDIXES	1261
Appendix A. Steering group	1263
Appendix B. Agenda of the workshop	1264
Appendix C. List of participants	1265

SESSION IV



ECONOMIC IMPACTS ON VESSELS AND SHORELINES



AN ECONOMIC PERSPECTIVE ON THE PROBLEM OF MARINE DEBRIS

Norman F. Meade, Kathleen M. Drazek, and Vernon R. Leeworthy
Office of Oceanography and Marine Assessment
National Ocean Service, NOAA
Rockville, Maryland 20852, U.S.A.

ABSTRACT

This paper examines the role of economic analysis in the development and implementation of an effective public policy to address the problem of marine debris. The economic theory of common property resources and other relevant aspects of natural resource and environmental economics are explained and used as a basis to critically review the economics literature on marine debris. Gaps in knowledge are identified and an economic data collection and research agenda is proposed.

INTRODUCTION

Many of the economic issues associated with the problem of marine debris are similar to those surrounding oil and hazardous substances pollution of the marine environment. The marine debris problem has not, however, received the same degree of attention by the research community as oil and hazardous substances pollution has. This is particularly true in the field of natural resource and environmental economics, as the review of the literature amply illustrates. Although a number of studies have shown that marine debris can have deleterious effects on marine life (Balazs 1985; Calkins 1985; Day et al. 1985; Bengston et al. 1988; Cooper et al. 1988), the current body of knowledge is insufficient to provide an assessment of the magnitude of the problem. And although research is continuing on the impacts of marine debris, neither does a coordinated effort exist to structure this research toward providing such an assessment, nor are efforts under way to ensure that research results are formulated in a way that will be useful for economic assessments.

This paper provides an overview of the economic aspects of the marine debris problem and suggests how economic analysis can play a role in finding effective and rational solutions. A research agenda is proposed to aid in quantifying the economic dimensions of the problem and assessing the effectiveness of economic incentives in achieving compliance with various laws and regulations.

THE ECONOMIC PERSPECTIVE

Absent significant economic incentives, compliance with environmental laws and regulations is usually low. It is true that education, moral suasion, and fear of punishment will stimulate many to abide by laws and regulation, but past experience has shown that these efforts alone will not significantly reduce noncompliance with environmental regulations. Here we discuss how an economist would approach analytically the problem of marine debris, including the issue of compliance with prohibitions on debris disposal.

Background

In order to show how economics can be used in analyzing the problem caused by marine debris, it is necessary to provide a brief description of the economic theory of natural resource and environmental economics. This will help clarify some of the concepts behind such familiar terms as market failure, economic efficiency, benefit-cost analysis, economic damage assessment, the value of environmental improvement, and cost effectiveness. These terms are related to methods for analyzing policy alternatives designed to correct problems in the way individuals use scarce natural resources (including environmental goods and services).

The literature on the problem of marine debris highlights a wide range of detrimental impacts on living and nonliving resources. These detrimental impacts are known, generally, in economic terms as external diseconomies (or simply "externalities" for ease of exposition). Externalities arise when the marketplace fails to balance competing uses of a resource so that a particular resource's value to society is maximized. Under ideal circumstances, competitive markets will consider all the costs and benefits of an activity, balance competing uses, and produce the maximum net benefit to society. Thorough study of the market failures which result in marine debris would undoubtedly lead to more effective solutions.

Common Property Resources and Nonmarket Goods

Two sources of market failure predominate in the natural resource and environmental economics literature: common property resources and nonmarket goods. One type is discussed in a classic article by Hardin (1968) who wrote of the "tragedy of the commons." Common property is overexploited because everyone has the right to use it, but no one has personal responsibility for it. Rivers, estuaries, and oceans are examples of common property. It is not surprising then that these bodies of water are overutilized as waste repositories, since dumpers do not have to pay the full social cost for their use. Given the rising, high cost of land-based disposal, we can expect pressure on these resources to continue.

Even if private property rights for natural resources exist, the second type of market failure occurs because markets cannot be easily organized for many environmental goods and services. They form a general category called "nonmarket goods and services." An example of the existence of market failure where private property rights exist is in the

market for wetlands. Many wetlands are privately owned but may be used in a nonoptimal way by the private owner because he or she cannot capture the many social (public) benefits produced, such as water recharge, storm protection, water purification, wildlife habitat, and fishery production. The wetland owner is unable to identify the beneficiaries or measure the amount of individual benefit for each of these services, therefore these services go unpriced and undervalued in actual market transactions. From the owner's point of view, he or she may maximize the value of wetlands by developing them, but from society's point of view wetlands may be misallocated since the value of nonmarket services is ignored.

The marine debris problem combines both types of market failure. Most of the resources affected, living and nonliving, are common property and have nonmarket values. Effective solutions to the marine debris problem must focus on resolving these two market failures. Implementing systems of private property rights in the rivers, estuaries, and oceans does not seem feasible. The solution to the common property resource problem has largely been government ownership and management. The government, it is often assumed, could represent and balance competing uses of resources if all the costs and benefits of the various activities were known. The government, acting as the private sole owner, could presumably maximize the value of its resources. However, experience has shown that such an outcome is not likely for a variety of reasons: lack of information, overlapping jurisdictions, conflicts of interests across jurisdictions, and the co-opting of politicians and managers by a particular interest group, to name a few.

Markets are vitally important sources of information on the value of goods and services. It is this aspect more than any other that leads to efficient outcomes from smoothly functioning markets. The costs and benefits of various courses of action are discovered through billions of private transactions. The major problem for nonmarket goods and services is the absence of quantifiable information about the costs and benefits of actions which affect them. Two broadly defined categories of nonmarket goods and services are expected to account for a major portion of the social costs of marine debris: recreational use value and intrinsic value of natural resources and the environment.

Recreational Use Value

Recreational use is generally recognized as second in importance only to human health as a beneficiary of water pollution control. Over the past 20 years, economists have been developing information collection and analytical techniques to estimate the recreational use value of natural resources. Survey sampling techniques and the use of questionnaires are the primary methods of information collection. Analytical techniques fall into two general categories; demand modeling and the use of direct valuation questions, e.g., contingent valuation approach. In demand modeling, individual expenditures on goods and services used in producing a recreational experience serve as proxies for actual market prices. In the contingent valuation approach, individuals are given a hypothetical situation defining the quantity and quality of the recreation experience. They are then asked how they would value in dollar terms a particular change in the quantity or

quality of a recreation resource. Some economists prefer the demand modeling approach because it is based on actual behavior; others prefer the contingent valuation method because of the flexibility it provides for addressing incremental environmental changes. Both have imperfections, and research on improved methods for estimating recreational use values continues.

Intrinsic Value

One of the value categories that is often excluded from estimates of the total economic value of nonmarket goods is referred to as intrinsic value. This term is used to define values that people place on natural resources that are independent of their present use. These values can be reduced by human activities that lower the quantity and quality of the resources in question. Such values appear to derive from a variety of motives including the desire to bequeath a legacy of natural resources such as clean oceans to future generations, or the sense of well-being that results from simply knowing that certain natural resources exist.

In the few empirical studies that have been completed to date, aggregate intrinsic values for unique natural resources have been shown to be quite large. As to the likely ratio of intrinsic values to use values, it is still too early to draw any firm conclusions. Most who have studied this issue agree that intrinsic values exist, but continue to debate how they can be measured accurately. The methods of collecting and analyzing data on intrinsic values closely follow the contingent valuation method used for recreational use values. Research on this important area of valuation is likely to intensify in the near future.

Efficient and Equitable Allocation of a Pollutant

Economic efficiency is one normative criterion for judging various policy outcomes. It is based on the maximization of the net social benefits to society from any activity (net benefits being equal to total social benefits minus total social costs). It is a normative criterion because there are an infinite number of economically efficient outcomes, each associated with a different distribution of wealth and income. A change in the distribution of wealth and income could change the benefits and costs of any activity and therefore the amount of the activity that is economically efficient. The distribution of wealth and income is another normative criterion used for judging policy outcomes and is commonly referred to as the equity or fairness criterion. Economists artificially separate the two criteria of economic efficiency and equity in order to make analysis tractable. Below, the concepts of economic costs and benefits and the economically efficient allocation of a pollutant are discussed. Following that, equity and another criterion related to efficiency, cost effectiveness, are presented.

Economic Costs

The fundamental economic measure of the cost of any action is its opportunity cost. This basic concept has an analogy in physics: two objects

cannot occupy the same space at the same time. In other words, one cannot undertake one activity without giving up something else. Opportunity cost, therefore, measures the value of the next best thing forgone in order to have the preferred choice. Social cost is simply measured by how much of some other thing is given up in order to have the preferred choice. For example, to estimate the full social cost of cleaning a marsh after an oil spill, one should count the opportunity costs of all the equipment, supplies, and wages paid to employees (using market prices), plus the nonmarket opportunity costs of any physical damage done (including those caused by the cleanup itself). Of course measuring the value of the aesthetic and biomass damage inflicted, since there is no market price established for them, is difficult.

Opportunity costs are incurred regardless of whether monetary transactions, or exchanges, take place. Both explicit costs, which show up in an accounts ledger, and implicit costs should be included in any full social cost accounting of a change in the quality or quantity of a natural resource. For example, the social cost of a beach littered with debris includes the cost of cleanup plus the lost enjoyment of the beach caused by the nonmarket aesthetic insult of the debris' presence until the cleanup is accomplished. Thus the social costs of any activity (beach litter) include the lost benefits from other activities impaired by that activity (beach use).

Economic Benefits

A benefit is the economic value of any good or service that provides utility or satisfaction to one or more individuals. Benefits enhance a person or group's well-being. They can be derived from the consumption of commodities such as offshore oil and gas, or fish, or from nonconsumptive enjoyment of a sunset or body surfing. Commodities, especially those valued in the competitive marketplace, where externalities do not exist, are much easier to measure because their prices are determined in arm's length exchanges which reflect the consumer's willingness to pay and the cost of all inputs used in their production.

Economic Efficiency

Economically efficient outcomes in the choice between competing activities are ones where net benefits (total benefits minus total costs) to society are maximized. When dealing with pollution, this concept is more easily understood by an equivalent formulation involving the minimization of two rather different types of costs: damage costs and control (or avoidance) costs. In the case of marine debris, damage costs would include such social costs as lost recreational use, intrinsic damage such as harm to pristine environments or marine mammals, and damages to ships from entanglement of propellers and steering gear. Control or avoidance costs include the cost of avoiding the pollution as well as the cost of removing or recycling the marine debris causing the harm.

The economically efficient outcome will occur at the quantity of marine debris corresponding to the point where the marginal control cost is equal to the marginal damage cost. This is shown in Figure 1 as point Q*.




Figure 1.--Efficient allocation of a pollutant--static case.

A nonzero optimum quantity of marine debris at point Q^* implies that there is some benefit from the use of products that end up as marine debris. Reducing the quantity of marine debris below Q^* would be inefficient from society's point of view because the social cost of reducing it by an additional unit would exceed the value of an additional unit of other goods and services otherwise damaged. The zero level of marine debris is not a socially desirable outcome in this case.

The above static analysis assumes that marine debris items are not persistent pollutants, that is, the pollutant does not have detrimental impacts over many time periods. Some forms of marine debris, however, are persistent pollutants. Even if all marine debris were controlled today, the amount accumulated in the environment would still have detrimental impacts for years to come. Because of the persistent nature of this type of pollutant, the analysis of the efficient pollutant level must take into account the intergenerational transfer of costs and benefits. In economics, we call this the dynamic efficiency criterion. Dynamic efficiency would be achieved at the pollutant level that maximizes the present value of net benefits over time. The mathematical formulation would be:

$$PV[B_1 \dots B_n] = \sum_{i=0}^n \frac{(B-C)_i}{(1+r)^i}$$

where B equals the total benefits of the goods that are produced jointly with the pollutant, i.e., marine debris; C the total cost of producing these goods plus the cost imposed on other goods and services impacted by the marine debris; i the time period; and r the social discount rate used

to make net benefits comparable across different time periods. The dynamically efficient allocation of a pollutant in this case has to satisfy the condition that the present value of the marginal net benefit from the last unit in period one equals the present value of the marginal benefit in each following period (Tietenberg 1988).

There is one interesting difference between the first efficiency outcome presented in Figure 1 (the static efficiency criteria) and the dynamically efficient outcome. In the dynamically efficient outcome, new marine debris after a certain amount of time must be eliminated. In the static outcome of Figure 1, Q^* , marine debris enters the environment each new time period. However, the dynamically efficient outcome recognizes that marine debris such as plastics causes damage over many periods. Thus, as marine debris continues to accumulate in the environment, not only the new but also the old marine debris is causing damage resulting in social costs. At some future time the old marine debris will have accumulated to a point where the costs are so high that economic efficiency requires the elimination of all new marine debris. That is, the point is reached where it is less costly to recycle all new marine debris or switch to cheaper substitutes.

Equity

As mentioned above, there are an infinite number of economically efficient allocations of marine debris depending upon the distribution of wealth and income. Wealth, broadly speaking, would include the amounts of both human and nonhuman capital a person owns. Human capital is a person's skills and abilities. Income is a flow from the stock of human and non-human capital. An increase in marine debris may result in an increase in the cost of beach visitation, since a person may have to travel further to get to a clean beach. This increase in cost can be thought of as a decrease in income available to the person to purchase other goods and services--an opportunity cost. Equity addresses the question of fairness in the distribution of net benefits from any activity. No generally accepted standards of fairness exist. Resolution of disputes over fairness are generally resolved in political or judicial processes. Implementation of policies that have high net benefits can fail because the benefits of the activity are concentrated in one region of the country and the costs in another. Unless the region that is disadvantaged is compensated for the added costs imposed by the policy, the policy may be defeated. There are several criteria that are generally used in evaluating the issue of equity. They are horizontal equity, vertical equity, and sustainability.

Horizontal equity occurs when people with equal incomes are treated equally. This can be used in judging the geographic fairness of a given policy. If people with comparable income levels in different parts of the country receive different net benefits, the horizontal equity criterion is violated.

Vertical equity deals with the treatment of unequals or those with different incomes. In assessing vertical equity, net benefits are distributed among income groups either progressively, regressively, or

proportionally. Distribution is said to be proportional if the net benefit received is proportional to income. It is said to be regressive if the net benefit represents a larger proportion of the income of the rich than of the poor, and is progressive if, as a proportion of their income, the poor receive a larger share than the rich. Since many of our societal programs are designed to aid the poor, it is usually assumed that regressive policies are bad. Some economic efficiency may be sacrificed to achieve greater equity.

The last criterion is sustainability. This involves intergenerational transfers of net benefits. As we have seen in the discussion of efficiency above, the marine debris problem can be characterized by intergenerational transfers because of the persistent nature of the pollutant. The sustainability criterion suggests that, at a minimum, future generations should be left no worse off than present generations.

Cost Effectiveness

A concept more closely related to the efficiency criterion of policy is the cost effectiveness approach. Under this approach it is recognized that, due to the lack of full and accurate information, determination of the optimal efficiency point is impossible. The cost effectiveness approach evaluates policies and management strategies as to the least costly way in which a given level of environmental quality can be achieved. In the case of persistent marine debris, the economically efficient solution may be an eventual ban on its use and disposal in the oceans altogether. However, compliance with such a ban would likely result in economic hardship for certain sectors of the economy and would be costly to enforce.

Laws and regulations that contain market-based incentive systems are, in theory, less costly than traditional regulatory approaches. Incentive systems use market forces to reduce pollution by requiring polluters to pay all or part of the social cost of their activity. They are penalized economically for high levels of pollution and are rewarded with lower fees for reduced levels of pollution. The laws and regulations that currently exist on marine debris do not contain market-based incentive systems to achieve compliance. This is an area where future research could pay big dividends.

Economic Impact

Many government officials appear more persuaded by the effects of their decisions and policies on sales, employment, and income, i.e., economic impact, than by efficiency, equity, sustainability, or cost effectiveness. Much of the time, concern about sales, employment, and income is expressed in terms of equity or fairness and reflects genuine concern for the health and welfare of people in the communities affected by various decisions and policies. However, economists would generally agree that maximization of sales, employment, and income are not preferable to economic efficiency as objectives of social policy, since irrational conclusions are often derived from analyses based upon maximization of economic impact. An example should help clarify this point.

Consider Figure 2, showing the demand and supply of commercially caught fish. The demand for commercially caught fish is shown in D_1 ; S_1 and S_2 are the supply of fish before and after pollution, respectively. Before pollution, consumers purchase Q_1 pounds of fish per time period at price P_1 per pound. Total sales revenue is equal to the area OP_1AQ_1 .

If pollution reduces the stock of fish, the supply curve shifts back to S_2 and consumers now purchase only Q_2 pounds per time period at the higher price, P_2 . Total revenue is now equal to the area OP_2BQ_2 . The problem with this analysis is that total revenue may have increased, decreased, or remained the same depending upon the price elasticity of demand. If demand is inelastic (a 10% increase in price will result in a <10% decrease in quantity demanded), then total revenue will increase. Thus, when demand is inelastic, if pollution reduces fish stocks it results in increases in total sales revenue.

Now consider the efficiency approach. Area $P_1P_1^*A$ measures the net value (consumer's surplus) associated with commercial fishing before the pollution. This would be a measure of the net benefits of commercial fishing to society. Now when pollution reduces the stocks, supply shifts to S_2 and the new consumer's surplus is equal to the area $P_2P_1^*B$, which is less than the area $P_1P_1^*A$ by the amount equal to the area $P_1P_2^*BA$. Thus, using the efficiency criterion, there is a net loss to society from the pollution injury to this commercial fishery. This loss would then be compared to the gains in consumer's surplus from the products that result in the pollution to determine if society gains or loses from their production.

Such comparisons are commonly known as benefit-cost analyses. They provide more comprehensive information to decisionmakers about the overall result of a given project or policy change than the rather incomplete picture conveyed by economic impact analyses. A benefit-cost analysis can help determine whether, for example, the social benefits of a specific set of policies to reduce marine debris outweigh their costs.

Categories of Social Cost

The following categories can be delineated as the major areas of known economic costs or externalities associated with marine debris:

- Commercial fisheries. Through what is called "ghost fishing," discarded or lost nets and other types of debris can entangle fish and reduce the quantity of various species and thereby impose costs on fishermen and consumers. Debris can also become entangled in fishermen's nets and either damage them or cause them to operate inefficiently.
- Ships. Debris can become entangled in the propellers and steering gear and can clog the water intake of vessels, thereby causing physical damage to ships of all types, including recreational fishing, cargo, military, and research vessels, and imposing repair and delay costs on their owners.




Figure 2.--Impact of pollution on the supply of commercially caught fish.

- Marine mammals, birds, and turtles. Through entanglement in and ingestion of plastics, we know that large numbers of birds and animals become injured and die, imposing costs on those members of society who obtain use value from these animals through viewing, hunting, and scientific research, or intrinsic values from the mere fact that these organisms exist.
- Recreation, such as beach use, hiking, camping, and picnicking. Debris causes aesthetic losses, as demonstrated by users who are willing to go to considerable expense to avoid it, such as through cleanup of beaches or extra travel to recreate in areas with less debris. Property owners in coastal areas may also suffer reductions in the value of their property if debris renders it less desirable from an aesthetic or recreational standpoint.
- Long-term impact. There could be other, as yet unknown, long-term impacts of marine debris on the health of humans and the biota which now, or may at some time in the future, impose unexpected costs on society.

The State of Economic Knowledge on Marine Debris

To date there have been only a handful of economic studies directed at the problem of marine debris. The present state of knowledge is reminiscent of what was known about the economics of oil spills and their prevention

some 20 years ago. A small number of studies have been conducted by state or local governments on the out-of-pocket costs but not necessarily the full opportunity costs of cleaning up small sections of beaches. There has been one detailed study on the effects of debris on individuals' willingness to pay for tourist accommodations in a small area of coastal Massachusetts several years ago (Wilman 1984). It revealed that overnight tourists did place a premium on reduced quantities of beach debris. However, the study ironically did not set out to measure the benefits of debris reduction, but rather the economic costs of oil spills on Cape Cod beaches. Since there were no actual oil spills there to study, the author used debris as a surrogate for the effect of oil on the value of beach recreation. There has also been one study on the costs of recycling shipboard plastic waste in the Port of Newport, Oregon (Recht 1988). It provides some useful information and anecdotes of what such a program entails from both a management and a cost standpoint. And finally, there has been one paper written on the types of economic incentives that might be applied to the problem of debris and what general types would likely be effective (Sutinen 1988). At present there are economic studies under way on some aspects of the debris problem that plagued the New England and mid-Atlantic coasts of the United States during the summers of 1987 and 1988.

In addition to the modest amount of economic research directed at the debris problem, there is some important complementary research being conducted on the value of various types of beach use, intrinsic values of natural resources, the costs and benefits of waste recycling programs, and the costs and marketability of degradable plastics. Results of such governmental and academic research programs can be found in the natural resource and environmental economics literature.

RESEARCH AGENDA

A review of the literature reveals that there is little known about the magnitude of the marine debris problem or of its social costs (or conversely, the benefits of a reduction in the quantity of debris). Justification of public programs to mitigate or eliminate these costs will require such estimates. But knowledge of these costs is only a first step. Laws and regulations require changing people's behavior to bring them into compliance. Market-based incentives will likely be the most cost-effective means of achieving compliance. Research is therefore needed on the relative effectiveness of various market based incentive programs in achieving compliance with various laws and regulations. Below is a list of suggested research projects that partially address both the issue of identifying the magnitude of the social costs of marine debris and various market-based incentive programs.

Social Costs

Aesthetics

Debris makes beaches and other recreational areas less attractive. Shorefront properties are also made less attractive, but whereas the loss in value of shorefront properties may show up in market transactions, the recreational values are nonmarket. Two studies are recommended to help understand the magnitude of this type of economic loss.

1. A study of the economic costs of debris on a specific set of beaches. This study would pick a set of beaches and investigate the economic value of lost services that would result from different levels of debris on the beach. The beaches chosen would ideally have wide regional representation. The study should be designed so that the methodology and the loss estimates could be expanded to other regions of the country.
2. A study of property value losses due to marine debris. Property value studies have been used by economists in estimating the economic damages from various environmental pollution problems. These techniques could easily be extended to the marine debris problem. Several regional studies should be conducted to show the effects throughout the nation.

Intrinsic Value

Debris traps and entangles fish and wildlife. Fish and wildlife also ingest various types of debris resulting in morbidity and mortality. This type of physical injury to the environment results in economic damage to individuals that value the right of fish and wildlife to exist or remain unharmed in pristine environments.

A study could be made of the economic cost incurred when individuals of some subpopulation of a noncommercial species (e.g., birds, mammals) become entangled in or ingest marine debris. This study could involve the threat of extinction or only the loss in social value when a small number of a species are lost or harmed. The study should be based on a national survey since many individuals outside coastal areas will experience this type of loss.

Fouling of Vessels and Fishing Gear

When vessels and their gear are impaired by contact with marine debris, there are two kinds of costs: a) the repair and replacement cost for the damaged gear and b) the opportunity cost of the vessel and gear when it is not in productive service. Commercial fishing or shipping impacts entail market losses, but for recreational boating, market and nonmarket losses must be considered. Two projects could be undertaken to quantify the incidence of impairment and the magnitude of costs.

1. Investigate the incidence of impairment for each of the following industry groups: commercial fishing, shipping, and recreational boating. Research should attempt to quantify the extent of the problem nationally and identify regions of critical concern.
2. Estimate the magnitude of costs for each of the three industry groups above. These could be small surveys among owners or operators in each of the industry groups. Areas

identified as representing the most severe problems should be used for each industry group.

Commercial and Recreational Fisheries

The greatest impact of marine debris on fish stocks is, apparently, the ghost fishing phenomenon. A secondary, but potentially large, impact is the possibility that consumer perception of contamination of fish stocks by marine debris can influence the demand and price of related fish products. This impact could extend to recreational fisheries because one of the main components of value in the recreational fishery is the consumption of fish.

1. Ghost fishing. Ghost fishing has an economic cost in terms of the wasted resource. For commercial fisheries it is the market value of the lost product, whereas for recreational fisheries it is the lost value due to lower catch rates. This project should involve both biologists and economists. Current economic research on the impact of catch rates on recreational fishing demand and value could be utilized in assessing the cost of ghost fishing.
2. The impact of perceived contamination on the price of and demand for fish. A project which collects and describes incidents of market effects (i.e., commercial fisheries only) from perceived contamination would provide at least some evidence of the economic costs of marine debris. A survey of the economics literature and of knowledgeable people to gather these incidents in the form of a research report should be conducted. Additional studies could follow, if warranted.

Compliance and Incentives

The greatest challenge in resolving the marine debris problem will be in finding and implementing the right mix of market-based incentives and enforcement to bring about compliance with various laws and regulations on the disposal of debris. The following projects would investigate the use of fees and incentives as part of the marine debris solution.

1. Deposits on the return of nondegradable products. The efficiency of deposits on beverage containers as a means of controlling land debris is well documented. This research project would investigate the potential for deposits for the return of plastic marine debris. It should focus on coastal states which have experience with deposit systems.
2. Fees on the use of nondegradable materials. Business firms and households are good at allocating scarce resources which they must pay for. Fees on plastic would be an incentive to substitute other materials. However, business firms must be treated differently from the household sector because

foreign firms could simply displace domestic firms. Foreign made products using a host of nondegradable but cheaper materials could replace domestically produced goods made of more expensive degradable materials. This project would investigate the feasibility of fees on potential debris in the marine environment.

3. Investigation of the economic gains that can accrue to a particular region as a consequence of consolidating waste handling facilities. New U.S. laws require that vessels bring their nondegradable waste to port. Ports are required to handle the solid waste. Within particular regions, it may be very costly for ports to handle all of the vessel-borne waste. An economic study of the costs of onshore waste handling would prepare ports for the resource demands and for setting port fees. When the costs differ among ports, there may be incentives to use different ports. Further, there are incentives to dump trash if fees are based on the amount of trash that is brought ashore.
4. Investigation of alternatives to traditional methods of compliance. Policies combining punishment and reward which partly subsidize the adoption of compliance techniques and impose clear penalties for the absence of compliance are used elsewhere in government regulation. This research program, would study compliance programs which include education, incentives, and penalties for a specific portion of the industry.

REFERENCES

- Balazs, G. H.
 1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Bengston, J. L., C. W. Fowler, H. Kajimura, R. Merrick, K. Yoshida, and S. Nomura.
 1988. Fur seal entanglement studies: Juvenile males and newly weaned pups, St. Paul Island, Alaska. In P. Kozloff and H. Kajimura (editors), Fur seal investigations, 1985, p. 34-57. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-146.
- Calkins, D. G.
 1985. Stellar sea lion entanglement in marine debris. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 308-314. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

- Cooper, R. A., H. A. Carr, and A. H. Hulbert.
1988. Manned submersible and ROV assessment of ghost gillnets on Jeffries and Stellwagen Banks, Gulf of Maine. U.S. Natl. Mar. Fish. Serv. and Mass. State Div. Mar. Fish., 17 p.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Hardin, G.
1968. The tragedy of the commons. Science 162:1243-1248.
- Recht, F.
1988. Report on a port-based project to reduce marine debris. Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. NWFAC Admin. Rep. H-88-13, 75 p.
- Sutinen, J. G.
1988. Remarks to the oceans of plastic workshop. In Oceans of plastic: A workshop on fisheries generated marine debris and derelict fishing gear, p. 50-61. Alaska Sea Grant College, Anchorage, AK.
- Tietenberg, T.
1988. Environmental and natural resource economics. Scott, Forsman and Co., Glenview, IL, p. 26-32.
- Wilman, E. A.
1984. External costs of coastal beach pollution: An hedonic approach. Resources for the Future, Inc., Wash., D.C., 194 p.

ESTIMATION OF DAMAGES TO FISHING VESSELS CAUSED BY
MARINE DEBRIS, BASED ON INSURANCE STATISTICS

Shuichi Takehama
Fishing Ground Environment Conservation Division
Fisheries Agency, the Government of Japan
Tokyo, Japan

ABSTRACT

An estimation has been made of the number of accidents, the amount of damage, and losses to and of fishing vessels caused by marine debris, based on data available from the insurance covering fishing vessels. Such accidents, damage, and losses caused by debris in the sea include those resulting from collision with drifting objects, entanglement of floating objects in the propeller blades, and clogging of the water intake for the engine cooling system. According to insurance statistics, losses attributed to the above-mentioned causes were ¥4.4 billion in 1985, whereas the losses and damage sustained by all the fishing vessels of <1,000 gross tons in Japan are estimated at ¥6.6 billion per year.

INTRODUCTION

Since ancient times, many different kinds of objects drifting in the sea have endangered ships. In the past, drifting wood and blocks of ice usually made up such driftage. More recently new kinds of debris including discarded plastic materials have emerged. Accidents and trouble caused by lost or discarded fishing nets and ropes becoming entangled in propeller blades, and the overheating of engines resulting from plastic debris clogging the water intake of an engine cooling system have been reported.

However, there are very limited statistical data available showing the number of vessels damaged by marine debris and the magnitude of the damage and losses. The reason for this lies in the fact that there are few systems for collecting such statistical data. On the other hand, at least part of such damage is covered by insurance, and by analyzing insurance-related data, it is possible to obtain approximate figures on the number of cases and the amount of damage.

In Japan, there is an insurance system for fishing vessels of 1,000 gross tons (GT) based on the Compensation Law Concerning Damages, etc., to Fishing Vessels. Established for the purpose of stabilizing the fishing industry which, is said to be subject to a great many dangers, this system

operates with a subsidy from the central government. As many as 60% of all fishing vessels of <1,000 GT are covered by this insurance system.

This insurance covers a wide range of damage to fishing vessels including damage such as sinking and fire, damage to the cargo including fish caught and stored on board, and loss of life of the crew. Also covered by the insurance are accidents resulting from marine debris.

This report illustrates the accidents and trouble sustained by fishing vessels in Japan as documented by a nationwide organization handling damage insurance (the Fishing Vessel Insurance Center, which is principally in charge of management of the insurance system) and estimates the magnitude of the losses and damage as well.

METHODS

The statistical data used in this report have come from two published reports relating to the damage to fishing vessels owing to accidents: Statistics of Fishing Vessel Insurance (hereinafter called the Insurance Statistics) and Report of Special Analyses on Accidents of Fishing Vessels Insured (hereinafter called the Special Report). The Insurance Statistics is issued annually, and the Special Report was compiled based on a detailed analysis of 1985 insurance data.

In the tables of the Insurance Statistics, damages sustained are classified according to those caused by bad weather such as heavy wind and rough seas, those caused by engine trouble such as a faulty lubrication system, those caused by human error in operating the ship and machines, those caused by drifting objects (this category is ambiguous but considered to be collision with drifting objects other than ice blocks), and those caused by foreign material tangled in the propellers. The cases to be discussed in this report will relate to damage caused by driftage, entanglement of foreign material in the propellers, and engine trouble resulting from trouble with the water cooling system.

Using the Special Report, more detailed analysis was possible. Accidents or trouble caused by drifting objects were classified into three categories: collision, cooling system trouble, and entanglement. As for the accidents or trouble with the cooling system, data indicating particular damaged points were also provided. As regards damage caused by drifting objects, the details given are those prepared especially for this report, obtained by reprocessing the original computer master tape used for the Special Report.

RESULTS

Fishing Vessel Insurance Statistics

Listed in Table 1 are the number of fishing vessels insured, the number of fishing vessels registered with the Fisheries Agency, the Government of Japan, and the percentage of insured fishing vessels (number of insured fishing vessels and number of registered fishing vessels). In

Table 1.--Number of fishing vessels registered, the number insured, and the ratio of vessels insured to total number of vessels (by gross tons (GT) and year) (Fishing Vessel Insurance Center 1985; Fisheries Agency of Japan 1975, 1982, 1983, 1984, 1985, 1986).

Vessels (GT)	1975	1982	1983	1984	1985	1986
Number of fishing vessels registered (1)						
<5	316,683	363,875	364,620	365,207	364,197	361,838
5-20	19,397	28,038	28,216	28,304	28,343	28,488
10-50	2,555	1,574	1,436	1,361	1,243	1,105
50-100	4,022	3,640	3,523	3,372	3,016	2,764
100-1,000	3,223	3,312	3,325	3,302	3,272	3,222
Total	345,880	400,439	401,120	401,546	400,071	397,417
Number of fishing vessels insured (2)						
<5	167,700	196,287	198,343	202,009	205,744	210,628
5-20	16,592	24,396	24,411	24,495	24,473	25,205
10-50	1,918	1,200	1,120	1,051	957	884
50-100	3,580	3,049	2,860	2,651	2,441	2,340
100-1,000	2,371	2,498	2,527	2,530	2,527	2,506
Total	192,161	227,430	229,261	232,736	236,142	241,563
Ratio of vessels insured to total vessels (2)/(1)						
<5	0.530	0.539	0.544	0.553	0.565	0.582
5-20	0.855	0.870	0.865	0.865	0.863	0.885
10-50	0.751	0.782	0.780	0.772	0.770	0.800
50-100	0.890	0.838	0.812	0.786	0.809	0.847
100-1,000	0.736	0.754	0.760	0.766	0.772	0.778
Total	0.556	0.568	0.572	0.580	0.590	0.608

1986, of the vessels smaller than 5 GT, >200,000 units were insured, or 58% of all the registered fishing vessels of that size group. (All fishing vessels in Japan must be registered when they are built.) The percentage of insured fishing vessels of 5 GT and greater is as high as about 80%, even though the total number of such ships is much smaller than the number of ships of <5 GT.

Figure 1-1 and Figure 2 show the frequency of accidents and the average amount of damage per accident, respectively, for the driftage, floating ice blocks, entanglement of the propeller blades, and trouble with the engine cooling system based on the Insurance Statistics.




Figure 1-1.--Change in frequency of accidents (number of accidents/number of vessels insured) by type of accident. (Source: Statistics of fishing vessel insurance. Fisheries Agency of Japan.)




Figure 1-2.--Change in frequency of accidents caused by drifting ice blocks by size class of vessel. (GT = gross tons). (Source: Statistics of fishing vessel insurance. Fisheries Agency of Japan.)




Figure 1-3.--Change in frequency of accidents or trouble with engine cooling system by size class of vessel (GT = gross tons). (Source: Statistics of fishing vessel insurance. Fisheries Agency of Japan.)




Figure 1-4.--Change in frequency of accidents caused by floating objects by size class of vessel (GT = gross tons). (Source: Statistics of fishing vessel insurance. Fisheries Agency of Japan.)




Figure 1-5.--Change in frequency of accidents caused by entanglement by size class of vessel (GT = gross tons). (Source: Statistics of fishing vessel insurance. Fisheries Agency of Japan.)




Figure 2.--Change in average cost of damage per accident by type of accident. Because the drifting ice category is so large, the cost shown in this figure has been divided by 10. (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)

The damage to the engine cooling system has a variety of causes including plastic bags. Damage by "drifting objects" has resulted from collision with objects floating in the sea, as described later. Damage by "entanglement" of fishing net and ropes in the propeller blades includes damage done not only by lost or discarded objects but also by fishing nets in actual use.

Except for damage done by "drifting objects," it was impossible to isolate the damage caused by only marine debris using the Insurance Statistics. These are, however, the only data available which enable us to review a historical change in accidents, and therefore, marine debris as a cause of the change is inferred by their use. In terms of frequency of accidents (number of accidents per number of fishing vessels insured), "drifting objects" is highest at about 10% (number of accident per number of insured vessels), which is more than twice the frequency attributable to other causes (Fig. 1-1). Conversely, the frequency of accidents is very low when it comes to the damage done by "floating ice," since such damage is limited to specific seasons and to specific areas of the sea. The frequency of accidents resulting from all categories shown has remained stable in the past 5 years. However, with regard to the accidents caused by drifting objects, this frequency is about 40% higher than it was in 1975, a percentage that has remained stable for the past 5 years.

Looking at the average cost per accident by type, "floating ice" stands out at ¥1 million or more suggesting that whereas its frequency of occurrence is low, the cost caused damage can be very great (Fig. 2). In contrast, accidents caused by drifting objects and entanglement in the propellers are low in cost, averaging around ¥100,000 per case.

With regard to differences in the frequency of accidents by size of fishing vessels, the frequency of "floating ice" is high in the case of ships of 100 GT or more, and is quite low with those of <5 GT. This is a natural consequence, because small ships rarely operate in a sea filled with drifting ice blocks during winter months (Fig. 1-2).

The frequency of accidents associated with the engine cooling system is highest for fishing vessels of 5-20 GT, next highest for those in the 20-100 GT class, and lowest for those of <5 GT. In any of these groups, the frequency of accidents tends to decline slightly over time (Fig. 1-3). The frequency of accidents caused by "floating objects," is highest with ships of 5 to 20 GT, followed by those of <5 GT and those 20 through 50 GT. It is the lowest with ships of 50 GT or larger. In most of those brackets, the frequency of accidents remains nearly the same. For vessels smaller than 5 GT, the frequency of accidents increased by 60% during the period from 1975 to 1982, and stabilized thereafter (Fig. 1-4).

In the entanglement of foreign materials in the propellers, too, the frequency is highest with vessels of 5 to 20 GT, followed by those <5 GT, 20 to 50 GT, and 50 to 100 GT. It is lowest with ships of 100 GT or larger. In terms of changes in frequency with time, ships in the of 20 to 50 GT and 50 to 100 GT classes are gradually increasing in number, whereas the frequency in the other brackets has remained stable (Fig. 1-5).

Special Report on Accidents of Insured Fishing Vessels

The purpose of the Special Report was to produce a detailed analysis based on data provided by the fishing vessel insurance. In order to do so, statements requesting payment were reprocessed to be collectively indicated on the form appearing in Figure 3. On this form, the types of accidents are classified (e.g., collision, fire, grounding, entanglement, engine trouble) as are the causes of the accidents (e.g., floating ice, drifting objects, inadequate watch), and by combining these categories, it is possible to determine the number of accidents of different kinds that were caused by floating objects and the cost as well. On the form, trouble with the engine cooling system is broken down into trouble resulting from plastic debris and that caused by other factors.

Accidents Caused by Floating Objects

Itemized in Table 2 by size of fishing vessel are the number of accidents classified by types of accidents caused by driftage and the amount of damage expressed in terms of money. Figures 4 and 5 show the frequency of accidents (the number of accidents divided by the number of insured fishing vessels) by type of accident and the average amount of damage per accident, respectively.

In 1985, there were a total of 32,8484 accidents resulting from drifting objects. There were 22,605 cases (69%) caused by collision, 5,809 cases (18%) with engine-related troubles, and 4,287 cases (13%) associated with entanglement. One hundred and forty-seven cases did not fall under any such classification (Table 2).

The cost of damage totaled ¥4.4 billion. It is said, in general, that the average such cost goes up as the size of the ship becomes larger. With ships <20 GT, the cost associated with engine trouble is the highest, and in the case of larger ships, the cost resulting from collision is the highest (Fig. 5).

In all size categories, accident frequencies are highest for those caused by collision, whereas the frequencies are lowest for engine trouble in all but the smallest ship size bracket. The frequency for the 5-20 GT category is highest in all types of accidents, whereas the frequency is low for size brackets of 50 GT or more, except for entanglement of foreign material in the propellers. The frequency of such entanglement is relatively high for 50 to 100 GT vessels, and low for those of <5 GT (Fig. 4).

Comparison of Accidents Caused by Marine Debris With Those of Other Causes

Figure 6 shows accidents caused by floating objects as a percentage of all accidents, by type of accident. The causes other than the driftage include entanglement with the fishing gears in actual use in the category of entanglement, and collision with submerged rocks in the category of collision. Collision as referred to here does not include ship-to-ship collision or grounding.

60年度 漁船保険事故分析調査表

(3) CAUSE OF ENGINE DAMAGE

(1) TYPE OF ACCIDENT

(2) MAIN CAUSE OF ACCIDENT

(1) Type of accident

(2) Main cause of accident

(3) Cause of engine damage

Collision with ships
 Collision with ice blocks
 Collision with others
 Grounding
 Capsizing
 Fire
 Missing
 Sinking
 Water intrusion
 Damage by rough sea
 Entanglement
 Explosion
 Theft
 Damage by lightning
 Engine trouble
 Machine trouble
 Other

Typhoon
 Other weather phenomenon
 Inadequate watch
 Autopilot system
 Floating ice
 Floating objects other than ice
 Careless handling of fire
 Improper ship maintenance
 Improper machine maintenance
 Intentional damage by others
 Improper ship mooring
 Improper fishing operation
 Improper ship operation
 Improper machine operation
 Other

Insufficient cooling resulting from clogging with plastic film.
 Insufficient cooling from other causes.
 Insufficient oil or deteriorated oil.
 Overload.
 Breakage of crank pin or bolt.
 Dropping of outboard engine.
 Other.

Figure 3.--Compilation form for the Special Report.

Table 2.--Number of accidents and damage caused by marine debris by type of accident and by vessel size (gross ton (GT)). (Number is actual number, and amount of total damage and damage per accident are shown in ¥1,000.) (Fishing Vessel Insurance Center 1985; Fisheries Agency of Japan 1975, 1982, 1983, 1984, 1985, 1986.)

Accident or damage	Vessels 0-5 GT			Vessels 5-20 GT			Vessels 20-50 GT		
	Number	Damage	D/N ^a	Number	Damage	D/N ^a	Number	Damage	D/N ^a
Collision	18,644	1,639,473	87.9	3,715	626,365	168.6	78	102,718	1,316.9
Entanglement	2,852	192,917	67.6	1,279	159,982	125.1	44	18,244	414.6
Engine trouble	4,773	814,839	170.7	940	363,436	386.6	35	20,521	586.3
Other	94	16,962	180.4	42	22,881	544.8	5	4,799	959.8
Total	26,363	2,664,191	101.1	5,976	1,172,664	196.2	162	146,282	903.0
Accident or damage	Vessels 50-100 GT			Vessels 100-1,000 GT			Total all vessels		
	Number	Damage	D/N ^a	Number	Damage	D/N ^a	Number	Damage	D/N ^a
Collision	90	132,083	1,467.6	78	158,264	2,029.0	2,605	2,658,903	117.6
Entanglement	82	26,217	319.7	30	40,153	1,338.4	4,287	437,513	102.1
Engine trouble	48	46,613	917.1	13	17,045	1,311.2	5,809	1,262,454	217.3
Other	2	862	431.0	4	30,557	7,639.3	147	76,061	517.4
Total	222	205,775	926.9	125	246,019	1,968.2	32,848	4,434,931	135.0

^aDamage divided by number.




Figure 4.--Frequency of accidents caused by floating objects by type of accident and by size class of vessel (GT = gross tons). (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)




Figure 5.--Average cost of damage per accident caused by floating objects by type of accident and by size class of vessel (GT = gross tons). (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)




Figure 6.--Percentage of the number of accidents caused by floating objects in relation to total number, by type of accident, and by size class of vessel (GT = gross tons). (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)

Percentagewise, engine trouble caused by floating objects is low on the whole, and the percentage decreases in reverse proportion to the size of ships.

For vessels <50 GT, the percentage of entanglement trouble caused by driftage increases with vessel size, but with the vessels >50 GT, the percentage tends to decrease.

In cases of collision, the percentage is the highest for vessels in the 5 to 20 GT category, and decreases as vessel size becomes larger.

It may be generalized that the percentage of damages attributable to drifting objects is lower in reverse proportion to the size of ships (Fig. 6).

Accidents or Trouble With the Engine Cooling System

The Special Report gives statistical data detailing the number of accidents or engine trouble and the amount (in yen) of damage resulting from improper engine cooling caused by plastic debris clogging the cooling water intake.

In 1985, there were 2,576 accidents with damage to engine cooling systems caused by plastic debris, which were covered by insurance. The cost of damages totaled ¥614 million. The frequency of accidents and the average cost per accident causing damages to the engines are shown in Figures 7 and 8, respectively. In those figures, causes of damage to the cooling system by other than plastic debris are added for reference. The frequency of accidents caused by plastic debris is lower for fishing vessels of larger size with the exception of those <5 GT. The average damage, on the other hand, increases with large ships.

There is slightly more engine trouble caused by factors other than plastic debris than trouble attributable to plastic debris, by size group of fishing vessels. The specific cause of these other accidents is not known. The cost per accident is nearly equal to or somewhat lower than that caused by plastic debris. According to Usui of the Fishing Vessel Insurance Center, who compiled the Special Report trouble with the cooling water systems occurs frequently as a result of clogging of the inlet ports for cooling water with drifting objects such as wood or grass. Since the average cost per accident is nearly the same, it is conceivable that accidents caused by other than plastic debris are similar to those caused by plastics. This seems to support Usui's statement.

Figure 9 shows the locations of main damage to the engines caused by a deficiency of the cooling system, which in turn was caused by plastic debris, as clarified in the Special Report. The damages to cylinder heads accounted for 57%; cylinder liners, 19%; and pistons, 14%.

DISCUSSION

The data of the Special Report differ somewhat from the Insurance Statistics for 1988. According to the Insurance Statistics, the number of insured vessels is 236,142, whereas the number given in the Special Report is 245,826, greater than the former by 9,700 ships or 4%. This difference is mostly in small ships, and therefore does not seem to adversely affect a comparison between the two sets of data.

As shown in Figure 10, in comparing the frequencies of accidents caused by collision with floating objects (category "floating objects" from the Insurance Statistics and "collision with floating objects" from the Special Report), they were found nearly the same in all ship size brackets.

The frequency of "entanglement" accidents given in the Insurance Statistics coincides well with the frequency of "entanglement" accidents in the Special Report. The frequency of "floating debris-related entanglement" in the Special Report is much lower when compared to the




Figure 7.--Frequency of accidents of engine trouble caused by plastic debris and other factors by size class of vessel. (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)




Figure 8.--Average cost per accident of engine trouble caused by plastic debris and other factors by size class of vessel. (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)

other two (Fig. 11). Because of this, "drifting objects" in Figure 1-1 represents changes in frequency of collisions with objects drifting in the sea, and "entanglement" in the same figure includes those accidents caused by other than floating debris, as previously stated. Further, Figure 11 shows that the percentage of entanglement accidents caused by other than floating debris becomes lower as the vessels become larger, when compared to entanglement accidents caused by floating debris. This is understandable, because human activities including fishing are much higher in the coastal areas than farther offshore and opportunities of encountering mooring ropes or fishing nets in actual use are greater in coastal waters. The density of marine debris distribution also lessens in the offshore waters.

The Special Report indicates that the frequencies of all three types of accidents (collision, entanglement, and engine trouble) show the same tendency, becoming lower as the vessel size become larger with the exception of vessels of <5 GT (Fig. 4). Such results are understandable, because the distribution density of drifting objects is high in the coastal waters in general, and small ships tend to operate in the vicinity of the coast. Several reasons can be conceived for the low frequency of accidents involving ships <5 GT. These are:

- Many boats with outboard engines are included in this category and their engine and water intake are easy to monitor, making it easier to detect the start of trouble such as trouble with the cooling system.
- Small vessels rarely operate at night.
- Small vessels are easier to monitor adequately than larger vessels.

According to the Special Report, the amounts of damages covered by insurance and resulting from collision, entanglement, engine trouble, and other accidents associated with objects floating in the sea were ¥2,659 million, ¥437 million, ¥1,262 million, and ¥76 million, respectively, for a total of ¥4,435 million. The total amount of such damage for all vessels in 1985 is estimated to be ¥6,608 million, a figure determined by extrapolation using ratios of the number of insured ships to the number of fishing vessels that were registered, by size (tonnage).

The estimated figure, however, is considered to be an overestimation. It is unrealistic to think that all the fishing vessels registered actually operated in that year. (Some are not in use any more and have not yet been removed from the register.) However, there are no statistics covering the number of fishing vessels that did operate during 1985. Furthermore, the statistical data used may have some problems in their characteristics since they were not prepared for this type of analysis, but they are considered to be pretty reliable. The actual size of the damage is thought to be somewhere between ¥4.4 billion and ¥6.6 billion. Such an amount is so huge that it calls for some review to determine its appropriateness. The total fishery production in 1985 by fishery management units with vessels of




Figure 9.--Engine part which is reported as damaged most severely in an accident of engine trouble caused by plastic debris. (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)




Figure 10.--Comparison of frequencies of accidents caused by "floating objects" from the Insurance Statistics, and "collision with floating objects" from the Special Report. (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.)




Figure 11.--Comparison of frequency of entanglement accidents.
 (Source: Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center. Statistics of fishing vessel insurance. Fisheries Agency of Japan.)

<1,000 GT was ¥2,165 billion. Realizing that the cost of running the fishing business in general is roughly 90% of the sales, and the total output is ¥1,949 billion, therefore damages costing ¥6.6 million, or 0.3% of the above-mentioned figure, do not seem to be unrealistic. It is, however, based solely on the available statistical data. No study has yet been made of the system used in the operation of the insurance, including confirming of the accidents. Also, no study has been made of the available ship accident reports. I invite comments and opinions from those involved in the Japanese fishing vessel insurance system.

ACKNOWLEDGMENT

In preparing this report, cooperation and assistance have been received from the following people. I would like to express appreciation to Masataka Seki and Michiko Yamagishi, Tokyo Software Development Company, Ltd.; Shigeki Fujita, Division of Fisheries Insurance, Fisheries Agency,

the Government of Japan; Akira Kito, Division of Fishing Ground Environment Conservation, Fisheries Agency, the Government of Japan; and Michio Usui, Fishing Vessel Insurance Center.

REFERENCES

Fishing Vessel Insurance Center.

1985. Report of special analysis on accidents of fishing vessels insured. Fishing Vessel Insurance Center.

Japan. Fisheries Agency.

1975. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1982. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1983. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1984. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1985. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1986. Statistics of fishing vessel insurance. Fisheries Agency of Japan.

1987 Summarized fishery statistics, 1987. Planning Division, Fisheries Agency of Japan.

n.d. Tables of statistics of fishing vessels. Fisheries Agency of Japan. No. 23, 35-39.

NEW YORK STATE MARINE DEBRIS PROGRAM

Roberta E. Weisbrod
New York State Department of Environmental Conservation
47-40 Twenty-First Street
Long Island City, New York 11101, U.S.A.

ABSTRACT

New York State has a multifaceted approach to prevention of marine debris. Regarding the sources of infectious waste-related debris, the state issued jointly with New Jersey a series of similar regulations which provided for a manifest tracking system. The state also recently passed additional legislation which increases the penalties for illegal disposal and removes the small quantities generator exemption; an additional US\$2 million was earmarked for enforcement.

The program to prevent the major portion of debris continues. The Department of Environmental Conservation works with Federal and local agencies to minimize contribution from such sources as combined sewer overflows and solid waste handling. The department also works with local environmental groups, education institutions, and the marine trades association in a public awareness campaign. The conclusions drawn from documented cleanups at eight beaches during Beach Cleanup Day, 8 October 1988, are discussed.

MEDICAL WASTES AND THE BEACH WASHUPS OF 1988: ISSUES AND IMPACTS

Kathryn D. Wagner*
Oceans and Environment Program
Office of Technology Assessment
Congress of the United States
Washington, D.C. 20510, U.S.A.

*Present address: Department of Political Science, Swarthmore College, 500 College Avenue, Swarthmore, PA 19081.

The views expressed in this paper do not necessarily represent the views of the Technology Assessment Board, the Technology Assessment Advisory Council, or their individual members.

ABSTRACT

The beach washups of medical wastes in 1988 resulted in beach closings, even if the extent of public health hazard posed by these wastes is not known with certainty. The nature and extent of economic impacts of the closings to the Long Island and New Jersey areas are assessed, based on available information.

Investigations to determine the sources of medical waste and other floatable marine debris along the shores of New York City, Long Island, and other nearby coastlines find that the primary sources are the Fresh Kills landfill (including barges transporting waste to it), marine transfer stations, combined sewer overflows, raw sewage discharges, and storm water outlets. Other sources, such as illegal dumping, probably contribute a smaller portion of floatables. This has important implications for whether some types of laws and programs being proposed and adopted, such as a manifest tracking system, will adequately address the problem of beach washups.

OVERVIEW

On 23 May 1988, a garbage slick nearly a mile in length along the shore of Ocean County, New Jersey, marked the season's first major washup of marine debris. Needles, syringes, and empty prescription bottles with New York addresses were among the floatable marine debris washed ashore. Beaches were closed as a result of this and similar incidents, including closing of more than 24 km (15 mi) of Long Island beaches from 6 to 8 July.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

Throughout the summer of 1988, national attention focused on reports of beach washups of medical wastes at various locations in the United States, much of the attention being given to washups on the east coast.

The east coast washups actually were about 10% or less "medical-related" waste. The largest single medical-related floatable was the insulin-type disposable syringe (New York State Department of Environmental Conservation (NYSDEC) 1988). Misconceptions about the quantities and nature of medical wastes washed ashore apparently resulted in part from the misidentification of items and from inaccurate media reports.

The degree of risk posed by medical wastes is not known. Proper handling, treatment, and disposal of these wastes are believed to minimize human health and environmental risks. Yet incidents resulting from careless or illegal disposal do pose aesthetic problems, and certainly help create public apprehension over current medical waste management practices. Aesthetic degradation and possible adverse health effects from medical wastes on public beaches are impacts difficult to measure and assess directly. Economic impacts resulting from the washups may be a more direct measure of their importance.

First, it is useful to define medical wastes and look at the broader context within which they are generated, managed, and regulated. Medical wastes include all infectious waste, hazardous (including low-level radioactive) wastes, and any other wastes generated from all types of health care institutions, including hospitals, clinics, doctors' (including dental and veterinary) offices, and medical laboratories (Lee 1988). The main focus of concern has been on the portion of medical wastes that are defined as infectious or "red bag" wastes, and how they are classified (e.g., as a solid, hazardous, or "special" waste) and regulated. The Centers for Disease Control and the U.S. Environmental Protection Agency (EPA) both designate as infectious pathological waste blood and blood products, contaminated sharps, and microbiological wastes (U.S. Congress 1988). The main sources of these wastes receiving attention are hospitals and other large facilities, but other sources of infectious wastes, such as sewage overflows, can also be significant contributors of environmental contamination.

The EPA reports that approximately 3.2 million tons of medical wastes from hospitals are generated each year, or about 2% of the total municipal solid waste stream. Currently, most generators of medical waste designate between 10 and 15% of it as infectious. The NYSDEC estimates that approximately 315 tons of medical wastes a day are generated by New York City's (NYC) 75 hospitals; and of this about 63 tons is infectious waste (NYSDEC 1988). Most of the medical waste washed ashore in the Greater NYC, Long Island, New Jersey, and even Rhode Island coastal areas is presumed to have emanated from the Greater NYC area. The actual amount of waste which washes ashore is not known with certainty, but is regarded as a large volume (NYSDEC 1988). At times, in some locations two pieces of medical debris per mile are found, while during a "garbage slick," garbage bags full of medical waste can be collected (NYSDEC 1988; Associated Press 1989).

Most infectious waste from hospitals is incinerated, while most noninfectious medical waste is landfilled. However, just as beach wash-up incidents raise public concern over current medical waste management practices, considerations of liability and worker safety lead some operators of municipal solid waste landfills and incinerators to refuse to take any medical wastes. As medical waste management becomes increasingly problematic for these types of reasons, an additional concern becomes the increased potential for illegal disposal.

The situation is further complicated by an uncertain and incongruous regulatory climate. Inconsistencies exist in the Federal guidelines for states regarding definitions and management options suggested for medical and infectious waste. Currently, no Federal regulations exist that comprehensively address the handling, transportation, treatment, and disposal of medical waste. This could change if the issue of medical wastes remains part of the current reauthorization effort for the Resource Conservation and Recovery Act (RCRA) or if the demonstration program of the Medical Waste Tracking Act (MWTA 42 U.S.C. 6901 et seq.) is expanded and extended in the future. As will be discussed below, other, specific types of management and enforcement actions may best address the issue of medical waste in beach washups. The Office of Technology Assessment (OTA), as part of a larger assessment of municipal solid waste, issued a separate background paper on medical waste management in October 1988 (U.S. Congress 1988). Reference should be made to that paper for a more detailed overview of medical waste management issues.

POSSIBLE ECONOMIC IMPACTS

The beach washups of 1988 in the Long Island and New Jersey areas had potential economic impacts of both revenue losses and costs. Revenue losses to the travel and tourism industry can result from declines in beach use, recreational fishing, and use of charter and party boats (R. L. Associates and U.S. Travel Data Center 1988; Thomas Conoscenti & Associates 1988; Ofiara and Brown 1989). Other possible economic impacts include increased beach maintenance and surveillance costs.

The focus here, based on very limited available information and in light of important caveats, is on the possible revenue losses to travel and tourism. Available information is suggestive of the types of short-term economic impacts which may be associated with the beach washups of 1988. It should be emphasized, however, that the information is not conclusive; some of it is anecdotal, most of the estimates of revenue losses are based on limited data (at most, for 2 or 3 years), and longer-term trends were not taken into account.

Causal links between the changes in beach use and tourism patterns and the beach washups and closings have not been carefully established, but are assumed in the estimates cited. Further, the methods used to estimate the changes and their impacts are not highly rigorous. Valid comparisons between losses to both New Jersey and Long Island coastal communities cannot be made given the different techniques used to estimate revenue losses.

The Long Island Tourism and Convention Commission reported a decline of 18% (i.e., an estimated 4.6 million fewer persons) in beach attendance in 1988 compared to 1987, and attributes this to the beach closings (Thomas Conoscenti & Associates 1988; Fey 1989). The commission also noted a decrease in attendance at all resorts and beaches (whether closed or not) as a result of the beach closings. Not surprisingly, decreased spending accompanied the lower beach attendance (Table 1). The difference between the 1987 tourism base and the estimated actual 1988 tourism base was \$921.2 million. The commission, however, calculated an estimated net loss of \$1.4 billion due to the beach closings in 1988, comparing the actual estimated tourism base of 1988 with that of an estimate of the industry base if it had grown in 1988 at the historical rate of 5.6% (Thomas Conoscenti & Associates 1988). The commission also reported that the actual net effect was likely to be considerably less than \$1.4 billion since it could be assumed that some of the tourists who did not visit the beaches probably participated in other activities on Long Island.

One part of the tourism and travel industry which may not be reflected in the calculations of the commission's survey (but is included in the marine recreational fishing category of the New Jersey estimates discussed below) is the charter and party boat businesses. A survey of NYC and Long Island charter and party boats owners found a 30 and 26% decline, respectively, in the number of passengers carried and trips conducted in 1988 compared to previous years (reportedly from 1985 through 1987) (DiLernia and Malchoff 1989). Floatables, including medical wastes, were considered by 60% of the party boat captains to be the most important issue affecting their business in 1988. Yet the respondents also agreed that a number of other factors threaten the profitability of the boat businesses, such as general marine pollution, a drop in fish stock abundance, and the high cost of operating vessels (DiLernia and Malchoff 1989).

One preliminary analysis of the impacts of the beach washups on all the beach towns in Monmouth, Ocean, Atlantic, and Cape May Counties, New Jersey, found that the overall range in beach attendance decrease from 1987 to 1988 was 7.9 to 34% (Ofiara and Brown 1989). (It should be noted that these investigators are completing their investigation and a more detailed version of their results will be available in April 1989.) Each of the four coastal counties experienced beach closings in 1987, but the beach attendance decrease between 1986 and 1987 ranged from 8.9 to 18.7%. The survey also indicated a 58% decline in beach attendance from 1985 to 1988 reported by seven New Jersey communities (Ofiara and Brown 1989).

According to the New Jersey Division of Travel and Tourism, an estimated 1.9 million fewer persons visited the New Jersey shore in 1988 than in 1987, a 22% drop in attendance (R. L. Associates and U.S. Travel Data Center 1988). In a 1988 survey of visitors, 22% considered themselves less likely to visit the New Jersey shore in 1989, with approximately the same number of respondents in this category as in 1987. Forty-four percent of this group of respondents identified pollution as their number one reason for not returning in 1989 (R. L. Associates and U.S. Travel Data Center 1988). (Pollution was in fact the single biggest reason given by those who indicated they would not be returning the following year to the New Jersey shore. This is a 10 percentage point increase over those who

Table 1.--Tourist and convention expenditures. Total Long Island, 1987 and 1988. (Source: Thomas Conoscenti & Associates, Inc. 1988.)

Visitors and their expenditures	1987	1988	Difference
Tourist and convention visitors (millions)	25.5	20.9	4.6
Expenditures (millions \$)			
-Lodging ^{a b}	368.9	332.2	36.7
-Food ^c	1,147.5	1,003.2	144.3
-Transportation ^d	255.0	219.5	35.5
-Entertainment	561.1	505.3	55.8
-Other ^e	1,009.2	908.8	100.4
Total	3,341.7	2,969.0	372.7
Annual total impact (millions \$)	7,685.9	6,828.7	857.2
Other direct summer activity (millions \$) ^f	589.0	525.0	64.0
Total tourist/convention industry (millions \$)	8,274.9	7,353.7	921.2

^aBased on 14,000 rooms.

^bAverage lodging rate (1987 = \$95/night; 1988 = \$100/night).

^cAverage \$45/day in 1987 and \$48/day in 1988.

^dIncludes day trips.

^eOther - e.g., retail sales.

^fVisitors to homeowners in summer.

had indicated in a survey the year before that they would be less likely to visit the shore in 1988 due to pollution. Yet about the same percentage (50% in 1987 and 47% in 1988) indicated that they were as likely to visit the shore the following year.)

The division also reports that a 9% decline in total expenditures occurred in 1988 at the New Jersey shore, a loss of approximately \$745.6 million. A more extensive analysis of revenue losses sustained by recreational fishing, beach use, and travel and tourism combined finds total estimated losses to range from \$545.9 million to \$2,022.85 million and, with all indirect effects included, to range from \$820.7 million to \$3,060.8 million (Ofiara and Brown 1989; Table 2).

It should be stressed again that these calculations are not comparable given the different methods used to derive them, and that the bases for them may be imprecise and have not been evaluated by the OTA. Yet, even with these qualifications, it appears that revenue losses have occurred. And it seems reasonable to assume that part of these losses was due to changes in beach use, recreational fishing and boating, and travel and tourism patterns which seem to have resulted primarily from the beach closings of 1988.

Table 2.--Aggregate estimated economic impacts to beach use, travel and tourism, and marine recreational fishing, New Jersey, 1988. All dollars are in 1987 dollars. (Source: Ofiara and Brown 1989.)

Category ^a	Trips (No.)	Economic benefits	Expenditures (\$1,000,000)	Gross value
Beach use				
Minimum	5,763,200	117.61	223.07	340.68
Maximum	24,493,600	499.83	948.05	1,447.87
Multiplier impacts				
Minimum		117.61	423.61	541.22
Maximum		499.83	1,800.35	2,300.18
Travel and tourism				
No. of businesses affected				
Minimum				395
Maximum				1,699
No. of lost jobs				
Minimum				9,553
Maximum				14,114
Lost wages (\$)				
Minimum				34.34
Maximum				147.79
Recreational fishing				
Minimum	1,332,600	88.28	82.60	170.88
Maximum	3,331,500	220.69	206.50	427.19
Multiplier impacts				
Minimum		88.28	156.86	245.14
Maximum		220.69	392.15	612.84
All activities				
Minimum		205.89	305.67	^b 545.90
Maximum		720.52	1,154.55	^b 2,022.85
Multiplier impacts				
Minimum		205.89	580.47	^b 820.70
Maximum		720.52	1,154.55	^b 3,060.80

^aMinimum refers to the minimum of the range. Maximum refers to the maximum of the range. Multiplier impacts are derived from the product of expenditures times 1.899; the New Jersey State multiplier associated with net output (includes value added), plus economic benefits.

^bThe sum of benefits, expenditures, and lost wages.

Even more limited information exists on the actual or estimated dollar impacts of various economic costs than on revenue losses for travel and tourism. For example, New Jersey spends approximately \$3 million annually on beach cleanups (New Jersey State 1987). A National Park Service official in Long Island indicates that the amount of money allocated for cleaning beaches and water quality testing has tripled, however, given the need for increased monitoring and surveillance since the beach wash-up problem arose in 1988 (J. Tanacredi, National Park Service, pers. commun. 14 February 1989). Figures are not readily available on these exact costs or how prevalent such increases are.

The actual washups may not have contained much medical waste, but the perception created by media reports that these wastes were appearing with frequency on beaches might have been a deterrent to beachgoers (R. L. Associates and U.S. Travel Data Center 1988; DiLernia and Malchoff 1989). It is not clear, even with a summer of fewer washups and the attendant lack of publicity, how quickly these economies will recover. Ofiara and Brown's (1989) review of previous studies indicates that there can be economic impacts (in some cases depressing fish prices for several years) from health advisories and subsequent media reporting of them.

BEACH WASHUPS: LOCATIONS AND SOURCES

In general, beaches closest to the sources of floatable marine debris (including medical waste floatable debris) are most likely to experience floatable strandings (New Jersey State 1987; NYSDEC 1988; Swanson 1988; Swanson and Zimmer 1989). Examination of potential sources and consideration of weather factors (i.e., winds and surface currents) seem to confirm this general relationship (NYSDEC 1988; Swanson 1988; Swanson and Zimmer 1989). There are a number of likely sources of the medical wastes and other materials in the beach washups along the NYC, Long Island, New Jersey, and other nearby shores. Table 3 lists the locations and dates of beach closings in the summer of 1988.

The weather appears to be an important factor in explaining the number of large beach washups in 1988. In 1988, as in 1976 (the last time long stretches of Long Island and other Greater NYC beaches closed due to pollution), a weather pattern of winds predominantly from one direction prevailed before the major washups (Swanson 1988; Spaulding et al. 1989; Swanson and Zimmer 1989). The most significant source of floatables is the Hudson/Raritan Estuary, and wind is the primary source of movement of floatables once they reach the bight. The prevailing south-southwesterly winds of early summer 1988 made Long Island beaches particularly susceptible to beach washups (Swanson 1988). A hindcast study for Long Island confirmed this relationship (NYSDEC 1988). The State of Rhode Island concluded that the New York Bight was also the probable source of the medical waste debris on its shores in 1988 (NYSDEC 1988; Spaulding et al. 1989).

Investigation showed that the primary sources of medical waste and other floatable marine debris along the shores of NYC, Long Island, and other nearby coastlines are the Fresh Kills landfill (including barges transporting waste to it), marine transfer stations, combined sewer

Table 3.--Summary of beach closings, 1988. (Source: New York State Department of Environmental Conservation 1988.)

County	Beach	Dates
Long Island		
Nassau	Nassau	7/6-7
	Long	7/6 (7/29 high bacteria)
	Jones	7/6, 7/8
	Lido	7/6
	Oyster Bay Town	7/7-8
	Gilgo	7/8
Suffolk	Robert Moses State Park	7/6-8
	Fire Island	7/7-8
	Babylon Town	7/8-10
	Smith Point State Park	7/10
	Quogue	7/23-27 (high bacteria)
	Shirley	7/29 (high bacteria)
New York City		
Queens	Rockaway	7/8, 7/26-28
	Jacob Riis Park	7/17-20
	Atlantic	7/17 (7/29 high bacteria)
Kings	Coney Island	7/12-13 (7/17 high bacteria)
	Brighton	7/12-13
	Manhattan	7/12-13
Richmond (Staten Island)	South	7/11 to close of season (9/3)
	Midland	7/10-8/18
	Great Kills	7/13-28 (8/9 high bacteria)
	Miller Field	7/13-25 (8/9 high bacteria)

overflows, raw sewage discharges, and storm-water outlets. Other sources such as illegal dumping probably contribute a smaller portion of floatables.

The NYSDEC's investigation into the sources of beach washups in 1988 concluded that medical-related wastes are sent to the Fresh Kills landfill, where some debris escapes into the water from a "hospital waste mooring" (NYSDEC 1988). Eight of nine municipal marine transfer stations (MTS's) and one private MTS currently operate in NYC. The NYC Department of Sanitation is responsible for off-loading wastes from trucks to barges for transfer to the landfills. Apparently, current loading practices cause spillage at the MTS's and there is not an effective system to remove such spillage (NYSDEC 1988).

The NYSDEC also reported that sewage treatment failures occurred prior to the beach washups in 1988, followed by rainstorms which flushed out floatable material collected in storm drains during a relatively dry period (NYSDEC 1988).

Although debris from the Fresh Kills landfill, sewage discharges (including combined sewage overflows), and MTS's are the most significant sources of floatables in the Greater NYC area, NYSDEC noted that "it is clear that medical-related waste has been disposed of illegally into the garbage and into the sewers [and that] these two sources contributed to the beach debris" (NYSDEC 1988). To date, illegal disposal of medical wastes appears to be a more significant problem on land than in the waterways, but it is possible that some illegal disposal directly into the water also occurs (NYSDEC 1988).

This information on the sources of medical waste in washups has important implications for whether some types of laws and programs being proposed and adopted to address the beach wash-up problem, such as a manifest tracking system, will be adequate. Another legitimate concern is that as regulation of medical wastes increases, disincentives for illegal disposal also need to be pursued, e.g., a manifest system or vigorous enforcement.

GOVERNMENTAL RESPONSES AND RELEVANT POLICY ISSUES

Whatever their actual aesthetic, social, and economic impacts may be, a critical issue is what governmental efforts if any will be effective in addressing the problem of beach washups of medical wastes. As noted above, currently no comprehensive Federal requirements exist for the management of medical wastes (U.S. Congress 1988). The MWTA of 1988 passed by Congress was in part an attempt to address the problems of beach washups of medical wastes and illegal disposal of medical wastes.

It is not clear, however, in light of the sources which appeared primarily responsible for the beach washups of 1988, that the "cradle-to-grave" type of manifest tracking system established by MWTA will have a significant impact on the washups of medical wastes. Other actions, such as improved waste management handling at marine transfer stations and at landfills in marine areas may more directly address the problem. Increased enforcement efforts appear prudent in any case, given the need to ensure that incentives for illegal disposal do not increase if the handling, transportation, treatment, and disposal of medical waste are increasingly regulated.

The MWTA establishes a demonstration tracking system (MWTA, Sections 11001-11003) and directs the EPA and another Federal agency to undertake studies of certain medical waste management issues (Sections 11008 and 11009). The intent is to develop a basis for determining whether and in what ways the Federal Government should regulate medical wastes. The MWTA specifically applies to Connecticut, New Jersey, New York, and the Great Lakes States (Section 11001). Any of the Great Lakes States may opt out of the demonstration program and any state can opt in; Connecticut, New Jersey, and New York can petition out if they have a program at least as

stringent as that of the Federal Government. Civil penalties of up to \$25,000 per day for each violation, criminal penalties of up to \$50,000 per day per violation, and in addition, jail terms of up to 5 years may be imposed in states implementing the tracking system (Section 11005).

On 24 March 1989, EPA established the 2-year pilot Federal tracking program authorized by MWTIA by publishing its "Standards for the Tracking and Management of Medical Waste; Interim Final Rule and Request for Comments" in the Federal Register (p. 12326-12395; 40 CFR Parts 22 and 259). Yet, as EPA points out in its press release of 13 March 1989:

"Many of the suspected sources of last summer's beach wash-up problems will not be affected by the new tracking system. Preliminary analyses of last summer's beach washups and additional EPA studies underway indicate that likely sources of the washups included improper handling of ordinary trash and sewer overflows which contain wastes from home health care and illegal drug use. To the extent that all of these sources contribute to environmental degradation, the problems will persist despite the new regulations (EPA 1989a)."

The manifest will not track medical-related wastes emanating from a number of the primary sources identified by the NYSDEC investigation. Interestingly, the cost of compliance with the requirements of the MWTIA, including the manifest system, is estimated by the EPA to increase the cost of medical wastes disposal by approximately \$0.08/lb on average (EPA 1989b). According to EPA (1989b), average annual compliance costs per facility range from about \$3,750 for hospitals to about \$70 for dentists. These figures and the per pound figure are considered to be low estimates by some waste industry officials.

In any case, New York State and New Jersey cooperatively adopted a tracking system in August 1988. It is not clear, however, whether either state will petition to opt out of the Federal program. A number of other governmental actions, including an interagency Floatables Action Plan for the New York Bight and action programs by individual states (e.g., New York State), have been initiated to address the problem of floatable medical wastes and other floatable debris in the Greater New York Harbor area (Molinari 1989; Weisbrod 1989).

The New York Bight Floatables Action Plan, a multiagency effort led by EPA Region II, is part of the New York Bight Restoration Plan. It includes such actions as studies of floatables in 1987 and 1988 and continued surveillance, regular cleanups at "key locations," other cleanup as necessary, and a communication network (Molinari 1989). In addition to expanding its public information program, NYSDEC's response to control beach washups of floatables includes combined sewer overflow abatement, improved operation and maintenance at sewer treatment plants (STP's), stricter controls at MTS's for the handling of solid waste, more stringent regulation of medical waste, and enhanced enforcement of medical and all solid waste regulations (Weisbrod 1989).

The NYSDEC anticipates that when new state medical waste regulations become effective, a capacity shortfall for medical waste disposal may result (Markell 1989). Older facilities may close if they anticipate that it will be too expensive to meet new regulations and, given the difficulties of siting new waste facilities of any type, incentives for illegal dumping could indirectly be fostered. For this reason, the state increased the criminal and civil penalties for medical waste violations in 1988 and plans an aggressive enforcement program (Markell 1989). Brooklyn District Attorney Elizabeth Holtzman supports strong enforcement efforts in the NYC area and actively prosecutes violators of existing medical waste management laws (Holtzman 1987, 1988).

Congress amended the Ocean Dumping Act (formally known as the Marine Protection, Research, and Sanctuaries Act of 1972, 33 U.S.C. Sections 1401 et seq.) in 1988 to increase the penalties for illegal disposal of medical wastes by public vessels. Some medical waste discovered along the coast of North Carolina and a few other locations was traced to discharges from U.S. Navy vessels (Associated Press 1988). According to the new amendments, civil penalties of not more than \$125,000 for each violation can be assessed by EPA for "engaging in activity involving the dumping of medical waste" as regulated by the law. Criminal penalties of not more than \$250,000, or imprisonment of not more than 5 years, or both, and possible forfeitures of property can also be imposed.

CONCLUSION

The beach washups of the summer of 1988 had a range of impacts (aesthetic, social, and economic) which may not have been precisely calculated, but did generate governmental responses to the appearance of medical waste on our beaches. Medical waste along our coasts also drew attention to a broader range of issues associated with medical waste management. Some of the specific programs initiated by state and local governments to address the beach washups of floatables may be most effective in the near term. The importance of the Federal demonstration tracking program for medical wastes in abating medical waste floatables in beach washups is not clear. Its significance to the improved management of all medical waste will need to be evaluated in the light of future regulatory programs. Nonetheless, using experience gained in regulating the hazardous and solid waste streams, there is opportunity for government at all levels to proceed in devising programs to manage medical waste wisely and efficiently in order to alleviate public concern, protect human health, and provide environmental protection.

REFERENCES

- Associated Press.
1988. Beach medical waste is claimed by the Navy. Associated Press, 9 Aug. 1988.
1989. Another summer of swimming among syringes? Associated Press, 19 Mar. 1989.

DiLernia, A., and M. Malchoff.

1989. 1988 survey of Long Island and New York City charter and party boat businesses. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.

Fey, G.

1989. Impact of environmental issues on tourism. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.

Holtzman, E.

1987. D. A. Holtzman announced indictment of medical waste company Press release, Brooklyn District Attorney's Office, Brooklyn, N.Y., 30 July 1987.

1988. Testimony at hearings before the U.S. House of Representatives, Committee on Small Business, Subcommittee on Regulation and Business Opportunities, hearing on health hazards posed in the generation, handling, and disposal of infectious wastes, 9 Aug. 1988.

Lee, M.

1988. CRS report for Congress: Infectious waste and beach closings. Congressional Research Service, Library of Congress, Wash. D.C., 9 Sept. 1988.

Markell, D.

1989. Infectious waste laws in New York State: An enforcement perspective. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.

Molinari, P.

1989. New York Bight Floatables Action Plan. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.

New Jersey State.

1987. New Jersey floatables study: Possible sources, transport, and beach survey results. New Jersey State.

New York State, Department of Environmental Conservation.

1988. Investigation: Sources of beach washups in 1988. Department of Environmental Conservation, New York State, Dec. 1988.


Ofiara, D., and B. Brown.

1989. Marine pollution events of 1988 and their effect on travel, tourism, and recreational activities in New Jersey. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.


- R. L. Associates, and U.S. Travel Data Center.
1988. The economic impact of visitors to the New Jersey shore the summer of 1988. Study conducted for the New Jersey Division of Travel and Tourism, Nov. 1988.
- Spaulding, M. et al.
1989. Hindcast of medical waste trajectories in southern New England waters. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.
- Swanson, R. L.
1988. The sources of infectious wastes. Paper presented at the Infectious Waste Disposal Conference, Wash., D.C.
- Swanson, R. L., and R. Zimmer.
1989. Overview of sources and marine transport of floatable wastes. Paper presented at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.
- Thomas Conoscenti & Associates, Inc.
1988. Long Island tourist/Convention estimates for 1987 and 1988. Prepared for the Long Island Tourism and Convention Commission.
- U.S. Congress, Office of Technology Assessment.
1988. Issues in medical waste management. Background paper October 1988. U.S. Congress, Office of Technology Assessment, Wash., D.C.
- U.S. Environmental Protection Agency.
1989a. EPA sets up pilot tracking program for medical waste. EPA Environmental News, 13 Mar. 1989.

1989b. Standards for the tracking and management of medical waste; interim final rule and request for comments. Federal Register, p. 12326-12395; 40 CFR 22, 259 (24 Mar. 1989).
- Weisbrod, R.
1989. Prevention and response: New York State actions. Presentation at the Floatables Wastes in the Ocean Conference, 21 Mar. 1989, State University of New York, Stonybrook, N.Y.

SESSION V



SOLUTIONS THROUGH TECHNOLOGY



THE PHILOSOPHY AND PRACTICE OF DEGRADABLE PLASTICS

Gerald Scott
Polymer Science
Department of Chemical Engineering and Applied Chemistry
Aston University
Birmingham B4 7ET, United Kingdom

ABSTRACT

A major advantage of plastics in packaging is their resistance to attack of microorganisms. After discard, however, this characteristic creates major problems in their disposal, since unlike nature's litter, they do not return to the biological cycle. Strategies for retaining the advantages of plastics during use but of triggering biodegradation after discard are discussed. Photobiodegradable plastics which have been used successfully as agricultural mulching film for many years are seen to be the potential solution to the marine debris problem.

THE ENVIRONMENTAL IMPACT OF PLASTICS

The growth of consumer packaging has been one of the most visible phenomena of the twentieth century. On balance, it has been a beneficial development which has facilitated the distribution and storage of perishable goods to the advantage of the international community.

The modern packaging industry has its roots in the petrochemical industry, and the cheap commodity plastics, polyethylene, polypropylene, polystyrene, and polyvinyl chloride, are the major polymeric materials currently in use. They have a number of advantages in common, of which the following are the more important:

- They are cheap and easy to fabricate into common items of packaging.
- They are resistant to water and microorganisms and are, therefore, able to protect perishable goods from biological attack.

However, the very characteristics which make plastics so useful in packaging cause considerable problems in their ultimate disposal. Unlike cellulosic packaging materials (paper, cardboards, and the cellulose-based plastics), the oil-based plastics do not biodegrade back to the carbon

cycle when discarded in the environment. This is why they present one of the most visible litter problems of the twentieth century. The persistence of plastics litter in recreational areas, where it gave maximum offense, became evident in the late 1960's. The response of the plastics industry at that time was to say that it was unlikely that technical solutions to this problem could be found because the characteristics of biodegradable plastics were "the antithesis of the nature of packaging materials" (Staudinger 1970). The Society of Chemical Industry in 1970 expressed its alternative strategy as follows:

" . . .the Society will seek to make common cause with all movements and organisations concerned with preventing environmental deriliction [sic] by littering" (Staudinger 1970).

This did not mean that they would encourage research into making plastics more biocompatible, but that they would encourage educational programs directed toward making the public more environmentally aware of the litter problem. This is still essentially the polymer industry's public stance today. The responsibility for plastics pollution is presented as that of the user, who must be educated into nonlittering habits (Claus 1987; O'Connell 1987; Society of the Plastics Industry 1988).

In the meantime, the problem continues to grow. There was a temporary respite in the mid-1970's, when it appeared that the oil crisis would lead to widespread recycling of plastics waste. This did not happen on any scale for technical and sociological reasons, and the problem became more severe even through the years of high polymer prices.

One of the earliest surveys of plastics pollution in the seas was carried out by the author in the early 1970's. Over a 3-year period, a fivefold increase in plastics litter was observed on a remote shoreline in northwest Scotland (Table 1) (Scott 1972a, 1975a). The conclusion drawn from this survey was that most of the plastic litter found on the seashore is seaborne and wind driven. The nature and location of the litter suggested that it came predominantly from shipping and not from local inhabitants or visitors. In this context, the good intentions of the educators were seen to be both misguided and misdirected. Commercial pressures to use the sea as a convenient "waste bin" have proved to be much more persuasive than homilies by The Tidy Britain Group, and even the threat of legislation has little effect due to the difficulty of policing this on the high seas.

This early evidence of sea littering has been confirmed by many subsequent studies (Dixon and Cooke 1977; Dixon 1978; Dixon and Dixon 1981; Fowler and Merrell 1986; Andrady 1987; Heneman 1988), and there is increasing evidence that plastics debris can kill birds and animals by ingestion and strangulation (Fowler and Merrell 1986; Andrady 1987; Heneman 1988).

Certain types of nonbiodegradable plastics waste have come in for most criticism over the years. The most visual and intrusive are the large polyethylene bags used for packaging agricultural and industrial products and domestic carrier bags, all of which float on the sea and accumulate on land. Even more aesthetically objectionable are the smaller items which

Table 1.--Accumulation of plastic litter at Strathaird Point, Isle of Skye, Scotland, in a 3-year period (number of packages per 50 yd).

Type of packaging (polymer) ^a	August 1971	August 1974
Detergent (LDPE)	7	5
Detergent (HDPE)	--	31
Bleach, sanitary fluid (HDPE)	15	49
Oil (HDPE)	4	18
Cosmetic (HDPE)	3	3
Carpet cleaner (HDPE)	--	5
Food (HIPS or ABS)	1	7
Table salt (HDPE)	--	10
Milk (HDPE)	--	16
Heavy gauge bags (LDPE)	2	6
Small transparent bags (LDPE)	--	29
Carrier bags (LDPE)	--	5
Heavy gauge sheets (LDPE)	6	15
Miscellaneous unidentified	3	14

^aLDPE - low density polyethylene, HDPE - high density polyethylene, HIPS - high impact polystyrene, ABS - acrylonitrile-butadiene-styrene copolymer.

originate from sewage disposal and which, although probably harmless compared with other components of sewage, cause great offense on beaches and in other environmentally sensitive areas (Johnson 1987). Six-pack collars, used for carrying beer and soft drink cans, have been particularly indicted as a cause of entanglement for birds and small animals, and discarded ropes and fishing nets are equally a cause of suffering and sometimes death to wildlife. These are all examples of litter which does not biodegrade, and although the polymers do degrade slowly under the influence of sunlight and oxygen and the erosive influence of the weather, these natural processes are not fast enough to eliminate the dangerous effects of man-made polymers in the environment.

Fortunately, a good deal of work in academic laboratories, particularly in the United Kingdom and Canada, had led to an understanding of the chemistry involved in the oxidative degradation of polymers. Associated with this was a fundamental understanding of antioxidant and ultraviolet (UV) stabilizer mechanisms which suggested the possibility of designing polymers with controlled outdoor stability.

THE ENVIRONMENTAL STABILITY OF POLYMERS

Biodegradability of Polymers

As has been discussed above, the main reason that the man-made polymers have assumed a position of such importance in the packaging

industry is because of their excellent water barrier properties. Since they are not readily penetrated by water, they act as an effective barrier, even in very thin films, to the attack of microorganisms. However, just because they are not accessible to microorganisms, they normally remain resistant to microbiological attack after discard in the environment.

All man-made polymers are not so hydrophobic as the carbon-chain polymers, however, and in general, the closer in structure polymers are to the natural polymers, the more biodegradable they become. Thus, the polyamides, which resemble the polypeptides in chemical structure, do absorb water and slowly biodegrade. In the case of polyurethane foams, biodegradation may take place quite rapidly because of the high internal surface area of the foam structure. Recently, man has been able to utilize nature's ability to synthesize and store within the biological cell certain types of polyester (Table 2) to produce a truly biodegradable polymer with physical and mechanical resemblances to the polyolefins (Lloyd 1987).

It is an unfortunate irony, however, that the nearer synthetic polymers approach the structure and properties of the natural polymers, the less useful they become as packaging materials because of the impairment of their barrier properties. Cellulose-based packaging has been largely abandoned over the years in favor of the hydrophobic polymers, and it is highly unlikely that the packaging industry would now be willing to return to less effective materials even if they could be produced at the same price as commodity plastics. It appears then that some other stratagem has to be sought to ensure that packaging materials are returned to the biological cycle when discarded in the outdoor environment.

Oxidative Degradation of Polymers

All organic polymers degrade due to the combined effects of oxygen, sunlight, and water by processes which do not, at least in the early stages, involve biological agencies (Scott 1965). They do so, however, at rates which differ by several orders of magnitude. Fluorinated polymers (e.g., Teflon) are in general the most resistant to environmental deterioration, and in the absence of light they can survive for many decades. Hydrocarbon polymers, and particularly the unsaturated rubbers, are much less resistant to oxidation, and even polyethylene, which on the basis of its structure should be chemically inert, does oxidize slowly unless protected against the effects of the environment. The small amounts of antioxidants which are added as processing stabilizers are normally sufficient to effectively stabilize polyethylene against the effects of oxidation in the absence of sunlight, but much more effective combinations of antioxidants and light stabilizers have to be used in order to give the polymer the durability required for use in outdoor applications (Scott 1979-88; Grassie and Scott 1985).

Oxidation of polymers leads to the formation of a variety of oxygen-containing functional groups as part of the polymer chain, of which the most important are hydroperoxides, carbonyl groups, alcohols, and carboxylic acids (Fig. 1) (Grassie and Scott 1985). These lead to the modification of the polymer surface, making it hydrophilic and allowing microorganisms to preferentially remove the oxygen functions. Figure 2

Table 2.--Commercially available degradable plastics.

Common description	Composition	Trade name	Manufacturer
Biodegradable polymers			
Poly (3-hydroxybutyrate-3-hydroxyvalerate)	Biosynthetic copolymer of 3-hydroxybutyric and 3-hydroxyvaleric acids	Biopol	ICI (United Kingdom)
Polymers containing a biodegradable filler^a			
Starch-filled polyethylene (Griffin process)	Physical blend of LDPE and starch	Bioplast Ecostar	Coloroll (United Kingdom) St. Lawrence Starch (Canada)
Photodegradable copolymers			
Ethylene-carbon monoxide copolymers		E/CO	DuPont (United States) Union Carbide (United States) Dow (United States)
Vinyl ketone copolymers (Guillet process)	Copolymers of ethylene, propylene, and styrene with a vinyl ketone	Ecolyte	EcoPlastics (Canada)
Photosensitizing and photoactivating additives			
Iron salts	Probably ferric stearate	PolyGrade	Ampercet (United States)
Aromatic ketones	Probably benzophenone with metal stearates	Not commercial	Princeton Polymer Lab. (United States)
Antioxidant photoactivator (Scott-Gilead process)	Ferric thiolates (sometimes with other metal thiolates (in polyethylene) (in polypropylene baler twine)	Plastor Greenplast Plastigone Ecoten ^b Litterless Cleanfield	Plastopil (Israel) Polydress (Germany) Enichem Agricoltura (Italy) Plastigone Technologies Inc. (United States) Amerplast (Finland) Plastigone Technologies Inc. (United States) American Brazilian Company (United States)

^aLDPE = low density polyethylene. ^bManufacture discontinued.




Figure 2.--Infrared spectra of high density polyethylene (HDPE) films with different histories: 1--with antioxidant after standing at ambient temperature for 1 year; 2--as 1 without antioxidant; 3--as 2 after standing for 3 years; 4--as 3 after treatment with an aerated medium inoculated with cultivated soil; 5--HDPE powder without antioxidant exposed to aerobic biodegradation for 2 years before molding to film with exclusion of air. (Grassie and Scott (1985) with permission, originally reproduced from a doctoral thesis by Dr. A. C. Albertsson with permission.)

Photooxidation of Polymers

Thermal oxidation of polymers is not a very controllable process. Although it can be catalyzed by cooxidation agents or transition metal ions, these are present in varying amounts of foodstuffs. However, autoxidation is readily inhibited by antioxidants, which are deliberately added to stabilize the polymer during manufacture (Grassie and Scott 1985). Furthermore, the addition of even a few percent of polymer-soluble coagents substantially alters the physical characteristics of the polyolefins (Scott 1988a, 1988b).

Photooxidation, by contrast, is a much more controllable process, since it is not appreciably affected by thermal antioxidants or contaminants. As early as 1971, it was suggested (Eggins et al. 1971) that controlled photooxidation was a potentially useful way of dealing with the problem of nonbiodegradable packaging litter. It offers the very considerable advantage over true biodegradation of the main polymer chain that, until photooxidation has occurred in the environment where the package has been discarded, its properties do not differ in any respect from conventional packaging made from the same polymer. The mechanism of photooxidation is essentially similar to that of thermal oxidation. The essential difference is the way in which the autoxidation chain reaction is initiated, which, in turn, depends on the presence of photoinitiators in the polymer (Grassie and Scott 1985).

Typical photoinitiators that have been used to sensitize the photodegradation of plastics in the outdoor environment after discard are listed in Table 2. Although many more have been reported in the patent literature, these are the only ones which have reached the marketplace. Companies producing them are also listed in Table 2.

Photodegradable plastics can be broadly classified into types:

- Copolymers in which the sensitizer, a carbonyl group, is built into the polymer.
- Conventional plastics to which the sensitizer is added, generally as a masterbatch and in some cases as a replacement for the usual processing stabilizer.

Copolymers

In the first approach, degradation occurs primarily by photolysis of the polymer backbone, leading to reduction in molecular weight and fragmentation of the polymer (Fig. 3). The Norrish Type II process is the predominant mechanism leading to chain scission, giving rise to ketone and vinyl groups at the end of the polymer chains (Grassie and Scott 1985). An early example of such a carbonyl-modified polymer was claimed in a patent by DuPont (Brubaker 1950), and this process has been commercialized relatively recently by DuPont and Union Carbide among others (Johnson 1987). The polymers embrittle rapidly and without any induction period (Harlen and Nicholas 1987; Statz and Dorris 1987), and if a period of




Figure 3.--Photolysis of in-chain carbonyl in polyethylene.

"safe" use is required, for example in storage or for a defined period out-of-doors, then the packaging has to be protected against UV irradiation. This kind of degradable plastic cannot, therefore, be used in agricultural protective film (mulch), where the polymer has to retain its strength for a well-defined period of time in sunlight but then has to photodegrade rapidly over a period of a few weeks. Rapid degradation appears to cease after all the in-chain carbonyl groups have photolyzed, and there is evidence (Harlen and Nicholas 1987) that the molecular weight decreases to a minimum and then increases again (Table 3). There is a similar minimum in the elongation to break, and these phenomena suggest that oxidation does not occur to any extent. This is consistent with the fact that if biodegradation does occur, it is extremely slow (Statz and Dorris 1987)--a considerable disadvantage of this type of product, since the fragmented products will still tend to accumulate in the environment.

In a more sophisticated approach to photodegradable copolymers, Guillet and his coworkers have copolymerized a variety of monomers, of which styrene, ethylene, and propylene are the most important, with vinyl ketones (Guillet 1973). These polymers have been commercialized under the name Ecolyte by Ecoplastics Ltd. (Redpath 1987). These polymers photolyze by essentially the same mechanism as the E/CO polymers, but both Guillet (Guillet et al. 1974; Jones et al. 1974) and Redpath (1987) have reported that the Ecolyte copolymers do biodegrade after fragmentation. This implies that the Norrish Type I process must play some part in initiating a conventional autoxidation chain reaction. However, the Ecolyte polymers suffer from the same disadvantage as the E/CO polymers in that they do not

Table 3.--Molecular weight changes in E/CO polymer (2.74% CO) on ultraviolet irradiation.

Exposure time, h	M_n	M_w
0	45,000	618,700
650	7,300	15,000
1,350	11,100	39,100

have a controllable induction period before rapid photodegradation commences.

Sensitizer Additives

In this approach, two main types of additive have been used. The first class falls broadly into the general class of triplet sensitizers, most importantly the benzophenones (Takahachi and Suzuki 1964) (see H. Omichi (1983) for a survey of the literature up to 1983). The mechanism of their action is summarized in Figure 4 and the related quinones act in the same way. Although the carbonyl sensitizers cause rapid photooxidation from the beginning of UV exposure, they autoretard rapidly in the case of




Figure 4.--Mechanism of photoinitiation and autoretardation by triplet sensitizers.





Figure 5.--Comparison of the photooxidation, as measured by carbonyl formation (at $1,710\text{ cm}^{-1}$), of low density polyethylene (LDPE) films containing different kinds of photoactivator at similar concentrations. 4,4'DMB, 4,4'dimethoxybenzophenone; Fe(III)AcAc, iron acetyl acetonate; Fe(III)DNDC, iron dinonyl dithiocarbamate; Fe(III)L2, iron complex of 4-methyl-2-hydroxyacetophenone oxime; Co(II)DNDC, cobalt dinonyl dithiocarbamate (UV stabilizer). (From Amin and Scott (1974) with permission.)

polyethylene and the photooxidation virtually ceases after a period of time (Fig. 5) (Takahachi and Suzuki 1964). This is because the hydrogen abstraction step in Figure 4 leads to a "stable" radical which is able to trap out the chain-carrying species formed in the photooxidation process. This has unfortunate consequences for the long-term oxidation and biodegradation of the polymer, and this, coupled with the lack of an appropriate time control mechanism, has resulted in no commercial developments with this type of sensitizer system.


The second class of photosensitizer additives is based on transition metal ion compounds and is the most important class in use today. Transition metal ions have been extensively studied as photosensitizers for polyolefins (Takahachi and Suzuki 1964; Mellor et al. 1973; Amin and Scott 1974; Chew et al. 1977). Many polymer-soluble metal carboxylates, notably Co^{3+} and Fe^{3+} , are powerful photoprooxidants which catalyze

photooxidation right from the beginning of UV irradiation (Fig. 5), and in the form of polymer-soluble carboxylates or acetylacetonates they cause melt degradation of the polymer during processing (Amin and Scott 1974). They cannot, therefore, be used alone in polymers in conventional processing operations because of their unfortunate effect on the melt stability of polymer and on the shelf-aging behavior of the fabricated product. Although conventional antioxidants improve processing and aging characteristics, they also interfere with the photosensitizing effect of the transition metal ions (Mellor et al. 1973).


The photoinitiating mechanism of the transition metal salts involves photolysis to give the reduced form of the metal ion and a free radical:



The anion radical readily abstracts a hydrogen atom from the polymer




Once formed, hydroperoxides control the subsequent oxidative degradation by the usual redox reactions with metal ions, e.g.,




Transition Metal Ion Antioxidant-Photosensitizers

Many metal complexes containing sulphur as a ligand are antioxidants and photostabilizers. Although this behavior is not limited to sulphur compounds, members of the latter class have gained a position of some importance as heat and light stabilizers for polyolefins (Scott 1965; Al-Malaika et al. 1983; Al-Malaika and Scott 1983). The dithiocarbamates (I) and the dithiophosphates (II) are representative of this class of stabilizers and exert their effect by destroying hydroperoxides by an



I



II

ionic mechanism (Al-Malaika et al. 1983; Al-Malaika and Scott 1983). One of their most striking and useful attributes is that they produce a well-defined and reproducible induction period during which the ligand is destroyed and at the end of which the metal ions which form part of the antioxidant are released and subsequently behave very much like the free transition metal ions described in the previous section (Fig. 5) (Scott 1965).

In 1971, Scott filed a patent based on the above concept in which antioxidant and photosensitizer properties were both contained in the same molecule, although this could be made in situ in the polymer by reacting the metal ion with the antioxidant (Scott 1971). The Fe(III) complexes of I and II are representative of this class of "delayed action" photoactivators. Not only do they replace conventional processing stabilizers by virtue of their antioxidant properties, but they are also effective heat stabilizers and short-term light stabilizers. The fact that the induction period to photooxidation could be controlled by varying the concentration of the metal complexes led in the early 1970's to the use of this system in agricultural mulching film, which requires a finely controlled lifetime before rapid photooxidation and biodegradation commences (Scott 1972b, 1972c, 1973a, 1973b, 1973c, 1975b, 1975c, 1976; Scott and Gilead 1978). This material was marketed as Plastor by Plastopil Hazorea, and subsequent development in collaboration with Gilead led to further patents (Scott and Gilead 1978) concerned with the fine control of the "safe" period, which is so essential for agricultural purposes (Scott and Gilead 1982).

The later developments involve the use of two component systems in which the length of the induction period is controlled by one metal thiolate and the rate of photooxidation by a second. The Scott-Gilead process is currently used in the commercial growing of soft fruits, vegetables, and some cereals in Italy, Germany, France, and the United States. In addition to the name Plastor, these products are sold under the trade names Greenplast (Enichem Agricoltura, Italy), Plastigone and Litterless (Plastigone Technologies Inc., United States). The process is also used in polypropylene binder twine by the American Brazilian Company in the United States under the trade name Cleanfield.

The use of the Scott-Gilead system in agriculture has established the reliability of this technology. Figure 6 shows it in use near the Dead Sea in Israel. At the end of the induction period, the polyethylene film photodegrades rapidly, and biodegradation is complete by the beginning of the following season. There has been no buildup of nonbiodegraded plastics on any of the sites where it has been used.

In parallel with the above developments, trials began on the use of the original iron thiolate system in packaging. This led to the manufacture of carrier bags in Finland in 1973 under the name Ecoten. Figures 7 and 8 illustrate the progression of the degradation of Ecoten carrier bags exposed at intervals out-of-doors in Birmingham (Scott 1976). The time delay of about 2 weeks of summer sunshine before rapid degradation commenced was introduced at the request of the user to safeguard against adventitious exposure to light of the packaging during use. A measure of the effectiveness of the iron thiolates as thermal antioxidants is the fact that 15 years after their manufacture, carrier bags made by this process and stored in the absence of light are still as strong as when they were first manufactured. This process is currently being evaluated in check-out bags (United States), carrier bags (United Kingdom), and six-pack collars (United States), all of which tend to end up as litter.



Figure 6.--Progress of photooxidation and biodegradation of Plastor (photobiodegradable) mulching film in Israel. 1--immediately after laying; 2--after cropping; 3--after ploughing. (Photographs by courtesy Plastopil Hazorea, Ltd.)



Figure 7.--Ecoten carrier bags exposed out-of-doors in Birmingham at 1-month intervals.




Figure 8.--Change in mechanical properties of Ecoten carrier bags exposed out-of-doors in Birmingham at 1-month intervals.

TECHNICAL REQUIREMENTS OF PHOTOBIODEGRADABLE PLASTICS

In spite of the successful use of photobiodegradable plastics in agriculture and to a lesser extent in packaging over the past 15 years, the packaging industry has been reluctant to accept them as a contribution toward the control of plastics litter. The arguments used against them are frequently couched in technical language, but are very often based on misconception rather than on technical fact and experience. The critics of degradable plastics are generally governed less by the published scientific evidence than by a reluctance to accept liability for the pollution resulting from the activities of their own industry. The pseudosociological arguments frequently used are intended as much for politicians as for the users of packaging.

The use of degradable plastics in agriculture was dictated by economic necessity. The same pressures are not evident in the use of degradable plastic in packaging, since rapid disintegration after use does not enhance the primary function of the package. Legislative bodies have, therefore, been reluctant to prohibit the use of nondegradable materials for purely aesthetic reasons. However, it is now clear that the situation is much more critical in the oceans, where there is a real threat to marine birds and animals. Experience in agriculture has unambiguously demonstrated that there are no economic or technical reasons why photobiodegradable plastics should not now be introduced into all bulk packaging. There is increasing evidence that the manufacturers and users of packaging are beginning to listen to the "green" movements, which advocate working with rather than against nature. The idea of returning waste plastics to the biological cycle is rapidly gaining popularity with the retailers of "organic" products and even with the large supermarkets.

However, packaging manufacturers are faced with a dilemma. The claims of the various degradation systems available are very difficult to check out. Furthermore, where they have been evaluated, the performance of some of the processes does not match up to the claims made for them. The terminology used is often confusing, and most technologists cannot distinguish between the subtleties of anaerobic and aerobic biodegradation, photodegradation, and photobiodegradation. There are at present no objective performance tests currently available which allow the user to compare the performance of degradable plastics. Until such criteria are available, it seems likely that the polymer industries will continue to argue on specious grounds against the introduction of more biocompatible, but also potentially more expensive, materials. In these circumstances, it seems unlikely that the polymer industries themselves will wish to fund the necessary research to establish the appropriate standards. The pressure for change must come from the users of packaging in association with the environmental pressure groups.

NET PACKAGING AND FISHING NETS

One of the most serious threats to marine life undoubtedly comes from ghost fishing, where both animals and fish become fatally entangled in

Table 4.--Combination effects of a photoactivator (FeDMC, 0.01 g/100 g) and a photostabilizer (NiDBC, variable) on the photooxidative stability of polypropylene.

Concentration of NiDBC, g/100 g	0	0.1	0.2	0.3	0.4
Embrittlement time, in hours	116	956	1,515	2,250	2,516

discarded nets. Much of this lethal debris arises accidentally by natural wear and tear on fishing gear, which has a finite life due to normal oxidative aging and biodegradation processes. Some of it, however, particularly in the packaging field, results from deliberate discard by the user. Most of the polymer used in these applications is polyolefin-based and so, as seen above, does not biodegrade rapidly unless sensitized to oxidation. An exceptionally wide range of lifetimes is thus required for netting, ranging from packaging, where ideally photobiodegradation should commence immediately on discard, to fishing net, where a service life of 5 years or more is required. This might at first sight appear to be an impossible achievement when coupled with the need for rapid photooxidation and biodegradation at the end of the useful life of the net. In practice, the Scott-Gilead system has been shown to be capable of providing this range of degradation times in polypropylene and high density polyethylene.

Table 4 shows that different combinations of photoactivator (FeDMC) and photoantioxidant (NiDBC) in polypropylene give a twenty-fivefold range of useful lifetimes (Scott and Gilead 1982), and more recent work has shown that this can be increased to a greater than fiftyfold range using a combination of different additives. In practice, this means that if a lifetime of 2 months is required for short-term packaging, then, using the same polymer but a different combination of the same additives, a lifetime of 8 years is technically feasible for fishing nets. This extraordinary lifetime control is accompanied in all cases by rapid photooxidation and biodegradation after embrittlement of the fiber. This process is already in commercial use in polypropylene binder twine (American Brazilian Company), where a lifetime of about 1 year is required followed by the rapid disappearance of the fiber from the field. It is fortunate that polypropylene, due to its low density, floats on the surface of water, where it is subjected to the combined effects of sunlight, oxygen, and microorganisms. This process is ideal for the protection of the environment from net packaging, ropes, and fishing nets which fail in service.

THE FUTURE

Polymer technologists have in the past been concerned to make sure that their products lasted as long as possible in the environment. It is increasingly being recognized that a more sophisticated approach is required in the future. Many products are required to last only as long as

they fulfill their useful function, and the chemistry of stabilization has now advanced to the point where predictable lifetime control of polymeric materials is not only feasible but is now well proven through the pioneering activities of Gilead and his coworkers in plasticulture (Scott and Gilead 1982; Gilead 1985; Gilead and Ennis 1987).

REFERENCES

- Al-Malaika, S., K. B. Chakraborty, and G. Scott.
1983. Developments in polymer stabilisation - 6. In G. Scott (editor), *Peroxidolylic antioxidants: Metal complexes containing sulphur ligands*, p. 73-120. Elsevier Appl. Sci. Publ.
- Al-Malaika, S., and G. Scott.
1983. Photostabilisation of polyolefins. In N. S. Allen (editor), *Degradation and stabilization of polyolefins*, p. 281-335. Elsevier Appl. Sci. Publ.
- Amin, M. E., and G. Scott.
1974. Photo-initiated oxidation of polyethylene; effect of photosensitisers. *Eur. Polym. J.* 10:1019-1028.
- Andrady, A. L.
1987. *Plastics in the marine environment. Symposium on degradable plastics*, p. 22-25. Society of the Plastic Industry, Wash., D.C.
- Brubaker, M. M.
1950. DuPont. U.S. Patent 2,495,286.
- Chew, C. H., L. M. Gan, and G. Scott.
1977. Mechanism of the photo-oxidation of polyethylene. *Eur. Polym. J.* 13:361-364.
- Claus, P.
1987. *Degradable plastics in Europe. Symposium on degradable plastics*, p. 4. Society of the Plastic Industry, Wash., D.C.
- Dixon, T. R.
1978. *Mar. Pollut. Bull.* 1:91.
- Dixon, T. R., and A. J. Cooke.
1977. Discarded containers on a Kent beach. *Mar. Pollut. Bull.* 8:105-109.
- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.
- Eggins, H. O. W., J. Mills, A. Holt, and G. Scott.
1971. Biodeterioration and biodegradation of synthetic polymers. In G. Sykes and F. A. Skinner (editors), *Microbial aspects of pollution*, p. 267-279. Acad. Press, Lond.

- Fowler, C. W., and T. R. Merrell.
1986. Victims of plastic technology. *Alaska Fish Game* 18(2):34-37.
- Gilead, D.
1985. Plastics that self-destruct. *Chemtech.*, May, p. 299-301.
- Gilead, D., and R. S. Ennis.
1987. A new, time-controlled, photodegradable plastic. *Symposium on degradable plastics*, p. 37-38. Society of the Plastic Industry, Wash., D.C.
- Grassie, N., and G. Scott.
1984. *Polymer degradation and stabilization*. Cambridge Univ. Press, Cambridge.
- Grassie, N., and G. Scott.
1985. *Polymer degradation and stabilization*. Cambridge Univ. Press, Cambridge, Chapters 1 and 4.
- Griffin, G. J. L.
1987. Degradable plastic films. *Symposium on degradable plastics*, p. 47-49. Society of the Plastic Industry, Wash., D.C.
- Guillet, J. E.
1973. *Polymers and ecological problems*. Plenum Press, N.Y.
- Guillet, J. E., T. W. Regulski, and T. B. McAneney.
1974. *Environ. Sci. Tech.* 8:923.
- Harlen, G. M., and A. Nicholas.
1987. Degradable ethylene-carbon monoxide copolymer. *Symposium on degradable plastics*, p. 14-17. Society of the Plastic Industry Wash., D.C.
- Heneman, B., and the Center for Environmental Education.
1988. Persistent marine debris in the North Sea, northwest Atlantic Ocean, wider Caribbean area, and the west coast of Baja California. Unpublished report to the Marine Mammal Commission and National Ocean Pollution Office, NOAA, Department of Commerce.
- Johnson, R.
1987. An SPI overview of degradable plastics. *Symposium on degradable plastics*, p. 6-12. Society of the Plastic Industry, Wash., D.C.
- Jones, P. H., D. Prasad, M. Hoskins, M. H. Morgan, and J. E. Guillet.
1974. *Environ. Sci. Technol.* 8:919.
- Lloyd, D. R.
1987. PHBV--biodegradable plastic. *Symposium on degradable plastics*, p. 19-21. Society of the Plastic Industry, Wash., D.C.

- Maddovar, W. J., and G. M. Chapman.
1987. Making plastics biodegradable using modified starch additions. Symposium on degradable plastics, p. 41-44. Society of the Plastic Industry, Wash., D.C.
- Mellor, D. C., A. B. Moir, and G. Scott.
1973. The effect of processing conditions on the UV stability of polyolefins. *Eur. Polym. J.* 9:219-225.
- O'Connell, C. E.
1987. Remarks of C. E. O'Connell. Symposium on degradable plastics, p. 1-2. Society of the Plastic Industry, Wash., D.C.
- Omichi, H.
1983. Photodegradable polyolefins. In N. S. Allen (editor), *Degradation and stabilisation of polyolefins*, p. 187-211. Elsevier Appl. Sci. Publ.
- Potts, J. E.
1982. The encyclopaedia of chemical technology. Section C5, p. 626-669. Wiley, N.Y.
- Redpath, A. E.
1987. Photodegradable controlled life-time plastics: A strategic environmental advantage for the plastics industry. Symposium on degradable plastics, p. 31-32. Society of the Plastic Industry, Wash., D.C.
- Scott, G.
1965. Atmospheric oxidation and antioxidants. Elsevier, Amsterdam.
1971. British Patent 1,356,107.
1972a. Plastics and coastal pollution. *Int. J. Environ. Stud.* 3:35-41.
1972b. Disposal of plastics. In Beadle (editor), *Plastic forming*, p. 161. Macmillan Engineering Evaluations.
1972c. Photodegradable plastics: An answer to pollution? *Design Engineering*, Oct., 77-79.
1973a. Delayed action photoactivator for polymers. In J. E. Guillet (editor), *Polymers and ecological problems*, p. 27. Plenum Press, N.Y.
1973b. Degradability of polymers and plastics. PRI preprints, Nov., Paper 6.
1973c. Improving the environment: Chemistry and plastics waste. *Britain* 9:267-272.

- 1975a. The growth of packaging litter. *Int. J. Environ. Stud.* 7:131-133.
- 1975b. Biological recycling of polymers. *Polymer Age* 6:54.
- 1975c. Polymers and pollution. In Penn and McAuliffe (editors), *Chemistry and pollution*. Macmillan Press.
1976. Time-controlled stabilisation of polyolefins. *J. Polym. Sci., Polym. Symp.* 57:357-374.
- 1988a. Migration and loss of antioxidants from polyethylene. *Food Additives and Contaminants*, 5(Supp. 1):421-432.
- 1988b. Potential toxicological problems associated with antioxidants. *Free Radical Res. Comm.* 5:141-147.
- Scott, G. (editor).
1979-88. *Developments in polymer stabilisation*. Elsevier Appl. Sci. Publ.
- Scott, G., and D. Gilead.
1978. British Patent 1,586,344.
1982. Time-controlled stabilisation of polymers. In G. Scott (editor), *Developments in polymer stabilisation - 5*, p. 71-106. Elsevier Appl. Sci. Publ.
1984. U.S. Patent 4,461,853.
1985. U.S. Patent 4,519,161.
- Society of the Plastics Industry.
1988.
- Statz, R. J., and M. C. Dorris.
1987. Photodegradable polyethylene. *Symposium on degradable plastics*, p. 51-55. Society of the Plastic Industry, Wash., D.C.
- Staudinger, J. J. P.
1970. Disposal of plastic waste and litter. *Society of Chemical Industry Monogr.* 35.
- Takahachi, T., and K. Suzuki.
1964. *Chem. High Polym. (Japan)* 21:494.

ENVIRONMENTAL DEGRADATION OF PLASTICS UNDER LAND AND MARINE EXPOSURE CONDITIONS

Anthony L. Andrady
Research Triangle Institute
Chemistry and Life Sciences
Research Triangle Park, North Carolina 27709, U.S.A.

ABSTRACT

Several types of thermoplastic and latex rubber materials commonly encountered in marine plastic debris were weathered outdoors in air and while floating in seawater, under North Carolina climatic conditions. The degradation of the different samples was monitored by tensile property determination.

In general, the various materials exposed outdoors in seawater tended to weather at a slower rate than the materials exposed outdoors in air. This retardation of weathering is probably a result of lack of heat buildup in samples exposed in seawater. Also, surface fouling on samples in seawater may have shielded them from light to some extent.

INTRODUCTION

In spite of their short history, synthetic polymers, particularly plastics, have gained wide popularity as the material of choice in a wide range of packaging, building, and other applications. With the current consumption of plastics reaching around 27.5 million metric tons (MT) (50 billion lb) in the United States (Modern Plastics 1988), plastics will without doubt continue to replace conventional materials such as glass, metal, wood, and paper in a variety of additional uses. The projected production in the year 2000 is expected to be 41.3 to 55.1 million MT (75 to 100 billion lb).

The popularity of plastics in packaging and other applications is attributed to the unique, useful characteristics of the material. These include light weight, excellent mechanical strength (tensile properties, tear resistance, and impact resistance), readily controllable and superior optical properties (clarity, gloss, and color), biological inertness, easy processability, low cost, and outstanding durability. Because they are synthetic materials, plastic compositions might be "tailor made," within limits, to obtain specific useful characteristics. As a response to an historically consistent consumer demand for stronger and longer lasting plastics, the industry has continually improved the durability of plastics, especially for outdoor exposure conditions.

The recent surge in the use of plastics as the material of choice in diverse applications is also reflected in uses of plastics at sea, particularly by the fishing industry. A very significant source of marine plastic debris is gear related. Introduction of plastics into the world's oceans started in the late 1940's with the changeover from natural fibers (jute, cotton, hemp) to synthetic polymer fibers in the construction of fishing gear (Uchida 1985). Today, nearly all the fishing gear used in developed countries is manufactured from durable synthetic materials (Klust 1973), and the commercial fishing industry is the prime source of plastics in the oceans.

In addition, passenger, freight, military, and research vessels, as well as beach users introduce plastic materials into the marine environment. Invariably, most of the non-gear-related plastic waste is discharged into the ocean as postconsumer waste from vessels or is washed into the ocean from the beach environment (Parker et al. 1987). The magnitude and the nature of this influx of plastics into the sea from such sources vary widely depending on the season of the year and the geographic region (Pruter 1987).

An inevitable consequence of increased usage of plastics, particularly in packaging applications, is the increased amounts of postconsumer plastic waste. The municipal waste stream (amounting to about >88,000 MT (160 million lb) annually in the United States) in urban areas now consists of about 7% postconsumer plastics, a figure that may increase to more than 9% by the year 2000 (Franklin Associates 1988). While accurate estimates of their lifetimes in the environment are not reliably known, plastics are perceived as being exceptionally persistent materials requiring hundreds of years of exposure to facilitate biodegradation.

Parallel estimates for the quantities of plastic waste in the world's oceans are not available. Estimates by several workers (Dahlberg et al. 1985; Pruter 1987), though probably very crude, allow an appreciation of the magnitude of waste at sea.

With a total annual world catch of about 45 million MT of fish (Parker et al. 1987), a substantial amount of plastic fishing gear is routinely introduced into the ocean. The estimates of worldwide losses of commercial gear vary from a low of about 750 MT tons annually (National Academy of Sciences (NAS) 1975) to as much as 75,000 MT/year (Merrell 1980). In addition to gear losses, fishing vessels also discharge "domestic" plastic waste. A 1986 estimate places the number of commercial vessels operating annually in the United States at 125,700 (Parker and Yang 1986). The world's fleet of fishing vessels is believed to discharge 23,000 tons of plastics annually into the sea (Horseman 1985).

A detailed discussion of the ecological concerns related to plastic debris at sea is beyond the scope of this paper. Several excellent reviews on the fate of plastic debris, the specific hazards posed by such debris in specific marine species, and the general impact of plastics on the populations of target species have been published (Day et al. 1985; Center for Environmental Education (CEE) 1987; Laist 1987).

Available evidence indicates entanglement by the debris and the ingestion of the debris to be the primary concerns with a variety of affected marine animals (including birds, turtles, marine mammals, and fish). These affected populations seem to seek out the debris (either mistaking it for prey or because of mere curiosity); such behavior leads to more fatalities than might be expected on the basis of random encounters with debris. The invariable association of either entangled fish or residual food in most of the plastic waste discharged into the sea also concentrates these species in the same geographic locations that have high incidences of plastic waste (Laist 1987). Recent declines in the natural populations of the Hawaiian monk seal, *Monachus schauinslandi*, by 4 to 8% per year have been attributed, at least in part, to entanglement in plastic waste (Fowler 1985, 1987).

PLASTICS AT SEA

Key Characteristics of Marine Plastic Waste

The potential negative impact of waste plastics on the marine resource depends upon the following key characteristics of the material.

- **Geometry.** Shape of the debris is important from the point of view of entanglement. Products such as six-pack rings and netting represent more of a potential hazard than an equivalent mass of the same polymer in the form of a laminate.
- **Durability.** The likelihood of encounter between a given item of marine debris and a marine animal depends upon the lifetime of the material. The duration available for the encounter is crucial in determining the potential hazard posed by the plastic material. Unfortunately, little information is available on the lifetime of plastics at sea. Lack of this information is a definite setback in the assessment of potential hazards posed by plastic waste.
- **Strength.** Strength of the debris material determines the likelihood that an entangled animal can escape. Alternatively, the possible obstruction of the gut in case of ingestion is less likely if the material is weak enough to mechanically fail during the ingestion process.
- **Toxicity.** Plastics, being undigestible macromolecules, cannot be absorbed through the gut lining. They are, therefore, not toxic materials. However, the plastics used in the fabrication of products may contain chemical additives which can be absorbed and assimilated.

Of these characteristics, lifetime is perhaps the most important. An attempt was therefore made in the present work to determine the relative lifetimes of some relevant debris items exposed on land and floating in seawater. The study will determine if the plastic materials floating on

seawater degrade at rates different from those exposed to the same natural weathering conditions, but in air.

Weathering of Plastics Under Marine Exposure Conditions

Sunlight-induced degradation is the principal mechanism of weathering of plastics outdoors. As sunlight is freely available to plastics floating in the sea, weathering might be expected to occur at sea at rates comparable to those on land. However, there are several reasons to expect the rate of degradation at sea to be different from that on land.

- High humidity is known to accelerate the rates of degradation of several classes of plastics (Davis and Sims 1983). This may be brought about by the "plasticizing" action of small quantities of sorbed water leading to increased accessibility of the matrix to atmospheric oxygen or by the leaching out of stabilizing additives from the formulation.
- Plastics exposed to sunlight outdoors undergo "heat buildup," a process which results in the plastic material reaching significantly higher temperatures than the surrounding air (Summers et al. 1983). The higher temperatures generally result in an acceleration of light-induced degradation and may even be high enough to induce significant thermooxidative degradation. Plastics at sea will not suffer from such heat buildup and may consequently undergo slower oxidative degradation and photodegradation.
- All materials exposed to the sea invariably undergo fouling (Fischer et al. 1984). In the initial stages of fouling, a biofilm forms on the surface of plastic. Gradual enrichment of the biofilm leads to a rich algal growth within it. Consequently, the biofilm becomes opaque, and the light available to the plastic for photodegradation is restricted. Thus, the rate of photodegradation at sea might be determined in part by the rate of fouling.
- Advanced stages of fouling are characterized by the colonization of the plastic surface by macrofoulants such as bryozoans. The weight of the macrofoulant and that of debris they entrap might even partially submerge the material. As the ultraviolet portion of sunlight is attenuated on passage through seawater, submerged plastics would necessarily undergo a slower rate of photodegradation. Microbe-rich foulant film may, however, also tend to accelerate the biodegradation process by providing a rich biotic population in contact with the plastic surface.

DEFINITIONS DEVELOPMENT

An important issue relating to the discussion of degradability and enhanced degradable plastics is that of definitions. The various terms are

used in a loose manner in the literature with no consistency. The need to develop adequate definitions has been pointed out recently by Andrady (1988) and others (General Accounting Office 1988). Definitions of the term "plastics" as it applies to the MARPOL Annex V Convention and the following terms relating to "degradability" are proposed as a starting point for further development.

For the purposes of MARPOL Annex V, the following definition of plastics has been proposed.

Plastic: A solid material which contains as an essential ingredient one or more synthetic organic high polymer; and is formed (shaped) during either manufacture of the polymer or the fabrication into a finished product by heat and/or pressure.

The proposed definition includes both rubber and plastics (plastic products as well as virgin resin pellets). Inorganic polymers such as glasses are excluded along with low molecular weight polymers which are not "high" polymers or solids. The latter includes polymeric waxes, varnishes, and lubricants. The definition excludes any polymers produced by living organisms, including cellulose, natural rubber, and bacterial polyesters. In the case of a composite material where one component is a polymer, the material is excluded if the polymer itself is a minor component not essential to the formulation.

Deterioration: Embrittlement and/or loss of physical integrity of a polymer regardless of the mechanism which brings about these changes.

The deterioration process might be the result of either a chemical or a physical process. Nonchemical deterioration of plastics plays an important role in environmental deterioration of plastics.

Degradation: Deterioration which results from a chemical process.

Degradation might be further subdivided, based on the agency causing the chemical change. "Photodegradation" refers to degradation brought about by light. "Biodegradation" is that due to living organisms. Degradation due to slow oxidation of the plastic (especially at elevated temperatures) is "thermooxidative degradation," while that due to the chemical action of water is hydrolytic degradation or "hydrolysis." To be consistent with the definition for "degradation" proposed herein, it is necessary to show that a loss in property (in this case, tensile properties) occurred and that such loss is a result of a chemical reaction. The term "degradation" has been used throughout this paper without establishing that the loss was due primarily to a chemical change. However, the chemical nature of the processes which result in strength loss in plastics exposed to outdoor environments are well known.

The nonchemical deterioration processes might be similarly subdivided. "Dissolution" (or swelling in water), in which no chemical changes take

place, might be viewed as a deterioration of the plastic in water (or an aqueous solution). "Biodeterioration" can take place when the plastic is attacked by borers (at sea) and rodents: The net result is the breakdown of a larger piece of plastic into small fragments. In "physical deterioration" a plastic loses strength due to purely physical phenomena (for instance due to freeze-thaw cycles).

This proposed scheme allows a facile classification of the related processes and makes a clear distinction between degradation which tends to chemically break down plastics, and nonchemical deterioration processes, which merely reduce the particle size of the plastic. The distinction is of obvious importance from an environmental point of view.

EXPERIMENTAL

Materials

The following plastic products, selected on the basis of reported composition of beach debris (CEE 1987), were included in this study:

- Polyethylene film (low density). Representative of the plastic used in six-pack rings, plastic bags, etc.
- Polypropylene strapping tape. Commercially available.
- Trawl netting material (orange and blue-green color).
- Latex rubber balloons. Commercial sample.
- Foamed polystyrene sheet. Commercial sample.
- Rapidly degradable polyethylene. Commercial sample.

Polyethylene bags are a well-known component of marine debris. The threat to marine turtles via ingestion of plastic bags has been reported (Balazs 1985). While strapping bands (usually made of polypropylene or polyester) are not a major component of the debris, they present a particularly severe threat of entanglement to marine mammals (Laist 1987). Trawl webbing is a major component of floating plastic debris in some regions (Pruter 1987). A fraction of the latex rubber balloons released in promotional events may eventually reach the oceans, where their ingestion may present a threat to turtles and other species.

Weathering and Sampling

Experiments were carried out at the exposure facility at Duke Marine Laboratory in Beaufort, North Carolina. Figure 1 shows the ambient temperature of the seawater during the period of exposure and the air temperatures for the area as recorded by the National Weather Service.




Figure 1.--Average monthly temperatures and rainfall for Morehead City, North Carolina for 1987.

The samples exposed on land were affixed with staples to a wooden platform and exposed horizontally on the flat roof of a laboratory building. These were backed by wood and were about 15.2 cm (6 in) from the roof surface. Another set of samples was exposed floating in a tank of seawater, with fresh seawater continuously flowing through the tank to maintain a depth of about 30.5-45.7 cm (12-18 in) of water at all times. Exposure in the tank as opposed to directly at sea has several associated advantages.

In the preliminary experiments carried out with samples directly floating in an enclosed section of sea, the samples tended to accumulate mud and debris on the surface due to tidal action. Exposure within the tank ensured minimal accumulation of soil and other floating debris on the sample while providing a fresh, clear, biologically active seawater medium. The experiment thus simulates the conditions best suited for rapid photodegradation. The exposure was carried out for a period of 1 year.

Sampling was carried out at the end of every second month for all samples. The exposed samples were placed in a black plastic bag and transported to the Research Triangle Institute for measurement of tensile properties. The samples exposed at sea were dried for about 3 h in an air oven at 50°C and were stored in the dark at ambient temperatures.

Tensile Testing

Measurement of tensile properties was carried out with an Instron Mechanical Tester, Model 1122 generally in accordance with ASTM D 638 (American Society for Testing and Materials) Tensile Properties of Plastics. No further preconditioning of the samples was done prior to testing. Air-powered grips were used to hold the samples. Smooth grip faces were used with the polyethylene samples; for latex balloons and strapping tape, a serrated face had to be used to avoid slippage. Table 1 gives the test parameters for various types of samples tested. In the case of trawl webbing and strapping tape, where the fibrous nature and surface markings, respectively, made it difficult to determine the true area of cross section, the load to break is reported.

RESULTS AND DISCUSSION

Polyethylene Film

Table 2 summarizes the tensile property data for the polyethylene film samples exposed on land and at sea. Both the sea and air samples exhibit an increase in tensile strength after 2 months. This increase may be due in part to relaxation of stresses frozen into the sample during processing, a common occurrence during early exposure of processed thermoplastics.

Figure 2 shows the variation of ultimate extension with exposure time. Clearly, the samples exposed floating on seawater degraded at a much slower rate. The samples exposed at sea showed a 12% loss in ultimate extension after 12 months while the air-exposed samples lost 95% of the ultimate extension after only 6 months. Ultimate extension is considered a more

Test parameter ^a	A	B	C	D	E
Beam capacity (kg)	1,000	1,000	1,000	1,000	1,000
Full scale load (kg)	10	20,100	5	200	2/5
Crosshead speed (mm/min)	100	100	20	100	50
Gauge length (cm)	5	1.5	4	4	3
Clamp	Pneumatic	Pneumatic	Pneumatic	Pneumatic	Pneumatic
Jaw face size (in)	1 × 1.5	1 × 1.5	1 × 1.5	1 × 1.5	1 × 1.5

^aA - Polyethylene film and degradable polyethylene, B - strapping tape, C - Styrofoam sheets, D - trawl netting, E - balloons.

appropriate parameter than tensile strength for measuring physical degradation since it reflects the brittleness and consequent tendency of the plastic to fragment. Statistical tests at the 0.05 level of significance showed no significant difference in mean ultimate extension values of the 2- and 12-month samples exposed at sea but did show a significant difference in mean ultimate extension of the 2- and 6-month samples exposed in air.

Polypropylene Tape

Table 3 gives the summary data relating to the weathering of polypropylene strapping tape. The formulation contained a filler and the material was highly anisotropic, easily tearing along its length. Material did not "neck" on extension but ruptured gradually. As the surface of the tape was not smooth enough (because of an embossed pattern on the surface) to obtain an accurate value for thickness, the maximum load rather than tensile strength is reported. Figure 3 illustrates the observed changes in ultimate extension. After 12 months, samples exposed on land had lost 90% of the initial ultimate extension, while samples exposed at sea had lost only 26% of the initial value. Thus, while some degradation does occur in samples exposed on seawater, it is much less pronounced than that for samples exposed in air for a comparable duration of exposure. This conclusion is further illustrated by testing for the statistical significance of differences in mean ultimate extensions at the 0.05 level. A significant difference exists in mean ultimate extension for the samples exposed in air for 0 and 12 months and for those exposed at sea for 0 and 12 months. As expected, a statistically significant difference was also found between the mean ultimate extensions of the 12-month samples exposed at sea and in air.

Trawl Netting

Tensile property data for net samples are given in Table 4. The data are reported as maximum load (kg), which often coincided with the ultimate load of the material, and ultimate extension.

Table 2.--Summary of data relating to weathering of low-density polyethylene film samples.

Duration (months)	Tensile strength (kg/cm ²)			Ultimate extension (%)			Number of samples
	Mean	S.D.	S.E.	Mean	S.D.	S.E.	
Samples exposed in air							
0	124.1	19.6	6.1	548	71	29	6
2	143.1	9.9	4.4	541	38	17	5
4	99.9	5.1	2.9	188	166	96	3
6	115.8	6.5	3.3	27	18	9	4
Samples exposed in seawater							
2	139.5	17.1	7.7	613	133	59	5
4	131.0	12.8	5.7	547	95	42	5
6	132.3	23.6	13.7	601	197	114	3
8	117.3	13.4	6.0	511	147	65	5
10	117.8	7.3	2.9	550	106	46	6
12	118.7	7.6	3.4	541	87	39	5

Table 3.--Summary of data relating to weathering of plastic strapping tape.

Duration (months)	Maximum load (kg)			Ultimate extension (%)			Number of samples
	Mean	S.D.	S.E.	Mean	S.D.	S.E.	
Samples exposed in air							
0	75.5	2.0	1.0	82	2	1	4
2	68.2	1.7	0.8	70	7	3	4
4	40.2	5.2	2.6	43	4	2	4
6	20.1	3.2	1.6	19	5	2	4
8	14.9	3.5	1.8	12	4	2	4
10	13.2	2.7	1.4	10	5	1	4
12	11.3	0.7	0.4	8	1	1	3
Samples exposed in seawater							
0	76.5	5.4	2.7	89	5	2	4
4	77.0	4.0	2.0	91	3	2	4
6	74.3	2.5	1.3	82	2	1	4
8	73.2	5.6	2.8	79	5	3	4
10	64.0	5.2	2.6	63	8	4	4
12	67.2	3.3	1.6	61	9	4	4

Note: Sample width was half the size of regular width of the tape.




Figure 2.--The variation of the mean ultimate extension of polyethylene films with the duration of exposure.




Figure 3.--The variation of the mean ultimate extension of polypropylene tapes with the duration of exposure.

Table 4.--Summary of data relating to weathering of trawl web material.

Duration (months)	Maximum load (kg)			Ultimate extension (%)			Number of samples
	Mean	S.D.	S.E.	Mean	S.D.	S.E.	
Orange-colored netting							
Samples exposed in air							
0	126	3.8	1.9	46.5	4.8	2.4	4
2	121	13.8	6.9	36.9	2.7	1.4	4
4	120	10.6	7.5	41.0	5.9	4.2	3
6	117	9.3	4.7	41.7	5.6	2.8	4
8	125	4.0	2.0	47.4	1.7	0.9	4
10	121	7.7	3.9	47.7	6.5	3.2	4
12	125	8.5	4.3	49.1	8.4	4.2	4
Samples exposed in seawater							
4	132	9.1	4.6	62.1	3.8	1.9	4
6	123	13.4	6.7	49.1	4.1	2.1	4
8	129	6.7	3.3	53.5	2.9	1.4	4
10	128	10.7	5.4	53.5	2.9	1.4	4
12	127	11.6	5.8	49.1	3.9	2.0	4
Blue-colored netting							
Samples exposed in air							
0	115	10.5	5.2	63.0	7.1	3.5	4
2	88	11.4	5.7	41.4	2.9	1.5	4
4	104	7.9	4.0	46.6	8.2	4.1	4
6	96	11.3	5.7	49.1	8.6	4.3	4
8	70	12.1	6.1	32.3	10.9	5.5	4
10	93	7.3	3.7	44.5	3.5	1.7	4
12	94	3.8	1.9	49.5	5.0	2.5	4
Samples exposed in seawater							
2	100	12.0	6.0	65.7	9.9	5.0	4
4	96	9.2	4.6	53.4	9.8	4.9	4
6	99	7.3	3.6	60.2	2.5	1.3	4
8	101	7.5	3.8	61.6	5.8	2.9	4
10	113	2.8	1.4	61.8	5.3	2.7	4
12	104	3.2	1.6	60.4	1.4	0.7	4

The changes in tensile properties obtained with trawl web samples on exposure in air and in seawater are less dramatic and are not shown in a figure. Netting intended for commercial fishing is compounded specifically for exceptional durability usually using hindered-amine and other light stabilizers. The data, however, did show that the blue netting is relatively more prone to outdoor degradation than is orange netting when exposed in air. While the difference is small, it is significant. On exposure in seawater, neither material underwent any significant change in maximum load up to 12 months of exposure (at which time the experiment was discontinued). The only conclusion that might be drawn from these samples is that they would persist longer in the environment, relative to the packaging materials and balloons tested.

Rubber Balloons

The strength and extensibility of the rubber balloons determine to a great extent the likelihood of the ingested material obstructing the air or gut passages of turtles. Retention of elasticity is of particular concern, as elastic materials are likely to be difficult to dislodge from the air passages or alimentary canals. Table 5 summarizes the tensile property data on balloons exposed under present experimental conditions. Figure 4 shows the variation of ultimate extension with duration of exposure. In air, the rubber lost 59% of its ultimate extension after only 2 months. For the same time at sea, the rubber lost only 11% of its ultimate extension. The balloons continued to retain their elasticity during exposure in seawater, with only a 48% loss after 12 months. In air, however, the balloons lost 94% of their ultimate extension after 6 months, beyond which time the samples were too weak and brittle to be tested. As with the plastic samples, the rate of degradation in seawater was much slower than that in air. The degree of hazard associated with a partially deteriorated balloon depends on the particle size which might be safely ingested by the target species. Such information on turtles and other relevant species is not available at the present time. However, the above results indicate that if the balloons do pose a hazard to marine life, they would, under present experimental conditions, be a threat for a relatively longer period of time at sea than on land.

Polystyrene Foam

In view of the abundance of polystyrene foam pieces in marine debris, the weathering behavior of polystyrene was particularly interesting. On exposure in air the foam underwent rapid yellowing, which apparently was a surface reaction. The sample exposed on seawater also underwent yellowing, although the algal fouling of the surfaces made it difficult to measure the extent of yellowing.

The yellowness index increased up to about the sixth month of exposure and decreased thereafter. However, the development of yellowness was also accompanied by embrittlement of the exposed surface. Over the exposure period of 1 year, a surface layer of up to half the original thickness became brittle enough to crumble on handling (and could be easily scraped out). Wind and rain are likely to remove at least some of the yellowed material during exposure. This may explain the reduction in the extent of yellowing at the longer exposure times.

Table 5.--Summary of data relating to weathering of latex rubber balloons.

Duration (months)	Tensile strength (kg/cm ²)			Ultimate extension (%)			Number of samples
	Mean	S.D.	S.E.	Mean	S.D.	S.E.	
Samples exposed in air							
0	96.7	7.2	3.6	986	100	50	5
2	3.6	1.9	0.9	405	184	92	4
4	1.9	--	--	140	--	--	2
6	1.4	--	--	63	--	--	2
^a 8	--	--	--	--	--	--	--
^a 10	--	--	--	--	--	--	--
Samples exposed in seawater							
2	22.7	3.4	1.5	874	107	48	5
4	21.5	5.4	2.4	727	75	34	5
6	16.0	3.1	1.5	611	69	34	4
8	14.0	3.6	1.8	600	87	44	4
10	18.3	3.5	1.7	719	74	37	4
12	9.1	1.0	0.6	513	26	15	3

^aToo brittle or weak to be tested.

The yellowness index of the degraded polystyrene foam correlates well ($r = 0.90$) with the tensile strength of the degraded material up to 6 months of exposure in air. Lack of such a correlation at longer exposure times is also possibly due to loss of embrittled yellow surface material (from rain, wind).

In fact, the thickness of the degraded (removable) yellow surface layer increased with duration of exposure for both sets of samples. The reduced thickness of the samples after the embrittled layer was scraped off is given below.

Duration Months)	Thickness of lower layer (cm)	
	Air	Seawater
0	0.418	0.418
2	0.349	0.221
4	0.308	0.164
6	0.234	0.168
8	0.217	0.229
10	0.214	0.155




Figure 4.--The variation of mean ultimate extension of latex rubber balloons with the duration of exposure.

The tensile strength can thus be calculated in two ways: based on original thickness, and based on the thickness of the unembrittled layer. Table 6 summarizes the tensile property data. If the oxidative degradation process was restricted to the yellowed brittle surface layer, the tensile strength of the underlying polystyrene would yield about the same tensile strength regardless of the duration of exposure. However, as seen in the table and in Figure 5, the tensile strength based on reduced thickness of the material also decreases with the duration of exposure. The lower unembrittled region is apparently accessible to the free radicals generated during the photo reaction.

Expanded polystyrene was the only type of plastic material tested where the rate of deterioration (of tensile properties) was faster at sea than on land. In air, the material requires an exposure of at least a year to decrease its tensile strength by 40%. Exposure in seawater reduces the tensile strength by over 60% in 4 months!

Under the present exposure conditions, the polystyrene foam material deteriorates relatively rapidly when exposed outdoors on seawater. This would lead to the breaking up of the material into smaller pieces fairly easily. Unlike most other plastic debris items, pieces of foamed polystyrene are not capable of entanglement. They might, however, be ingested by a variety of species, especially when covered with foulants. Effects of ingestion of weathered polystyrene foam material are not known.

Table 6.--Summary of data relating to weathering of expanded-extruded polystyrene.

Duration (months)	Tensile strength ^a (kg/cm ²)			Tensile strength ^b (kg/cm ²)			Ultimate extension (%)			Number of samples
	Mean	S.D.	S.E.	Mean	S.D.	S.E.	Mean	S.D.	S.E.	
Samples exposed in air										
0	3.89	0.50	0.21	3.89	0.50	0.20	3.9	1.70	0.07	6
2	4.31	0.34	0.17	5.16	0.40	0.20	3.5	0.29	0.14	4
4	3.46	0.59	0.29	4.70	0.80	0.40	3.9	0.32	0.16	4
6	2.45	0.27	0.19	4.37	0.48	0.24	2.9	0.13	0.06	4
8	2.39	0.27	0.19	4.60	0.52	0.26	3.2	0.24	0.12	4
10	2.61	0.14	0.07	5.09	0.27	0.14	3.2	0.24	0.12	4
12	2.37	0.27	0.19	4.53	0.51	0.25	3.3	0.32	0.16	4
Samples exposed in seawater										
2	2.88	0.21	0.09	5.45	0.40	0.18	4.6	0.80	0.36	5
4	1.13	0.71	0.36	5.50	1.82	0.91	4.1	1.60	0.80	4
6	1.09	0.17	0.09	3.20	0.44	0.22	2.2	0.88	0.44	4
8	1.22	0.29	0.13	2.22	0.54	0.25	1.9	0.29	0.13	5
10	0.69	0.09	0.04	2.13	0.22	0.11	1.6	0.24	0.12	4

^aTensile strength calculated using the initial area of cross section.

^bTensile strength calculated using the area of cross section based on residual unembrittled layer.

Enhanced Photodegradable Polyethylene (Ethylene Carbon Monoxide Copolymer)

The weathering behavior of the enhanced degradable polyethylene was quite different from that of the polyethylene homopolymer sample (Table 7). As might be expected, the samples exposed in air rapidly degraded, losing nearly 99% of the initial value of mean ultimate extension within 6 weeks of exposure (Fig. 6). The tensile strength decreased slowly reaching to about 50% of the initial value in the same period of exposure. At this stage samples were embrittled and too weak to be tested.

The samples exposed in seawater also degraded rapidly on exposure losing nearly 95% of the initial value of mean ultimate elongation in about 6 weeks. However, the material did not reach the same stage of final embrittlement obtained with samples in air until after 14 weeks of exposure. The mechanism leading to a plateau in the mean extension values from about the sixth to fourteenth week of exposure is not understood. But it is of little practical consequence. It is clear that under the experimental conditions of the study, the enhanced photodegradable polyethylene performs




Figure 5.--The variation of mean tensile strength of expanded extruded polystyrene with the duration of exposure.




Figure 6.--The variation of mean ultimate extension of enhanced degradable polyethylene sheets with the duration of exposure.

Table 7.--Summary of data on outdoor weathering of enhanced photodegradable six-pack ring material (LDPE).

Duration (weeks)	Tensile strength (kg/cm ²)		Ultimate extension (%)	
	Mean	S.E.	Mean	S.E.
Exposure in air				
0	160.4	1.0	398	3.4
1	122.4	2.1	35.6	8.3
2	128.0	2.4	21.0	4.4
3	134.1	2.4	16.8	1.4
4	104.7	8.0	10.7	2.6
5	86.3	6.9	5.7	0.8
Exposure in seawater				
0	160.4	1.0	398	3.4
1	112.0	3.7	145.6	23.3
2	112.3	1.0	42.4	3.3
3	115.1	0.8	42.3	5.4
4	120.4	0.9	25.8	4.1
5	116.5	2.6	44.1	16.7
6	120.2	12.6	19.1	1.0
7	122.9	0.8	21.0	3.1
8	121.1	0.4	17.71	0.5
9	119.7	2.8	18.4	1.4
10	122.6	0.5	18.1	0.6
11	122.8	0.5	22.6	5.1
12	116.3	5.7	11.3	1.3
13	119.1	14.0	19.4	0.8
14	73.77	21.8	13.9	2.2
15	58.9	6.6	6.9	1.7

satisfactorily at sea for all practical purposes. The initial rates of decrease in the mean tensile properties obtained with exposure in seawater, are somewhat slower than those obtained with exposure on land.

SUMMARY FINDINGS

Table 8 illustrates the general findings of the exposure study by a comparison of tensile properties before and after exposure in air and in seawater for the different samples. Data relating to a single duration of exposure are shown to illustrate the general trend observed. Except for the netting, rates of degradation for the samples in seawater were much

Table 8.--Comparison of weathering data for exposure on land and at sea.

Sample	Duration of exposure (months)	Percent decrease in the mean value of tensile property			
		Air		Seawater	
		Strength ^a	Extension	Strength ^a	Extension
Polyethylene film	6	6.6	95.1	No change	No change
Polypropylene tape	12	85.0	90.2	11.0	31.5
Latex balloons	6	98.6	93.6	83.5	38.0
Expanded polystyrene	10	32.9	18.0	82.3	65.2
Netting	12	No change	No change	No change	No change
Rapidly degradable polyethylene	1.25	46.2	98.6	27.1	88.9

^aThe percentages reported as based on the maximum load in the case of netting and polypropylene type materials.

slower than the degradation rates on land. Netting material did not show any significant variation in tensile properties due to the type or duration of exposure. Enhanced degradable polyethylene six-pack ring material degraded in about the same time scale under both air and seawater exposure.

The marked retardation of the weathering process observed in some types of plastic materials floating in seawater might be attributed to: (a) differences in heat buildup and (b) fouling of samples in seawater.

A significant fraction of the sunlight impinging on a plastic surface is absorbed by the material as heat. Depending on the nature of the plastic, the velocity of the air around it, and the temperature difference between the plastic and the surroundings, this absorbed energy maintains the plastic at a temperature higher than that of the surrounding air (Summers et al. 1983). Plastics exposed in air undergo heat buildup easily. The effect is even more pronounced in the present samples, which were exposed on a thermally insulating wood surface. Under such conditions, the heat buildup is likely to be higher than for the case of exposure on soil, thus simulating weathering under "worst case" conditions and accelerating the degradation process in the samples exposed in air. Samples exposed in seawater, however, are held at near ambient temperature leading to slower rates of degradation.

Samples floating on seawater underwent extensive fouling during the exposure. Foulants were mostly algae except for several *Balanus* sp. found on samples exposed for over 8 months. The experimental method used in the present study (containment of samples in a shallow tank) is likely to have reduced the extent of fouling and prevented the settlement of debris (or

"silting") on the sample surface. Fouling retards the photodegradation by restricting the light available to the plastic.

CONCLUSIONS

Under present experimental conditions, low density polyethylene film, polypropylene tape, and latex rubber balloon samples were found to significantly degrade when exposed in air outdoors. Marked decreases in ultimate extension were obtained in 1 year of exposure. Similar samples exposed at the same site, but floating on seawater, degraded at a significantly slower rate during the same period of time.

The lower rates of degradation might primarily be attributed to lack of heat buildup in samples exposed on seawater. Biofouling of sample surface leading to reduced light availability may also have decreased the rate of weathering.

Foamed polystyrene degraded at a faster rate in seawater than in air when exposed outdoors.

Enhanced degradable six-pack ring polyethylene degraded at nearly the same rate when exposed in air and in seawater.

ACKNOWLEDGMENTS

This work was supported by the National Marine and Fisheries Service (NMFS), NOAA. The technical assistance and guidance provided by James M. Coe (Alaska Fisheries Science Center, NMFS) is acknowledged with thanks.

REFERENCES

- Andrady, A. L.
1988. Plastics that degrade faster. *ASTM Standardization News*, p. 46.
- Balazs, G. H.
1985. Impact of ocean debris on marine turtles: Entanglement and ingestion. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 387-429. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Center for Environmental Education.
1987. *Plastics in the ocean: More than a litter problem*. Center for Environmental Education, Wash., D.C., 128 p.
- Dahlberg, M. L., and R. H. Day.
1985. Observations of man-made objects on the surface of the North Pacific Ocean. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 198-212. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

- Davis, A., and D. Sims.
1983. Weathering of polymers. Appl. Sci. Publ., Lond.
- Day, R. H., D. H. S. Wehle, and F. C. Coleman.
1985. Ingestion of plastic pollutants by marine birds. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 344-386. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.
- Fischer, C. E., V. J. Castelli, S. D. Rogers, and H. R. Bleile.
1984. Technology for control of marine biofouling. In J. D. Costlow and R. C. Tipper (editors), Marine biodeterioration: An interdisciplinary study. Naval Inst. Press, Annapolis, MD.
- Fowler, C. W.
1985. An evaluation of the role of entanglement in the population dynamics of northern fur seals on the Pribilof Islands. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 291-307. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

1987. Marine debris and northern fur seals. A case study. Mar. Pollut. Bull. 18:326-335.
- Franklin Associates.
1988. Characterization of municipal solid waste in the United States 1960-2000. Franklin Associates.
- General Accounting Office.
1988. Degradable plastics. Standards, research and development. General Accounting Office/RCED-88-208, pl.
- Horseman, P. V.
1985. Garbage kills. BBC Wildlife, Aug., p. 391-393.
- Klust, G.
1973. Netting material for fishing gear. Food and Agriculture Organization/Fishing News Books Ltd., England.
- Laist, D.
1987. Overview of the biological effects of lost and discarded plastic debris in the marine environment. Mar. Pollut. Bull. 18:319-326.
- Merrell, T. R., Jr.
1980. Accumulation of plastic litter on beaches of Amchitka Island, Alaska. Mar. Environ. Res. 3:171-184.
- Modern Plastics.
1988. U.S. resin sales by process and market. Modern Plastics, January 1988, p. 96.

National Academy of Sciences.

1975. Marine litter. In *Assessing potential ocean pollutants*, p. 405-438. A report of the study panel on assessing potential ocean pollutants to the Ocean Affairs Board, Commission on Natural Resources, National Academy of Sciences, National Research Council Wash., D.C.

Parker N. R., S. C. Hunter, and R. J. Yang.

1987. Development of methodology to reduce the disposal of non-degradable refuse into the marine environment. Final report on NOAA Contract 85-ABC-00203.

Parker, N. R., and R. J. Yang.

1986. Development of methodology to reduce the disposal of non-degradable refuse into the marine environment. Paper presented at the 6th International Ocean Disposal Symposium, 21-25 April 1986, Pacific Grove, CA.

Pruter, A. T.

1987. Sources, quantities and distribution of persistent plastics in the marine environment. *Mar. Pollut. Bull.* 18:305-310.

Summers, J. W., E. Rabinovitch, and G. J. Queensbury.

1983. Predicting heat buildup due to the sun's energy. *J. Vinyl Technol.* 5:110.

Uchida, R. N.

1985. The types and estimated amounts of fish net deployed in the North Pacific. In R. S. Shomura and H. O. Yoshida (editors), *Proceedings of the Workshop on the Fate and Impact of Marine Debris*, 26-29 November 1984, Honolulu, Hawaii, p. 37-108. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

RECYCLING OF MARINE PLASTICS DEBRIS THROUGH MELT REPROCESSING:
A CASE FOR LOST OR ABANDONED FISHING GEAR

M. Xanthos, P. G. Kelleher, A. Patel
Polymer Processing Institute at
Stevens Institute of Technology-Castle Point on the Hudson
Hoboken, New Jersey 07030 U.S.A.

and

W. G. Gordon
New Jersey Marine Sciences Consortium
Sandy Hook Field Station
Fort Hancock, New Jersey 07732, U.S.A.

ABSTRACT

In attempts to offer a technological solution to the marine plastic debris problem, the Polymer Processing Institute in collaboration with the New Jersey Marine Sciences Consortium began in January 1989 a 2-year research program that focuses on the collection, characterization, and subsequent recycling of unusable plastic fishing gear. The present paper contains initial data developed during the first quarter of 1989 (Phase I) on netting containing nylon-6, nylon-6,6, and polypropylene plastics. Molecular weight determination and thermal analysis suggest that none of the nylons are badly degraded; thus, the nets could be reprocessed without difficulties in the recycling extruder. However, the presence of fiber coatings (e.g., "green enamel," "tar") was found to complicate the polymer characterization and to interfere with the recycling process. The fact that these coatings cannot be easily removed suggests that further work is required to determine their possible adverse effects on the processing and product characteristics. The present paper also provides a complete description of the work planned for the two successive phases of the project.

INTRODUCTION

Among the various marine plastic debris items, abandoned or lost fishing gear is a major pollutant adversely affecting marine life and disturbing the ecological equilibrium. Recent estimates of the National Academy of Sciences bring the amount of fishing gear lost in the sea to about 135,000 metric tons/annum (Lautenberg 1987). Many marine species die from entanglement in such plastic debris. It is believed that 30,000

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

northern fur seals, *Callorhinus ursinus*, and at least 250,000 birds die each year in nets laid by salmon fishermen, as well as in other debris. The problem appears to become increasingly important as a result of the recent deployment in the North Pacific of huge drifting nets stretching up to 30-50 km (Shabecoff 1989). If these lightweight monofilament nets are discarded or lost, they do not degrade but continue to drift unless retrieved.

The Polymer Processing Institute, in a joint effort with the New Jersey Marine Sciences Consortium (NJMSC), began in January 1989 a 2-year program having as an ultimate goal the development of cost-effective methods for complying with the prohibition of marine pollution. The project is funded by the Saltonstall-Kennedy Act and will attempt to offer a technological solution to the marine plastic debris problem. Its specific objective is to establish incentives for recycling lost or abandoned fishing gear such as nets, lines, and ropes by demonstrating that these materials can be converted into useful products when melt reprocessed.

DESCRIPTION OF THE RESEARCH PROJECT

This project consists of two phases: Phase I, completed in January 1990, concentrates on the collection, on a national level, of discarded fibrous fishing gear, followed by separation, identification, and subsequent chemical and physical characterization of the plastics involved. Preliminary data indicate that the major types of plastics to be recovered are nylon-6, nylon-6,6, and polypropylene (PP), all possibly contaminated or degraded as a result of prolonged exposure in the marine environment. The extent of environmental degradation and its effect on rheological and mechanical properties will dictate the design of the melt reprocessing experiments of the next phase.

In Phase II of the project (to be completed in its second year), various processing methods will be evaluated with the final objective of producing molding materials with properties equivalent or superior to those of virgin resins. Reprocessing will be conducted in a twin-screw extruder and will involve blending of the recovered netting with virgin resin, blending with other recycled resins of different chemical structure, or chemical modification. It is expected that these methods will upgrade the properties of the recovered plastics and produce value-added compounds. At this stage of the project, potential markets for the recycled product will be identified as well as companies interested in establishing facilities for recycling and reprocessing. Based on the analysis of the collection and separation data, the possible financial incentives that will encourage recycling will be estimated.

EXPERIMENTAL RESULTS

Characteristics of Collected Fishing Gear

Samples of used and new fishing nets were supplied by NJMSC. The used samples that were collected during the first quarter of Phase I varied

widely in characteristics such as net mesh size, filament diameter, type of coating, degree of exposure, polymer type, and site of collection. All samples were classified and characterized as shown in Table 1. The majority of the collected nets were relatively clean, with minimum visible gross contamination from either organic or inorganic matter.

Identification of Plastics and Contaminants

The identification of plastics and their contaminants was undertaken in order to establish the proper reprocessing conditions and to minimize any possible adverse effects of the contaminants on processing and long-term polymer stability.

Identification methods used for plastics were infrared (IR) spectroscopy, melting temperature determination, and solubility measurements (the latter only for nylons). The preferred analytical method was IR spectroscopy, since the presence of coatings (e.g., "green dip," "tar") often interfered with the melting temperature and solubility determinations. The IR spectra were obtained on films cast on a AgCl plate from solutions of 90% formic acid (nylons) or hot xylene (polypropylene). Table 2 summarizes the identification results for all plastics. It is shown that nylon-6 and nylon-6,6 are the materials of choice for nets, whereas polypropylene is used for ropes and lines. It is interesting to note that the green-dipped twine in 88-2 contained both nylon-6 and nylon-6,6 fibers; by contrast, the attached repair portion was only nondipped nylon-6,6 fibers.

A thorough analysis and identification of contaminants was not completed at this stage of the project. Some surface elements that were identified by combined scanning electron microscopy and energy dispersive x-ray analysis included silicon and iron on the 88-2 fibers. These could correspond to sand and rust, respectively. Also, the "green dip" on 88-2 and 89-2 and the tar coating of 89-1 contain infusible cross-linked organic materials that turn black (degrade?) when heated near the melting temperature of nylons. The coatings level was not precisely determined. Preliminary thermogravimetric data suggest that the green dip content is at least 3-5%, whereas the tar content is at least 1% of the weight of the fiber.

Separation of Individual Plastics and Removal of Contaminants

Separation of the individual identified plastics present in the nets or the net and line combinations was not considered necessary at this stage of the project, since the proximity in chemical structures and melting points could allow these polymers to be eventually processed as blends (e.g., nylon-6 with nylon-6,6 in sample 88-2, nylon-6 with PP in sample 88-3). Floats, buoys, and lead-containing lines were the only articles that were removed. With respect to contaminant removal, the insolubility of the "green dip" and "tar" coatings in either water or organic solvents appears to preclude their economical separation from the fibers. Also, any water treatment of the fibers by slurring under agitation does not seem to be beneficial in improving their thermooxidative stability, as certain

Table 1.--Classification and description of used and new fishing gear received in 1988 and first quarter 1989.

Designation	Type or description	Color	Mesh (cm)/ diameter (mm)	Other information
Used				
88-1	Fyke(?) net with braided line attached; ca. 14 kg	Dark brown	Net: 7.5/1.2 Line: --/6.3	Brittle, dusty, badly degraded.
88-2	Trawl net, apparently dipped in green "enamel" and repaired with non-dipped net; ca. 50 kg	Dipped part: green or faded green Repaired part: off-white	8.3-10.5/3	Collected in New Jersey; variable in color, mesh and diameter; tar patches.
88-3	Monofilament gillnet with braided line attached; ca. 10 kg	Net: natural Line: green	Net: 8.3/0.37 Line: --/6.3	Collected in New Jersey.
89-1	Trawl net [30-ft flat otter] dipped in tar; ca. 25 kg	Black	4.5/1.2	Bought in Connecticut; used once per month for 1 year, stored in shed, repaired.
89-2	Trawl net [30-ft flat otter] dipped in green enamel; ca. 25 kg	Light green	3.8/1.2	Bought in Mississippi; used twice per month for 1 year, stored in shed.
New				
89-3-V	Twine net sample	Off-white	3.8/1.6	None.
89-4-V	Multifilament net sample	Light green	12.7/0.8	None.
89-5-V	Braided net sample	Light green	20/2.5	None.
89-6-V	Twine net sample	Blue	3.4/1.6	None.

Table 2.--Identification of polymers (by infrared spectroscopy, and solubility and melting temperature measurements) in used and new fishing gear.

Gear designation	Type of polymers
Used	
88-1	
Net	Not identified
Line	Not identified
88-2	
Green net	Nylon-6 and nylon-6,6
Faded green net	Nylon-6 and nylon-6,6
Repair net	Nylon-6,6
88-3	
Net	Nylon-6
Line	Polypropylene
89-1	
Net	Nylon-6
89-2	
Net	Nylon-6
New	
89-3-V	Nylon-6,6
89-4-V	Nylon-6
89-5-V	Polypropylene
89-6-V	Polypropylene

experiments showed. Thus, the only treatment prior to characterization was the removal of loose surface debris by air jetting.

Characterization of the Recovered Nylon Plastics

The characterization of the recovered plastics could provide valuable information on the extent of their degradation and their thermooxidative stability during reprocessing.

Table 3 summarizes the intrinsic viscosity and molecular weight data of all nylon polymers including three virgin materials in pellet form. Since the exact molecular weight of a fiber prior to exposure is not known, the extent of degradation can be only approximated. Thus, by comparing the intrinsic viscosities of the recovered polymers with those in the new nets and certain virgin pellets, one can conclude that with the possible exception of the "88-2 faded green" none of the other used nets showed extensive degradation of molecular weight. Thermal analysis (Table 4) by differential scanning calorimetry (DSC) and thermal stability investigations (Table 5) by thermogravimetry (TGA) tend to support the above conclusion. Differences in fusion and crystallization behavior between the fibers in the used or new net and the commercial pellets reflect largely the presence of

Table 3.--Intrinsic viscosity and molecular weight of nylon polymers: comparison of used and new fishing nets with commercial polymer pellets.

Sample designation	Intrinsic viscosity η^a (dl/g)	Molecular weight (M_v)	
		Nylon-6	Nylon-6,6
Used nets			
88-2			
Green	1.45	^b 43,840	or 21,480
Faded green	1.07	^b 30,380	or 14,140
Repair	1.31		18,710
88-3	1.52	46,520	
89-1	1.42	42,660	
89-2	1.31	38,840	
New nets			
89-3-V	1.44		21,460
89-4-V	1.61	50,090	
Pellets			
Zytel 101-F	1.16		15,760
Capron 8207-F	1.47	44,730	
Capron BHS-D	1.70	53,440	

^a η - Measured in 90% formic acid solution at 25°C.

^bFigures given refer to the individual polymers since both nylons are present in these nets in unknown ratios.

Table 4.--Thermal analysis (differential scanning calorimetry (DSC)) of nylon polymers: comparison of used and new fishing nets with commercial polymer pellets. Note: Heating, cooling, and reheating at 20°C/min; fusion data on second heating.

Polymer	Fusion temperature onset/maximum (°C)	Heat of fusion (cal/g)	Crystallization temperature onset/maximum (°C)	Heat of crystallization (cal/g)
Nylon-6				
Used nets				
88-2				
Green	211/222	10.7	187/181	-12.1
Faded green	212/223	10.9	187/182	-12.4
88-3	202/215	10.4	173/164	-11.4
89-1	208/221	8.3	181/174	-10.4
89-2	195/200	10.2	184/176	-12.0
New nets				
89-4-V	205/220	9.9	186/171	-12.1
Pellets				
Capron 8207-F	209/222	12.8	180/168	-13.4
Capron BHS-D	212/226	11.4	172/163	-10.7
Nylon-6,6				
Used nets				
88-2				
Repair	253/263	16.2	231/227	-12.7
New nets				
89-3-V	250/259	11.8	229/222	-12.9
Pellets				
Zytel 101-F	256/265	11.3	226/214	-14.0

coatings, rather than extensive polymer degradation. For example, coatings and contaminants or their residues appear to act as nucleating agents, as evidenced by the higher crystallization temperatures of the polymers in all the coated used and new nets. Also, extensive weight losses in TGA are probably associated with the early decomposition of the coatings and contaminants and not with that of the degraded polymer itself (with the possible exception of the "88-2 faded green" sample). Table 5 also shows that the thermal stability of the samples is practically independent of the type of test atmosphere (nitrogen versus oxygen).

Table 5.--Thermal stability (thermogravimetry (TGA)) of nylon polymers: comparison of used and new fishing nets with commercial polymer pellets.

Polymer	Standard TGA ^a				Isothermal TGA ^a		
	Onset decomposition temperature (°C)		Temperature at 10% weight retention (°C)		Isothermal temperature (°C)	Weight % decrease after 15 min	
	Nitrogen	Oxygen ^b	Nitrogen	Oxygen ^b			Nitrogen
Nylon-6							
Used nets							
88-2							
Green	429	NM	560	NM	225	4.03	4.84
Faded green	396	NM	673	NM	230	1.35	1.48
88-3	445	NM	513	NM	225	0.33	0.42
89-1	459	NM	519	NM	230	0.91	1.06
89-2	426	NM	525	NM	230	2.71	3.49
New nets							
89-4-V	459	NM	513	NM	230	0.26	0.38
Pellets							
Capron 8207-F	466	453	513	497	225	0.25	0.25
Capron BHS-D	471	NM	506	NM	230	0.62	0.67
Nylon-6,6							
Used nets							
88-2							
Repair	449	NM	530	NM	270	1.18	4.31
New nets							
89-3-V	451	NM	514	NM	270	1.23	1.75
Pellets							
Zytel 101-F	439	436	503	512	270	0.33	0.58

^aStandard TGA at 40°C/min; in isothermal TGA temperature reached within 2 min.

^bNM indicates not measured.

Experiments on Net Size Reduction and Extrusion Reprocessing

Attempts to chop net 88-2 into smaller pieces in a laboratory granulator were not successful even in the presence of dry ice. However, it was possible to feed continuously the same net into the hopper of a single-screw extruder and produce an extrudate that could be pelletized. The extrudate was black in color, presumably as a result of the decomposition of the fiber coating. Further experiments are under way in order to produce solid particles with dimensions suitable for feeding in polymer processing machinery such as extruders or injection molding machines.

CONCLUSIONS

The identification results on samples of unusable fishing gear collected during the first quarter of 1989 indicate that the major plastic involved are nylon-6, nylon-6,6, and to a lesser extent polypropylene. The initial characterization results of the exposed nylons are encouraging: it appears that none of the nylons are badly degraded in terms of reduced molecular weight or loss of thermooxidative stability. This could mean satisfactory processing characteristics during remelting of the net in the recycling extruder. The presence of fiber coatings that cannot be easily removed tends to complicate not only identification and characterization, but also the reprocessing behavior of the nylon nets. Our future efforts will include the complete analysis and characterization of typical commercial fiber coatings for a better understanding of their effects on processing. Other research areas addressed during this phase of the project include the full characterization of the collected exposed polypropylene gear and the investigation of net size reduction methods.

ACKNOWLEDGMENT

This research is supported by National Oceanic and Atmospheric Administration Grant No. NA 89 AA-H-SK005.

REFERENCES

- Lautenberg, F. [Senator]
1987. Introducing the Plastic Pollution Control Act of 1987.
Congressional Record-Senate, S269S, March 3, 1987.
- Shabecoff, P.
1989. The New York Times, March 21, 1989, p. C1.

CONTROL OF PLASTIC WASTES ABOARD NAVAL SHIPS AT SEA

Craig S. Alig
David Taylor Research Center Code 2834
Department of the Navy
Annapolis, Maryland 21402, U.S.A.

Larry Koss
Office of Chief of Naval Operations
Department of the Navy
Washington, D.C. 20350, U.S.A.

Tom Scarano
Naval Sea Systems Command
Department of the Navy
Washington, D.C. 20362, U.S.A.

and

Fred Chitty
Supply Operations and Policy
Commander in Chief, Atlantic Fleet
Department of the Navy
Norfolk, Virginia 23511, U.S.A.

ABSTRACT

The U.S. Navy is taking a proactive approach to comply with the prohibition on the at-sea discharge of plastics mandated by the Marine Plastic Pollution Research and Control Act of 1987. For U.S. naval ships, space and weight constraints, high crew densities, and mission requirements create unique solid waste management problems.

In pursuit of a zero discharge goal without significant adverse impact, processes, activities, operations, and systems to reduce plastics discharges are being identified, evaluated, and documented. Operational, supply, and technology-oriented solutions are now being demonstrated. Seven ships have been nominated by Commander in Chief U.S. Atlantic and Commander in Chief U.S. Pacific Fleets to participate in a plastics waste reduction demonstration project. Each ship was asked to develop its own instructions and procedures to eliminate the discharge of plastic wastes and to implement those instructions in a manner consistent with the operational requirements and mission of the ship.

Thus far, a submarine tender, a frigate, a destroyer, and two submarines have participated. Scientists and engineers from the David Taylor Research Center have collected waste generation rate and characterization data and have monitored and documented lessons learned. Naval Supply Systems Command has provided support for the demonstrations by recommending substitutes for plastic products, and new waste processing systems being developed by the Naval Sea Systems Command have been evaluated. Procedures for both the source separation of plastic and non-plastic wastes and the separation of food-contaminated plastic waste from non-food-contaminated plastic waste have been very successful aboard each of the demonstration ships. Plastic wastes have been stored and returned to port. A new Navy-developed vertical trash compactor has been successfully used to process plastic and nonplastic waste separately, and a pulper has been successfully used to process large volumes of degradable, nonplastic waste for ocean discharge.

SOLID WASTE: THE NAVY SHIPBOARD PROBLEM

For as long as ships have sailed the oceans, waste has been thrown overboard at sea. This practice continued unchallenged for centuries. The ocean's vast size and powerful assimilative capacity easily absorbed ship-generated waste with no apparent adverse impact. For many years, the waste consisted of simple degradable materials; later, the waste included metal, which sank. However, the relatively recent development of synthetic materials such as plastic, which float and persist in the marine environment when thrown into the sea, has changed the perception that there is no harm in discharging ship-generated trash at sea.

The visible evidence of ships' discharges now points an accusing finger at the maritime industry and the military for polluting the oceans, even though a great proportion of trash on the beach originates from sources ashore. Initially, the concern over marine debris, whether floating at sea or washed onto the beach, was because it offended our sense of aesthetics. However, as plastic became more pervasive in its application, other problems developed. Plastic line began to foul ships' propellers, and drifting plastic sheeting clogged ships' seawater intakes.

Floating marine debris also presented a unique problem for warships--it compromised security. Buoyant bags of trash establish a trail of floating waste which can betray a ship's location. Floating waste can be recovered more easily, enabling adversaries to gain information from the items contained therein.

Regulations of the U.S. Navy prohibit the discharge of any trash within 25 nmi of any shore, and require that all trash be weighted before disposal at sea to ensure that it sinks. However, it is difficult for shipboard personnel to consistently package or process waste for negative buoyancy without the use of special equipment.

In 1970, when the nation finally confronted the environmental crisis and sweeping clean air and water legislation was established, the Commanders in Chief of the Atlantic and Pacific Fleets recognized the need to develop strategies and technologies to deal with the solid waste problem. A comprehensive Naval Shipboard Refuse Study (NSRS) revealed that each person afloat generated about 1.4 kg (3.05 lb) of solid waste per day (Table 1). (Note that less than five thousandths of a kilogram per person per day was plastic waste.)

However, warships at sea had no holding capacity to store waste on board. Waste storage also created health and sanitation hazards and fire control problems. Overboard disposal by Navy ships continued to be the practice.

Recognizing that an alternative to overboard disposal was necessary, the Naval Sea Systems Command set out to find suitable solid waste processing systems. Initially, their goal was to process the degradable waste so it would not float when discharged and to compact the intrinsically heavy, inert material so it would sink to the bottom. This would eliminate the problems caused by floating debris. At the time, plastic waste was not viewed as a serious problem because of the small amounts generated aboard ship.

What seemed a simple concept in the early 1970's proved to be extremely difficult to execute. Commercially available equipment could not meet the rigorous requirements imposed by the Navy (Table 2). Dozens of candidates were evaluated at the Navy's David Taylor Research Center, but none could satisfy the demands of a warship. The first real equipment successes began when industry teamed with the increasing experience of the Navy, and a family of Navy-model food waste disposers was developed.

During the late 1970's, the Navy's engineering communities at the Naval Sea Systems Command and David Taylor Research Center initiated a long-term shipboard solid waste control research and development program.

RISING TIDE OF MARINE PLASTICS AND U.S. REACTION

Public concern over marine debris magnified enormously in the 1980's because of the terrible impact that synthetic material, particularly plastic, was having on marine life. The amount of floating marine debris continued to increase, creating more beach litter and overwhelming waterfront communities struggling to maintain a high-quality beach environment.

The increase in floating marine debris corresponded to the increased use of plastic products in the home, industry, and marketplace. Plastic and synthetic products found their way aboard maritime ships--and then overboard. Comparing studies conducted in 1971 and 1987, Table 1 shows an approximate twentyfold increase in Navy shipboard plastic waste. No prohibitions existed in the early 1980's against the discharge of shipboard-generated waste once a ship was beyond 3 nmi of the shoreline.

Table 1.--Generation of naval shipboard solid waste
(kg (lb) per person per day).

Item	Naval shipboard refuse study - 1971	Naval shipboard refuse study - 1987
Plastic	0.004 (0.01)	0.095 (0.21)
Food waste	0.603 (1.33)	0.580 (1.28)
Glass	0.008 (0.02)	0.059 (0.13)
Metal	0.299 (0.66)	0.186 (0.41)
Rubber	0.004 (0.01)	0.004 (0.01)
Paper, other	0.463 (1.02)	0.503 (1.11)
Total	1.38 (3.05)	1.43 (3.15)

Table 2.--Criteria for naval shipboard
waste processing equipment.

Parameter	Requirement
Reliability	High
Manpower	Low
Safety	Extremely safe
Space needed and weight	Low
Simplicity	Extremely simple
Ability to withstand shock and vibration	Rugged beyond belief

During the past few years, the deadly impact that synthetic plastic materials have on marine sea life has been graphically documented and widely publicized by environmental organizations. Countless photographs document the deadly consequences of ingestion of plastic by birds, turtles, and marine mammals, and their entanglement in synthetic fishing line and nets and in plastic sheeting.

Clearly, the Navy is not a contributor to the deadliest form of marine plastic--the synthetic rope of fishing nets. The low number of Navy ships at sea compared to the commercial fishing and merchant fleets makes the Navy a minor contributor (ca. 2.5%) to the total plastics waste problem. However, the high population density aboard naval vessels and the plastic waste discharged daily from each ship at sea adds up over time. As a role model, the Navy must demonstrate leadership with an intensive effort to eliminate the discharge of floating marine debris.

The International Convention for the Prevention of Pollution from Ships (commonly known as MARPOL) was the first comprehensive agreement to

control marine pollution worldwide. The MARPOL was drafted in 1973 (MARPOL 73) and updated in 1978 (MARPOL 73/78). The MARPOL 73/78 included Annex V as an option which would prohibit ships from discharging plastic wastes at sea. The convention was ratified in 1980 with its protocol of Annexes 1 and 2 to eliminate the discharge of oil at sea; yet, it was 1987 before the international outcry against plastic waste forced leaders around the world to take action on Annex V.

In the fall of 1987, the U.S. Senate gave its unanimous consent to the ratification of Annex V to MARPOL 73/78. The 29 signatory nations represented over 50% of the world's merchant fleet tonnage. Annex V dictates that no vessel from a signatory nation may dispose of any plastics into the sea or dump floating solid waste within 25 nmi of any shoreline. Annex V also prohibits the discharge of any solid waste (except ground food waste) into special areas such as the Baltic or Mediterranean Seas.

Congress passed enabling legislation immediately after ratification of Annex V--the Marine Plastic Pollution Research and Control Act of 1987 (Public Law 100-220), which took effect 1 January 1989 for all maritime vessels, and takes effect 1 January 1994 for Navy ships. Congress recognized that full compliance in 5 years would be extremely difficult for the Navy because of the time required to complete development of and to procure and install the appropriate equipment on about 500 ships. Therefore, Congress required that the Navy report in 3 years on progress made toward full compliance, with the expectation that the compliance deadline could be extended if warranted.

In October 1987, the Assistant Secretary of the Navy for Shipbuilding and Logistics created an ad hoc advisory committee on plastics. For 7 months, the committee met and traveled to Navy research centers and supply depots, and visited several naval ships of various types. In June 1988 it delivered its final report to the Assistant Secretary. The report contained 42 specific recommendations for the Navy to meet its solid waste management goals by 1994. The recommendations were divided into four categories: technology, operations, supply, and education.

CRITICAL NAVY ISSUES

In formulating a plan to achieve full compliance with P.L. 100-220, a number of critical issues had to be addressed.

- How do we separate the plastic waste, which comprises 7% by weight of all the solid waste generated, without creating labor-intensive efforts, which could negatively affect crew morale?
- Where do we install solid waste processing equipment aboard a military vessel so that it is centrally located, efficiently arranged, and minimizes the crew's labor burden? (While the food waste (0.58 kg (1.28 lb) per person per day) can be discharged directly overboard at sea when processed through galley or scullery garbage grinders, the remaining solid

waste (0.85 kg (1.87 lb) per person per day) must be transported and processed for disposal or storage. For example, 900 kg (close to 1 ton) of waste per day is generated at dozens of different rates and locations aboard a 1,000-man ship. It must be carried by hand to processing centers and then carried elsewhere for disposal or long-term storage.)

- Where do we find space to store solid waste? (There is very little designated trash storage space aboard a warship, and there are no unused spaces that can readily be made available. However, since plastic waste cannot be discharged overboard, space must be found without creating fire, health, or sanitation hazards, and without reducing the quality of life aboard ship. Full regulatory compliance demands that equipment be developed specifically to process plastic waste for volume reduction and sanitation.)
- How do we reduce the quantity of plastic waste generated aboard ship? (Alternatives are available for products such as polyethylene trash bags and polystyrene coffee cups. However, plastic is widely used for packaging, and often is the most cost-effective material for that application, especially food products. It has taken years to develop and implement plastics that are efficient and economical (e.g., shrink wrap). Material and product substitutions that perform as well and are as economical may require a long-term search.) Realistically, plastic waste may best be managed by accepting its continued use and developing a plastic waste processor which, together with recycling, will allow us to control the plastic waste storage problem aboard ship.

NAVY PLAN FOR FULL COMPLIANCE

The Navy's approach to full compliance with MARPOL Annex V and P.L. 100-220 contains four parts. They are technology initiatives, operational changes, substitutes for plastic products, and education. We may think of "people" as the fifth part of our approach. Unfortunately, the human side is sometimes the most difficult to specify, predict, and control.

Technology Initiatives

In the context of naval ships, technology refers to the equipment that will be installed to provide each ship with part of the capability required for onboard management of solid and plastic wastes. It is important to understand the rationale behind solid waste technology initiatives before the details can be presented.

First, our shipboard technology initiatives reflect the need to comply with all of the requirements of Annex V, which includes managing the total solid waste stream and prohibiting the discharge of plastics. The complexities associated with shipboard equipment installation force us to consider

the total solution to the problem rather than small areas at a time. This approach seems necessary also if we are to achieve our goal and implement timely solutions at a reasonable cost. Therefore, the Navy's technology program places the same emphasis on solid waste management as on plastics discharge prohibition.

Second, a "generic" solution for a "typical" ship would be reassuring; however, many unique solutions are needed to satisfy the multiplicity of ship designs and operating scenarios. The Navy may install solid waste management equipment on approximately 500 ships that fall into about 60 different ship classes! Additionally, surface ships may carry as few as 200 people or as many as 6,000, and submarines have requirements that are entirely unique.

Third, naval ships differ significantly from commercial vessels and will find it more difficult to comply with Annex V because their population density usually is much higher. A 305-m (1,000-ft) naval ship may have a crew of 6,000, while the same size oil tanker may have a crew of 40. Obviously, the contributing population determines the quantity of plastic and other solid waste produced. While the maritime fleet has similar problems with large population densities on cruise ships, they differ significantly from Navy ships in their mission, purpose, and time at sea. Furthermore, Navy ships have no occupational specialty to manage solid waste.

The Navy is developing three shipboard systems that will be major factors in our compliance with Annex V: a vertical trash compactor, a solid waste pulper, and a plastic waste processor. Each system will be of a single size with a fixed capacity, making it easier to train operators and obtain parts necessary for repair and maintenance. Larger ships may require multiple units.

Presently, onboard incineration of solid and plastic wastes does not play a major role in the Navy's plan to comply with Annex V, because this type of burning emits potentially toxic and corrosive waste products in its exhaust gases and ash. Additionally, our experience with conventional marine solid waste incinerators has shown that suitable, high-capacity incinerators that meet the requirements of Table 2 are unavailable. However, we are investigating advanced thermal destruction technologies for limited use on ships operating under unique conditions; in some cases, this may be the best alternative to achieve ultimate, at-sea volume reduction of solid waste.

Shipboard Vertical Trash Compactor

The Navy's research and development program of trash compactors began in 1979. Our objective was to develop a machine that was reliable, easy to operate, sanitary, safe, and would allow ships to meet environmental regulations for the discharge of solid waste. We found it difficult to achieve negative buoyancy in trash that was compacted into a degradable container. Finally, we were able to ensure that the container would sink by spraying seawater into the compaction chamber, then using high compaction pressure

to force the water into the pores of the trash to displace the air. A preproduction prototype, such as the one shown in Figure 1, is undergoing technical evaluation aboard a Navy destroyer and has met with outstanding success in the past year.

The Navy shipboard vertical trash compactor was designed to meet Navy standards for maintainability, reliability, safety, shock, vibration, structure-borne and airborne noise, electromagnetic compatibility, and habitability. It can process solid waste composed of glass bottles, metal cans, paper products, and other nonindustrial and nonhazardous waste into 20.4-kg (45-lb) trash slugs. The slugs are contained in cloth bags that can be hand carried. The compactor provides extended-time trash storage, trash slugs that sink without added metal weights, continuous safety checks, and fully automated operation controlled by a programmable logic controller. The most critical parts of the compactor are made from materials that are corrosion resistant; the compactor can be disassembled for movement and installation aboard ship. Its vertical configuration results in a footprint only 0.6 x 1.8 m (2 x 6 ft). Volume reductions greater than 5:1 were achieved when processing plastic waste for storage aboard ship. The first of these units should be delivered to the fleet within 2 years.

Shipboard Solid Waste Pulper

Pulping solid waste is not new to the Navy. One class of ship has been successful in using commercial pulpers to process mixed solid waste. Galley and scullery food waste disposers are actually small pulpers. Used as a shipboard solid waste processing method, pulping mixes waste with water, reduces the size of the solids, and creates a wet pulp or slurry, which can be pumped directly overboard. Pulped waste is more readily biodegradable than unpulped waste, tends to be negatively or neutrally buoyant, and disperses rapidly when discharged.

Typically, pulpers operate as follows. Waste enters a large tank through a feed chute and is mixed with water. The water softens and saturates the waste material so that it can be reduced in size more easily. The mixture forms a vortex caused by the rotation of a cutting mechanism located in the bottom of the tank. The pumping action of the cutting mechanism draws trash down the vortex to the bottom of the tank where it is cut and pressed through a perforated sizing ring before being discharged as a slurry. The Navy shipboard pulper is being designed to process paper, food waste, and fiberboard. A conceptual design of the pulper appears as Figure 2. Metal, glass, plastic, and cloth are considered nonpulpable, although the design of the pulper makes it highly resistant to damage from accidental insertion of these items. It will process about 75% of the total solid waste generated aboard a ship at a rate of 262 kg (600 lb) of mixed solid waste per hour. Our experience demonstrates that these pulpers are extremely reliable and simple to operate. The first units could be delivered to the fleet within 4 years.




Figure 1.--Shipboard vertical trash compactor.




Figure 2.--Shipboard solid waste pulper.

Plastic Waste Processor

Plastic waste comprises about 7% by weight of the total solid waste generated on board a Navy ship. Of that quantity, 50% is contaminated with food waste. Storage of food-contaminated waste aboard ship requires significant volume reductions along with sterilization or similar treatment to control noxious odors.

The Navy's shipboard plastic waste processor is in the early stage of development. While the exact design and configuration are uncertain, Figure 3 depicts our developmental objective. The densified, sanitized block of waste plastic will be suitable for long-term shipboard storage until it can be off-loaded ashore. We anticipate a 30:1 volume reduction and an end product that is recyclable. Our goal is to have the first units delivered to the fleet within 6 years.

Shipboard Equipment Configurations

To plan the installation of our solid waste management equipment, naval ships are grouped into four categories--small, medium, and large surface ships, and submarines. The conceptual plan calls for compactors only to be installed on small surface ships; compactors and plastic waste processors on the mid-sized ships; and compactors, plastic waste processors, and pulpers on larger ships. Some larger ships may require more than one of each system to ensure maximum efficiency. This plan assumes that personnel on board each ship will separate plastic manually at its source and will use their food waste disposers.

On smaller surface ships, all solid waste (except food) will be processed through the compactor. Negatively buoyant, nonplastic slugs will be stored on board for shore disposal or for overboard discharge where permissible. The compactor can process an all-plastic slug which provides at least a 5:1 volume reduction.

Medium-sized surface ships can accommodate a compactor and a plastic waste processor. The compactor will process all of the solid waste except separated plastics and food waste. Nonplastic slugs will be stored on board for disposal ashore or for overboard discharge. The plastic waste processor will process all plastics including food-contaminated waste, and the densified (30:1) and sanitized plastic will be stored on board for disposal ashore.

All three systems will be installed on larger surface ships in single or multiple units, depending on the need and the space available. Each system will be targeted to a specific segment of the solid waste stream to ensure maximum utilization and efficiency. Separated plastic waste will be processed for storage and shore disposal. The remainder of the solid waste stream will be separated at its source into pulpables (e.g., paper, fiberboard, and light wood) and nonpulpables (e.g., glass and metal cans). The compactor will process nonpulpables into negatively buoyant slugs for overboard discharge when the ship is 25 nmi from the shoreline. Pulpables will be processed by the pulper when the ship is at least 12 nmi from shore




Figure 3.--Shipboard plastic waste processor.

and pumped overboard. Larger ships are most able to maximize the use of waste management technology and minimize the amount of waste retained on board for shore disposal.

Because of limited space and other constraints, the current Navy plan does not include the development and installation of new solid and plastic waste equipment for submarines. However, submarines will comply with Annex V to the maximum extent possible by using material substitutions to minimize the generation of plastic waste; by source separation and onboard storage of plastic waste that is not food-contaminated; and by continuing the current practice of compaction, weighting for negative buoyancy, and overboard discharge of the remaining solid waste.

Operational Constraints

The Navy's response to the need for operational changes met with four major constraints.

1. *Manpower.*--An 81-h work week is standard for sailors operating at sea, and expectations of what a crew member is to accomplish during a shift leaves no time for special handling of plastic waste. The establishment of additional jobs to fulfill this need is not probable because no funding is available for such a position, and there is no berthing space on the ships for additional crew members.
2. *Space.*--Space is at a premium aboard even the largest naval ships. The typical seagoing merchant ship is considerably larger than the average naval surface combatant, yet its crew numbers 30 compared to 300 sailors aboard a Navy ship. While aircraft carriers are the largest ships in the fleet, their crew exceeds 6,000, thus making their population density similar to that of our smaller surface ships. The cramped quarters aboard our ships leave little space for the installation of new equipment.
3. *Quality of life.*--It is critical that the Navy maintain the quality of life at sea as a top priority to enable us to continue to attract high quality personnel. Routine operations involve 7 to 25 continuous days at sea, with 45 days not unusual. Long deployments may require 80 to 150 continuous days at sea. Thus, it is imperative that each sailor have a clean, healthy, and safe place in which to live. Controlling plastic waste cannot be allowed to negatively affect quality of life aboard ship.
4. *Financial constraints.*--Congress has not allocated additional funds for the increased operating costs that will be incurred during implementation of the procedures to prevent plastic pollution. In fact, budgets are being cut and daily operating funds are scarce. In forecasting the operational changes ahead, funding concerns force us to note that the

cost of paper coffee cups is double that of polystyrene cups, and paper trash can liners are quadruple the price of those made of plastic.

Hence, the Navy must reduce plastic pollution in the marine environment while operating within these realities. We must change the way we process solid waste, yet minimize the increase in manpower and financial resources required and maintain the high quality of life aboard ship.

Substitutes for Plastic Products

Many items contribute to the problems of plastic waste aboard ship, most of which cannot be controlled. A destroyer with a crew of approximately 300 was used as one of our plastic waste reduction demonstration study ships. During one 16-day period, the David Taylor Research Center study team inventoried 6,179 individual pieces of plastic waste. "Miscellaneous" sources, those not attributable to berthing, work center, or food service areas, accounted for 25% of the waste. The two most numerous items were six-pack rings and trash bags; however, frozen meat packaging, food wrap, and food containers accounted for 14 of the remaining top 23 categories of plastic waste collected. Acceptable, nonplastic substitutes for these items will not be available for many years, if ever.

However, nonplastic substitutes are available for some of the plastic items. Ships have found nonplastic substitutes for coffee cups and stirrers, tableware, and trash can liners. While plastic bags are still used to collect and hold plastic waste for disposal ashore, paper bags are now specified for collection and at-sea disposal of nonplastic solid waste.

New procedures will eliminate some disposable plastic products. For example, food waste can be disposed in food pulpers or garbage grinders, which eliminates serious storage problems caused by the collection of food waste in plastic bags. This practice also requires less manpower; the daily garbage does not have to be carried from the galley to the fantail and dumped overboard for disposal.

Education

All shipboard personnel must be educated on the hazard that waste plastic poses to marine life and on the procedures necessary for effective shipboard solid waste management. Navy personnel have become increasingly aware of the potential adverse impact that shipboard operations have on the environment. Shipboard solid waste separation management must become a task that each crew member accepts as part of the ship's routine operation.

IMMEDIATE REDUCTIONS IN PLASTICS DISPOSAL AT SEA: SHIPBOARD PLASTIC WASTE REDUCTION DEMONSTRATIONS

The 5-year implementation period does not afford us the time to conduct paper studies. While the research and development required for technical solutions have been accelerated and are moving forward as quickly

as possible, we must implement changes in waste management and disposal practices now, not because Congress expects it of us, but because it is environmentally expedient.

Changing shipboard operational procedures was the single viable alternative to effect the expeditious reduction of plastic wastes discharged at sea. Our focus was on the segregation of plastic wastes from other solid wastes, and its short-term storage aboard ship. Many of the recommendations in the report of the ad hoc advisory committee on plastics were directed toward achieving this objective. Shipboard plastic waste reduction demonstration studies were conducted aboard five surface ships and two submarines to test and evaluate each recommendation. Studies on two additional surface ships are planned. Thus far, the demonstration studies have illustrated several major points:

- Navy shipboard plastic waste represents about 7% by weight of the total solid waste stream; 50% of that waste originates in food service areas.
- Plastic waste generation is nearly constant across ship classes, ranging between 45 and 90 g (0.1 and 0.2 lb) per person per day.
- Separation of the plastic from other waste at its source was most effective and required the least effort. Trash cans or bags labeled "Plastic Waste Only" were essential. In most living and work spaces they would hold a 1- to 2-week accumulation of plastic waste.
- Onboard storage of plastic contaminated with food waste was limited to approximately 3 days before the noxious odor began to affect the quality of life and it posed a threat to health and sanitation conditions.
- Uncontaminated plastic waste can be stored on board up to 3 weeks; the originating work center seems to be the most appropriate choice for storage on most ships. The plastic waste was collected and placed in pier-side dumpsters when a ship returned to port.

POLICY GUIDANCE

The commanders of the Atlantic and Pacific fleets have issued policy guidance to the commanding officers of the ships. Each ship must separate and store plastic that is not contaminated with food waste for at least the first 20 days of any underway period, and longer if space allows. Plastic that is contaminated with food waste must be held on board for the last 3 days of any underway period. Plastic waste stored on board will be off-loaded in port. If retention compromises the health, safety, or combat readiness of the ship and those aboard, properly packaged and negatively buoyant plastic waste may be discharged overboard when the ship is beyond 50 nmi from any shoreline. Such disposal must be approved by the

commanding officer and a log entry made indicating the time and position of overboard discharge. Noncompliance with the dumping policy must be reported in writing. This policy became effective 15 January 1989 for ships in the Atlantic fleet, and 1 March 1989 for Pacific fleet ships. The effective dates were chosen to coincide with the issuance of education packages for each ship. Each package includes the "Ship's Guide for the Management of Plastic Waste at Sea."

Fleet implementation of the new procedures will produce an immediate reduction in the total plastic waste discharged overboard by an astounding 70%! The aggressive action exhibited by our operational forces affords our engineering community time to develop and install solid waste management equipment aboard ships that will promote still further reductions. Also, the Navy's supply community will use this opportunity to research alternative products to reduce the amount of plastic used on board ship. The U.S. Navy stands firmly committed to achieving full compliance with all environmental regulations.

SHIPBOARD WASTE DISPOSAL: TAKING OUT
THE TRASH UNDER THE NEW RULES

Lissa A. Martinez
7107 Cedar Avenue
Takoma Park, Maryland 20912, U.S.A.

ABSTRACT

In 1988, the Society of Naval Architects and Marine Engineers Panel M-17, Disposal of Shipboard Wastes, convened two workshops to encourage open discussion of the diverse options available to comply with MARPOL 73/78 Annex V (Garbage). This paper reviews the available engineering, operational, and managerial changes useful in implementing Annex V and outlines the "good marine practice" suggested by the discussions in Panel M-17. Examples of approaches actually adopted by different commercial operators will be offered.

INTRODUCTION

A new marine pollution prevention regime, MARPOL 73/78 Annex V (Garbage), came into force internationally on 31 December 1988 (International Maritime Organization (IMO) 1978; U.S. Congress 1987). Annex V requires ship operators to change the way shipboard garbage is handled and immediately bans discarding plastic materials anywhere in the sea (Whitehead 1988). Disposing of shipboard garbage properly matters more to the company, the sailor, and the national authorities, because Annex V changes the long-accepted maritime practice of tossing garbage into the sea. However, taking out the trash under the new rules means more than stopping sailors from chucking everything over the side. Annex V is no antilitter campaign, but is part of a fundamental shift in the way ship crews and managers operate (Horsman 1982; Vauk and Schrey 1987). Making the transition to a commercial fleet that is able to obey the MARPOL Annex V will take a combination of changing how people have usually done things and providing them with the tools they need to do things differently.

Implementing Annex V has become the job of ship designers, ship operators, and maritime environmental specialists who have the expertise in shipboard systems design and operation. A ship is a small place, of a fixed size, occupied by people, cargo, and a lot of machinery. Rarely is a ship built with spare space or operated with extra people. Under these common constraints, a change in one shipboard activity often has a consequence in another activity. In this paper, the author examines the implementation of the MARPOL Annex V in the merchant fleet from the

perspective of the marine technical professionals, to convey a sense of the "good marine practice" they need to select, install, and operate solutions to comply with Annex V.

**Background: The Role of the Society of
Naval Architects and Marine Engineers**

The Society of Naval Architects and Marine Engineers (SNAME) is a U.S. organization of ship designers, ship builders, and ship operators. One standing technical panel of the society is Panel M-17 (Disposal of Shipboard Wastes), which is made up of professionals who work in engineering and management to enable ships to meet the legal requirements for environmental protection (SNAME 1982). In 1988, the SNAME Panel on Disposal of Shipboard Wastes convened two shirt-sleeves workshops on "The Shipboard Engineering and Environmental Aspects of Implementing MARPOL 73/78 Annex V (Garbage)" to encourage open discussion of the diverse options available to comply with Annex V. The first panel workshop on 18 July 1988 was hosted by the Office of the Chief Scientist, National Oceanic and Atmospheric Administration (NOAA). By popular demand, a second meeting was held on 12 October 1988, hosted by the Waste Combustion Equipment Council of the National Solid Waste Management Association (NSWMA).

The Panel M-17 workshops have been lively and useful exchanges of information and opinion. More than 80 people attended, including waste disposal firms representatives, port authority representatives, fleet operators, marine engineers and designers, environmental lawyers and regulators, and supply officers, all of whom have work to do to implement Annex V. The meeting participants provided useful information about shipboard and shoreside waste disposal equipment, local port implementation needs, disposal costs, U.S. Coast Guard regulatory proposals, and ways to design a compliance alternative that make sense for individual ships (Martinez 1989).

Enlightened self-interest helped motivate such a free exchange of information. Each fleet had to comply with the new international convention by 31 December 1988, with little lead time to order equipment or change vessel operations. Domestically, the Coast Guard had only a year after Congress passed the new law to draft and issue new rules that apply to all boats, ships, and oil rigs operating in the waters of the United States. No regulations were in place and many in the merchant marine were uncertain what would satisfy the authorities or how to do it. It seemed a good idea to sit down and talk about what we faced.

Background: MARPOL Implementation Philosophies

The MARPOL 73/78 Annex V (Garbage) is the third pollution prevention regime to be imposed on the world merchant fleet. Official shorthand for the International Convention for the Prevention of Pollution from Ships 1973, as amended by the Protocol of 1978, the MARPOL 73/78 contains five annexes that address particular types of ship-source marine pollution. The first annex implemented addressed oil pollution and the second addressed chemical cargo wastes. However, Annex V is philosophically different in

its approach, and maritime professionals need to understand that philosophical difference.

The MARPOL 73/78 Annex I (Oil) and Annex II (Bulk Chemicals) prescribed the way to comply, including which equipment to use and what procedures to adopt. In addition, it was clear where the responsibility for day-to-day compliance rested. The wastes involved come from cargo tanks, which are the responsibility of the deck officers, or from machinery spaces, which are the responsibility of the engineers. Enforcement and compliance were rigidly defined, and neither the Coast Guard nor ship operators had much leeway from the very start. Finally, not all vessels had to comply by the same date, because internationally agreed upon schedules gave older vessels more time to come into compliance.

The MARPOL 73/78 Annex V (Garbage) is so different that it has taken a while to get used to it. Annex V mandates that overboard disposal must change and that plastic disposal must cease, but implementing Annex V relies on none of the methods which were so central to the previous pollution conventions. Annex V instead:

- applies to ship-generated garbage, regardless of the source, and clearly includes hotel and galley services, which are the responsibility of the stewards;
- does not require specific new equipment in either ships or ports;
- does not tell ship operators or port authorities how to comply; and
- applies to all vessels in all waters immediately, with no delayed implementation schedules for existing vessels.

These are both the strengths and the weaknesses of the entry into force of Annex V. On the positive side, operators are not shackled to a technologically rigid solution and are free to develop their own best way to comply with the new requirements. Also on the positive side, implementing Annex V is not just the engineer's or deck officer's responsibility, but is a shared responsibility across the ship's crew and the shoreside supporting organization. This allows an operator to experiment and make incremental changes that can yield a compliance solution tailored to the way the ship operates (U.S. Department of Commerce 1988). On the negative side, the entire fleet had to comply by 31 December 1988, which meant that few people had the experience in solving the problem aboard ships and few were familiar enough with the annex to protect themselves from the uninformed speculation that was circulating, and everyone wanted equipment or advice at the same time.

TACKLING THE PROBLEM

Table 1, the "MARPOL 73/78 Annex V Summary of Garbage Discharge Restrictions," (U.S. Department of Transportation 1988b, 1988c) sets out

Table 1.--The MARPOL 73/78 Annex V summary of garbage disposal restrictions.

Garbage type	All vessels except offshore platforms and associated vessels		Offshore platforms and associated vessels ^b
	Outside special areas	In special areas ^a	
Plastics--includes synthetic ropes and fishing nets and plastic bags	Disposal prohibited	Disposal prohibited	Disposal prohibited.
Floating dunnage, lining and packing materials	Disposal prohibited 25 nmi from nearest land	Disposal prohibited	Disposal prohibited.
Paper, rags, glass, metal, bottles, crockery, and similar refuse ^c	Disposal prohibited <25 nmi from nearest land	Disposal prohibited	Disposal prohibited.
Food waste not comminuted or ground	Disposal prohibited <12 nmi from nearest land	Disposal prohibited	Disposal prohibited.
Food waste comminuted or ground ^c	Disposal prohibited <3 nmi from nearest land	Disposal prohibited <12 nmi from nearest land	Disposal prohibited <12 nmi from nearest land.
Mixed refuse types	(d)	(d)	(d)

^aSpecial areas are the Mediterranean, Baltic, Red and Black Seas, and Persian Gulf areas.
^bOffshore platforms and associated vessels includes all fixed or floating platforms engaged in exploration or exploitation and associated offshore processing of seabed mineral resources, and all vessels alongside or within 500 m of such platforms.
^cComminuted or ground garbage must be able to pass through a screen with a mesh size no larger than 25 mm.
^dWhen garbage is mixed with other harmful substances having different disposal or discharge requirements, the more stringent disposal requirements shall apply.

the different classes of wastes that are now regulated and where their disposal is now restricted. As stated previously, the saving grace of Annex V is that no technology is mandated. This injects a little breathing room into the transition and removes any cause for alarm if you cannot retrofit equipment by the entry-into-force date. The open "philosophy" for compliance gives a company more freedom to design its own "right" way, but also compels a designer to consider more alternatives. It means that the role of the marine engineer in implementing Annex V is different. A company cannot delegate the task to an engineer, as was done with Annexes I and II, and expect everything to fall into place neatly. It's not that the engineer cannot deliver a technical solution, it is rather that the solution is not in the hardware.

Diverse ways are available to comply with Annex V, exercising both managerial and technical prerogative. "Management" changes can greatly reduce the amount of waste generated and reduce the size of the "engineering" solution needed. Some options affect vessel operations that are not the usual jurisdiction of the marine engineers, such as provisioning the accommodations and securing the cargo. A good starting point for evaluating the situation is to read the Marine Environment Protection Committee (MEPC) Guidelines for the Implementation of Annex V (Garbage) (IMO 1988b). This document has been written to introduce the merchant mariner and the maritime designers to the problem-solving method that is best used in handling shipboard wastes. It works well, and the credit goes to the drafters of the document for producing a practical and usable text. The guidelines are well regarded by the designers and operators who have had the opportunity to use them. Better yet, the guidelines are amenable to modifications as operators and authorities gain experience in implementing Annex V.

In the United States, marine engineers and designers have had only partial information available for fulfilling the marine engineering tasks related to Annex V. Each compliance decision has consequences for the way the ship operates, and some apparently simple solutions affect shipboard operations and costs more severely than do some apparently more complex solutions.

Shipboard sanitation and safety must be safeguarded when selecting an installation or retrofit (Signorino 1988). Panel M-17 convened to become familiar with the solid waste management practices in port facilities and with the state of the art of types of equipment used to process or destroy the wastes now addressed in the Annex V (Garbage) requirements. Very few of us, even in the Panel M-17 community of specialists, had previously studied the ship's garbage, Annex V, garbage equipment available for shipboard or shoreside use, or how to safely retrofit garbage equipment on existing ships. Some options use expensive and unfamiliar equipment, such as package incinerators, large capacity compactors, or pulpers. A quick review of the decision and technical options follows.

Estimating the Waste Stream

Before the problem can be solved, some estimate has to be devised of how much garbage will be handled. If possible, detailed inventories can be

done, but many operators are unable to spend the time and money on gathering such information before compliance is required. The discussion in the M-17 meeting reviewed the various ways that people have selected to estimate the ship-generated garbage needing treatment. The following ways have been used with some success:

- The Coast Guard regulatory docket for the implementation of the MARPOL Annex V includes a study that creates a unit called the GBE or "40-gallon garbage bag equivalent" (U.S. Department of Transportation 1988a).
- The U.S. Navy 1988 inventory of shipboard waste yielded numbers of 1.4 kg(3 lb)/person/day of garbage which includes 0.5 kg (1 lb) of food waste and 0.9 kg (0.2 lb) of plastic. This is a twentyfold increase in the average amount of plastic since the 1971 inventory.
- The waste disposal industry categorizes "incinerable waste" according to its heat release by the following classification (Norske Hydro 1988). Note that these categories presume no presorting of waste (Table 2).

Characterizing the Waste Stream

Along the way, the types of shipboard activities that generate the plastic waste will be identified. Each solution to Annex V is simplified if the shipboard waste streams are kept separate, rather than being mixed. Clean plastic can be kept separate from the food-contaminated plastics and both can be collected separately from the other waste that can still be discharged at sea. But making waste separation work requires some cooperation from the crew members or passengers on the ship. It was suggested that SNAME help develop some simple crew and officer training sessions on waste source separation, to motivate and inform people about how it simplified the overall Annex V solution.

The Zero Equipment Option

Having evaluated the amounts, sources, and types of plastic being used and discarded as a result of shipboard operations, the ship operator can implement a few shipboard and shoreside managerial options to comply with Annex V. Those actions require no equipment installations, but affect shoreside company practices and shipboard crew practices.

Change the Purchasing of Ship Supplies

A quick scan of most ships identifies where plastics and other problem materials are used (Cavaliere 1988). In a number of uses, plastic has become the material of choice because it is safer to work with and is unbreakable. Other uses aboard ship, however, are convenience uses, just as are the uses of plastic ashore. It is similar to converting a shoreside business or home away from using the plastics that are so easily available.

Table 2.--Incinerable waste categorized according to its heat release.

Type	Classification of solid combustible waste materials	KJ/kg	Incom- bustible solids %	Moisture content %	Kg/dm ³
0	TRASH, a mixture of highly combustible waste--paper, cardboard cartons, wood boxes, and combustible floor sweepings from commercial and industrial activities. Contains up to 10% by weight of plastic bags, coated paper, treated corrugated cardboard, oily rags, and plastic or rubber scraps.	20,000	5	10	0.15-0.2
1	RUBBISH, a mixture of combustible waste--paper, cardboard cartons, wood scrap, foliage, and combustible floor sweepings from commercial and industrial activities. Contains up to 20% by weight of restaurant or cafeteria waste, but little or no treated papers, plastic, or rubber wastes.	15,000	10	25	0.15-0.2
2	REFUSE, an approximately even mixture of rubbish and garbage by weight, common to apartment and residential occupancy.	10,000	7	50	0.2-0.3

When a plastic item loses its disposable advantage, it loses most of its purchase appeal. Sometimes, an item that is currently purchased can be done without. A plastic item can be purchased in another material, a disposable item can be eliminated in favor of buying a reusable version that needs to be washed. A formal inventory may not be needed as much as a scavenger hunt for the unnoticed plastics that then become candidates for elimination.

Establish Who Is Responsible Aboard the Ship

Every cause needs a champion. In the U.S. merchant marine, jobs are commonly defined rigidly as deck, engine, or steward. Tackling Annex V implementation falls in no single department and requires the participation of all personnel or passengers aboard. Each ship should designate a specific person to be responsible for shepherding the entire ship into compliance with Annex V.

Port Reception Facilities

Send It Ashore, But Where?

Each port must provide reception facilities for shipboard garbage. If Annex V creates uncertainties for ship operators, the port operators are just as uncertain about what to do. In the past, it has been difficult and expensive for some ports to provide the reception facilities required by the previous MARPOL annexes, so ports are not thrilled by the obligation to provide "adequate" garbage reception facilities. The task is difficult for the port, which can only guess at (1) the number of ships bringing in foreign "food wastes" that will need Animal and Plant Health Inspection Service (APHIS) certified disposal, and (2) the amount of plastic-contaminated waste that will now be brought ashore to add to the port community's shoreside waste stream.

Many port cities are already straining to deal with their own municipal garbage problems, and adding more ship-generated garbage to the local landfill is not an easy thing to sell. The sentiment of those attending the M-17 meetings seemed to be that most U.S. ports are unprepared to meet the reception facility requirement and the ship operator will still be left "holding the bag."

Some German ports have already imposed a vessel fee, whether the vessel uses the port garbage service or not. At least one U.S. port is considering the same action (Nightingale 1988). Such a fee would be about \$150 or more per ship per port call. Such mandatory fees affect the ship operators on those routes, because they may lessen the incentive to install extensive garbage handling equipment on board the ship.

When contracting for disposal services, the usual measurements are tons or cubic yards, because that is how the hauler is charged at the landfill (NSWMA 1988).

Animal and Plant Health Inspection Service Requirements

The port reception facilities must include APHIS waste-handling facilities. The APHIS restrictions are intended to prevent the introduction of foreign plagues, such as foot-and-mouth disease, into the United States (U.S. Department of Agriculture undated). Any organic wastes that have possibly been contaminated by contact with foreign foodstuffs or foreign livestock are subject to quarantine and can be handled only by authorized APHIS contractors or APHIS personnel themselves. The wastes taken off the ship must be totally sterilized, either through autoclaving or by incineration, and the remaining matter must be securely landfilled. The APHIS requirements are not new and are not changed by the Annex V regulations. Much of the plastic wastes coming from ships as a result of the Annex V regime, however, will be food packaging or food serving articles and is subject to the APHIS restrictions. That volume of waste may increase greatly, especially in the interim compliance periods, when vessel operators may prefer to store rather than treat the plastic wastes.

All APHIS wastes must be stored separated from other garbage to avoid contamination, and when off-loaded, it must be delivered to a certified facility for proper sterilization or incineration. All transport must meet strict quarantine requirements. As a result, APHIS waste is expensive to handle. A ship may be billed by the pound of APHIS waste handled per pick up and frequency of pick up, since transport to an APHIS-certified facility is regulated. Ship officers and crew should make all efforts to keep the APHIS waste separated from regular garbage that does not require quarantine. Otherwise, the APHIS inspector, who makes the final decision as to which wastes must be quarantined, may require much larger amounts of ship's garbage to be quarantined, at the expense of the ship operator. The APHIS wastes should be stored on board the ship in leakproof containers until removal. There is no approved container, and it was the opinion of the waste handlers at the meeting that a Rubbermaid Roughneck, such as is used for curbside garbage pickup, was probably sufficient.

There are currently no more than 43 APHIS facilities in the area of U.S. ports. Wastes may not be transported through rural areas, which makes reaching some remote marine terminals almost impossible. Ship operators should contact their local APHIS officers immediately to get the details about any existing or planned APHIS-certified facilities in the vicinity of the ports where they anticipate needing APHIS wastes handled. It was also suggested that operators inquire of their shipping agents or ship management agents what kind of services the agents can provide.

Recently, APHIS administrators have brought up a new concern about handling compacted wastes. The APHIS regulations for steam sterilizing foreign garbage are based on experience with handling fresh, uncompacted wastes such as are off-loaded from an international airline flight. The autoclaving procedures depend on killing temperatures penetrating the core of the mass of garbage, and a half hour has generally proved effective with a margin of safety. However, well-compacted wastes are, by definition, more dense, and the APHIS has no confidence that a half hour of steam exposure will penetrate the core of the garbage mass. Practically, this means

that a ship operator should now be cautious about compacting APHIS wastes as well, because the savings in storage may be offset by a higher cost for APHIS disposal. The APHIS is likely to have to recalibrate the autoclaving time to compensate for the degree of compaction of the wastes (i.e., 10 to 1, 20 to 1, 40 to 1) in order to ensure that the steam penetrates to the core and kills. Any longer interval of autoclaving is certainly going to increase the operating costs of the autoclave and the price of APHIS disposal for the compacted wastes.

The APHIS-quarantined wastes are not the same as "infectious" wastes. It is important not to confuse the two, because it is much more expensive to dispose of infectious wastes (i.e., hospital wastes). Plus, there is so much public outcry over the recent well-publicized incidents of hospital waste washing ashore on eastern U.S. beaches that no ship operator should invite trouble by using the term "infectious waste" when he means APHIS-quarantined waste.

The Shipboard Equipment Option

Though Annex V requires no equipment to be installed on a ship, many operators will choose to add garbage handling equipment such as compactors, comminuters, or incinerators. Each piece of equipment installed, whether new or retrofit, needs to operate safely and effectively and not create any disease hazards for the shipboard personnel. In all cases, the tradeoffs need to be identified before expensive decisions are made. In the MEPC Guidelines, Section 5, "Shipboard Equipment for Processing Garbage" requests ". . . information on the development and use of shipboard. . ." comminuters, compactors, and marine garbage incinerators. This is a genuine request for a technical exchange and is another of the provisions unique to Annex V implementation.

Comminuters: Specifications and Installation Needs

Comminuters are a type of heavy-duty garbage grinder. Though not required by Annex V, comminuters are mentioned in both Annex V and in the proposed U.S. regulations. They are useful primarily for handling galley and scullery wastes that can be discharged in the zone between 3 and 12 nmi offshore (or anywhere within a special area). Comminuters must reduce the wastes to pass through a screen 25-mm (1-in) square. Such equipment is available for galley installation and works well.

Storage tanks for comminuted food wastes were discussed briefly. Such a tank allows a ship's steward to continue comminuting food wastes while within 3 nmi, but avoid discharging it into restricted waters. If used, the tanks must be installed so as to be easily flushed clean. Tank materials must be able to withstand potential corrosion from rotting food slurries and must be adequately vented to prevent anaerobic conditions in the tank.

Comminuted food wastes should never be flushed to black water (sewage) holding tanks or to marine sanitation devices (shipboard sewage treatment plants). Food wastes cannot be adequately degraded by the microorganisms

in the systems and can overload the aerobic capacity of the tanks and make the whole system go septic. Such a ghastly mess must be avoided at all costs.

Pulpers

The SOMAT Corporation makes a pulper for use on U.S. Navy ships that works like a comminuter, but further processes the slurried waste to press out the water and dry the waste material enough to make it easier to burn or store. One unit can separate out plastics because the plastics float in the pulping chamber while the other wastes pass through. These devices are about the size of a washing machine.

Compactors: Specifications and Installation Needs

Garbage compactors, used on some ships, have had mixed success. Many of the original units were never intended to be installed on a rolling, heaving ship or be used in the salt-laden sea atmosphere. Purpose-built units for marine installation are now readily available, however, and they fare better. The principal reason for using a compactor is clear: garbage storage takes less room. However, the stored compacted garbage, especially food-contaminated plastics, can "ripen" to a stinky mess if not properly isolated and disinfected. The M-17 discussion raised the following points:

- Hygiene for stored wastes needs to be guaranteed, since the accumulated wastes will otherwise rot and attract vermin. The U.S. Public Health Service (PHS) has standards for shipboard sanitation that must not be compromised. To prevent rotting, food wastes may need to be frozen or at least refrigerated in the 40°F cold room until disposal ashore. This may cut into the steward's storage space.
- Though compactors are usually not large, using them requires organizing the garbage collection and installing them requires identifying enough space for storing the compacted garbage as well. Both the compactor space and the storage space should have adequate space drains and hose washdown fixtures. The discharge of the "compactor juice" created is not regulated, as far as anyone present knew.
- Compactors may be used with unsorted garbage or with separated waste streams. It may be worthwhile for a ship to install more than one compactor, if one is used principally for the APHIS wastes generated from the galley and the other is used for all other plastic-containing wastes generated on board the ship that do not need quarantine. One fleet operator suggested that compactors were also useful for baling dry wastes to be recycled.

Incinerators: Specification, Selection, and Use

Using an incinerator for plastic wastes enables the ship crew to destroy the troublesome wastes rather than hold them and rely exclusively

on port reception facilities or shoreside waste haulers. Ship operators consider incinerators in a tradeoff study against other compliance options in the implementation of the MARPOL Annex V. The Annex V rules require only compliance, and the ship operator will want to know as much as possible about the consequences of installing an incinerator before making the decision.

Incinerators available for shipboard use differ significantly from each other and cannot be considered all the same. They differ in number of combustion chambers, rate of waste feed, form of waste feed, auxiliary fuel required, pretreatment or waste separation needed, amount of operator training needed, auxiliary equipment or ventilation needed, retrofit installation difficulties, and other ship-specific parameters.

Before installing an incinerator, a ship operator must know what needs to be incinerated and in what amounts. Some wastes require shredding or similar pretreatment before incineration. Other wastes require more fuel to destroy than makes sense, so those wastes (e.g., metal scraps) should be separated ahead of time. Some wastes (e.g., glass) cannot be incinerated in some incinerators, but are handled by others. No single size vessel "needs" an incinerator. Ship owners are going to make this type of decision based more on how many people are on the ship and how the ship operates.

There are no technical standards for shipboard incinerators under Annex V. Neither the IMO nor the U.S. Environmental Protection Agency have set effluent or emission standards for the ashes, residues, or stack gases. The American Society for Testing and Materials (ASTM) Committee D-34 Waste Disposal is the proper group to develop performance and effluent standards, but there is no activity in D-34 to develop standards for incinerators of any size. The last attempt to develop such a standard failed due to a lack of agreement on what was acceptable.

In the United States, the operation of incinerators of a size suitable for shipboard use is not regulated by the Federal Government. At the state and port level, the regulation of small incinerators in communities ashore varies greatly. A shipboard incinerator might be subject to local incinerator restrictions if the incinerator is used while the ship is in port, just as ships operating in some California ports have to burn different fuels in order to meet the local air quality restrictions.

There are no IMO, United States, or Coast Guard residence time or minimum temperature standards for the combustion chamber used in shipboard incinerators. The Waste Combustion Equipment Council is working on an "industry standard" for incinerator performance. The classification society Germanischer Lloyd has developed regulations (Germanischer Lloyd 1987) and the Norwegian ship classification society Det Norske Veritas also has regulations for the equipment Det Norske Veritas (1980). In the United States, a shipboard incinerator construction standard and a selection guide are being developed under ASTM Committee F-25 Shipbuilding. When that is completed and accepted by ASTM, the Coast Guard may accept it as a technical standard. Until then, the Coast Guard is constrained to regulate

incinerators according to its existing marine engineering regulations on auxiliary boilers: control systems, flameout protections, space ventilation, enough room around the installation, and fire protection.

I am skeptical about using garbage incinerators for destroying shipboard plastics. The Canadian experience with municipal incinerators seems to have fallen short of what incineration might promise, because operating the plants perfectly is so crucial to the environmental effectiveness of the technology (Mohr 1988). Shipboard incinerators, unless carefully tested and tended, may only make the plastics prohibition another shell game by dumping dangerous substances into the sea via the air and the ash.

Integrated Waste Collection, Treatment, and Disposal

Some firms have developed totally integrated ship waste handling systems, and these have been installed on a number of passenger vessels. Successfully operating these systems requires that the crew learn how the system operates and uses the parts of the system to their best advantage. Unfortunately, some have proved too easy to ruin when silverware is thrown into the incinerator or the shredder is fed a full dose of bed linens.

EXAMPLES FROM THE FLEET

How have ship operators actually responded? Many organizations are still trying to make cost-effective decisions about how to comply with the new Annex V requirements. Some examples can be given, but the identities of the fleets have been removed because this information is largely anecdotal and companies may still decide to change their approach as they develop permanent compliance. The compliance approaches presented are eliminating shipboard plastics, installing compactors, installing fuel-fired incinerators for select waste streams, and installing an incinerator for all shipboard wastes.

Eliminating Plastics

Company A operates chemical carriers in the domestic trade of the United States. Most of the voyages are between refineries in the Gulf of Mexico and tank farms along the eastern seaboard. A separate portion of the fleet services the west coast of the United States. The crews make short voyages with frequent port stops, but no foreign trips. For this firm, APHIS restrictions pose no problem. However, because the Gulf of Mexico is part of a trade route, the company has to think now about what might be needed to comply with a special area designation in the Gulf of Mexico. Company A began changing its supply procurement practices in 1987 to eliminate plastics and disposable goods wherever possible. Polystyrene coffee cups were banned and heavyweight paper cups were purchased instead. Crew support for eliminating plastics has been strong, because the trash problem in the Gulf of Mexico is apparent as they sail her waters. Further material substitutions will be made, such as asking suppliers to deliver maintenance supplies in metal cans rather than plastic containers. The company has no clear idea how much plastic waste they can eliminate from the ships' garbage, but they want to do as much as they can with replacing

materials before locking themselves into an engineering solution. Equipment may be limited to galley comminuters and compactors, which fit more easily into the vessels' operations than do incinerators.

Compactors

Company B operates tankers with few people aboard along trade routes that bring the vessels into U.S. ports frequently. Company B chose to install compactors, after having previously considered installing incinerators on each vessel. The compactors are intended to be an interim compliance solution and Company B has not ruled out installing incinerators. First, however, the company wants to see what the wastes are on the ships, how they can be changed to nonplastic materials, and what standards for incinerator performance are developed by ASTM or IMO.

Incinerators for Selected Wastes

Company C ships carry general cargo and have a relatively small number of people working on board any vessel. They operate an irregular trade route around the Pacific with no guarantee of port facilities in some of the less frequently visited ports. Company C chose to install purpose-built diesel-fired incinerators to handle their waste while at sea and free them from relying on the uncertainties of the foreign port facilities. The managers and operators of the vessels appear satisfied with the units, which are regularly used.

Incinerator Installed for All Shipboard Wastes

Company D has tankers, so the ships have relatively few crew members, no passengers, and a steady load of maintenance activities with the probability of small oil leaks around machinery and deck piping fixtures. They service relatively remote terminals where it would be difficult to arrange for APHIS waste disposal. This firm elected to retrofit a large incinerator, so that the crew could destroy all the shipboard wastes without relying on port facilities for any garbage disposal. The experience of the fleet operators has been that the system works well so long as a high temperature is attained in the combustion chamber. One unanticipated limiting factor has been the disposal of dirty oil sorbent pads. The material of the pad itself burns nicely, but it also melts if the combustion chamber is not hot enough, and the melting pad material can pool and leak out the air inlets of the incinerator. The temperature of the chamber must be high enough when the pads enter the chamber to take the waste directly to combustion.

FUTURE MARPOL ANNEX V DEVELOPMENTS: SPECIAL AREAS

The term "special area" means an area where no dumping of garbage is allowed (U.S. Department of Transportation 1988a). An important point is that the requirements of each special area do not go into effect until all the national authorities bordering on the proposed special area officially notify the IMO that adequate reception facilities are available. Only then

does IMO send out a global notice of the special area designation, and 1 year later the designation goes into effect. At this point, only the Baltic Sea has met that requirement. The Baltic Sea will be a special area as of October 1989. The North Sea has also been proposed and the border countries are filing notices with IMO. The formal designation of the North Sea as a special areas under Annex V will occur in the fall of 1989. The Gulf of Mexico, bordered by Mexico, the United States, and Cuba, has been suggested as a special area under the MARPOL Annex V (IMO 1988a). However, neither Cuba nor Mexico are signatories to the MARPOL convention and it is not possible for IMO to designate a special area without the advance consent of all nations surrounding the proposed area. The problems that face those who favor designating the Gulf of Mexico as a special area point up the limitations of MARPOL Annex V, even after its entry into force.

SUGGESTIONS FOR FUTURE DEVELOPMENT

Educating Passengers and Tourists

Passenger ships have been the source of a lot of garbage tossed into the sea (Smock 1988). Citizen outreach should include a campaign to tell potential passenger ship customers about MARPOL Annex V and their role in ensuring its success. Environmental professionals interested in eliminating marine debris should target the travel industry and the vacationing public with information emphasizing the benefits of the changes in shipboard handling of plastic materials. Passenger ship operators will be affected by this pollution prevention annex far more than they have been affected by any previous MARPOL annex. As "hotels" in a highly competitive travel and leisure market, passenger ship operators must have some assurance that complying with Annex V can be accomplished with a minimum of disruption to the carefree atmosphere that they try to provide to the vacationing passengers. Recent practice on many short cruises and "party boats" operating out of U.S. ports has been to use disposable materials, principally plastics, for many food and drink services. On longer voyages, passengers bring a variety of personal products for their own use. In North America at least, travellers can buy shampoos, razors, and deodorants in convenient unbreakable small plastic packages that are expected to be discarded when empty. Till now, there was no reason for a ship passenger to think twice about bringing aboard plastic containers. This may be the first time a person realizes the toothpaste tube is plastic. The MARPOL Annex V changes all that, and the travelling public should be encouraged via the travel magazines and other literature to learn to leave the plastic disposables ashore or to expect to keep them until returning to shore. The travelling public is increasingly sophisticated and will probably be glad to make such small changes, if it protects the pristine open sea that they desire.

Switching to nonplastic items will be a bigger adjustment on these vessels, because it will affect how service is delivered to the fare-paying passengers. First, fare-paying passengers cannot be compelled to sort their garbage, as crew members can, so the ship operator must devise a shipboard system that either handles all the collected waste without sorting, or depends on the crew members to separate the trash after it is

picked up from throughout the ship. Second, replacing plastics in some instances will increase costs directly. For example, paper cups cost two or three times as much as expanded polystyrene cups cost. A passenger ship operator should be rewarded, not penalized, for moving away from the disposable plastics that are cheaper to use. Passengers and environmental organizations should praise the successful ship operators and challenge the others to do as well.

Develop Plastic Melters for Port Facilities or for Ships

This promising technology is already available for shoreside use, though it has not been widely used in the United States. The process reduces the volume of the typical plastic trash by a factor of 40, creating extruded blocks of plastic that are also likely to meet the hygiene and vermin-killing standards of the APHS and PHS for shipboard sanitation. Plastic treated by a "melter" does not have to be sorted by material types. On the other hand, the plastic handled this way is not being recycled as such; it is simply being reduced in volume and must still be returned to shore for disposal.

Burning plastics may not be necessary, if a manageable melter for plastic wastes can be devised. A small unit might be useful for a passenger vessel or a small terminal, and a larger unit might be very useful for a remotely located terminal that has no easy way to handle the plastic wastes that are delivered by ships. The U.S. Navy is researching the development of a shipboard piece of equipment for use aboard its larger vessels, but that work is not expected to produce a prototype for several years.

Repackage With Selective Plastics

All plastics are not equal (Society of the Plastics Industry 1988). Incineration creates different combustion products, for example. Polyvinyl chloride polymer contains chlorine atoms, so that incinerating these materials is guaranteed to release hydrochloric acid. Other plastics, such as polyethylene or polypropylene, burn with less hazardous combustion products. In addition, plastics melt at different temperatures and react differently in waste treatment processes. So, the operators may gain some advantage in waste handling by changing from one kind of plastic to another. By investigating the types of plastic used in the products brought aboard ship, the ship operator can retain the advantages of using plastic in some products, but the disposal problems can be simplified, both at the ship and in the ports (Council on Plastics and Packaging in the Environment 1988).

In some senses, repackaging means rethinking away from disposable items to items that have a longer life span. Some shoreside restaurants have replaced plastic plates with plastic or wicker baskets that are lined with paper napkins for each serving. Yes, more paper napkins get used, but the waste bin has much less plastic debris in it.

Switch to Degradable Plastic

Some new plastic resins are being marketed as degradable, either because they photodegrade in ambient light or because they biodegrade when microbes decompose certain starches or celluloses used in their manufacture. These resins and any products made from them are considered plastic under Annex V and the Coast Guard regulations.

Refining Waste Handling to Large-Scale System

Now the APHIS system is set up to handle daily operations at airports, but it is much less prepared to handle daily operations at seaports. The APHIS organization must make the transition to be able to handle a larger volume of wastes, without the extraordinary arrangements that ship operators are having to deal with today. That may require more APHIS inspectors and will certainly require more APHIS-certified facilities accessible to arriving ships throughout the country.

Long-Range Prospects--Ten Years Down the Road

Ten years from 31 December 1988 should find the MARPOL 73/78 Annex V (Garbage) transition complete. At some time between now and then, ports and ships will learn how to manage shipboard garbage in a way that satisfies the annex, safeguards the sanitation and hygiene of the ship, safeguards the port country from animal or plant pestilence, and compensates the disposal firms without bankrupting the ship operators. After all, the shift in shipboard garbage disposal is not happening in isolation. Concurrently, shoreside communities and industries are recognizing that the present disposal practices for municipal garbage must change and that the popular use of "disposable" products leaves a permanent legacy ashore, just as it does at sea. The changes that the annex demands of the ship operators may soon be mirrored ashore. As more people confront the same garbage handling problems in their businesses and homes, engineers and designers will have more incentive to develop efficient products and processes, which may then offer a better alternative to the ship operators than those that exist now. Ten years from now, taking out the garbage will be very different, whether you are at sea or on land.

ACKNOWLEDGMENTS

I thank David Cottingham, Office of Chief Scientist, NOAA, Washington, D.C.; the Waste Combustion Equipment Council, NSWMA, Washington, D.C.; and the Ships Machinery Committee, SNAME, Jersey City, NJ.

REFERENCES

Cavaliere, L. A.

1988. Trash management at sea - A summary of Annex V to MARPOL, The International Convention for the Prevention of Pollution from Ships, presented to the International Ship Suppliers Association, September 27, 1988.

Council on Plastics and Packaging in the Environment.

1988. Society of the Plastics Industry fact sheet, plastic packaging recycling. Council on Plastics and Packaging in the Environment.

Det Norske Veritas.

1980. Boilers, pressure vessels, thermal-oil installations and incinerators. In Rules for the classification of steel ships, machinery and systems. Part 4 Chapter 3, January 1980. Det Norske Veritas.

Germanischer Lloyd.

1987. Regulations for the design, equipment and testing of waste incinerating systems on seagoing ships. July 1987. Germanischer Lloyd.

Horsman, P. V.

1982. The amount of garbage pollution from merchant ships. Mar. Pollut. Bull. 13:167-169.

International Maritime Organization.

1978. Annex V, Regulations for the prevention of pollution by garbage from ships. In the International Convention for the Prevention of Pollution from Ships, 1973, as modified by the Protocol of 1978 relating thereto (MARPOL 73/78).

1988a. Acceptance by the Government of the United States, optional annexes, including the understanding to make "every reasonable effort" to make the Gulf of Mexico a "special area" under MARPOL 73/78 Annex V. Submitted by the United States to the International Maritime Organization, issued as Int. Maritime Organ. Doc. PMP Circ. 72, 11 January 1988.

1988b. Marine environment protection committee, guidelines for the implementation of Annex V, regulations for the prevention of pollution by garbage from ships [includes the full text of Annex V as an appendix]. Especially section 5 "Shipboard equipment for processing garbage", dated September 8, 1988. Issued as Doc. No. MEPC 26/WP.6.

Martinez, L.A.

1989. Compliance deadline for MARPOL 73/78 Annex V, in technical and research activities section. Mar. Tech. 26:1, p. xxvii.

Mohr, M.

1988. The burning question. Incinerate or recycle: What should we do with our garbage? Harrowsmith 78, March/April.

National Solid Waste Management Association.

1988. The 1987 tip fee survey - Last year's rise was biggest ever. 1730 Rhode Island Ave., N.W., Suite 1000, Wash., D.C. National Solid Waste Management Association.

Nightingale, D. E. B.

1988. Port of Seattle weighs options for handling MARPOL V waste. World Wide Shipping, October/November.

Norske Hydro.

1988. Classification of solid combustibile waste materials. Sales literature for shipboard waste disposal system including incinerators. Norske Hydro.

Signorino, R.

1988. Office of Maritime Safety Standards Programs, Occupational Safety and Health Administration, U.S. Department of Labor. (pers. commun. to L. Martinez as chair of M-17), July.

Smock, D.

1988. Are shipboard plastics all washed up? Plastics World, September.

Society of Naval Architects and Marine Engineers.

1982. Guide for the disposal of shipboard wastes. Prepared by Panel M-17 (Disposal of Shipboard Wastes) of the Ship's Machinery Committee, January 1, 1982. Tech. Res. Bull. 3-33. [Available from SNAME, 601 Pavonia Avenue, Jersey City, NJ.]

Society of the Plastics Industry.

1988. Voluntary plastic container coding system. Society of the Plastics Industry.

U.S. Congress.

1987. Title II - Marine Plastic Pollution Research and Control Act of 1987. Public Law 100-220, One Hundredth Congress of the United States of America. U.S. Congress.

U.S. Department of Agriculture.

Undated. Animal and plant health inspection service regulations. Title 9, Code of Federal Regulations, Part 94, Section 94.5 Garbage: regulations on storage and movement on certain means of conveyance. U.S. Department of Agriculture.

U.S. Department of Commerce.

1988. National Oceanic and Atmospheric Administration. Office of Marine Operations Instruction 9593. Subject: Pollution Abatement and Control, with Appendix B, Pollution Control Equipment Survey, issued January 6. U.S. Department of Commerce.

U.S. Department of Transportation.

1988a. Transportation Systems Center. Economic analysis of designating the Gulf of Mexico as a "special area" under Annex V of the MARPOL Protocol (draft final report). July 22, 1988. U.S. Department of Transportation.

- 1988b. United States Coast Guard. 33 CFR 151 and 158 Regulations implementing the pollution prevention requirements of Annex V of MARPOL 73/78; advance notice of proposed rulemaking. Federal Register, Part IV, p. 23884 through 23895, Friday, June 24, 1988. U.S. Department of Transportation.
- 1988c. United States Coast Guard. 33 CFR 151, 155, and 158. 46 CFR Part 25. Regulations implementing the pollution prevention provisions of Annex V of MARPOL 73/78; notice of proposed rulemaking, Federal Register, Part V, p. 43622 through 44617. Thursday, 27 October 1988.
- Vauk, G. J. M., and E. Schrey.
1987. Litter pollution from ships in the German Bight. Mar. Pollut. Bull. 18:316-318.
- Whitehead, J. R.
1988. Reducing plastic pollution in the marine environment: The U.S. Coast Guard and implementation of Annex V of MARPOL 73/78. In Proceedings of Oceans '88, Baltimore, MD, November 1, 1988.

LOW TECHNOLOGY (BURN BARREL) DISPOSAL OF
SHIPBOARD-GENERATED (MARPOL V) WASTES

Terrill J. Chang*
SCS Engineers
Redmond, Washington 98053, U.S.A.

*Present address: c/o Camp Dresser & McKee, One Woodward Avenue, Ste.
1500, Detroit, MI 48226.

ABSTRACT

To help ensure more widespread compliance with marine disposal laws, alternatives are needed that are applicable to a variety of wastes, but are less costly than using marine incinerators. Burn barrels are low technology burners for disposing of MARPOL wastes at sea. They are not considered state-of-the-art combustion devices, but they are a practical, technically feasible alternative. They comply with existing environmental and marine regulations and are currently in use. Design and operational guidelines for optimizing safety and combustion in burn barrels were developed.

INTRODUCTION AND REGULATORY FRAMEWORK

Traditionally, shipboard waste has been dumped at sea without regard for the impact of the waste on marine life or navigation. Marine debris is recognized as a growing problem, threatening marine life, beaches, and vessel safety worldwide. Public Law 100-220, The Marine Plastic Research and Control Act (effective 30 December 1988) domestically implements MARPOL Annex V. It restricts at-sea discharge of garbage to certain zones and bans all at-sea disposal of plastics. In addition, it requires ports to have available suitable waste reception facilities.

Shoreside disposal of certain wastes is restricted by U.S. Department of Agriculture (USDA) regulations for controlling plant and animal diseases and pests. Wastes contaminated by food from foreign ports (except Canada) must be enclosed in leakproof containers and brought ashore under USDA supervision. These wastes must then be disposed of via incineration, steam sterilization, or grinding into a sewer. Thus, existing regulations make disposal of shipboard wastes more difficult both at sea and ashore.

ALTERNATIVES FOR DISPOSAL OF SHIPBOARD WASTES

Under MARPOL and USDA regulations, vessel operators are faced with the following disposal alternatives, each of which has drawbacks:

- Incineration. True marine incinerators, those with combustion air fans, are expensive, moderately complicated, and occupy valuable deck space.
- Grinding. This process is generally suitable only for food wastes and cannot be used to dispose of plastics at sea.
- Compaction. Requires wastes to be stored, using valuable space. It creates possible odor problems and a potential for morale and aesthetics problems as the crew must work in close proximity to stored garbage.
- Onshore disposal (per local solid waste or USDA regulations). Drawbacks are similar to compaction, but uncompacted garbage requires even more space aboard ship.
- Biodegradable plastics. This may apply to some packaging materials; biodegradable rope and nets are not yet available; degradation products may be toxic or otherwise harmful to marine life.
- Burn barrels (low-technology burners). These may produce excessive harmful emissions or pose safety and fire hazards.

BURN BARRELS--ONE POSSIBLE SOLUTION

No single technology or disposal method can solve the marine debris problem by itself. A variety of technologies are necessary to accommodate the range of wastes produced, vessel and crew sizes, trip durations, and missions. To help ensure more widespread compliance with marine disposal laws, alternatives are needed that are applicable to a variety of wastes, but are less costly than using marine incinerators.

It has been reported that a variety of vessels are currently utilizing low-technology burn barrels to dispose of their wastes at sea. Burn barrels are simple (typically, a 208.2-L (55-gal) drum with some holes cut in the sides), "low tech," and similar to those barrels commonly used to burn household trash during the 1950's. The Marine Entanglement Program, National Marine Fisheries Service, NOAA, retained SCS Engineers to evaluate the technical feasibility, safety, and potential environmental impacts of using burn barrels to dispose of shipboard wastes.

It should be stressed that the burn barrel technique is not being advocated. However, under present regulatory authority, such technology is permissible and actively being utilized. Thus, the goal of the investigation was to provide guidelines for burn barrels that would enable legal

disposal of shipboard wastes while protecting the environment, shipboard personnel, and the vessel itself.

Design Guidelines

Because actual construction and testing of a burn barrel were beyond the scope of this project, guidelines for the design, construction, and operation of burn barrels were developed (Fig. 1). A primary consideration was to optimize combustion, that is, to make the barrel act like an incinerator as far as practical in a unit with no moving parts. Types of wastes to be burned, environmental regulations, operator safety, and fire prevention were also evaluated. The burn barrel should be large enough to burn the expected volume of waste in a reasonable time, but without occupying too much deck space.


As shown in the schematic, a complete burn barrel installation should have the following features:

- Combustion chamber (208.2-L (55-gal), 16-gauge steel, Department of Transportation standard 17C drum) located inside an 321.7 L (85-gal) steel overpack drum.
- Combustion air inlets for underfire and overfire air.
- Air gap to cool combustion chamber and preheat combustion air.
- Spark arrester, grate, ash scoop or pan, rain cap.
- Suitable anchoring and insulation.
- Adequate clearances from all combustible surfaces.
- Location: aft and downwind.
- Nearby fire hose or extinguisher and first-aid kit.

Operational Guidelines

Operational guidelines for burn barrels are largely a matter of common sense. Primary concerns are good combustion and fire safety.

- Burn during calm sea conditions, avoiding rainy weather.
- Build the fire with loosely stacked paper and wood kindling, not with flammable liquids. Add plastics in small amounts to ensure burning rather than melting.
- Avoid potential explosives such as liquid-filled bottles and aerosol cans.



SCHEMATIC OF BURN BARREL

NOT TO SCALE

PRELIMINARY DRAWING
NOT FOR CONSTRUCTION

Figure 1.--Schematic of a burn barrel.

- Agitate the wastes with a metal poker for more complete combustion.
- Don't cook food over the fire.
- Wear safety goggles or glasses while operating the burn barrel.
- Avoid inhaling burn barrel smoke and fumes, which might contain hydrochloric acid from burning plastics.

ENVIRONMENTAL CONSIDERATIONS

Ash Disposal

The ash resulting from proper burn barrel operations should consist of only sand, dirt, metal, and glass (all of which do not burn); small amounts of unburned carbon (similar to charcoal); and small globules of melted plastic. The latter are still considered to be plastic and cannot be legally disposed of at sea. Since separating the plastic is inconvenient, the entire supply of ash should be stored in a metal container and disposed of properly ashore.

Air Emissions

The U.S. Environmental Protection Agency currently has no air pollution regulations that apply to burn barrels, and has no plans to promulgate any. State and local air pollution district regulations vary widely, and some of these may affect the use of burn barrels in certain coastal jurisdictions.

Actual air emissions from a burn barrel were not tested, but major components are expected to be water vapor, carbon dioxide, particulates (smoke), carbon monoxide, small amounts of hydrogen chloride (from chlorinated plastics and salt air), and various products of incomplete combustions.

The combustion conditions in burn barrels are more similar to open burning than to an incinerator. Furthermore, they lack air pollution control equipment to clean emissions. On the other hand, due to the small quantities of wastes per burn barrel, airborne emissions are expected to be modest. Air quality impacts from burn barrels operated on the open ocean are not anticipated to be significant. However, emissions testing of burn barrels is warranted.

COST CONSIDERATIONS

A burn barrel is expected to cost approximately \$500, while steam sterilization of wastes to meet USDA regulations is estimated to cost about 30¢/lb. Assuming a 5-day trip with a 30-person crew generating 1.64 kg (4.4 lb) per person per day, it would cost about \$200 to dispose of their wastes via steam sterilization. If a burn barrel were used instead, it would pay

for itself in only 2.5 trips. Alternatively, a marine incinerator would cost upwards of \$20,000.

CONCLUSION

While they are not considered to be state-of-the-art combustion devices, burn barrels are a practical and technically feasible alternative. When properly used, they appear to comply with existing environmental and marine regulations. It is believed that they can provide a safe, convenient, and low-cost alternative to either onshore disposal or incineration of shipboard-generated wastes.

PROVIDING REFUSE RECEPTION FACILITIES AND MORE:
THE PORT'S ROLE IN THE MARINE DEBRIS SOLUTION

Fran Recht and Sidney Lasseigne
Pacific States Marine Fisheries Commission
Marine Debris and Habitat Project
Depoe Bay, Oregon 97341, U.S.A.

ABSTRACT

A marine debris pilot project was conducted by the Port of Newport, Oregon, in response to the requirements of MARPOL Annex V. Project findings are summarized. The authors also document what has been learned from the preliminary application of these findings to several other west coast U.S. ports. They discuss the different aspects of refuse collection and recycling that should be examined by a port when gearing up for adequate dockside refuse disposal.

The technical part of dealing with marine refuse in a west coast U.S. port is seldom difficult. Initiating action can be-- many ports and mariner groups do not consider marine debris a high priority item. However, those with port responsibility have responded positively to outside encouragement and minimal assistance.

INTRODUCTION

If mariners are expected to return plastics and other refuse to port they must have convenient dockside refuse disposal available to them. The U.S. Coast Guard has acknowledged this when writing the regulations to implement the provisions of MARPOL Annex V. Regulations require that all commercial ports and docks, no matter their size, provide refuse reception facilities.

While it may seem logical to expect all docks to have garbage containers, similar laws designed to control ocean pollution (e.g., those regulating oil or sewage disposal) have not succeeded because they failed to assure that facilities were universally available. Pollution containment facilities were required only of ports meeting some minimum size requirements. Mariners have often been uncertain where oil or sewage could be off-loaded.

Fortunately, this will not be the case with garbage. A fisherman off-loading fish at the processing house will be also able to off-load a

sack of plastics or a used net. A fisherman from California, calling for the first time at a port in Alaska, will know that he can find a place to deposit trash and that the port must accept his old cable.

With these regulations then, ports and docks have been handed a much-expanded if not new role as garbage collectors--a role that many ports are unprepared to handle. Realizing this lack of preparedness and the crucial role that ports would play in solving the marine debris problem, the Marine Entanglement Research Program, of the National Marine Fisheries Service (NMFS), National Oceanic and Atmospheric Administration (NOAA) provided funding for a pilot port project. Between January 1987 and May 1988, the Port of Newport, Newport, Oregon, conducted a demonstration program which addressed questions about port refuse disposal facility needs especially for fishermen and boaters, costs and cost recovery, and mariner education. A report, a detailed reference guide, and a videotape summary resulted from that pilot program and are available through the Marine Debris Information Offices, NOAA.

Work was begun in November 1988 to apply the findings of the demonstration project to ports and fishing groups on the west coast of the United States. It continued through June 1990. This work was coordinated by the Pacific States Marine Fisheries Commission with funding from a Saltonstall-Kennedy fisheries development grant. Various methods were used: Ports and fishing groups were sent written information and educational resources, marine debris exhibits were featured at trade expositions, and workshops and seminars were conducted with port groups, extension agents, and educators. Additionally, eight target ports (two in each of the west coast states) were visited periodically to provide them with direct assistance in assessing their refuse disposal needs and options, enlisting local mariner support, and encouraging community awareness. Targeted areas were chosen with the help of the commercial fishing industry and the Sea Grant marine agents. They were areas with active commercial and recreational fishing activity where need was shown and mariner and port support was considered likely. These port areas will in turn serve as examples for and be able to assist surrounding areas on the basis of their experiences. The target port areas were Petersburg and Homer, Alaska; Anacortes and Westport, Washington; Astoria and Coos Bay, Oregon; and Eureka and Morro Bay, California.

This paper will summarize the findings of the pilot port program. It will then mention what has been learned in applying the pilot project experiences to other port areas on the west coast of the United States.

PILOT PORT PROJECT SERVES AS MODEL

While no port is likely to be thrilled with the news that it is legally obligated to accept refuse from vessels, pilot project experiences indicated that ports can benefit by examining and improving their refuse handling system and by becoming involved in educating their customers about the marine debris problem. Some of the positive results of the Port of Newport pilot project were:

- a high and voluntary (Annex V had not yet been implemented) return of refuse back to shore;
- reduced port refuse disposal costs, despite the much-increased volume of refuse being returned;
- an improved rapport of the port with the fishing community;
- the pride of ownership mariners, port workers, and community members felt in the project; and
- frequent, positive media attention focused on the port and fishermen's efforts.

The acceptance and continuity of the marine refuse disposal project at the port resulted from:

- the convenience and comprehensiveness of the refuse reception facilities;
- the use of a simple, low-cost recycling system which reduced refuse disposal costs and port labor involvement;
- port management and worker support for recycling efforts; and
- fishermen's cooperation with port refuse and recycling efforts.

The support of commercial fishermen and other mariners for ocean cleanup efforts and for increasing community awareness of the marine debris problem resulted from:

- involvement of respected mariners and community members in the project advisory group and their willingness to take an active role in promoting awareness among their peers;
- direct contact with mariners, port workers, and community members (such as is achieved by conducting an oral survey, asking for ideas about improving the refuse system, or by organizing a beach cleanup) as a means of provoking thought and creating a sense of involvement;
- the widespread dispersal of a variety of educational and promotional materials such as brochures, decals, posters, displays, and slide and videotaped shows; and
- frequent local media reports of port, mariner, and community efforts to deal with the marine debris problem.

In work with other ports these approaches are emphasized and have been found to be generally adaptable.

APPLICATION OF PILOT PROJECT FINDINGS TO OTHER PORTS

The pilot port project differed from the situation to be encountered in most ports. The pilot project provided both focus and funds for marine debris work at the Port of Newport, minimizing port risk and financial restrictions and allowing the hiring of a full time manager, who was able to concentrate solely on marine debris work and plan and implement activities related both to facility development and education.

Though Annex V implementation might focus port attention on refuse facility work, ports we have seen seem generally unconcerned about this regulation's enforcement and have not made compliance a high priority. Most have not planned or budgeted for refuse system changes, nor assigned anyone to be in charge of carrying out necessary work. An active role in mariner and public education has not been considered by most ports.

What then of the Newport experience can be applied to other ports? Can the factors that made Newport's project successful--the convenient and cost-effective refuse disposal system which emphasized recycling, the mariner cooperation and support, and the community involvement--be counted on in other areas? What changes have been observed in work with ports that differ in physical size and layout, types of vessels and clientele served, type of equipment and services offered, and political organization?

Beginning Work With a Port

Given the general willingness of a port to look at its refuse handling situation in light of Annex V, the approach of holding a meeting to bring together the port, mariners, and refuse handlers has been found to be applicable and effective. Though it is unlikely that such a meeting will be initiated by either the port or the mariner group, we have found it easy to arrange by initiating the idea with these parties and asking the Sea Grant marine agents (or other community organizers) for help in identifying and inviting participants, arranging meeting facilities, and helping to moderate the discussion. The preliminary meeting is used to inform participants of the marine debris problem and the applicable laws, encourage the discussion of refuse service needs and options, and plan the necessary improvements.

In order to encourage the port to commit to making changes, it is important that mariner concern about the issue be evidenced by attendance at the meeting and participation in discussions. Mariner input to the port can be obtained at the meeting if the moderator will ask specific questions to draw out mariner discussion. It may be important for the moderator to discuss needs, ideas, and problems with mariners beforehand in order to foster honest discussion. The advisory group, if one can be formed, or an interested mariner group will need to follow through and continue to encourage the port to accomplish the suggested changes. While we have found that fostering the formation of an active advisory board is difficult unless a specific marine debris or solid waste project is defined, it is not an impossibility if a local person takes the initiative to keep things going, define a role for the advisory group, and call the meetings.

It is also effective for the moderator or a port official to ask about resources or specific commitments from the refuse haulers or recyclers at this preliminary meeting. (For example, What size bins can you provide? How frequently will the containers be emptied?) However, even with these details worked out, it is imperative that someone at the port be assigned to follow through to make the necessary arrangements.

In ports where either the port or the refuse service is operated under the jurisdiction of the municipality or city, special attention needs to be paid to informing these entities of mariner needs. City managers, engineers, and public works employees should be involved in the preliminary meetings with mariners and port officials in order to avoid subsequent problems and misunderstandings.

Gaining Port Interest

The attitude and interest of the port or dock manager or harbormaster regarding the marine debris problem and the new law will be the key determinants of progress regarding port refuse facility needs.

Many ports consider themselves to be already providing adequate refuse reception facilities, even though dumpsters may regularly overflow or be inaccessible, and mariners have no place to dispose of large refuse items. These ports need to be "pushed." That push can come from a manager who becomes especially interested in this marine debris issue. It is not uncommon to find such people, especially when the potential for more efficient refuse service and improved port-customer relations become apparent. Interested managers have quickly influenced change by establishing informal refuse reception areas (i.e., by beginning piles for wood, metal, and net), by speaking to mariners about new laws and port plans, by increasing the size of refuse containers, by pursuing recycling options, by including plans for marine debris facility improvements in their grant applications, and by encouraging education about marine debris in public schools.

We have found port and harbormaster groups quite interested in having their members informed of the new laws, refuse handling options, and available resources, and have found them quite willing to distribute written information and promotional materials to their members and arrange for presentations before their groups. We have also found harbormasters involved in marine debris work willing to share their experiences with their peers at port association meetings. These short presentations are extremely influential and generate many "hands on"-type questions, result in new cost-saving ideas, and foster a positive attitude toward attempting refuse system changes.

The biggest stumbling blocks we have encountered are not technical or financial, but attitudinal and political: managers who care only to meet the letter of the law, or who are unconcerned with user relations; city and port tensions that restrict port autonomy and the ability and willingness of the port to make financial commitments or changes or to solve problems (e.g., lack of cooperation between city and port in prosecuting the non-mariner citizens who use port dumpsters).

Assessing a Refuse System for Adequacy and Convenience

Analysis of the existing refuse handling system and port layout is useful in identifying problem areas and needs specific to that port's situation. It is important to critically examine refuse can or dumpster placement (for convenience and visibility), refuse container capacity, and emptying schedules to identify their adequacy in serving mariners and in handling refuse loads especially during the high-use times. The availability of carts, hoists, and forklifts for moving refuse may also be a determinant of convenience.

A one-time walk-around assessment is not adequate, however. Observations must be made during the various busy seasons of the port and these observations should be "reality checked." Resident mariners and port workers can be valuable resources for determining problem areas and times, but only if specific questions are asked. "When do refuse containers overflow and where?" gives much better information than asking, "Is the refuse system adequate?" It is also important to observe what mariners or the port actually do, despite what one might be told. For example, if a refuse container is located far from the access ramps to the vessels, notice whether the mariners actually use it, even if the port or mariners themselves report that they do.

Negotiating Refuse System Options

Often better service, additional service, and lower costs can be negotiated in port meetings with refuse haulers. In Astoria, Oregon, the port meeting with refuse company officials resulted in the willingness of the refuse hauler to back farther out on the dock to service a container. The Petersburg, Alaska, harbor district, under city jurisdiction, is meeting with city officials, who also operate the refuse service, to negotiate charges to account for the dumping of household refuse in port containers and for emptying half-filled dumpsters.

Determining Whether Recycling Will Work

A recycling system is a viable and cost-saving option for ports in which the following conditions exist:

- Ports are convinced that there are substantial benefits to recycling: recycling saves them significant refuse disposal costs, expands their refuse reception capacity, or provides a welcome service to mariners.
- There are operating recycling systems in the area and nearby markets for recycled goods. (If markets are far away, recycling is still a possibility if nonprofit groups can arrange with shipping companies to waive their backhauling charges.)
- Recyclers or community organizations will come to the port to haul away the collected materials without port involvement.

(Ports generally have shown little interest in collecting recyclables if they must also haul them away.)

- Port concerns about recycling can be addressed. Lack of familiarity with recycling, concern about system efficiency, and uncertainty about recycling markets are deterrents to port recycling interest. Ports need to be assured that the materials they accumulate will not be difficult to get rid of. Ports need to be willing to experiment with recycling.
- Recycling provides mariners increased convenience and benefit. Mariners will use recycling areas if they are provided at the point of disposal and if they are clearly signed. It helps if mariners are familiar with a recycling system or concept (e.g., fishermen in Coos Bay, Oregon, were aware of the Newport system; cardboard was already being recycled in the town of Bellingham, Washington).

Anticipating Recycling Potential and Possibilities

Refuse container contents can be examined and mariners queried to determine the types of waste materials generated and the potential for cost savings through recycling. Speaking to mariners will also indicate their level of awareness about recycling and their interest.

Visual examination of refuse container contents from port to port indicates that the amounts of metal, wood, paper, glass, and gear items found are quite variable. However, we have found that most all commercial fishing ports examined to date could realize cost savings by collecting and separating cardboard items for recycling. The recycling of this item is an easy and impressive first step and may stimulate further interest in recycling. (This was Newport's experience as well as that in Coos Bay, Oregon, where the harbormaster noted with much enthusiasm, "It works!")

In most ports serving the larger commercial fishing vessels, the establishment of a central area to both collect and store wood, cable, metal, and net items (for recycling or giving away) can provide a convenience to mariners and also encourage the proper disposal of particularly large items. Ports have often started these areas informally by stacking materials on pallets or in old containers, or by simply making distinct piles of different types of materials and posting signs. All those that have done so are impressed by how quickly additional materials get placed on the piles by mariners. Even hand-painted signs are effective in encouraging the proper sorting of refuse items. Ports that serve a primarily recreational vessel fleet will probably not find a need for such a central area, unless infrequent pickup of recyclable materials necessitates storage.

We have found that wherever there has been an established recycling program in a local area, it is relatively easy for the port to tie into it. Port contact with the recyclers has always resulted in their willingness to cooperate with the ports in establishing a workable recycling system and hauling schedule. Recycling containers need not be elaborate or expensive

and can usually be built by the port or acquired. Refuse or recycling companies often provide free recycling bins, while donations of the bins can often be obtained from local businesses, from a city (e.g., they may have surplus bins), or from restaurants (208.2-L (55-gal) drums). School shop classes may also be willing to fabricate them. Ports may find it awkward to ask for such donations, and such requests may be easier and more effective if they come from a local mariner or a citizen.

Recycling containers must be able to be emptied easily by the recycler and must be clearly designated so as not to accumulate trash. Bright colors and clear signage are essential. The color blue is being used by most west coast ports for their recycling containers, with the idea that coast-wide consistency will make mariner recognition easier.

Seine and trawl nets are in demand by fishermen and the general public and will be removed from an accessible collection area if it is signed.

Though nets are made of materials with recycling potential (nylon, polyethylene, and polypropylene), used-net recycling efforts are still experimental in nature in the United States, and the details of collection, transportation, and market value of the nets are still undetermined.

At least 20 ports on the west coast have begun recycling programs. Table 1 summarizes these efforts.

Beginning a Port Recycling Program

Though ports may not set up a recycling system on their own, most welcome assistance in getting one started. The following steps are usually followed to establish a recycling program:

1. Assess refuse materials generated at the port.
2. Determine which materials are accepted or collected by area recyclers.
3. Explore demand and markets for uncollected but recyclable goods, e.g., fish nets.
4. Work with mariner groups and port to design a system that will be utilized.
5. Order or make recycling signs.
6. Order or make recycling bins or designate reception areas.
7. Implement the recycling plan, placing signs and bins.
8. Inform mariner groups of the bins and encourage their proper use. Inform the media.

Table 1.--West coast port recycling systems.^a (Note: Used oil is also recycled at all these marinas.)

Port and contact	Type of materials collected	Collection methods	Other notes
Newport, OR Bud Shoemaker (503) 265-7758	Cardboard, metal, wood Troll wire Metal, cable, wood, nets	Wood fish bins 208.2 L (55 gal) barrel Reception area, barge	System's 3d year Compounds/screens Refuse volume reduced one-third.
Charleston, OR Don Yost (503) 888-2548	Cardboard, metal, wood Cable, nets	Wood fish bins Reception area	8 h/month labor saves \$200/month due to cardboard reduction alone.
Astoria, OR Bill Cook (503) 325-8279	Cardboard, metal, wood Cable, nets, some plastic	Reception area Reception area	New refuse area Completed December 1989.
Westport, WA Karl Wallin (206) 533-9528	Cardboard, plastic, lines Glass, cable, nets	Wood fish bins Wood fish bins	System in place October 1989 Located dockside and near office, launch ramp, net repair.
Anacortes, WA Dale Fowler (206) 293-0694	Metal, cable, nets, wood	Reception area	Got free advertising of net availability.
Bellingham, WA Art Choat (206) 676-2500	Aluminum, cardboard Wood, metal, nets	Wood fish bins Reception area	Recycling project assisted by Washington Sea Grant.
Friday Harbor, WA Bart Mathews (206) 378-3688	Glass (white, brown, green) Aluminum	Commercial 15.3 m ³ (20 yd ³) recycling bin, provided and hauled by garbage company	Recycling system so well-used had to enlarge and automate system.

Table 1.--Continued.

Port and contact	Type of materials collected	Collection methods	Other notes
Everett, WA Karen Bukis (206) 259-6001	Cardboard, aluminum, mixed paper, glass plastics (in 1991)	208.2-L (55-gal) barrels obtained from yogurt company for \$8 each	Three barrels in each of 13 loca- tions. Started April 1989, in first 10 months recycled 2,270 kg (5,000 lb) paper, 1,816 kg (4,000 lb) glass, 454 kg (1,000 lb) aluminum. Garbage bill reduced \$7,500 from previous year.
Ilwaco, WA Bob Petersen (206) 642-3144	Aluminum cans Nets, cable, wood, metal	Wire mesh cans Reception area	Cans benefit baseball team.
Port Townsend, WA Andrea Fontenot (206) 385-2355	Cans, papers, glass	113.5-L (30-gal) garbage cans	Five roofed "environmental centers" built summer 1989. Part of water- front revitalization plan.
Seattle, WA Marla Kleiven (206) 728-3394	Glass, aluminum, newspaper	208.2-L (55-gal) drums	In 11 cedar-fenced refuse areas at recreational marina.
Greg Money (206) 728-3395	All types recyclables, including plastics, paper	3.1 m ³ (4 yd ³) dumpster	Three locations at commercial fish- ing marina for comingled materials. Port pays to have hauled but less than for trash.
Sequim, WA Jan Hardin (206) 683-9898	Cardboard Glass, aluminum	Wood fish bins Wood fish bins	In place Planned for 1990.
Half Moon Bay, CA Bob McMahon (415) 726-5727	Cardboard, paper, glass Plastic jugs, aluminum	Rubbermaid 2-wheeled 208.2-L (55-gal) carts, transferred weekly to storage bins	Pilot study by Coastal Resources Center. Handbook available December 1990 from (916) 323-3508.

Table 1.--Continued.

Port and contact	Type of materials collected	Collection methods	Other notes
Oakland, CA area ^b Calvin Young (415) 891-3912	Aluminum, glass, plastic beverage bottles	Mini-house structures contain two 208.2-L (55-gal) barrels. PVC tube prevents removal of deposited materials	Materials for each of 27 units cost \$200. Emptied twice/week even in winter. In operation since April 1989.
Kodiak, AK George McCorkle (907) 486-5438	Aluminum, plastics/rubber Wood, metal, nets/rope Paper/cardboard Batteries	3.1 m ³ (4-yd ³) dumpsters Separate collection area	Seven dumpsters in each of two MARPOL stations. Operating since February 1990.

^aOther systems planned: Santa Cruz, CA, Steve Scheiblaue (408) 475-6161; Chula Vista, CA, Becky Clark, 550 Marina Parkway, Chula Vista, CA 92010; San Diego, CA area (a number of marinas); Libby Lucus (619) 235-0281; Edmonds, WA, Bill Stevens (206) 457-4505; Port Angeles, WA, Chuck Faires (206) 457-4505; Crescent City, CA, Rich Taylor (707) 464-6174.

^bSix marinas.

9. Monitor the recycling system to make sure materials are hauled on time and problems are resolved.

The Port Role in Mariner and Community Education

Most ports will play a limited but important role in stimulating mariner awareness and involvement in the marine debris solution. Ports welcome and even solicit help from mariner groups, community groups, and recyclers to inform mariners of the refuse disposal law and to stimulate interest. The ports are willing to make notices about the law available to mariners and, when they are provided, will display posters and distribute stickers and brochures. A limited number of ports, fish processors, and dock facilities actively seek out information to pass on to mariners.

Ports may be willing to meet with their user groups to discuss Annex V regulations, but they are not likely to coordinate or organize such meetings on their own. Likewise, though most ports welcome press coverage about their marine debris efforts and do not object to public service announcements that say "brought to you by your local port," they are not likely to seek out media contacts.

A few interested harbormasters have actively sought support for marine debris cleanup efforts. The harbormaster in Petersburg, Alaska, spent a whole day going from boat to boat to explain the law to mariners and his need to raise moorage rates in anticipation of the extra refuse load. The harbormaster in Astoria, Oregon, showed marine debris promotional materials to a solid waste committee. He has asked for their assistance in contacting mariners on the docks and has inspired them to go into the schools to inform children. In Avila Beach, California, the port manager, a former teacher, is developing a marine debris educational package for school children.

Fishing Industry Support

Our experience shows that support from commercial fishing groups can be expected. Fishing industry groups are interested in solving the marine debris problem and are willing to become involved in marine debris work.

When supplied with information about Annex V, industry groups and Sea Grant marine agents will promote awareness of the issue through newsletters. They will discuss the regulations at fishing group meetings and encourage affirmative action among members. Industry groups have also been willing to promote marine debris awareness at expositions by handing out materials, displaying posters, talking with mariners, and by making display space available for marine debris pictures.

When prompted or during a meeting, fishermen as well as Sea Grant agents are willing to talk to their port harbormasters about refuse disposal needs and ideas. Many individual fishermen have personal interest in this issue, and their support is an essential factor influencing the attitudes of others, promoting peer action, and pushing for port changes. Some fishermen, when provided information and promotional items may also

act on their own to bring this material to the attention of their peers, other mariner groups, and the schools.

Taking Action

Action related to the marine debris problem is unlikely unless someone takes charge. Someone must define and plan activities that address the marine debris problem, must organize and facilitate meetings, and must delegate the work. Do not assume that a port employee will take on or be assigned the lead role. If that level of involvement is not likely from the port, look for other individuals and groups for the port to work with.

Groups that have already been organized around beach cleanups or recycling are logical groups with which to work. They may be able to research or put together a port recycling program, coordinate a special promotional event, conduct mariner surveys, or organize a harbor cleanup or educational campaign. They are often willing to distribute information and show readymade video tape or slide presentations. The reference guide resulting from the Newport pilot project outlines such an outreach program, and emphasizes approaches such as involvement of port employees and mariners in program design and trouble shooting. This involvement has been effective in assuring the support of a marine debris program.

Groups already taking part in boater education efforts (e.g., the U.S. Power Squadrons and Coast Guard Auxiliary) can also provide support. They may be willing to incorporate marine debris information into their classes and during their contacts with mariners.

Other groups which may be interested in marine debris cleanup and recycling activities are environmental groups, community, senior, and scouting groups, and school science classes. They are most likely to act when they have readymade materials to give out or use. Interest and action are encouraged simply by making brochures, stickers, posters, curriculum materials, photo displays, and slide or video programs available to groups and teachers.

CONCLUSION

The technical part of dealing with marine refuse in a port is usually simple. Refuse containers are readily available in various sizes, hauling schedules or container sizes can be adjusted to meet increased demand, and recycling can often be used to decrease refuse disposal costs.

What is difficult is the initiation of action. What has been most apparent in work with the eight target ports, and with other ports on the west coast, is that while there is general support and interest by the port and mariner groups in the marine debris problem, it is not a high priority action item.

Most ports and mariner groups are not likely to take action on their own. However, if ports are approached, offered assistance, and encouraged (even by providing a simple catalyst such as information, notices, posters,

or brochures) much can be accomplished. All ports have been willing to attend and even help organize meetings which bring them together with mariners, refuse haulers, and others. Once port officials are stimulated to begin (by the development of personal interest, by gaining the tangible assistance of supportive mariners or community groups, by guilt, or by seeing the examples set by other ports) we found that many then initiate activities on their own.

This essential outside push can be provided by a mariner, citizen, recycler, or community group willing to do some research regarding refuse and recycling options, organize meetings or activities, and encourage the awareness of mariners. States, counties, and cities may be able to designate part of an employee's time for such port assistance through their solid waste, environmental quality, boating, or fishing departments. If such a person can organize the readily available assistance of the fisheries groups and the ports, tangible progress on the marine debris problem will soon be noted.

Further information on the NMFS-sponsored pilot port marine debris project is available in two reports: "Report on a port-based project to reduce marine debris" and "Dealing with Annex V--reference guide for ports." The former report describes the project in detail. The latter presents guidelines and resources that resulted from the pilot project. A video tape about the Newport project called "A marine refuse disposal project" has also been completed. All three resources are available from NOAA's Marine Debris Information Office, 1725 DeSales Street, N.W., Suite 500, Washington, D.C., 20036, U.S.A., phone (202) 429-5609.

DISPOSITION AND RECYCLING OF PLASTIC PRODUCTS INCLUDING USED NETS

Masahito Aizawa and Akinori Satou
Oceanography Office
N.S. Environmental Science Consultant Co., Ltd.
2-9-14, Toranomom, Minato-ku, Tokyo, 105 Japan

ABSTRACT

Disposition and recycling technologies of various plastic wastes are reviewed. The technologies employed vary by material and type of waste.

Case studies were carried out at Akkeshi and Wakkanai ports in Hokkaido, which are major base ports of Japanese salmon fishing vessels. Some used nets are recycled into nylon pellets; others are incinerated or landfilled.

BACKGROUND

Recycling technologies in Japan have been developed because Japan is poor in some natural resources. Japan must import most raw materials needed to produce industrial goods, and there exists national recognition of the necessity to make the utmost of raw materials through recycling and other means.

Japan is also a small and densely populated country. It is not easy to find places for waste disposal, and the environment tends to be susceptible to damage from waste disposal practices. Thus, Japan finds it necessary to develop effective recycling technologies. To date, effective recycling technologies have been developed mostly by small- and medium-scale enterprises, rather than by public research institutes. Thus, these enterprises tend to be small in scale and of great variety.

There are two principal methods of recycling used fishing nets-- pelletization and reutilization. Both methods have problems. Pelletization as a recycling business requires profit stabilization, while reutilization of nets requires further development of recycling methods in order to increase demand.

PLASTIC CONSUMPTION IN JAPAN

The output of primary products of plastics in Japan reached 4.77 million tons (MT) in 1987, an increase of about 12% over the 4-year period

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

from 1984. Responsible for this increased production is the continuous development of new plastic products, progressive development of synthetic technologies of resins, and the easy-to-process nature and low prices of plastics.

The lifespan of plastic products from marketing to disposal varies according to the resins and their products. The life of packaging materials and disposable containers, which together account for some 40% of plastic products, averages 1 to 2 years; on the other hand, sundry goods for household use and toys last for 3 to 5 years. As a result, 55% of plastic products are discarded within 5 years after their production. Pipes and other plastic products used in construction last for more than 10 years, but they account for only 20% of the total.

The most notable characteristics of plastic products are durability and low prices. The low prices, however, have brought about the "throw-away" practice; thus, the useful life of the products themselves is much shorter than that of plastic as a substance. The present problem is that the disposal of plastic is difficult, the difficulty stemming from its high durability.

CASE STUDIES OF RECYCLING WASTE PLASTICS

Introduced here are some Japanese technologies for recycling waste plastics. In recycling waste plastics, effective classification and separation of various plastics are essential to maintain good quality of recycled products and to lower production costs.

Plastic products are being used in almost every field, often in combination with wood, steel, or other materials. As a result, discarded plastics are mixed in with other waste material. Moreover, even if plastics are grouped by kinds at the time of refuse collection, they are usually covered with dust, earth, or sand, in which case additional processing is required.

Various techniques for classifying municipal refuse have been developed. They include techniques developed under the "Stardust '80 Project" conducted by the Agency of Industrial Science and Technology of the Ministry of International Trade and Industry in 1972-82 with the aim of turning such refuse into new resources.

There are two ways to recycle waste plastics--one is using it in new plastic products and the other is reducing the plastic as a material. Although in both cases quality control is a difficult task, this has developed into a new business in which 88 companies are already engaged throughout Japan. Recycled plastic products include manhole covers, piles, and flowerpots. The Ministry of International Trade and Industry has established the Japan Industrial Standard as guidelines for uses and quality control of such recycled products.

The pyrolytical method converts organic substances of high molecular weight into those of low molecular weight by use of heat energy. Suitable

substances for this process are waste plastics, used tires, and waste oils. As opposed to incineration, pyrolysis needs not only strictly controlled operations, but must also meet two conditions:

- Waste plastics as materials must be even in quality and available steadily and in large quantity.
- A reliable supply system of recycled products like oil and gas must be established.

TECHNOLOGY FOR SORTING MUNICIPAL WASTES (STARDUST '80 PROJECT)

This system is designed to sort municipal refuse into two or three groups by applying both the pulverizing and sorting processes simultaneously. These processes utilize the principle that different materials can be crushed into different sizes in accordance with their resistance to shock, compression, and shearing stress.

One such group is composed of kitchen garbage, fragile paper, and other goods which can be easily reduced to small pieces and passed through small screen holes. Another group is made up of plastics, metals, fibers, stronger paper products (cardboard paper and laminated paper), and other things which cannot be broken easily and are extracted from the machine almost in their original sizes. When compared with conventional pulverizing machines, this system has the following advantages:

- Pulverization and sorting are possible in one process with a small amount of power.
- As metals, plastics, and other items can be selectively broken almost in their original forms, the system faces less wear and therefore entails lower maintenance costs.
- Owing to its slow speed, the device produces less noise, vibration, and dust.
- A special anticoiling device prevents long objects from wrapping around the rotating shaft.
- It is possible to easily discharge overloading objects such as metal blocks, bulk waste, etc., out of the system through adjustment of its rotating speed.

TECHNOLOGY FOR SORTING AGRICULTURAL FILM

This system is capable of sorting and utilizing soil-smearred agricultural films without resorting to a washing process.

Films are first roughly crushed (about 40 mm) and then sent to a heavy, oil-fired stirring dryer where the material is dried with 110°C hot air. Soil and sand are loosened from the films and discharged from the

screen at the bottom of the drying chamber. Iron elements are eliminated by the use of magnets. The films are further crushed and once again dried (air drying) so that their quality is improved. After this, iron pieces are again removed by magnet and the films recovered in the form of fluff.

Recovery costs, excluding those of pelletization, amount to about ¥37.5/kg.

TECHNOLOGY FOR RECYCLING FOAMED STYROL FISH BOXES

Foamed styrol is used to make fish boxes and package cushioning, the latter used extensively in business offices and households.

Fish boxes are first used in fishing ports and later accumulated in markets and retail stores. Of the fish boxes handled in central wholesale markets throughout Japan in 1979, 42% were incinerated and about 22% were recycled.

Fish boxes to be recycled must be relatively clean. After being cut, they are heated and melted at 270°-280°C and molded into blocks weighing 8 kg each. These plastic blocks are exported through chemical companies to Southeast Asia, where they are used as raw materials for toys and re-exported to Japan, the United States, and elsewhere.

Such boxes were initially sold at ¥25/kg but their current price is ¥5/kg; the decline resulting from the price competition with virgin materials produced at low cost reflecting the current stagnant oil prices.

RECYCLING WASTE PLASTICS INTO ARTIFICIAL FISH REEFS

In Japan, the establishment of artificial habitats for fish has been promoted under a long-term national project designed to increase fish resources along the coasts.

Man-made waste plastics fish reefs are made of recycled polyethylene, polypropylene, and other polyolefin plastics derived from the plastic films used for packaging foods and other purposes.

In order to strengthen artificial fish reefs, various measures are taken, such as mixing ferro-oxide into the melted plastic to substantially increase density and building structural-use steel pipes into props. Between 1972 and 1988, a total of 250,000 m³ of fish reefs weighing about 18,000 MT, were installed on the seabeds around the Japanese coasts. These artificial reefs have the following characteristics:

- There is no limit to their durability.
- As they are made of assembling units, it is easy to adjust their shapes and sizes.
- Easy to assemble, the time needed for installation is short.

- As they do not exude alkali elements, a variety of seaweed and shellfish can safely attach to the artificial reef.

TECHNOLOGY FOR RECYCLING FISHING BOATS OF FIBER-REINFORCED PLASTICS

About 1965, fiber-reinforced plastics (FRP) began to be used for building fishing boats. Easy to mold, lightweight, durable, and economical, FRP fishing boats have been used widely in the coastal fisheries. In Japan the number of powered FRP boats reached 290,000 in 1984. The earliest of these vessels have reached "retirement age," and safe disposal is now a major problem.

In 1985 and 1986, the Fisheries Agency developed an FRP boat disposal system. The disposal procedure is discussed below.

Chosen for the experiment was an FRP vessel from the set net fishery; the vessel weighed 3.42 MT and measured $9.86 \times 2.6 \times 0.59$ m. The vessel was first crushed by a hammer crusher into 10 to 20 cm³ pieces. These pieces were carbonized in a batch-type pyrolysis incinerator which measured 3.4 cm³. Waste tires were used as fuel for the carbonization process.

The plastics decomposed in the pyrolysis were gasified and recovered in the form of oils, which were further separated into water, tar, pitch, and waxes and stored as reprocessed oils. Glass fiber residue was burned in the pyrolysis furnace.

Recovered were the following substances:

- Oils--For reasons of smell, low calories, and irregular viscosity, the recovered oils were traded as products below C-class fuel oil (heavy oil).
- Glass fibers--Recovered glass fibers were utilized as composite plastics for compound plastics called AMC or FMC. Pending further research, they may be usable as molded products for automobiles. Their utilization as powdered glass may also be possible.
- Carbons--They are a mixture of products from FRP and waste tires. It is possible to market those sifted through an 80-mesh screen.

TECHNOLOGIES FOR RECYCLING AND INCINERATING USED FISHING NETS

Used fishing nets can be utilized in their original form for applications other than fishing or they can be reprocessed into basic material and used to produce other things.

Recycling of Used Fishing Nets in Their Original Forms

There is a wide range of purposes for used fishing nets other than fishing. These include:

Collection of Planktonic Larvae of Scallops

When used for catching planktonic larvae of scallops, one-sixth of a unit (about 100 m/unit) of nylon monofilament driftnet used in the salmon fishery is packed in one onion bag. Salmon driftnets are used and not squid driftnets; the latter are unsuitable because they are smeared with squid ink. Scallop fishermen buy used nets from net-weaving companies at a price of about ¥1,600 per unit (including shipping costs) in Hakodate, Hokkaido.

Protection of Trees From Deer

In 1986, in order to protect trees in the training plantation of Kyoto University in Shibecha, Hokkaido, a reforested area within the plantation was surrounded with nets 1.4 m high that had been used previously for salmon fishing. Before such protection, 79% of the 86 trees in the area had been damaged by deer, *Cervus nippon esoensis*. No further damage has been reported since the installation of the nets.

Although such protective nets can be considered effective for preventing deer from harming trees in Hokkaido, this method is costly and labor intensive. In Chiba Prefecture located east of Tokyo, a deer was reported to have died when its antlers became entangled in a net.

Drying Tangle

In Japan, tangle is usually dried on pebbles after it is gathered from the sea. In some regions, trawl nets are used instead as drying racks.

Antibird Nets at Garbage Collection Sites

In Wakkanai, Hokkaido, used trawl nets cut into 2-m squares are spread over garbage cans at community garbage collection sites so that crows cannot scatter the garbage.

Reprocessing of Used Fishing Nets

Since 1988, three factories (two in Akkeshi and one in Hakodate) have been in operation pelletizing used nylon monofilament salmon and squid driftnets. The companies collect nets from various ports in Hokkaido. The pelletizing process is similar at all the companies and total production is an estimated 700 MT a year. The pellets are sold to chemical companies at a price of about ¥210/kg (1988) for the manufacture of electrical equipment for automobiles, bicycle saddles, electric fans, and other kinds of goods. When compared to virgin pellets, however, the tensile strength of recycled pellets is reported to be weaker by 10%.

One of the problems concerning pelletization lies in the great effort required in washing nets or otherwise removing impurities from them. Trawl nets are more likely to retain soil and trash at their webbings due to the method of fishing. As far as we are informed, there is practically no reprocessing of trawl nets at this time, presumably due to the cost of preparation.

Economically, the competitive advantage of reprocessed pellets over virgin pellets is very unstable; the price of the latter being subject to the price of petroleum.

Incineration of Used Fishing Nets

In Japan, as a means of reducing the volume of bulky fishing nets, high heat incineration, which is effective in minimizing harmful substances and ash, has been under study. Under consideration are batch-type pyrolysis furnaces (0.7 to 50 m³ in size) placed underground and designed to attain complete combustion through use of water-gas reactions.

This furnace is constructed so that combustion may take place using an air intake from above. This also makes it possible to adjust the supply of air so as to prevent unburned soot and dust from being discharged. The chimney flue helps combustion by serving as a secondary furnace.

When burned, plastics generally give off black smoke and, in some cases, harmful gases. No noxious smell or black smoke is produced, however, in this particular type of incinerator, even if used fishing nets, tires, foamed styrol, or lumps of plastics are burned.

RECYCLING OF USED FISHING NETS IN HOKKAIDO

Information regarding recycling of used fishing nets in Hokkaido was obtained from oral surveys conducted among staffs at government offices, fishing operators, net-weaving companies, and other related industries in Akkeshi and Wakkanai in 1988. Akkeshi is one of the five main ports in Hokkaido engaged in medium-sized salmon driftnet fishing operations. It is also known as a port for the squid driftnet fishery. Wakkanai, on the other hand, is a port mainly supporting a trawl fishery.

Akkeshi

Use of Fishing Nets

As of 1987, fishing operators in Akkeshi owned 13 fishing boats authorized for use in salmon and trout fishing with medium-sized driftnets. Six of them were also engaged in driftnet fishing for squid. In addition, 10 other boats from other ports landed their catch at Akkeshi; these vessels were not involved with the disposition of fishing nets.

Medium-sized salmon driftnets in use at Akkeshi mostly measure about 100 m in length. When the salmon and trout fishing season using medium-sized nets comes to an end around July, some nets are used again for squid

catching and then scrapped. Nets aboard boats used exclusively for salmon and trout fishing are transported to net-weaving companies. About 50% of the nets need no repairs and are used again the following season; 30 to 40% of the remaining nets are repaired and used again. About 20% of the salmon and trout fishing nets are scrapped each year, and almost all nets used to catch both salmon and squid are scrapped after the cuttlefish season ends. The lifespan of nets averages from 2.5 to 3 years.

Disposal and Recycling of Fishing Nets

For lack of landfill space and to avoid pollution, Akkeshi Town does not incinerate used fishing nets or discard them in landfills, although it discards foamed plastics like fish boxes as "incombustible refuse." Fishing operators burn their ropes and buoys separately.

In Akkeshi, there are two companies that reprocess nylon monofilament salmon and squid driftnets into pellets. Used nets are either brought to the processors by fishing operators and net-weaving companies or gathered without charge at port collection sites. Nets purchased from net makers and repairmen cost the processors ¥20/kg. It is estimated that about 600 to 900 MT of used nets are collected within and from outside Akkeshi annually. Among them, those unfit for recycling are incinerated by the reprocessors. The rest are melted and pelletized into an estimated 450 MT at the two reprocessing companies. Before 1987, some used fishing nets were exported through trading firms to China and Taiwan to be made into valves and other products. Such exports are not carried out today because of the unfavorable prices of petroleum.

One of the two pellet processors in Akkeshi is capable of producing 1.5 MT a day in a 9-h operation, but such facilities are in operation for only about one-third of the year. The company spent ¥14 million on a washing machine and another ¥13.5 million on a machine for melting and pelletizing plastics, with the total investments estimated to be some ¥60 million, including the land and building. The monthly production of 15 MT of pellets requires ¥740,000, including ¥200,000 for fuel and electricity and ¥520,000 for 100 man-days of labor.

Wakkanai

Use of Fishing Nets

As of 1987, Wakkanai had 19 boats which were allowed to fish in the offshore trawl fishery. On an average there are about 10 fishing nets for each boat; usually 2 or 3 nets are loaded on the boats, and also on board are 1 or 2 nets to be repaired. The remainder of the net supply is stored in warehouses or other locations of the fishing operators or net-weaving companies.

Damaged nets are repaired aboard the fishing boats when the damage is slight. Major repairs are made either in the fishing operators' own net-weaving shops or shops of independent net-weaving companies. Repairs usually become necessary after six or seven fishing operations; repairs

mostly concern the dislocation of rope from net mesh. Net webbing usually lasts for 2 years. Torn webbing is usually not mended, but replaced with a new section of webbing. Fishing operators spend about ¥10 million each year for net-repairing purposes.

Scrapping of Fishing Nets

Scraps of trawl webbing are accumulated during the process of repairing by net-weaving companies and fishing operators. About 15 to 18 MT of scrapped webbing is accumulated by net-weaving companies; the total excludes webbing used for heating purposes during winter. The amount of scrapped webbing accumulated by fishing operators is estimated at 17 MT per year.

Disposal and Recycling of Fishing Nets

Scrapped nets not used for heating or recycling are brought to garbage disposal plants in Wakkanai. An estimated 700 MT of fishing materials are brought to landfills annually, accounting for 1.6% of the municipal total in 1987. Of the 700 MT, nets made up 80% and ropes, 20%.


Some of the scrapped nets are incinerated at establishments of net makers and fishing operators. Here, stoves used as heaters during winter burn nets exclusively (or together with waste oil). This is probably the most popular method of used trawl net recycling practiced in Wakkanai.

Between 1982 and 1987, a producer of animal feed in Wakkanai used scrapped trawl nets as an auxiliary fuel in its coal boiler. To operate the boiler the company burned 30 used 12-MT truck tires every day and burned 2 to 3 MT of scrapped fishing nets every year. The feed producer obtained these nets free of charge at net-repairing shops.


ACKNOWLEDGMENTS

We express our deepest thanks to Shuichi Takehama and Nobuyuki Yagi, assistant director and officer, respectively, of the Ground Environment Conservation Division, the Fisheries Agency, Government of Japan, and to others who have cooperated in preparing this report.

SESSION VI



SOLUTIONS THROUGH LAW AND POLICY



STATUS OF THE U.S. ENVIRONMENTAL PROTECTION AGENCY
MARINE DEBRIS ACTIVITIES AND PROGRAMS

David P. Redford
Office of Marine and Estuarine Protection
U.S. Environmental Protection Agency
Washington, D.C. 20460, U.S.A.

ABSTRACT

The U.S. Environmental Protection Agency (EPA) issued on 18 January 1989 an Interim National Coastal and Marine Policy (NCMP), which addresses marine debris as one aspect of a number of marine degradation problems. This policy states that one of EPA's goals is the recovery of full use of shores, beaches, and water by reducing sources of plastics, floatables, and debris.

The EPA's objectives include:

1. To control disposal of medical wastes and aggressively enforce laws to protect the public from exposure to them.
2. To accelerate efforts to identify the sources of floatables, debris, and plastics.
3. To control these materials through new technologies and substitutes, permitting, and enforcement, as well as aggressive source reduction and waste minimization programs.

This paper summarizes the status and progress of many EPA activities and programs aimed at addressing and implementing the marine debris aspects of the NCMP.

DISCUSSION

Plastic debris in the marine environment has been shown to kill or harm marine life, damage vessels, and cause aesthetic and economic damage to beach communities.

The Final Report of the Interagency Task Force on Persistent Marine Debris (1988) described the potential sources of marine debris and separated these sources into two categories: ocean sources and land-based sources. Ocean sources include commercial fishing vessels, recreational

boating, merchant vessels, cruise ships, military and research vessels, and offshore oil rigs and supply vessels. Land-based sources include plastic manufacturing and processing activities, combined sewer overflows (CSO's) and sewage treatment plants, solid waste management practices, and litter. Debris from these sources washes up on U.S. beaches and clogs our waterways. Last summer's problems on the northeastern U.S. beaches show the large-scale impact that marine debris can have.

On 18 January 1989, the U.S. Environmental Protection Agency (EPA) issued an interim National Coastal and Marine Policy (EPA 1989b), which included the agency's position on marine debris.

The EPA is taking action against many of the sources of marine debris described above and the problems they pose. These actions include

- assessment of the items that make up marine debris and their sources,
- assessment of the specific items released to the environment from CSO's and storm sewers,
- medical waste tracking,
- enforcement,
- public education,
- cleanup plans.

Some of EPA's activities are described below in relation to the authority under which they are being implemented.

MARINE PLASTIC POLLUTION RESEARCH AND CONTROL ACT OF 1987

The Marine Plastic Pollution Research and Control Act (P.L. 100-220) (MPPRCA) contains several sections pertaining to plastics and marine debris. Among these are:

1. The development of regulations by the U.S. Coast Guard implementing the requirements of MARPOL Annex V. These regulations set forth the restrictions for disposal of garbage from ships and include prohibition of the disposal of plastics.
2. An assessment of the effects of plastic materials on the marine environment by the Secretary of Commerce.
3. The development of a joint public education program by EPA, the U.S. National Oceanic and Atmospheric Administration (NOAA), and the Secretary of Transportation.

4. An assessment of the problems associated with plastic debris in the New York Bight by the EPA.
5. An assessment by the EPA of methods to reduce plastic pollution.

Activities being carried on by EPA under the requirements of MPPRCA are described below.

New York Bight Report to Congress

The EPA has prepared a report to Congress on plastics in the New York Bight (EPA 1989d), describing the types and sources of floatable debris, the fate of floatable material, the recreational activities, control programs, and recommendations for research, monitoring, and control. The recommendations presented in the report include:

1. Monitoring and surveillance
 - Monitor sources.
 - Monitor beaches and waterways.
2. Local control measures
 - Improve combined sewer overflows.
 - Increase street cleaning.
 - Remove debris from beaches and waterways.
 - Encourage volunteer cleanup activities.
 - Develop public awareness and education campaigns.
3. Research
 - Study the rates of plastic degradation.
 - Study degradable plastics and potential applications.
 - Develop cost-effective recycling approaches.

Public Awareness and Citizen Monitoring

The EPA is working with the NOAA, the Coast Guard, and other agencies in developing citizen monitoring and cleanup patrols and a public awareness program on marine debris. To date, this has included the financial support for this conference, the Second International Conference on Marine Debris held in Hawaii, financial support for the 1988 beach cleanups conducted during Coastweeks by the Center for Environmental Education (now the Center for Marine Conservation (CMC)), and sponsorship of an upcoming informal

roundtable meeting on marine debris. The EPA will be using the results of the Hawaii conference and the roundtable meeting to plan future national-level public education activities.

These future activities will also be developed using the results of the Washington State Marine Debris Task Force studies, where EPA is working with the State of Washington and many Federal, state, and local groups to combat local marine debris problems. This task force, which was set up by Washington State on its own initiative, has pulled together Federal, state, and local interests and published a report (Marine Plastic Debris Task Force 1988). The report makes a variety of recommendations for controlling, monitoring, and removing marine debris. The recommendations already implemented, such as a state information coordinator, derelict fish net removal, and a citizen monitoring program, will be evaluated by EPA as potential activities to be included in the national program to be coordinated with other Federal agencies.

New York Bight Restoration Plan

A New York Harbor Floatables Action Plan (EPA 1989a) was developed as part of the New York Bight Restoration Plan. This floatables action plan describes the routine and responsive monitoring and cleanup activities that will occur in New York Harbor to remove floating debris that may cause environmental, aesthetic, and economic damage. The plan includes:

1. routine monitoring and cleanup of debris (particularly during times of extreme high tides or major storm events) using vessels, helicopters, and planes to monitor and specially designed vessels to cleanup debris;
2. development of a reporting and action network to report sightings of debris;
3. a contingency plan using the reporting network to activate the cleanup vessels to respond to reports of debris sightings.

Methods to Reduce Plastic Pollution

The EPA is preparing a report, due to Congress in the summer of 1989, which describes:

- the production of and major markets for plastics in the United States;
- the generation of plastic waste in the United States;
- the impacts of plastics on the management of solid wastes in the United States including transportation, disposal in landfills, and incineration;

- the sources, fates, and effects of plastics in the marine environment;
- an evaluation of potential solutions such as source reduction, use of degradable plastics, recycling, and substitution; and
- recommendations for action.

In assessing the potential sources and fates of plastics for this report to Congress, EPA has funded several field investigations. One such study was the 1988 CMC beach cleanups previously mentioned. These data will help determine the distribution of plastic articles on our nation's beaches and will be useful in determining which items are most prevalent and where EPA must focus more effort. The 1988 data and data from previous beach cleanups will be a useful baseline in estimating the effect of the U.S. regulations to implement MARPOL Annex V and other mitigation efforts. Future beach surveys will be used in an attempt to indicate trends and possibly show the effectiveness of the new controls on shipboard sources of plastic debris. It may still be difficult, however, to determine which plastic articles found on beaches come from shore-based sources and which come from vessels.

To determine the variability of plastic items in inshore waters and to assess the harbors as potential sources of debris in other coastal areas, the EPA has also conducted field investigations of the floating debris in several U.S. harbors through a contract to Battelle Ocean Sciences of Duxbury, Massachusetts (December 1988-February 1989). Harbors surveyed include Boston, New York, Philadelphia, Baltimore, Miami, Seattle, Tacoma, San Francisco, and Oakland. These surveys were conducted by towing surface nets through slicks of floating debris and then sorting, identifying, and counting the contents. Plastic items 0.3 mm and larger were studied, yielding information on not only the larger plastic items readily observed on beaches but also on the smaller plastic pellets and pieces which are not so obvious to a casual observer. The final EPA report of these surveys is currently being prepared. A summary of preliminary results is presented by Trulli et al. (1990). The preliminary findings of these surveys indicates that the presence of certain plastic items can be directly linked to the presence of CSO's where storm water and street runoff combine with sewage at times of heavy rain and are discharged directly into receiving water.

The apparent correlation between certain debris items and the presence of CSO's is the basis for a new study, being designed by Battelle Ocean Sciences, where the materials released from several of the CSO's in two cities, Philadelphia and Boston, will be identified and quantified. To do this, EPA will be placing nets around the outfalls from selected CSO's in these cities (May-June 1989) to collect the materials exiting the system during dry weather and during rainy conditions. The results of this study will directly address CSO's as a source of debris, and will be used by the agency in determining the appropriate actions to be taken under the Clean Water Act. Storm drains will also be sampled during this study to determine their contribution to marine debris.

LITTER FROM VESSEL TRANSPORT OF SOLID WASTE

The Shore Protection Act (P.L. 100-688, Sections 4001-4204) provides for controls on operations relating to the vessel transport of certain solid wastes (i.e., municipal or commercial waste) so that these wastes are not deposited in coastal waters.

The EPA is developing guidance with the Coast Guard to minimize deposition of solid wastes into coastal waters during loading, transporting, or unloading. A permit and enforcement program is being developed by the U.S. Department of Transportation such that all vessels transporting solid wastes would require a permit from the Coast Guard. It is estimated that about 400 vessels will need permits--about 100 in the New York Harbor area and 300 in the Gulf of Mexico, plus a few in other locations.

Under this act, EPA is also preparing a report to Congress describing the need and effectiveness of a tracking system for vessels transporting wastes in U.S. waters. Such a tracking system could be used to monitor the movement of wastes in U.S. waters and provide the agency with a mechanism for assuring that wastes are not illegally discharged.

In New York Harbor, an area where large volumes of trash are transported by barges, the State and City of New York have strengthened the requirements for trash barges. The implementation of these requirements is expected to minimize the loss of trash during the marine transport process. The new requirements include:

- limitations on load heights in barges;
- the placement of booms around marine transfer facilities;
- the use of scavenger vessels at marine transfer facilities to collect trash which falls into the water;
- the use of covers over barges to keep the wind from blowing trash off barges.

MEDICAL WASTE--TRACKING AND DISPOSAL

The Medical Waste Tracking Act of 1988 (P.L. 100-582) requires EPA to develop a 2-year demonstration program for the tracking of medical wastes. This program would track the generation and movement of medical wastes from "cradle to grave."

The EPA has prepared an Interim Final Regulation (EPA 1989e) which describes a tracking system to be used by the states participating in the demonstration program. Medical wastes from generators of 22.7 kg (50 lb) or greater per month would have to track the movement of their wastes to the point of final disposition. Generators of less than 22.7 kg (<50 lb) per month have the same requirements but need to keep logs for their medical wastes rather than initiate tracking forms. It is anticipated that

this tracking system will allow EPA to monitor the movement of medical wastes from the point of generation to final disposal and will assure that the wastes are not disposed of illegally where they may endanger public health or degrade the marine environment.

The U.S. Public Vessel Medical Waste Anti-Dumping Act of 1988 (P.L. 100-688 Sections 3101-3105) requires that all public vessels have a management plan for medical wastes on board ship and prohibits the disposal of these wastes at sea except in times of national emergency. The EPA is distributing guidance to all EPA programs that operate boats and ships. The guidance states EPA's policy that all medical wastes generated on board an EPA vessel will be stored in a secure area on board for disposal at an appropriate shore-based facility upon return from sea. No medical wastes can be thrown overboard. Other Federal agencies are also required by the act to issue similar guidance.

COMBINED SEWER OVERFLOWS AND STORM SEWERS

The Clean Water Act (33 U.S.C. 1251 et seq.) requires EPA to regulate discharges from municipal and industrial outfalls into U.S. waters. Under this authority, the agency recently developed a strategy for enforcing the provisions of the act regarding CSO's (EPA 1989c), which are significant sources of plastic street debris entering the marine environment.

The strategy requires that all CSO's be identified and categorized according to their status of compliance with technology and water quality-based regulations. There are about 1,200 combined sewer systems in the United States serving an estimated population of 43 million. States will be required to develop a state-wide strategy for the development and implementation of measures to reduce pollutant discharges from CSO's.

The EPA strategy sets forth three objectives:

1. to ensure that all CSO discharges occur only as a result of wet weather;
2. to bring all wet weather CSO discharge points into compliance with the technology-based requirements of the Clean Water Act and applicable state water quality standards; and
3. to minimize water quality, aquatic biota, and human health impacts from wet weather overflows that do occur.

The strategy confirms that CSO's are point sources independent of the treatment works and reaffirms that both technology-based and water quality-based requirements apply to CSO's. The strategy emphasizes that CSO point sources which are discharging without a permit are unlawful and must be issued permits or eliminated.

The agency has also proposed regulations which will describe the effluent requirements for storm sewers (EPA 1988). By controlling

the effluent from CSO's and storm drains, the agency hopes to significantly curtail the discharge of sewage and sewage-related plastics and street litter into the marine environment at times of heavy rain. To implement the types of controls available for long-term solutions will require major resource expenditures.

ENFORCEMENT ACTIONS

The Ocean Dumping Regulations (40 CFR parts 220-229), which implement a section of the Marine Protection, Research, and Sanctuaries Act (MPRSA) (33 U.S.C. 1401 et seq.), were promulgated in 1977. These regulations prohibit the transport for the purpose of dumping into the ocean of any "persistent inert synthetic or natural materials which may float or remain in suspension in the ocean in such a manner that they may interfere materially with fishing, navigation, or other legitimate uses of the ocean." Activities involving transport for the purpose of disposal at sea are regulated under this act, and permits granted by the agency prohibit the dumping of floatable plastics. Any activity involving the transport of floatable plastics or debris out to sea for the purpose of dumping is illegal, and subject to the fines and penalties described under the act.

Under this authority, EPA has initiated enforcement actions against Nassau County and the City of Long Beach, in New York, and National Seatrade, Inc., for the disposal of sludge containing plastics at the 170.5-km (106-mi) sewage sludge disposal site.

The agency is seeking a total of \$100,000 in fines for these violations of the MPRSA. The county, city, and National Seatrade, which barges treated sewage sludge, are charged with one violation which occurred in September 1988 when EPA staff observed floatables being dumped along with sewage sludge from the county and city at the 170.5-km (106-mi) site. The proposed civil penalty for this violation is \$50,000. The city is also charged in a second count with sending floatables along with its sludge to be ocean dumped in December 1988. The proposed penalty for this violation is \$50,000.

CONCLUSION

The EPA is taking steps to identify and control sources of marine debris, to cleanup existing marine debris, and to involve and educate the public through cooperation with other Federal, state, and local groups. Through a combined effort, EPA hopes to greatly reduce the volume of marine debris generated and its impacts. The EPA believes that one of the primary steps in reaching this goal is to control nondegradable debris before it becomes marine debris--that is, to control it at its source. This should include source reduction and recycling as well as general pollution prevention. Wastes must be managed correctly not only by transport and disposal companies but also by the consumer. Litter thrown into the streets or directly into waterways does have an impact on the marine environment. Plastic items flushed down toilets can, in some cities, end up in the ocean through combined sewer overflows or treatment plant shutdowns. The actions being taken today by EPA as well as other Federal, state, and local

agencies will have a beneficial impact on the environment. These actions must be supplemented, however, by consumers. Proper handling of discarded items by consumers is essential for the trash and sewage collection systems to work. It is the responsibility of all Federal, state, and local agencies to get this message out and build a national/world environmental ethic, or we will be forever cleaning up our environment rather than enjoying and benefiting from it.

REFERENCES

- Interagency Task Force on Persistent Marine Debris.
1988. Final report. Interagency Task Force on Persistent Marine Debris, 170 p.
- Marine Plastic Debris Task Force.
1988. Marine plastic debris action plan for Washington State. Marine Plastic Debris Task Force, Washington State Department of Natural Resources, 2d ed., December 1988, var pag.
- Trulli, W. R., H. K. Trulli, and D. P. Redford.
1990. Characterization of marine debris in selected harbors of the United States. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]
- U.S. Environmental Protection Agency.
1988. National pollution discharge elimination system permit application regulations for storm water discharges; proposed regulations. 53 FR 49416 (Dec. 7, 1988).
- 1989a. Action plan for addressing floatable debris in the New York Bight. Prepared by Battelle Ocean Sciences for Environmental Protection Agency Region II, 15 p.
- 1989b. Interim coastal and marine policy. U.S. Environmental Protection Agency, Wash., D.C.
- 1989c. Interim final national combined sewer overflow control strategy. U.S. Environmental Protection Agency, Wash. D.C.
- 1989d. Report to Congress on the New York Bight Plastics Study, Section 2302, Subtitle C of Marine Plastic Pollution, Research and Control Act of 1987, Title II of the United States-Japan Fishery Agreement Approval Act of 1987, P.L. 100-220, 25 p.
- 1989e. Standards for the tracking and management of medical waste; interim final rule and request for comments. 40 CFR parts 22 and 259.

INTERNATIONAL REGULATIONS FOR THE PREVENTION AND
CONTROL OF POLLUTION BY DEBRIS FROM SHIPS

David T. Edwards and Edward Rymarz
International Maritime Organization
London SE1 7SR, United Kingdom

(The views expressed herein are those of the authors and are not necessarily those of the International Maritime Organization.)

ABSTRACT

The amount of solid wastes entering the world's oceans each year is estimated to be in millions of metric tons. The sea floor, water surface, and beaches of the Earth's marine environment are littered with man-made materials. Studies show that most of the refuse washed up on beaches comes from ships. The International Maritime Organization, being responsible for the prevention and control of marine pollution from ships, adopted in 1973 the International Convention for the Prevention of Pollution from Ships. This convention, as modified by the 1978 Protocol hereafter referred to as MARPOL 73/78, applies to any ship of any type whatsoever. It covers all aspects of intentional pollution from ships by oil (Annex I), noxious substances carried in bulk (Annex II), or in packaged forms (Annex III), sewage (Annex IV), and garbage (Annex V).

The Annex V of MARPOL 73/78, entitled "Regulations for the Prevention of Pollution by Garbage from Ships," entered into force on 31 December 1988. In order to assist governments in developing and enacting domestic laws which would implement Annex V, the Marine Environment Protection Committee of the International Maritime Organization developed the Guidelines for the Implementation of Annex V of MARPOL 73/78. The guidelines are discussed in this paper.

INTRODUCTION

A considerable amount of debris originating from ships is disposed of at sea. Garbage from ships has traditionally been dumped into the sea as a matter of course and in proportion to the amount of similar wastes poured into the sea each year from the land. The quantities of waste disposed of in this way in the past are not considered excessive.

Today, however, the situation is very different. One reason is the growing everyday use of substances such as plastics which are nonbiodegradable: Once thrown into the sea, they are extremely persistent and potentially harmful if ingested by seabirds and marine mammals. Also, the aesthetic quality of coastlines and beaches has been devalued by the accumulation of such wastes.

Studies carried out in the United Kingdom by The Tidy Britain Group in 1978-79 showed that most of the refuse washed up on British beaches came from ships. Of some 20,000 items recovered, 42% were plastic (mostly containers of household products such as disinfectants, bleach, and washing-up liquid) and 22% metal. Of the latter, more than two-thirds turned out to be drink cans (International Maritime Organization (IMO) 1988b).

The amount of wastes generated by ships can be prodigious. A study carried out by a marine biologist at Newcastle University in 1982 showed that during a 44-day voyage, the 46 members of a merchant ship's crew dumped overboard 320 cardboard or paper boxes, 370 plastic beer-can holders, 162 crisp packets, 19 plastic bags, 2 plastic drums, 240 bottles, 5 glasses, 5,176 cans, and 2 metal drums. Not surprisingly perhaps, the same survey showed that of 600 ships entering ports in South Wales in 1977-78, only 13.5% used the waste disposal facilities provided on shore (Jones et al. 1981).

More recently, a survey carried out in the German Bight showed that the minimum amount of refuse drifting between Helgoland and the Elbe estuary is approximately 8.5 million pieces annually with a weight of 1,300 metric tons (MT), and 95% of the refuse found could be attributed to shipping (Vauk et al. 1987). In the Mediterranean, litter generated by shipping and oil drilling has been estimated at 325×10^6 kg/year. The total amount of litter generated in the world's oceans each year, assuming all ships' solid wastes are disposed of overboard, has been estimated at 6.5×10^6 MT/annum (Dixon and Dixon 1983).

According to the Center for Environmental Education (1986), the worldwide rate of disposal of domestic litter from merchant ships has been estimated at 110,000 MT, 0.7% of which is plastic. The amount of cargo-associated wastes including dunnage, shoring, pallets, wires, and covers is estimated at 5.6 million MT per year. Our considerations are limited to prevention of marine pollution from ships.

REGULATIONS DEVELOPED BY THE INTERNATIONAL MARITIME ORGANIZATION

To mitigate and control or even eliminate operational discharge from ships into the sea of all kinds of wastes including disposal of garbage, the IMO adopted in 1973 the International Convention for the Prevention of Pollution from Ships. This convention as modified by the 1978 Protocol (MARPOL 73/78) applies to any ship of any type whatsoever including hydrofoil boats, air-cushion vehicles, submersibles, floating rafts, and fixed or floating platforms operating in the marine environment. It covers all

aspects of intentional pollution from ships by oil (Annex I), noxious substances carried in bulk (Annex II) or in packaged forms (Annex III), sewage (Annex IV), and garbage (Annex V). (Annex V is attached as Appendix A.)

As of 8 February 1989, 55 states--the combined merchant fleets of which constitute 80.92% of gross tonnage of the world's merchant shipping--ratified the convention.

The Annexes III, IV, and V of MARPOL 73/78 are so-called "optional annexes." Each of them enters into force 12 months after the date on which at least 15 states with a combined merchant fleet of at least 50% of the gross tonnage of the world's merchant shipping have accepted it. With respect to Annex V, these conditions were fulfilled in December 1987, and subsequently, in accordance with the provisions of article 15(2) of the MARPOL Convention, this annex entered into force on 31 December 1988.

Thirty-nine states--the combined merchant fleets of which constitute 56.6% of gross tonnage of the world's merchant shipping--ratified Annex V. As of 8 February 1989, the states that have accepted Annex V are: Algeria, Antigua and Barbuda, Austria, Belgium, China, Colombia, Cote d'Ivoire, Czechoslovakia, Democratic People's Republic of Korea, Denmark, Egypt, Finland, France, Gabon, German Democratic Republic, Federal Republic of Germany, Greece, Hungary, Italy, Japan, Lebanon, Marshall Islands, Netherlands, Norway, Oman, Panama, Peru, Poland, Portugal, St. Vincent and Grenadines, Suriname, Sweden, Tunisia, Tuvalu, United Kingdom, U.S.S.R., United States, Uruguay, and Yugoslavia.

ANNEX V

The purpose of Annex V, in particular, is to prevent pollution caused by dumping into the sea all kinds of solid waste garbage, which includes inter alia all plastics, synthetic ropes, synthetic fishing nets, and plastic garbage bags. Disposal into the sea of all plastics is completely prohibited in all areas. However, some other forms of garbage may be disposed of at sea under strictly controlled conditions.

Dunnage, lining, and packing materials can only be disposed of at sea more than 25 nmi from land. Food wastes and all other garbage (including paper products, rags, glass, metal, bottles, and crockery) cannot be disposed of at sea within 12 nmi of land. Even stricter controls apply in sea areas called "special areas." For the purpose of Annex V, the special areas are: The Mediterranean Sea Area, the Baltic Sea Area, the Black Sea Area, the Red Sea Area, and the Gulfs Area. The Gulfs Area means the sea area located northwest of the rhumb line between Ras al Hadd (lat. 22°30'N, long. 59°48'E) and Ras al Fasteh (lat. 25°04'N, long. 61°25'E).

In order for ships to fully comply with the requirements on disposal of garbage within the special areas, they should be provided at ports and terminals in that area with facilities for the reception of garbage, without causing undue delay to ships.

Therefore, the government of each party to the convention whose coastline borders a special area is obliged to ensure that as soon as possible in all ports within the area, adequate reception facilities will be provided in accordance with Annex V regulations, taking into account the special needs of ships operating therein.

The IMO, after receiving notification from all governments of the states bordering a special area where adequate reception facilities have been provided, establishes the date when the requirements of the regulations on special areas shall take effect.

So far, only the states of the Baltic Sea Area (Denmark, Finland, German Democratic Republic, Federal Republic of Germany, Poland, Sweden, and U.S.S.R.) have notified IMO that requirements concerning provision of adequate reception facilities in that area have been fulfilled. The IMO notified all parties to the convention that 1 October 1989 was the established date, and starting on that date, the requirements on disposal of garbage within the Baltic Sea Area took effect (Resolution Marine Environment Protection Committee (MEPC) 31(26) adopted on 9 September 1988).

The establishment of special requirements for disposal of garbage from fixed and floating platforms is incorporated in Annex V. The disposal of any materials regulated by Annex V is completely prohibited from fixed and floating platforms which are engaged in exploration, exploitation, and associated offshore processing of seabed mineral resources, and from all other ships when alongside or within 500 m of such platforms.

The exception is the disposal into the sea of food wastes, which are permitted when they have been passed through a comminuter or grinder, but only when such fixed or floating platforms are located more than 12 nmi from land and all other ships are alongside or within 500 m of such platforms.

Limitations on the discharge of garbage from ships, as specified in Annex V, are summarized in Table 1.

GUIDELINES FOR THE IMPLEMENTATION OF ANNEX V OF MARPOL 73/78

The MEPC of the IMO at consecutive sessions in 1987 and 1988 considered the problem of implementation of Annex V. The working groups on optional annexes, under the chairmanship of T. A. Wastler for the 24th and 25th sessions and D. B. Pascoe for the 26th session, both chairmen from the United States, worked on the development of the Guidelines for the Implementation of Annex V of MARPOL 73/78, which were approved at the 26th session of MEPC, 5-9 September 1988 (IMO 1988a).

The main objectives of these guidelines are to (1) assist governments which have ratified Annex V in developing and enacting domestic laws which give force to and implement Annex V; (2) assist vessel operators in complying with the requirements set forth in Annex V and domestic laws;

Table 1.--Summary of at-sea garbage disposal regulations, MARPOL, Annex V.

Garbage type	All ships except platforms ^a		Offshore platforms ^a
	Outside special areas	In special areas ^b	
Plastics--includes synthetic ropes and fishing nets and plastic garbage bags	Disposal prohibited	Disposal prohibited	Disposal prohibited.
Floating dunnage, lining, and packing materials	>25 nmi from nearest land	Disposal prohibited	Disposal prohibited.
Paper, rags, glass, metal, bottles, crockery, and similar refuse	>12 nmi from nearest land	Disposal prohibited	Disposal prohibited.
All other garbage including paper rags, glass, etc., comminuted or ground	>3 nmi from nearest land	Disposal prohibited	Disposal prohibited.
Food waste not comminuted or ground	>12 nmi from nearest land	>12 nmi from nearest land	Disposal prohibited.
Food waste comminuted or ground ^c	>3 nmi from nearest land	>12 nmi from nearest land	>12 nmi from nearest land.
Mixed refuse types	(d)	(d)	(d)

^aOffshore platforms and associated ships include all fixed or floating platforms engaged in exploration or exploitation of seabed mineral resources, and all ships alongside or within 500 m of such platforms.

^bGarbage disposal regulations for special areas shall take effect in accordance with regulation 5(4)(b) of Annex V.

^cComminuted or ground garbage must be able to pass through a screen with mesh size no larger than 25 mm.

^dWhen garbage is mixed with other harmful substances having different disposal or discharge requirements, the most stringent disposal requirements shall apply.

Note: The Baltic Sea Special Area Disposal Regulations take effect on 1 October 1989.

and (3) assist port and terminal operators in assessing the need for and providing adequate reception facilities for garbage generated on different types of ships.

The guidelines are divided into seven categories that provide a general framework upon which governments will be able to formulate programs for education and training of seafarers and others who must comply with the regulations; methods of reducing shipboard generation of garbage; shipboard garbage handling and storage procedures; shipboard equipment for processing garbage; estimation of the amounts of ship-generated garbage delivered to port; and actions to ensure compliance with the regulations.

Recognizing that Annex V regulations promote waste management systems for ships, and that ships vary tremendously in size, mission, complement, and capability, these guidelines include a range of waste management options that may be combined in many ways to facilitate compliance with Annex V. Further, recognizing that waste management technology for ships is in an early stage of development, it is recommended that governments and the IMO continue to gather information and review these guidelines periodically.

The convention provides definitions for terms used throughout these guidelines which establish the scope of Annex V requirements. These definitions are incorporated in Section 1 as follows:

- "Food waste" is any spoiled or unspoiled victual substance such as fruits, vegetables, dairy products, poultry, meat products, food scraps, food particles, and all other material contaminated by such waste generated aboard ship, principally in the galley and dining areas.
- "Plastic" is any solid material which contains as an essential ingredient one or more synthetic organic high polymers and which is formed (shaped) by heat and/or pressure during either manufacture of the polymer or its fabrication into a finished product. Plastics have material properties ranging from hard and brittle to soft and elastic. Plastics are used for a variety of marine purposes including, but not limited to, packaging (vapor-proof barriers, bottles, containers, liners), ship construction (fiberglass and laminated structures, siding, piping, insulation, flooring, carpets, fabrics, paints and finishes, adhesives, electrical and electronic components), disposable eating utensils and cups, bags, sheeting, floats, fishing nets, strapping bands, and rope and line.
- "Domestic waste" includes all types of food wastes and wastes generated in the living spaces on board ship.
- "Cargo-associated waste" is all materials which have become wastes as a result of use for cargo stowage and handling on board ship. Cargo-associated waste includes but is not

limited to dunnage, shoring, pallets, lining and packing materials, plywood, paper, cardboard, wire, and steel strapping.

- "Maintenance waste" is materials collected by the engine and deck departments while maintaining and operating the vessel, such as soot, machinery deposits, scraped paint, deck sweeping, wiping wastes, and rags.
- "Operational waste" is all cargo-associated waste and maintenance waste, and cargo residues defined as garbage in "Cargo residues" (see below).
- "Dishwater" is the residue from the manual or automatic washing of dishes and cooking utensils which have been precleaned to the extent that any food particles adhering to them would not normally interfere with the operation of automatic dishwashers. "Gray water" is drainage from dishwater, shower, laundry, bath, and washbasin drains and does not include drainage from toilets, urinals, hospitals, and animal spaces or drainage from cargo spaces.
- "Oily rags" are rags which have been saturated with oil as controlled in Annex I to the convention. "Contaminated rags" are rags which have been saturated with a substance defined as a harmful substance in the other annexes to the convention.
- "Cargo residues," for the purposes of these guidelines, are defined as remnants of any cargo material on board that cannot be placed in proper cargo holds (loading excess and spillage) or which remain in cargo holds and elsewhere after unloading procedures are completed (unloading residual and spillage). Cargo residues are expected to be in small quantities.
- "Fishing gear" is defined as any physical device or part thereof or combination of items that may be placed on or in the water with the intended purpose of capturing, or controlling for subsequent capture, living marine or freshwater organisms.
- "Seafarer," for the purposes of these guidelines, means anyone who goes to sea in a ship for any purpose including, but not limited to, transport of goods and services, exploration, exploitation and associated offshore processing of seabed mineral resources, fishing, and recreation.

In order to clarify the application of Annex V, the following comments and explanations were added.

- Dishwater and gray water are not included as garbage in the context of Annex V.
- Ash and clinkers from shipboard incinerators and coal burning boilers are operational wastes in the meaning of Annex V regulation 1(1) and therefore are included in the term "all other garbage" in the meaning of Annex V Regulations 3(1)(b)(ii) and 5(2)(a)(ii).
- Cargo residues are to be treated as "garbage" under Annex V except when those residues are substances defined or listed under the other annexes to the convention.
- Cargo residues of all other substances are not explicitly excluded from disposal as "garbage" under the overall definition of garbage in Annex V. However, certain of these substances may pose harm to the marine environment and because of their possible safety hazards may not be suitable for disposal at reception facilities equipped to handle general garbage. The disposal of such cargo residues should be based on the physical, chemical, and biological properties of the substance and may require special handling not normally provided by garbage reception facilities.

The remaining sections of the guidelines follow.

Training, Education, and Information

The definition of "ships" used in the convention requires these guidelines address not only the professional and commercial maritime community but also the noncommercial seafaring population as sources of pollution of the sea by garbage. Uniform programs in the field of training and education will make a valuable contribution to raising the level of the seafarers' compliance with Annex V, thereby ensuring compliance with the convention. Accordingly, governments should develop and undertake training, education, and public information programs suited for all seafaring communities under their jurisdictions.

Governments may exchange and maintain information relevant to compliance with Annex V regulations through the IMO. Accordingly, governments are encouraged to provide the organization with the following:

- Technical information on shipboard waste management methods such as recycling, incineration, compaction, sorting and sanitation systems, packaging, and provisioning methods.
- Copies of current domestic laws and regulations relating to the prevention of pollution of the sea by garbage.
- Educational materials developed to raise the level of compliance with Annex V. Contributions of this type include printed materials, posters, brochures, photographs, audio-

and videotapes, and films as well as synopses of training programs, seminars, and formal curriculums.

- Information and reports on the nature and extent of marine debris found along beaches and in coastal waters under their respective jurisdictions. In order to assess the effectiveness of Annex V, these studies should provide details on amounts, distribution, sources, and impacts of marine debris.

Governments are encouraged to amend their maritime certification examinations and requirements, as appropriate, to include a knowledge of duties imposed by national and international law regarding the control of pollution of the sea by garbage.

Governments are recommended to require all ships of their registry to permanently post a summary declaration stating the prohibition and restrictions for discharging garbage from ships under Annex V and the penalties for failure to comply. It is suggested this declaration be placed on a placard at least 12.5 x 20 cm made of durable material and fixed in a conspicuous place in galley spaces, mess deck, wardroom, bridge, main deck, and other areas of the ship, as appropriate. The placard should be printed in the language or languages understood by the crew and passengers.

Governments are encouraged to have maritime colleges and technical institutes under their jurisdiction develop or augment curriculums to include both the legal duties and the technical options for handling ship-generated garbage available to professional seafarers. These curriculums should also include information on environmental impacts of garbage. Suggested topics are listed below:

- Garbage in the marine environment, sources, types, and impacts.
- National and international laws related to or impinging upon shipboard waste management.
- Health and sanitation considerations related to the storage, handling, and transfer of ship-generated garbage.
- Current technology for onboard and shoreside processing of ship-generated garbage.
- Provisioning options, materials, and procedures to minimize the generation of garbage aboard ship.

Professional associations and societies of ship officers, engineers, naval architects, shipowners and managers, and seamen are encouraged to ensure their members' competency regarding the handling of ship-generated garbage.

Vessel and reception facility operators should establish training programs for personnel operating and maintaining garbage reception or processing equipment. It is suggested that the programs include instruction on what constitutes garbage and the applicable regulations for handling and disposing of it. Such training should be reviewed annually.

Generalized public information programs are needed to provide information to nonprofessional seafarers and others concerned with the health and stability of the marine environment regarding the impacts of garbage at sea. Governments and involved commercial organizations are encouraged to utilize the organization's library and to exchange resources and materials as appropriate to initiate internal and external public awareness programs.

Methods for delivering this information include radio and television, articles in periodicals and trade journals, voluntary public projects such as beach cleanup days and adopt-a-beach programs, public statements by high government officials, posters, brochures, conferences and symposia, cooperative research and development, voluntary product labeling, and teaching materials for public schools.

Audiences include recreational boaters and fishermen, port and terminal operators, coastal communities, ship supply industries, shipbuilders, waste management industries, plastic manufacturers and fabricators, trade associations, educators, and government.

The subjects addressed in these programs are recommended to include the responsibilities of citizens under national and international law; options for handling garbage at sea and upon return to shore; known sources and types of garbage; impacts of plastic debris on seabirds, fish, marine mammals, sea turtles, and ship operations; impacts on coastal tourist trade; current actions by governments and private organizations; and sources of additional information.

Minimizing the Amount of Potential Garbage

All ship operators should minimize the taking aboard of potential garbage and onboard generation of garbage.

Domestic wastes may be minimized through proper provisioning practices. Ship operators and governments should encourage ship's suppliers and provisioners to consider their products in terms of the garbage they generate. Options available to decrease the amount of domestic waste generated aboard ship include the following:

- Bulk packaging of consumable items. This may result in the creation of less waste, but factors such as inadequate shelf life once a container is opened must be considered to avoid increasing wastes.
- Reusable packaging and containers. This can decrease the amount of garbage being generated. Use of disposable cups,

utensils, dishes, towels and rags, and other convenience items should be limited.

- Provisions packaged in or made of materials other than disposable plastic. These should be selected to replenish ship supplies unless a reusable plastic alternate is available.

Operational waste generation is specific to individual ship activities and cargoes. It is recommended that manufacturers, shippers, ship operators, and governments consider the garbage associated with various categories of cargoes and take actions as needed to minimize their generation. Suggested actions are listed below.

- Replace disposable plastic sheeting used for cargo protection with permanent, reusable covering materials.
- Adopt stowage systems and methods that reuse coverings, dunnage, shoring, lining, and packing materials.
- Dispose of dunnage, lining, and packaging materials generated in port during cargo discharge at the port reception facilities and do not retain it on board for discharge at sea.

It may, in certain cases, be difficult for port reception facilities to handle cargo residues. They are usually created through inefficiencies in loading, unloading, and onboard handling, and it is, therefore, recommended that cargo be unloaded as efficiently as possible in order to avoid or minimize cargo residues.

Spillage of the cargo during transfer operations should be carefully controlled, both on board and from dockside. Spillage typically occurs in port. It should be completely cleaned up prior to sailing and either delivered into the intended cargo space or into the port reception facility. Shipboard areas where spillage is most common should be protected so that the residues are easily recovered.

Fishing gear once discharged becomes a harmful substance. Fishing vessel operators, their organizations, and their respective governments are encouraged to undertake such research, technology development, and regulations as may be necessary to minimize the probability of loss and maximize the probability of recovery of fishing gear from the ocean. It is recommended that fishing vessel operators record and report the loss and recovery of fishing gear. Techniques both to minimize the amount of fishing gear lost in the ocean and to maximize its recovery are listed below.

- Operators and associations of fishing vessels using untended, fixed, or drifting gear are encouraged to develop information exchanges with such other ship traffic as may be necessary to minimize accidental encounters between ships and gear.

Governments are encouraged to assist in the development of information systems where necessary.

- Fishery managers are encouraged to consider the probability of encounters between ship traffic and fishing gear when establishing seasons, areas, and gear-type regulations.
- Fishery managers and fishing vessel operators and associations are encouraged to utilize gear identification systems which provide information such as vessel name, registration number, and nationality. Such systems may be useful to promote reporting, recovery, and return of lost gear.
- Fishing vessel operators are encouraged to document positions and reasons for loss of their gear. To reduce the potential of entanglement and ghost fishing (capture of marine life by discharged fishing gear), benthic traps, trawls, and gillnets could be designed to have degradable panels or sections made of natural fiber twine, wood, or wire.
- Governments are encouraged to consider the development of technology for more effective fishing gear identification systems.

Governments are encouraged to undertake research and technology development to minimize potential garbage and its impacts on the marine environment. Following are suggested areas for such study:

- Development of recycling technology and systems for synthetic materials returned to shore as garbage.
- Development of technology for degradable synthetic materials to replace current plastic products, as appropriate. In this connection, governments should also study the impacts on the environment of the products of degradation of such new materials.

Shipboard Garbage Handling and Storage Procedures

Compliance with limitations on the discharge of garbage from ships as specified in Annex V requires personnel, equipment, and procedures for collecting, sorting, processing, storing, and disposing of garbage. Economic and procedural considerations associated with these activities include storage space requirements, sanitation, equipment and personnel costs, and in-port garbage service charges.

In complying with the provisions of Annex V, the most appropriate procedures for handling and storing garbage on ships will vary depending on factors such as the type and size of the ship, the area of operation (e.g., distance from nearest land), shipboard garbage processing equipment and storage space, crew size, duration of voyage, and regulations and reception

facilities at ports of call. Proper handling and storage minimize shipboard storage space requirements and enable efficient transfer of retained garbage to port reception facilities.

To ensure that the most effective and efficient handling and storage procedures are followed, it is recommended that vessel operators develop waste management plans that can be incorporated into crew and vessel operating manuals. Such plans should include the appointment of an environmental control officer. The manuals should identify crew responsibilities and procedures for all aspects of handling and storing garbage aboard the ship. Procedures for handling ship-generated garbage can be divided into four phases: Collection, processing, storage, and disposal. A generalized waste management plan for handling and disposal of ship-generated garbage is presented in Table 2. Specific procedures for each phase are discussed below.

Collection


Procedures for collecting garbage generated aboard ship should be based on consideration of what can be discarded overboard while en route. To reduce or avoid the need for sorting after collection, it is recommended that three categories of distinctively marked garbage receptacles be provided to receive garbage as it is generated. These separate receptacles (e.g., cans, bags, or bins) would receive (1) plastics and plastics mixed with nonplastic garbage, (2) food wastes (which includes materials contaminated by such wastes), and (3) other garbage which can be disposed of at sea. Receptacles for each of the three categories of garbage should be clearly marked and distinguishable by color, graphics, shape, size, or location. These receptacles should be provided in appropriate spaces throughout the ship (e.g., the engine room, mess deck, wardroom, galley, and other living or working spaces), and all crew members and passengers should be advised of what garbage should and should not be discarded in them. Crew responsibilities should be assigned for collecting or emptying these receptacles and taking the garbage to the appropriate processing or storage location. Use of such a system will facilitate subsequent shipboard processing and minimize the amount of garbage which must be stored aboard ship for return to port.

Separate cans or bags could be provided for receiving and storing glass, metal, plastics, paper, or other items which can be recycled. To encourage crew members to deposit such items in provided receptacles, proceeds generated from their return might be added to a ship's recreational fund.

Plastics and plastics mixed with nonplastic garbage.--Plastic garbage must be retained aboard ship for discharge at port reception facilities unless reduced to ash by incineration. When plastic garbage is not separated from other garbage, the mixture must be treated as if it were all plastic.

Synthetic fishing net and line scraps generated by the repair or operation of fishing gear may not be discarded at sea and should be

Table 2.--Options for shipboard handling and disposal of garbage.



collected in a manner that avoids its loss overboard. Such material may be incinerated, compacted, or stored along with other plastic waste, or it may be preferable to keep it separate from other types of garbage if it has strong odor or great volume.

Food wastes.--Some governments have regulations for controlling human, plant, and animal diseases that may be carried by foreign food wastes and materials that have been associated with them (e.g., food packaging and disposable eating utensils). These regulations may require incinerating, sterilizing, or other special treatment of garbage to destroy possible pest and disease organisms. Such garbage should be kept separate from other garbage and preferably retained for disposal in port in accordance with laws of the receiving country. With regard to such garbage, governments are reminded of their obligation to assure the provision of adequate reception facilities. Precautions must be taken to ensure that plastics contaminated by food wastes (e.g., plastic food wrappers) are not discharged at sea with other food wastes.

Other garbage.--Garbage in this category includes, but is not limited to, paper products, rags, glass, metal bottles, crockery, dunnage, lining, and packing materials. Vessels may find it desirable to separate dunnage, lining, and packing material, which will float, since this material is subject to a different discharge limit than other garbage in this category (Table 1). Such garbage should be kept separate from other garbage and preferably retained for disposal in port.

Oily rags and contaminated rags must be kept on board and discharged to a port reception facility or incinerated.

Recovery of garbage at sea.--Fishermen and other seafarers who recover derelict fishing gear and other persistent garbage during routine operations are encouraged to retain this material for disposal on shore. If lost pots or traps are recovered and space is not available for storage, fishermen and other seafarers are encouraged to remove and transport any line and webbing to port for disposal and return the bare frames to the water or, minimally, to cut open the traps to keep them from continuing to trap marine life.

Seafarers are further encouraged to recover other persistent garbage from the sea as opportunities arise and prudent practice permits.

Processing

Depending on factors such as the type of ship, area of operation, and size of crew, ships may be equipped with incinerators, compactors, comminuters, or other devices for shipboard garbage processing (see "Shipboard Equipment for Processing Garbage" below). Members of the crew should be assigned responsibility for operating this equipment on a schedule commensurate with ship needs. In selecting appropriate processing procedures, consider that compactors, incinerators, comminuters, and other such devices have a number of advantages, such as making it possible to discharge certain garbage at sea, reducing shipboard space requirements for storing

garbage, making it easier to off-load garbage in port, and enhancing assimilation of garbage discharged into the marine environment.

It should be noted that special rules on incineration may be established by authorities in some ports and may exist in some special areas. Incineration of the following items requires special precaution due to the potentially harmful environmental and health effects from combustion of by-products: Hazardous materials (e.g., scraped paint, impregnated wood) and certain types of plastics (e.g., polyvinyl chloride-based plastics). The problems of combustion of by-products are discussed in the section entitled "Shipboard Equipment for Processing Garbage."

Ships operating primarily in special areas or within 3 nmi of the nearest land should choose between storage of either compacted or uncompact materials for off-loading at port reception facilities or incineration with retention of ash and clinkers. This is the most restrictive situation in that no discharge is permitted.

Compactors make garbage easier to store, to transfer to port reception facilities, and to dispose of at sea when discharge limitations permit. In the latter case, compacted garbage also may sink more readily, which would reduce aesthetic impacts in coastal waters and along beaches and perhaps reduce the likelihood of marine life ingesting or otherwise interacting with discharged materials.

Ships operating primarily beyond 3 nmi from the nearest land are encouraged to install and use comminuters to grind food wastes to a particle size capable of passing through a screen with openings no larger than 25 mm. Although larger food scraps may be discharged beyond 12 nmi, it is recommended that comminuters be used even outside this limit because they hasten assimilation into the marine environment. Because food wastes comminuted with plastics cannot be discharged at sea, all plastic materials must be removed before food wastes are ground.

Storage

Garbage collected from living and working areas throughout the ship should be delivered to designated processing or storage locations. Garbage that must be returned to port for disposal may require long-term storage depending on the length of the voyage or arrangements for off-loading (e.g., transferring garbage to an offshore vessel for incineration or subsequent transfer ashore). Garbage which may be discarded overboard may require short-term storage or no storage. In all cases, garbage should be stored in a manner which avoids health and safety hazards. The following points should be considered when selecting procedures for storing garbage:

- Ships should use separate cans, drums, boxes, bags, or other containers for short-term (disposal garbage) and trip-long (nondisposable garbage) storage. Short-term storage would be appropriate for holding otherwise disposable garbage while a ship is passing through a restricted discharge area.

- Sufficient storage space and equipment (e.g., cans, drums, bags, or other containers) should be provided. Where space is limited, vessel operators are encouraged to install compactors or incinerators. To the extent possible, all processed and unprocessed garbage which must be stored for any length of time should be in tight, securely covered containers.
- Food wastes and associated garbage which are returned to port and which may carry diseases or pests should be stored in tightly covered containers and be kept separate from garbage which does not contain such food wastes. Both types of garbage should be stored in separate, clearly marked containers to avoid incorrect disposal and treatment on land.
- Waste fishing gear can be stored on deck if materials have strong odors or if their size is too great to permit storage elsewhere on the ship. In cases where gear is fouled with marine growth or dead organisms, it may be reasonable to tow gear behind the vessel for a time to wash it out before storing. If it cannot be recovered by the vessel, the appropriate coastal state should be notified of its location.
- Disinfection and both preventive and remedial pest control methods should be applied regularly in garbage storage areas.

Disposal

Although disposal is possible under Annex V, discharge of garbage to port reception facilities should be given first priority. Disposal of ship-generated garbage must be done in a manner consistent with the regulations summarized in Table 1. When disposing of garbage, the following points should be considered:

- Garbage which may be disposed of at sea can simply be discharged overboard. Disposal of uncompacted garbage is convenient, but results in a maximum number of floating objects which may reach shore even when discharged beyond 25 nmi from the nearest land. Compacted garbage is more likely to sink and thus less likely to pose aesthetic problems. If necessary and possible, weights should be added to promote sinking. Compacted bales of garbage should be discharged over deep water (50 m or more) to prevent rapid loss of their structural integrity due to wave action and currents.
- Floating cargo-associated waste that is not plastic or otherwise regulated under other MARPOL annexes may be discharged beyond 25 nmi from the nearest land. Cargo-associated waste that will sink and is not plastic or otherwise regulated may be discharged beyond 12 nmi from the nearest land. Most cargo-associated waste is generated during the loading and unloading process, usually at

dockside. It is recommended that every effort be made to deliver these wastes to the nearest port reception facility prior to the ship's departure.

- Maintenance wastes are generated more or less steadily during the course of routine ship operations. In some cases, maintenance wastes may be contaminated with substances, such as oil or toxic chemicals, controlled under other annexes or other pollution control laws. In such cases, the more stringent disposal requirements take precedence.
- To ensure timely transfer of large quantities of ship-generated garbage to port reception facilities, it is essential for ships or their agents to make arrangements well in advance for garbage reception. At the same time, disposal needs should be identified in order to make arrangements for garbage requiring special handling or other necessary arrangements. Special disposal needs might include off-loading food wastes and associated garbage which may carry certain disease or pest organisms, or unusually large, heavy, or odorous derelict fishing gear.

Shipboard Equipment for Processing Garbage

The range of options for garbage handling aboard ships depends largely upon costs, personnel limitations, generation rate, capacity, vessel configuration, and traffic patterns. The type of equipment available to address various facets of shipboard garbage handling include incinerators, compactors, comminuters, and their associated hardware.

Grinding or Comminution

When not in a special area, the discharge of comminuted food wastes and all other comminuted garbage (except plastics and floatable dunnage, lining, and packing materials) may be permitted under Regulation 3(1)(c) of Annex V beyond 3 nmi from the nearest land. It is recommended that such comminuted or ground garbage not be discharged into a ship's sewage treatment system unless the system is approved for treating such garbage. Furthermore, it should not be stored in bottoms or tanks containing oily wastes. Such actions can result in faulty operation of sewage treatment or oily water separator equipment and can cause sanitary problems for crew members and passengers. Options for grinding or comminution include the following:

- A wide variety of food waste grinders are available in the market and are commonly fitted in most modern ships' galleys. These food waste grinders produce a slurry of food particles and water that washes easily through the required 25 mm screen. Output ranges from 10 to 250 L/min. It is recommended that the discharge from shipboard comminuters be directed into a holding tank when the vessel is operating within an area where discharge is prohibited.

- Size reduction of certain other garbage items can be achieved by shredding or crushing, and machines for carrying out this process are available for use on board ships.

Information on the development and use of comminuters for garbage aboard ships should be forwarded to the organization.

Compaction

Table 3 gives compaction information for various types of garbage.

Most garbage can be compacted; the exceptions include unground plastics, fiber- and paperboard, bulky cargo containers, and thick metal items. Pressurized containers should not be compacted since they present an explosion hazard.

Compaction can reduce the volume of garbage into bags, boxes, or briquettes. When these compacted slugs are equally formed and structurally strong, they can be piled up in building block form; this permits the most efficient use of space in the storage compartments. The compaction ratio for normal mixed shipboard garbage may range as high as 12:1.

Some of the available compactors have options such as sanitizing, deodorizing, adjustable compaction ratios, bagging in plastic or paper, boxing in cardboard (with or without plastic or waxed paper lining), and baling. Paper or cardboard tends to become soaked and weakened by moisture in the garbage during long periods of onboard storage. There have also been problems due to the generation of gas and pressure which can explode tight plastic bags.

If grinding machines are used prior to compaction, the compaction ratio can be increased and the storage space decreased.

A compactor should be installed in a compartment with adequate room for operating and maintaining the unit and storing trash to be processed. The compartment should be located adjacent to the areas of food processing and commissary storerooms. If not already required by regulations, it is recommended that the space have freshwater washdown service, coamings, deck drains, adequate ventilation, and hand or automatic fixed fire fighting equipment.

Information on the development and use of shipboard compactors should be forwarded to the organization.

Incineration

Compared to the technology of land-based incineration, the state of the art in marine incinerators is not highly advanced, primarily because the technology has not yet been subject to constraints on air emissions or on the types of materials that could be incinerated. Marine incinerators in current use are predominately designed for intermittent operation and hand stoking, and typically do not include any provisions for air pollution

Table 3.--Compaction guide for shipboard-generated garbage.

Typical examples	Special handling by vessel personnel before compaction	Compaction characteristics			Onboard storage space
		Rate of alteration	Retention of compacted form	Density of compaction form	
Metal food and beverage containers, glass, small wood pieces	None	Very rapid	Almost 100%	High	Minimum
Comminuted plastics, fiber and paper-board	Minor--reduce material to size for feed, minimal manual labor	Rapid	Approximately 80%	Medium	Minimum
Small metal drums, uncomminuted cargo packing, large pieces of wood	Moderate--longer manual labor time required to size material for feed	Slow	Approximately 50%	Relatively low	Moderate
Uncomminuted plastics	Major--very long manual labor time to size material for feed; usually impractical	Very slow	Less than 10%	Very low	Maximum
Bulky metal cargo containers, thick metal items	Impractical for shipboard compaction not feasible	Not applicable	Not applicable	Not applicable	Maximum

control. Control of air pollution is normally required in many ports in the world. Prior to using an incinerator while in port, permission may be required from the port authority concerned. In general, the use of shipboard garbage incinerators in ports in or near urban areas should be discouraged, as their use will add to possible air pollution in these areas.

Table 4 is a guide for incineration of garbage, including combustibility, reduction of volume, residual materials, exhaust, onboard storage space, and any required special handling by vessel personnel. With the exception of metal and glass, most garbage is amenable to incineration.

In contrast to land-based incinerators, shipboard incinerators must be as compact as practicable, and with operating personnel at a premium, automatic operation is desirable. Most shipboard incinerators are designed for intermittent operation: The waste is charged to the incinerator, firing is started, and combustion typically lasts for 3 to 6 h.

Commercial marine incinerators currently available vary greatly in size, have natural or induced draft, and are hand-fired. It should be noted that incinerator ratings are usually quoted on the basis of heat input rate rather than on a weight charged basis because of the variability of the heat content in the wastes. Some modern incinerators are designed for continuous firing and can handle simultaneous disposal of nearly all shipboard waste.

Some of the advantages of the most advanced incinerators are: They operate under negative pressure, they are highly reliable since they have few moving parts, they require minimal operator skill, they are light in weight, and they have low exhaust and external skin temperatures.

Some of the disadvantages of incinerators are: The possible hazardous nature of the ash or vapor; dirty operation; excessive labor required for charging, stoking, and ash removal; and the probability of not meeting air pollution regulations imposed in certain harbors. Some of these disadvantages can be remedied by automatic equipment for charging, stoking, and discharging ash into the sea outside areas where such discharge is prohibited. The additional equipment to perform these automatic functions requires more installation space.

The incineration of predominantly plastic wastes, considered under some circumstances in complying with Annex V, requires more air and much higher temperatures for complete destruction. If plastics are to be burned in a safe manner, the incinerator should be suitable for the purpose; otherwise, the following problems can result:

- Depending on the type of plastic and conditions of combustion, some toxic gases can be generated in the exhaust stream, including vaporized hydrochloric (HCl) and hydrocyanic (HCN) acids. These and other intermediary products of plastic combustion can be extremely dangerous.

Table 4.---Incineration^a guide for shipboard-generated garbage.

Typical examples	Incineration characteristics				Onboard storage space	
	Special handling by vessel personnel before incineration	Combustibility	Reduction of volume	Residual		Exhaust
Paper packaging, food and beverage containers	Minor--easy to feed into hopper	High	Over 95%	Powder ash	Possibly smoky and not hazardous	Minimum.
Fiber and paperboard	Minor--reduce material to size for feed; minimum manual labor	High	Over 95%	Powder ash	Possibly smoky and not hazardous	Minimum.
Plastic packaging, food and beverage containers	Minor--easy to feed into hopper	High	Over 95%	Powder ash	Possibly smoky and hazardous based on incinerator design	Minimum.
Plastic sheeting, netting, rope, and bulk material	Moderate manual labor time for size reduction	High	Over 95%	Powder ash	Possibly smoky and hazardous based on incinerator design	Minimum.
Rubber hoses and bulk pieces	Major manual labor time for size reduction	High	Over 95%	Powder ash	Possibly smoky and hazardous based on incinerator design	Minimum.
Metal food and beverage containers	Minor--easy to feed into hopper	Low	Less 10%	Slag	Possibly smoky and not hazardous	Moderate.
Metal cargo, bulky containers, thick metal items	Major manual labor time for size reduction (not easily incinerated)	Very low	Less 5%	Large metal fragments and slag	Possibly smoky and not hazardous	Maximum.
Glass food and beverage containers	Minor--easy to feed low into hopper	Low	Less	Slag	Possibly smoky and not hazardous	
Wood, cargo containers and large wood scraps	Moderate manual labor time for size reduction	High	Over 95%	Powder ash	Possibly smoky and not hazardous	Minimum.

^aCheck local rules for possible restrictions.

- The ash from the combustion of some plastic products may contain heavy metal or other residues which can be toxic and should, therefore, not be discharged into the sea. Such ash should be retained on board where possible, and discharged at port reception facilities.
- The high temperatures generated during incineration of primarily plastic wastes may damage some garbage incinerators.

Plastic incineration requires 3 to 10 times more combustion air than average municipal refuse. If the proper level of oxygen is not supplied, high levels of soot will form in the exhaust stream.

Certain ship classification societies have established requirements for the operation or construction of incinerators. The International Association of Classification Societies can provide information as to such requirements.

Information on the development and utilization of marine garbage incinerator systems for shipboard use should be forwarded to the organization.

Port Reception Facilities for Garbage

Governments are urged to initiate at the earliest opportunity studies into the provision of reception facilities at ports in their respective countries. They should carry out the studies in close cooperation with port authorities and other local authorities responsible for garbage handling. Such studies should include information such as port-by-port listing of available garbage reception facilities, the types of garbage they are equipped to handle (e.g., food wastes contaminated with foreign disease or pest organisms, large pieces of derelict fishing gear, or refuse and operational wastes only), their capacities, and any special procedures required to use them. Governments should provide the results of their studies to the organization for inclusion in the Annex V library.

While selecting the most appropriate type of reception facility for a particular port, consideration should be given to several alternative methods available. In this regard, floating plants such as barges or self-propelled ships might be considered more effective for collection of garbage in a particular location than land-based facilities.

The equipment for treatment and disposal of garbage is a significant factor in determining the adequacy of a reception facility. It not only provides a measure of the time required to complete the process, but it also is the primary means for ensuring that ultimate disposal of the garbage is environmentally safe.

Governments, in assessing the adequacy of reception facilities, should also consider the technological problems associated with the treatment and disposal of garbage received from ships. Although the establishment of

waste management standards is not within the scope of the convention, governments should take responsible actions within their national programs to consider such standards.

The methodology for determining the adequacy of a reception facility should be based on the needs of each type of ship as well as the number and types of ships using the port. The size and location of a port should be considered in determining adequacy. Emphasis should also be given to calculating the quantities of ship-generated garbage not discharged into the sea in accordance with the provisions of Regulations 3, 4, and 5 of Annex V.

Vessel, port, and terminal operators should consider the following when determining quantities of garbage on a per ship basis:

- type of garbage,
- ship type and design,
- ship operating route,
- number of persons on board,
- duration of voyage,
- time spent in areas where discharge into the sea is prohibited or restricted, and
- time spent in port.

It should be noted that reception procedures may differ, and port reception may require onboard separation of food wastes (e.g., raw meat because of risk of animal diseases), cargo-associated waste, and domestic and maintenance waste.

The purpose of these guidelines will be attained if they provide the necessary stimulus to governments to initiate and continue studies of reception facilities and of treatment and disposal technology. Information on developments in this respect should be forwarded to the organization.

Ensuring Compliance With Annex V

Recognizing that direct enforcement of Annex V regulations, particularly at sea, is difficult to accomplish, governments are encouraged to consider not only restrictive and punitive measures but also the removal of any disincentives, creation of positive incentives, and the development of voluntary measures within the regulated community when developing programs and domestic legislation to ensure compliance with Annex V.

Enforcement

Governments should encourage their flag vessels to advise them of ports in foreign countries party to Annex V which do not have port reception facilities for garbage. This will provide a basis for advising responsible governments of possible problems and calling IMO's attention to possible infractions. An acceptable reporting format is in Appendix B.

Governments should establish a documentation system (e.g., letters or certificates) for ports and terminals under their jurisdiction having adequate facilities for receiving ship-generated garbage. Periodic inspection of the reception facilities is recommended.

Governments should identify appropriate enforcement agencies providing legal authority, adequate training, funding, and equipment to incorporate the enforcement of Annex V regulations into their responsibilities. In those cases where customs or agricultural officials are responsible for receiving and inspecting garbage, governments should ensure that the necessary inspections are facilitated as much as possible.

Governments should consider, where applicable, the use of garbage discharge reporting systems for ships (e.g., existing ship's deck logbook or record books). Such logs should, at a minimum, document the date, time, location by latitude and longitude or name of port, type of garbage (e.g., food, refuse, cargo-associated waste, or maintenance waste), and estimated amount of garbage discharged. Particular attention should be given to the reporting of:

- the loss of fishing gear,
- the discharge of cargo residues,
- any discharge in special areas,
- discharge at port reception facilities, and
- discharge of garbage at sea.

The issue of documents or receipts by port reception facilities might also assist the reporting system.

Encouraging and Facilitating Compliance

The augmentation of port reception facilities to serve ship traffic without undue delay or inconvenience may require capital investment from port and terminal operators as well as the waste management companies serving those ports. Governments are encouraged to evaluate means within their authority to lessen this impact, thereby helping to ensure that garbage delivered to port is actually received and disposed of properly at reasonable costs or without charging special fees to individual ships. Such means include, but are not limited to:

- tax incentives;
- loan guarantees;
- public vessel business preference;
- special funds to assist in problem situations such as remote ports with no land-based waste management systems to receive ships' garbage;
- government subsidies and special funds to help defray the cost of a bounty program for lost, abandoned, or discarded fishing gear or other persistent garbage. The program would make appropriate payments to persons who retrieve such fishing gear or any persistent garbage other than their own from marine waters under the jurisdiction of a government.

The installation of shipboard garbage processing equipment would facilitate compliance with Annex V and lessen the burden on port reception facilities to process garbage for disposal. Therefore, governments should consider actions to encourage the installation of certain types of garbage processing equipment on ships operating under their flag. For example, programs to lessen costs of purchasing and installing compactors, incinerators, and comminuters during construction of new ships would be very helpful.

Governments are encouraged to consider the economic impacts of domestic regulations intended to force compliance with Annex V. Unrealistic regulations may lead to higher levels of noncompliance than would an education program without specific regulatory requirements beyond Annex V itself. Due to the highly variable nature of ship operations and configurations, it seems appropriate to maintain the highest possible level of flexibility in domestic regulations to permit ships the greatest range of options in complying with Annex V.

Governments are encouraged to support research and development of technology that will simplify compliance with Annex V regulations for ships and ports. This research should concentrate on:

- shipboard waste-handling systems;
- ship provision innovations to minimize garbage generation;
- loading and unloading technology to minimize dunnage, spillage, and cargo residues; and
- new ship construction design to facilitate garbage management and transfer.

Governments are encouraged to work within the organization to develop port reception systems that simplify the transfer of garbage for international vessels.

Voluntary Measures

Governments are encouraged to assist ship operators' and seafarers' organizations in developing resolutions, bylaws, and other internal mechanisms that will encourage compliance with Annex V regulations. These groups include seamen's and officers' unions, associations of ship owners and insurers and classification societies, and pilot associations and fishermen's organizations.

Governments are encouraged to assist and support where possible the development of internal systems to promote compliance with Annex V in port authorities and associations, terminal operators' organizations, stevedores' and longshoremen's unions, and land-based waste management authorities.

CONCLUSIONS

The legal framework for international cooperation in the protection of the marine environment from pollution caused by disposal into the sea of garbage from ships was given effect with the entry into force of Annex V of MARPOL 73/78 on 31 December 1988 (Marine Environment Protection Committee of the International Maritime Organization 1988).

First priority was given to the uniform implementation of Annex V through the development and enactment of enabling domestic laws. It is envisaged that the guidelines described in this paper will facilitate this process and improve the effectiveness of these measures to prevent pollution of the seas by garbage from ships.

The next step will necessarily be the encouragement of the widest possible acceptance and implementation of Annex V by states. The IMO Secretariat has integrated implementation of Annex V into its technical assistance Program for the Protection of Marine Environment. It can, therefore, be expected that global and regional workshops and seminars will be organized by IMO in cooperation with interested international and regional organizations with the aim of familiarizing experts, in particular those from developing countries, with the regulations and guidelines and providing advice on their implementation.

In 1981, the IMO assembly of its member states adopted Resolution A.500(XIII). The crux of this resolution is the agreement that it is undesirable to amend existing conventions unless they have been in force for a reasonable period of time and experience has been gained regarding the costs to the maritime industry of their operation and the burden on the legislative and administrative resources of IMO member states. The principle endorsed is that new conventions or amendments to existing conventions should be considered on the basis of a "clear and well-documented demonstration of compelling need."

After further practical experience has been gained with the implementation and enforcement of Annex V, there will doubtless be suggestions coming forth on strengthening the provisions of the annex and on further elaborating the guidelines. These suggested changes are likely to include:

- a requirement that individual waste management plans be developed for particular categories of vessels;
- a requirement that log entries be made for all waste garbage disposal practices to facilitate port control;
- requirements concerning standards of the garbage handling equipment to be installed on ships;
- a requirement concerning garbage separation on board ships; and
- guidelines on incineration of garbage on board ships.

REFERENCES

- Center for Environmental Education.
1986. Use and disposal of non-biodegradable plastics in the marine and Great Lakes environments. Center for Environmental Education, Wash., D.C.
- Dixon, T. R., and T. J. Dixon.
1983. Marine litter surveillance on the North Atlantic Ocean shores of Portugal and the Western Isles of Scotland. Marine Litter Research Programme, Stage 5, Keep Britain Tidy Group.
- International Maritime Organization.
1988a. Guidelines for the implementation of Annex of MARPOL 73/78. (Sales No.: 5388.09E.)

1988b. Strategy for the protection of the marine environment.
- Jones, R. E., S. G. Pine, M. C. Ferriter, and F. E. M. Ferriter.
1981. Sea land litter. Report of two surveys conducted by the Keep Wales Tidy Campaign into marine waste disposal practices of merchant shipping and pleasurecraft on the South Wales coastline.
- Marine Environment Protection Committee of the International Maritime Organization.
1988. Marine Environment Protection Committee Resolution 31(26) adopted on 9 September 1988 and Implementation of Annexes V and IV of MARPOL 73/78. 26th Session, agenda item 10.
- Vauk, G., G. Dahlmann, E. Hartwig, J. C. Ranger, B. Reineking, E. Schrey, E. Vauk-Hentzelt.
1987. Recording of oil victims on the German North Sea coast and results of oil analyses as well as investigations of burdening of the German Bight by ship's refuse. A report.

APPENDIX A

ANNEX V

REGULATIONS FOR THE PREVENTION OF POLLUTION
BY GARBAGE FROM SHIPS

Regulation 1

Definitions

For the purposes of this Annex:

(1) "Garbage" means all kinds of victual, domestic and operational waste excluding fresh fish and parts thereof, generated during the normal operation of the ship and liable to be disposed of continuously or periodically except those substances which are defined or listed in other Annexes to the present Convention.

(2) "Nearest land". The term "from the nearest land" means from the baseline from which the territorial sea of the territory in question is established in accordance with international law except that, for the purposes of the present Convention "from the nearest land" off the north eastern coast of Australia shall mean from a line drawn from a point on the coast of Australia in

latitude 11°00' South, longitude 142°08' East to a point in
latitude 10°35' South,
longitude 141°55' East, thence to a point latitude 10°00' South,
longitude 142°00' East, thence to a point latitude 9°10' South,
longitude 143°52' East, thence to a point latitude 9°00' South,
longitude 144°30' East, thence to a point latitude 13°00' South,
longitude 144°00' East, thence to a point latitude 15°00' South,
longitude 146°00' East, thence to a point latitude 18°00' South,
longitude 147°00' East, thence to a point latitude 21°00' South,
longitude 153°00' East, thence to a point on the coast of Australia
in latitude 24°42' South, longitude 153°15' East.

(3) "Special area" means a sea area where for recognized technical reasons in relation to its oceanographical and ecological condition and to the particular character of its traffic the adoption of special mandatory methods for the prevention of sea pollution by garbage is required. Special areas shall include those listed in Regulation 5 of this Annex.

Regulation 2

Application

The provisions of this Annex shall apply to all ships.

Regulation 3

Disposal of Garbage Outside Special Areas

- (1) Subject to the provisions of Regulations 4, 5, and 6 of this Annex:
- (a) the disposal into the sea of all plastics, including but not limited to synthetic ropes, synthetic fishing nets and plastic garbage bags is prohibited;
 - (b) the disposal into the sea of the following garbage shall be made as far as practicable from the nearest land but in any case is prohibited if the distance from the nearest land is less than:
 - (i) 25 nautical miles for dunnage, lining and packing materials which will float;
 - (ii) 12 nautical miles for food wastes and all other garbage including paper products, rags, glass, metal, bottles, crockery and similar refuse;
 - (c) disposal into the sea of garbage specified in sub-paragraph (b)(ii) of this Regulation may be permitted when it has passed through a comminuter or grinder and made as far as practicable from the nearest land but in any case is prohibited if the distance from the nearest land is less than 3 nautical miles. Such comminuted or ground garbage shall be capable of passing through a screen with openings no greater than 25 millimetres.
- (2) When the garbage is mixed with other discharges having different disposal or discharge requirements the more stringent requirements shall apply.

Regulation 4

Special Requirements for Disposal of Garbage

- (1) Subject to the provisions of paragraph (2) of this Regulation, the disposal of any materials regulated by this Annex is prohibited from fixed or floating platforms engaged in the exploration, exploitation and associated offshore processing of seabed mineral resources, and from all other ships when alongside or within 500 metres of such platforms.
- (2) The disposal into the sea of food wastes may be permitted when they have been passed through a comminuter or grinder from such fixed or floating platforms located more than 12 nautical miles from land and all other ships when alongside or within 500 metres of such platforms. Such comminuted or ground food wastes shall be capable of passing through a screen with openings no greater than 25 millimetres.

Regulation 5

Disposal of Garbage Within Special Areas

(1) For the purposes of this Annex the special areas are the Mediterranean Sea area, the Baltic Sea area, the Black Sea area, the Red Sea area and the "Gulfs area" which are defined as follows:

- (a) The Mediterranean Sea area means the Mediterranean Sea proper including the gulfs and seas therein with the boundary between the Mediterranean and the Black Sea constituted by the 41°N parallel and bounded to the west by the Straits of Gibraltar at the meridian of 5°36'W.
- (b) The Baltic Sea area means the Baltic Sea proper with the Gulf of Bothnia and the Gulf of Finland and the entrance to the Baltic Sea bounded by the parallel of the Skaw in the Skagerrak at 57°44.8'N.
- (c) The Black Sea area means the Black Sea proper with the boundary between the Mediterranean and the Black Sea constituted by the parallel 41°N.
- (d) The Red Sea area means the Red Sea proper including the Gulfs of Suez and Aqaba bounded at the south by the rhumb line between Ras si Ane (12°8.5'N, 43°19.6'E) and Husn Murad (12°40.4'N, 43°30.2'E).
- (e) The "Gulfs area" means the sea area located north west of the rhumb line between Ras al Hadd (22°30'N, 59°48'E) and Ras al FasteH (25°04'N, 61°25'E).

(2) Subject to the provisions of Regulation 6 of this Annex:

- (a) disposal into the sea of the following is prohibited:
 - (i) all plastics, including but not limited to synthetic ropes, synthetic fishing nets and plastic garbage bags; and
 - (ii) all other garbage, including paper products, rags, glass, metal, bottles, crockery, dunnage, lining and packing materials;
- (b) disposal into the sea of food wastes shall be made as far as practicable from land, but in any case not less than 12 nautical miles from the nearest land.

(3) When the garbage is mixed with other discharges having different disposal or discharge requirements the more stringent requirements shall apply.

(4) Reception facilities within special areas:

- (a) The Government of each Party to the Convention, the coastline of which borders a special area undertakes to ensure that as soon as possible in all ports within a special area, adequate reception facilities are provided in accordance with Regulation 7 of this Annex, taking into account the special needs of ships operating in these areas.
- (b) The Government of each Party concerned shall notify the Organization of the measures taken pursuant to sub-paragraph (a) of this Regulation. Upon receipt of sufficient notifications the Organization shall establish a date from which the requirements of this Regulation in respect of the area in question shall take effect. The Organization shall notify all Parties of the date so established no less than twelve months in advance of that date.
- (c) After the date so established, ships calling also at ports in these special areas where such facilities are not yet available, shall fully comply with the requirements of this Regulation.

Regulation 6

Exceptions

Regulations 3, 4 and 5 of this Annex shall not apply to:

- (a) the disposal of garbage from a ship necessary for the purpose of securing the safety of a ship and those on board or saving life at sea; or
- (b) the escape of garbage resulting from damage to a ship or its equipment provided all reasonable precautions have been taken before and after the occurrence of the damage, for the purpose of preventing or minimizing the escape; or
- (c) the accidental loss of synthetic fishing nets or synthetic material incidental to the repair of such nets, provided that all reasonable precautions have been taken to prevent such loss.

Regulation 7

Reception Facilities

- (1) The Government of each Party to the Convention undertakes to ensure the provision of facilities at ports and terminals for the reception of garbage, without causing undue delay to ships, and according to the needs of the ships using them.
- (2) The Government of each Party shall notify the Organization for transmission to the Parties concerned of all cases where the facilities provided under this Regulation are alleged to be inadequate.

APPENDIX B

FORM FOR REPORTING ALLEGED INADEQUACY OF PORT
RECEPTION FACILITIES FOR GARBAGE

1. Country
Name of port or area
Location in the port (e.g., berth/terminal/jetty)
Date of incident
2. Type and amount of garbage for discharge to facility:
 - a. Total amount (m³):
Food waste
Cargo-associated waste
Maintenance waste
Other
 - b. Amount not accepted by the facility (m³):
Food waste
Cargo-associated waste
Maintenance waste
Other
3. Special problems encountered:
Undue delay
Inconvenient locality of facilities
Unreasonable charges for use of facilities
Use of facility not technically possible
Special national regulations
Other
4. Remarks: For example, information received from port authorities
or operators of reception facilities, reasons for
nonacceptance (2.b above).
5. Ship's particulars:
Name of ship
Owner or operator
Distinctive number or letters
Port of registry
Number of persons on board

Date of completion of form

Signature of master

REDRESSING THE PROBLEM OF PERSISTENT MARINE DEBRIS
THROUGH LAW AND PUBLIC POLICY: OPPORTUNITIES AND PITFALLS

Michael J. Bean
Wildlife Program
Environmental Defense Fund
Washington, D.C. 20036, U.S.A.

ABSTRACT

This paper explores a variety of legal and institutional authorities for redressing the problems of persistent plastic debris in ocean environments. A major focus is on the recently ratified Annex V of the International Convention for the Prevention of Pollution by Ships (MARPOL) and its implementing legislation in the United States, the Marine Plastic Pollution Research and Control Act of 1987. The U.S. Coast Guard's new implementing regulations are described and critiqued, and important problems in the initial implementation of these new requirements analyzed.

The discussion of MARPOL and its initial implementation will introduce two related concerns. The first is the potential for solutions to the persistent debris problem to create an entirely different set of environmental problems. For example, the apparent intention of many shipping interests to turn to on-board incineration as the preferred means of complying with MARPOL's prohibition against the disposal at sea of plastics creates potential new problems in the form of hazardous air emissions and disposal of ash at sea. The limited legal authorities for responding to those potential problems under the Resource Conservation and Recovery Act, section 311 of the Clean Water Act, and other laws are described.

A further example is the fact that the enormous public attention to the problems associated with persistent plastic debris in the oceans and elsewhere has stimulated much interest in requiring "degradable" plastics before very much is known about the environmental hazards of the products of degradation and the impact of degradability upon efforts to recycle plastics. Some of the recently enacted laws and pending legislation relating to degradable plastics are reviewed.

The second consideration raised by the discussion of MARPOL is the difficulty of addressing the problem of marine plastic pollution as a problem separate from the larger problem of the

proliferating use of plastics generally. The paper concludes with an appraisal of opportunities to redress marine plastic pollution by public policy measures that touch upon that problem only indirectly in the context of designing sensible overall solid waste disposal programs. Some of the key public policy issues before Congress and local legislative bodies with important implications for plastics will be addressed.

INTRODUCTION

The first international conference on marine debris (Shomura and Yoshida 1985) was a major catalyst for a startling array of legislative and public policy initiatives to address a suddenly critical environmental problem that few had previously considered to be a problem at all. This paper examines some of the more important of these initiatives, pointing out their potential for reducing the marine debris problem, their limitations, and their possible pitfalls in terms of exacerbating other environmental problems. Finally, while the focus of the original symposium was on plastics in the marine environment, it has become increasingly clear that plastics present a variety of environmental problems, not simply in the marine environment, but elsewhere as well. As a result of this fact, this paper argues that the most significant benefits to the marine environment may come about as a result of measures that are aimed at addressing the broader set of problems for which plastics generally are responsible.

MARPOL ANNEX V

Undoubtedly, the most significant public policy advance since the 1984 conference pertains to Annex V of the International Convention for the Prevention of Pollution from Ships, better known simply as MARPOL. As a result of the subsequent ratification of MARPOL Annex V by the United States, the Soviet Union, and a number of other countries, that international agreement banning the disposal at sea of vessel-generated plastic waste came into force on 31 December 1988. In the United States, legislation to implement Annex V and, in certain respects, to impose duties beyond those of Annex V itself, was enacted in late 1987. Most of the duties imposed by that legislation, the Marine Plastic Pollution Research and Control Act of 1987 (MPPRCA), were to take effect concurrently with the coming into force internationally of MARPOL Annex V. However, the U.S. Coast Guard has thus far published only proposed regulations to implement the new legislation. Until final regulations are promulgated, the duties imposed by MARPOL Annex V and its U.S. implementing legislation are in a sort of regulatory limbo.

Before turning to the issues that have slowed the promulgation of implementing regulations, let us review first what MARPOL Annex V is intended to do. Put simply, MARPOL Annex V, in tandem with the London Dumping Convention, makes unlawful the deliberate disposal at sea of persistent plastics. The MARPOL Annex V picks up where the London Dumping Convention leaves off; the latter prohibits the dumping of plastics from

sources other than the routine operation of the vessel itself; MARPOL Annex V, on the other hand, restricts the disposal of vessel-generated wastes at sea, requiring that certain types of waste be disposed of beyond specified distances from land and prohibiting altogether the disposal at sea of other types of waste, including all plastics. Since, from time immemorial, it has been the nearly universal practice of ships at sea to dispose of their own wastes overboard, and since plastics represent a rapidly growing fraction of the shipboard waste stream, MARPOL Annex V has the potential to reduce a significant source of marine plastic debris. How significant this source is for the most serious environmental problems associated with marine plastic debris and how much of MARPOL'S potential to cut into this source will actually be realized are still very much matters of conjecture.

In retrospect, considering the dramatic changes that MARPOL Annex V was intended to effect in the long established practices of ships at sea, it is remarkable how little controversy attended the U.S. ratification of Annex V and enactment of domestic implementing legislation for it. No significant opposition to either measure was voiced at congressional hearings, there were no contested floor amendments offered in either house of Congress, and both houses approved the legislation by unrecorded voice vote, indicating the absence of any serious opposition or controversy. Either the interests that were to be affected by the legislation had carefully considered its potential impact and concluded that those impacts were acceptable, or those interests simply failed to take notice that legislation with potentially far-reaching consequences was zipping through Congress with uncharacteristic speed. The subsequent anxieties expressed by parts of the shipping industry during the course of Coast Guard rule making incline me toward the latter explanation.

The first noteworthy aspect of MARPOL Annex V is its extraordinary scope. Prior to the coming into force of Annex V, MARPOL affected only a few large ships, the oil and chemical tankers whose operations are subject to regulation under Annexes I or II, the mandatory MARPOL annexes. Annex V, on the other hand, applies literally to "all ships," save for warships and other government-owned or operated ships being used for government noncommercial service. Thus, not just major merchant vessels, but also commercial fishing vessels, cruise liners, recreational craft, and even rowboats and canoes are subject to Annex V's proscriptions. Moreover, the U.S. domestic implementing legislation goes beyond Annex V's already broad scope by requiring military and other government vessels to comply fully within 5 years after MARPOL enters into force (i.e., by 31 December 1993).

The domestic implementing legislation expands Annex V's geographic scope as well. Whereas Annex V applies only to ships at sea, the MPPRCA applies to vessels in U.S. internal waters as well. Both foreign and U.S. vessels are subject to the U.S. law while in navigable waters or the exclusive economic zone (EEZ) of the United States. United States vessels remain subject to the proscriptions of the U.S. law wherever they may be, even beyond the U.S. EEZ. With respect to penalties for violations, MARPOL itself requires only that these "be adequate in severity to discourage violations." The U.S. implementing legislation fixes very high maximum criminal and civil penalties for violations of Annex V; indeed, they are

the same as those for violations of Annex I or II involving oil or noxious liquid substances. A knowing violation of any of the MARPOL annexes, the requirements of the implementing legislation, or regulations adopted thereunder may be punished by a fine of up to \$50,000 per offense and up to 5 years in prison. A civil penalty of up to \$25,000 may also be assessed for any violation of the above.

Clearly, MARPOL Annex V and its implementing legislation in the United States represent, on paper at least, a major commitment to eliminating at least one source of marine plastic debris. How will it work in practice? MARPOL Annex V does not tell a vessel operator what he should do with ship-generated plastic waste--it only tells him that he cannot dispose of it in the ocean. The Coast Guard, charged with developing regulations to implement Annex V and the MPPRCA, has not sought through those regulations to steer vessel operators in any particular direction. Rather, it too leaves up to each individual operator the decision of what to do with plastic waste.

Implicitly, there is in Annex V a sort of presumption that the best solution is to off-load any plastic waste in port. This is because Annex V not only restricts the trash disposal practices of ships, but also requires that there be adequate reception facilities for garbage at ports and terminals. There is a rather obvious ambiguity here, in that most nonplastic garbage can still be disposed of at sea. Thus, it is unclear whether to be "adequate" a reception facility must be capable of handling only ship-generated plastic waste (the only waste that cannot be disposed of at sea), or whether it must be capable of handling the much larger volume of other garbage that might--or might not--be brought back to port.

On this issue, the Coast Guard's proposed regulations basically punt. The proposed regulations include a recommended "worksheet" for estimating the likely quantity of garbage that a port or terminal may be expected to handle, but then go on to disavow the likely accuracy of the formulas in the worksheet and any intention to rely upon those formulas in determining whether a port or terminal has met its obligation to provide adequate reception facilities. Indeed, the proposed regulations rely upon a system of self-certification, and require such self-certification for only a limited number of ports, principally those that already receive ships subject to Annexes I or II. Moreover, the proposed regulations do not even require that in certifying its reception facilities as adequate, a port or terminal identify what those facilities are.

VESSEL RESPONSE: INCINERATION AT SEA

Now, if you are a ship owner, you might at this point begin to get a bit nervous. The Coast Guard's proposed regulations were published on 27 October; Annex V was scheduled to come into force just 9 weeks later. As of that time it would no longer be lawful to dispose of plastic trash at sea, yet the availability of adequate reception facilities in port to handle any trash brought back to port had to be taken on faith. It was very clear that many of the larger shipping interests did not have that faith. In a number of public and private forums connected with the Coast

Guard rule making, two sentiments were very clearly expressed by these interests. The first was that 31 December 1988, was simply too soon to expect effective compliance with Annex V's new requirements for both ships and ports. The second was that in order to guard against the contingency that ports and terminals might in fact not have the ability to handle ship-generated garbage expeditiously or at all, larger ships would have to take care of their waste disposal problems by themselves by installing shipboard incinerators. It was rather remarkable, given the expense, potential danger, and generally primitive technology of shipboard incinerators, that installation of incinerators was the first alternative many shipping interests chose to explore, while reduction or elimination of optional plastics aboard ship seemed scarcely to have been considered.

It is unclear at this juncture just how much use of shipboard incineration will be stimulated by MARPOL Annex V, but it is very clear that it has stimulated a great deal of consideration of that option as a way of complying with the annex. Whether at-sea incineration of plastics represents a net environmental gain or merely solves one environmental problem by creating another is an open question. The Coast Guard, in its notice of proposed rule making, acknowledged that "proper disposal of incinerator ash has not been fully studied," but advised that for purposes of Annex V, ash is to be treated as operational waste that can be disposed of in the ocean beyond 3 nmi from shore. Rather more cautious advice can be found in the Guidelines for the Implementation of Annex V adopted by the Marine Environment Protection Committee of the International Maritime Organization in September 1988. Those guidelines, which are advisory only, state that the "ash from the combustion of some plastic products may contain heavy metal or other residues which can be toxic and should therefore not be discharged into the sea. Such ashes should be retained on board, where possible, and discharged at port reception facilities."

Even the more cautionary advice found in the guidelines probably understates the likelihood that incinerator ash will contain toxic heavy metals. About half of the tests that have been done of mixed bottom and fly ash, and virtually all of the tests of fly ash alone from municipal incinerators have found levels of lead or cadmium or both that exceed U.S. Environmental Protection Agency (EPA) criteria for designating such material as "hazardous waste" under the Federal Resource Conservation and Recovery Act (RCRA) (Denison and Silbergeld 1988). Plastics, some of which use lead or cadmium as stabilizers and colorants, are believed to be a major source of both of those heavy metals in municipal incinerator ash. Indeed, a recent study indicates that plastics account for 71% of the lead and 88% of the cadmium in the combustible portion of the municipal solid waste stream (Franklin Associates 1989).

The above percentages are all the more remarkable, given that plastics represent only about 7% (by weight) of the municipal solid waste stream. There is every reason to expect plastics to comprise an even greater percentage of the waste stream fed into an onboard incinerator, since it is only plastics that cannot otherwise be disposed of at sea. Thus, while lead and cadmium levels are generally high enough in municipal incinerator ash to be considered hazardous, even higher levels appear quite likely in

ash from shipboard incinerators. If that ash were brought to shore, it might have to be disposed of as hazardous waste under the RCRA. Since RCRA's reach does not extend beyond 3 nmi from shore, however, neither it nor MARPOL prohibits the disposal of the same ash directly into the sea. Section 311 of the Clean Water Act authorizes the EPA to prohibit the discharge of hazardous substances into the contiguous zone of the United States, but incinerator ash is not yet among the substances designated as hazardous under this authority.

Incineration of plastics at sea presents not only the problem of disposing of ash, but also that of controlling atmospheric emissions. Land-based incinerators are typically equipped with sophisticated technology to capture harmful flue gases and reduce particulate emissions. Emissions from land-based incinerators are closely regulated under authority of the Clean Air Act. Onboard incinerators, because of their smaller size, shorter stacks, and other limitations, are unlikely to be equipped with any sort of emission control equipment; in any event, inasmuch as vessels at sea are outside the scope of the Clean Air Act, they will not be required to control their emissions. Yet, the incineration of some plastics, in particular polyvinyl chloride, produces highly toxic, corrosive gases.

Whether widespread conversion to at-sea incineration of plastics will create a new set of environmental problems is open to debate. What is clear is at least the potential for solutions to one problem to become themselves the source of another problem. Just as the long distance transport of acid rain-causing sulphur oxides was the unforeseen result of building higher and higher emission stacks to reduce local air pollution problems, so too might some of the solutions to the environmental problems of plastic debris cause other, largely unforeseen problems.

DEGRADABLE PLASTICS

Let me illustrate this with a discussion of a very hot topic-- "degradable" plastics. The concern with the problem of marine plastic pollution has helped trigger what can only be described as an avalanche of interest among public policy makers in the subject of degradable plastics. The MPPRCA of 1987 does not merely implement MARPOL Annex V. It also directs EPA to carry out a study of the adverse environmental effects of plastics generally, and to evaluate the feasibility of making products that present a particular hazard to the environment from "degradable plastics materials." Less than a year later, in October 1988, Congress passed another law, Public Law No. 100-556, requiring EPA, within 2 years thereafter, to issue regulations requiring that "plastic ring carriers" be made of "naturally degradable material" unless doing so is infeasible or will result in byproducts of degradation that present a greater threat to the environment. Only a month earlier, Congress included in the Defense Department Authorization Act, Public Law No. 100-456, a provision (section 352) directing the Secretary of Defense to report to Congress by 1 March 1990 his recommendations concerning the substitution of "degradable plastic items for nondegradable plastic items" used by the military. The Senate also passed Senate Resolution 412, a nonbinding resolution

expressing the sense of the Senate that EPA should encourage the use of biodegradable plastic bags and other items through its regulatory and informational programs.

In addition to these enacted measures, there were also a number of other bills introduced in the last Congress that would have mandated either the study of, or use of, degradable plastics for a variety of purposes. In the new Congress, interest will almost certainly be even higher. At the state level, degradable plastics are required for beverage ring carriers in at least 17 states and for still other products in a number of states. Corn-growing states in particular have an interest in pushing degradable plastic requirements, since at least some of the technologies for producing degradable plastic utilize corn starch as the ingredient that imparts degradability. Thus, it came as no great surprise to me to learn recently that the incoming chairman of the National Governor's Association, the Governor of Iowa, has degradable plastics near the top of the environmental agenda that he wants the association to pursue.

The rush to impose degradable plastic requirements has far outpaced any reasonable understanding of the extent to which degradable plastics are actually likely to contribute to the solution of current environmental problems and the likelihood that they might exacerbate others or create altogether new ones. Some of the promoters of degradable plastics have argued that products that might otherwise entangle marine animals will be so weakened that they will be readily broken. One seldom hears acknowledged, however, that those same products, and others that never presented any entanglement threat, may, by virtue of their gradual breakdown into multiple fragments, be far more likely to contribute to the problem of ingestion than they would have done had they remained intact. As for the chemical products of degradation, the discussions to date have rarely included mention of the fact that heavy metals and other toxic chemicals are commonly used as plasticizers, stabilizers, catalysts, and colorants in a wide variety of plastics. While a plastic product remains intact, these are relatively inaccessible to the environment. As that product degrades into ever smaller pieces and shorter polymer chains, however, these same chemicals are likely to become much more accessible to the environment. My point is not to say that, on balance, degradable plastics represent a greater environmental threat than nondegradable plastics; rather, it is only to say that no one yet knows what trade-offs are involved, yet the rush by many legislators to embrace degradable plastics has nearly become a stampede.

If degradable plastics are an unlikely panacea for the problem of marine plastic pollution, what other measures beyond MARPOL itself offer some hope of redressing the problem? The limitations of MARPOL Annex V are apparent. Even assuming widespread voluntary compliance with, and effective enforcement of, its prohibitions--and such assumptions are not easily indulged--MARPOL Annex V at best only addresses a fraction of the problem. It does not touch at all the problem of lost or damaged fishing gear, or marine plastic waste that can be traced back to beach litter, storm water runoff, sewage disposal, spills associated with marine transfer of municipal waste, factory discharges into inland waters and estuaries,

and other essentially land-based sources. All water runs downhill, and at the bottom of all those hills lies the ocean. Add to this diversity of sources of marine plastic pollution the fact that the incredible penetration of the market by more and more plastic products continues unabated, and it is not difficult to look upon the task of protecting the oceans from plastic debris as a job suited for Sisyphus.

PLASTICS IN THE SOLID WASTE STREAM

What public policy alternatives exist for making additional inroads into the marine plastic pollution problem? At least part of the answer, it seems likely, will derive from the eventual recognition that the problem of marine plastic debris is but one facet of a much larger set of problems stemming from the growing abundance of plastics in the solid waste stream.

Plastics present a number of unique problems, not just for living marine organisms, but for human communities struggling with the growing solid waste crisis. As already noted, plastics are primary contributors of some of the heavy metals in municipal incinerator ash that may cause that ash to be treated--and very expensively--as hazardous waste. In addition, the very low level of plastics recycling frustrates effective solid waste management programs.

Increasingly, many communities are recognizing that the most economical and environmentally sound response to the growing solid waste problem is to include an aggressive recycling component in their solid waste management programs. Yet, plastics in the waste stream are, by and large, replacing the very glass and metal containers that enjoy some of the highest recycling rates. Pressed on one side by the prospect of diminishing landfill capacity and on the other by the high cost of building and operating incinerators, many states and local governments are enacting laws that prescribe some minimum level of recycling of the overall municipal solid waste stream--typically at least 25% and sometimes twice that. Achieving those prescribed rates will be difficult unless either the amount of plastic in that stream is limited or the extent of plastics recycling dramatically increases. A number of communities are already trying to accomplish the former by prohibiting certain types of plastic packaging or by imposing differential taxes on certain products, with a higher tax on those made from materials not readily recycled. Suffolk County, New York, is an example. Its law prohibiting plastic bags and certain fast food packaging is serving as something of a test for how far local communities can go in banning plastic products. The plastics industry does not take this lightly. It has challenged the Suffolk County law in a New York trial court in a case entitled *Society of the Plastics Industry v. County of Suffolk*; the Environmental Defense Fund and the Natural Resources Defense Council have intervened in support of the county law.

Laws like that of Suffolk County are likely to become increasingly common unless dramatic strides in the recycling of plastics are made very soon. Of the more than 9 million MT (20 billion lb) of plastic in the municipal solid waste stream in 1986, only 68,000 MT (150 million lb), or

<1%, was recycled (EPA 1989). By contrast, more than 26% of paper and paperboard, and more than 50% of aluminum cans are recycled (EPA 1989). Virtually the only plastic product with a more than negligible recycling rate is the PET (polyethylene terephthalate) soft-drink bottle. However, returnable deposit legislation, which currently exists in only 10 states, represents the only significant means through which plastic bottles are currently being collected for recycling (EPA 1989). In Michigan, which has a required 10 cent deposit, the recovery rate is nearly 90%. In states without mandatory deposit legislation, there is virtually no recycling of plastic bottles. Thus, if the plastics industry wanted to give a real boost to recycling, it would break ranks with the bottling industry and support state mandatory deposit legislation.

In the long run, the amount of plastic debris in the marine environment is likely to be a function not merely of the waste disposal practices of vessels at sea, but of the laws and public policy measures that influence how much further growth of the plastics market will occur and how plastics in the solid waste stream generally are treated. Measures to mandate or encourage reusable or recyclable products, to limit product packaging, or to encourage nonplastic alternatives may ultimately contribute as much or more to the solution of the marine plastic debris problem as measures aimed directly at marine industries.

REFERENCES

- Denison, R. A., and E. K. Silbergeld.
1988. Risks of municipal solid waste incineration. *Risk Analysis* 8(33).
- Environmental Protection Agency.
1989. Draft report to Congress under the Marine Plastic Pollution Control Act.
- Franklin Associates, Ltd.
1989. Characterization of products containing lead and cadmium in municipal solid waste in the United States, 1970 to 2000, a report to the Environmental Protection Agency.
- Shomura, R. S., and H. O. Yoshida (editors).
1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

IMPLEMENTATION AND ENFORCEMENT OF ANNEX V
OF MARPOL 73/78 IN THE UNITED STATES

Raymond E. Seebald
Commandant (G-MPS-1)
U.S. Coast Guard Headquarters
Washington, D.C. 20593, U.S.A.

ABSTRACT

A new, bold international law entered into placed on 31 December 1988 in response to the threat to the marine environment posed by marine debris. Annex V of MARPOL 73/78, as it is known, is the first international agreement to attempt to reduce the amounts of plastics and other garbage waste discharged into the oceans from vessels. The U.S. Coast Guard's role was to interpret broad international guidelines and directives by Congress as stated in the Marine Plastic Pollution Research and Control Act of 1987, and develop practical enforceable regulations. These regulations which were developed reflect both the Coast Guard's expertise in enforcing Annexes I and II of MARPOL 73/78, and the numerous comments received from industry and the environmental and public sectors. The resulting performance-based regulations left the details of compliance up to vessels, which were required to restrict their discharge of garbage, and to the terminals and ports, which were required to provide reception facilities for the garbage wastes. Vessels have employed numerous different strategies to comply with the regulations. These strategies range from different operational methods to the installation of large garbage-treating equipment, based on the route, passenger load, type of trade, and size of vessel. Port facilities have applied an equally diverse response to compliance.

The Coast Guard's enforcement of these regulations involves a multiagency effort. The Department of Agriculture and the National Marine Fisheries Service, NOAA, conduct enforcement in areas of their own influence and expertise. The Coast Guard expects that these enforcement strategies together with educational efforts by local captains of the port should result in a high compliance rate on the part of the shipping community.

THE COAST GUARD'S ANNEX V COMPLIANCE REPORT: A CASE STUDY

Carl A. Crampton
U.S. Coast Guard (G-MPS-3)
Washington, D.C. 20593, U.S.A.

ABSTRACT

This case study documents the U.S. Coast Guard process for accomplishing the congressional mandate for a compliance report concerning the implementation of Annex V (Regulations for the Prevention of Pollution by Garbage from Ships) of MARPOL 73/78. Annex V was implemented by Title II of Public Law 100-220, the Marine Plastic Pollution Research and Control Act of 1987.

The study follows the development of an empirical methodology for assessing compliance with new regulations that implement Annex V. The goal is a report that measures the amount of Annex V garbage being brought ashore both before and after Annex V became effective. As a secondary objective, the Coast Guard would like to assess the effects of Annex V regulations on the coastal and marine environments. The project involves an evaluation of current information and its statistical validity as a preimplementation baseline. The evaluation is then used to develop a baseline estimate for the amount of Annex V garbage being generated. Once that information has been collected, a methodology for measuring Annex V compliance and for assessing the effects of Annex V regulations on the coastal and marine environments will be developed.

The Coast Guard must follow specific congressional intent for the compilation of this report. Congress required in section 2201, Title II of PL 100-220, that within 1 year from the date of enactment of the act, and every 2 years thereafter for a period of 6 years, the Coast Guard, in consultation with the Secretary of Agriculture and the Secretary of Commerce, report to them on compliance with Annex V in U.S. waters, including the waters of the U.S. exclusive economic zone. The report is to include a description of the enforcement mechanisms in place and an assessment of the need for additional enforcement authority. It must also address the extent to which garbage reception facilities have been made available at the ports, and the mechanisms used by the Coast Guard to ensure that these facilities are made available as required by Annex V.

In preparing this report, the Coast Guard must assess the extent to which vessels dispose of floatable dunnage materials beyond 25 nmi from shore and the extent to which they wash ashore, and recommend whether Annex V should be amended to prohibit the disposal of these materials at sea. The report must also include a detailed assessment of the fines levied by foreign nations for violations of Annexes I, II, and V committed by foreign flag vessels in the U.S. exclusive economic zone. Finally, the report must summarize the education efforts undertaken to inform the public of the problem of pollution of the marine environment by improperly disposed garbage.

HISTORY OF ENACTMENT

MARPOL 73/78

The International Convention for the Prevention of Pollution from Ships, 1973, as amended by the Protocol of 1978 (MARPOL 73/78), established baseline levels of practice in attempting to mitigate or eliminate damage to the environment by pollution from vessels. The MARPOL 73/78 is built of five annexes. Each annex is concerned with preventing pollution of the world's oceans by a different product or group of products. Annex I is concerned with preventing oil pollution, Annex II with the noxious liquid substances or chemicals, Annex III with the release of packaged hazardous materials, Annex IV deals with the prevention of pollution by sewage, and Annex V with the prevention of pollution by garbage. Annexes I and II are mandatory for signatories, while Annex III, IV, and V are optional, only becoming effective 1 year from the date when at least 15 nations have ratified them. The 15 nations must represent a cumulative total of 50% of world shipping tonnage.

The Act to Prevent Pollution From Ships

Domestic legislation was needed to implement the convention and its annexes in the United States. The MARPOL 73/78 was incorporated into U.S. law when the Act to Prevent Pollution from Ships was passed by the Congress of the United States in 1980. (For a more complete description of this process, see Whitehead 1988.) The Act to Prevent Pollution from Ships was codified in Title 33, United States Code, Sections 1901-1910.

The Marine Plastic Pollution Research and Control Act of 1987

Scientific and environmental forums during the 1970's and 1980's highlighted the amount and impact of garbage from ships on the world's oceans. Although ocean disposal of any type of garbage presents the potential for environmental damage, plastics seemed to be the most immediate problem. The characteristics of plastics which make them useful: lightweight, strong, and persistent, make them deadly to the ocean's biota. Title II of Public Law 100-220, the Marine Plastic Pollution Research and Control Act (MPPRCA) of 1987, implemented Annex V of MARPOL 73/78. Annex V prohibits the discharge of any plastic materials from ships at sea, limits the locations

where other garbage may be discharged, and requires signatories to provide reception facilities for the discharge of ships' garbage. Title II includes additional requirements for Annex V's implementation in the United States. One of those requirements is a report on compliance with the new law's requirements.

COMPLIANCE REPORT REQUIREMENTS

Provisions in the Law

Congress mandated in the law that "within 1 year after the effective date of this section, and biennially thereafter for a period of 6 years, the Secretary of the Department in which the U.S. Coast Guard is operating, in consultation with the Secretary of Agriculture and the Secretary of Commerce, shall report to the Congress regarding compliance with Annex V to the International Convention for the Prevention of Pollution from Ships, 1973, in United States waters" (P.L. 100-220).

The Congressional Record elaborated on this reporting requirement (U.S. Congress 1987). The report is required to include a description of the enforcement mechanisms in place and an assessment of the need for additional enforcement authority. It is to address the extent to which garbage reception facilities have been made available at the ports, and the mechanism used by the U.S. Coast Guard to ensure that these facilities are made available as required by Annex V. In preparing this report, the Coast Guard is to assess the extent to which vessels dispose of floatable dunnage materials beyond 25 nmi from shore and the extent to which they wash ashore, and to recommend whether Annex V should be amended to prohibit the disposal of these materials at sea. The report is to provide a detailed assessment of the fines levied by signatory nations for violations of Annexes I, II, and V by their vessels in the U.S. exclusive economic zone. Finally, the report is required to summarize the education efforts undertaken to inform the public of the problem of pollution of the marine environment by garbage that is improperly disposed of.

The Coast Guard is to solicit the advice of interested parties, including the shipping industry, ports, the commercial fishing industry, environmental groups, waste handling firms, and recreational boaters, in preparing this report. The input of industry and environmental groups is to be solicited through existing forums such as the Workshop on National Marine Pollution Problems and Needs conducted by the National Marine Pollution Planning Office of the U.S. National Oceanic and Atmospheric Administration (NOAA); the Marine Debris Roundtable, a national discussion group informally established by NOAA; the National Committee for the Prevention of Pollution, a subcommittee of the Shipping Coordinating Committee, which meets in advance of meetings of the Marine Environmental Protection Committee of the International Maritime Organization; and the National Boating Safety Advisory Council, an advisory committee to the Coast Guard that considers matters affecting recreational boating.

Completing the Feedback Loop

The Coast Guard wants to present a complete picture to Congress in the report. The current report requirements include information on violations, facilities available, and other enforcement information. However, it does not require relative measures against which enforcement data can be viewed. The Coast Guard compiled information on various vessel classes and their contribution to the garbage load in the oceans as a required portion of the regulatory process. It is logical that the report should examine compliance by the different vessel classes already identified. Compliance and regulatory effectiveness are related, and the information would be valuable to this and other Coast Guard regulatory efforts.

The regulation's effect on the environment is another item selected for examination by the report. The law's intended result is the elimination of ship-generated plastics from the waters of the United States, and a reduction in ship-generated garbage. Measurements of amounts of plastic and other garbage in the marine environment seem a necessary part of a compliance report. The effects of the law will be seen by evaluating compliance and environmental effects.

INITIAL PARAMETERS

Constraints

The plan for the required and desired portions of the report filters through a screen of constraints. The first of these is that no additional funds or other resources were allocated by Congress for implementation, enforcement, or for reporting compliance. The compliance report is to be developed using such resources as can be reprogrammed within the Coast Guard. What makes this constraint more difficult to work with is the routine shortage of Coast Guard operating funds. The lack of resources is amplified by any regulatory project's ability to compete within the Coast Guard's needs for operational missions. This is particularly true when those missions include drug interdiction efforts and Maritime Defense Zone planning. Another constraint is the internal human resources available to complete the task. Those responsible for developing the regulation and enforcement plans cannot devote the necessary time to the project without seriously impairing their ability to perform other necessary functions. A final constraint is the lack of information on the amounts, constituents, and sources of garbage in the waters of the world. Estimates and inferences are all that are currently available.

Needs

The report will need baseline information from which to estimate compliance. Knowledge of the amount of garbage, especially plastic garbage, presently in the ocean is necessary to gauge the effectiveness of the regulatory program. The effects of plastic and other garbage on the environment prior to the regulation will also have to be assessed. Measurements of current compliance and garbage generation by vessel types need to be made or estimated if trends are to be analyzed.

The Coast Guard is interested in assistance in developing the feedback-loop items necessary for placing in perspective the measures of compliance that Congress had identified. Questions concerning data collection needs and possible measures of effectiveness need to be answered. The report's time constraints make additional resource input necessary. In this instance the resource is expertise in evaluative design.

WORK PLAN

Need for Action

The Coast Guard is utilizing an incremental approach for the reporting process because of the short time allotted for the entire regulatory package, including the compliance report. The incremental approach allows general planning for completion of the regulatory project, while permitting changes in detail if new techniques or resources are identified. The method for compilation of the report has not been determined, but the end points for completion, as well as resource parameters for efforts to complete the report, are known. The Coast Guard began by identifying areas in which additional information and expertise were needed. It was decided that these areas should be addressed first. Members of the Transportation System Center (TSC), U.S. Department of Transportation, were retained for this effort.

Initial Studies

The first phase involved the search for existing information and studies. The Coast Guard was especially interested in data on the quality of the environment prior to enactment of the MPPRCA of 1987. However, it was also important to identify sources of empirical data for follow-on studies. Continuing information-gathering efforts or studies would be highly beneficial, as the data gathered would continue through the period in which the Coast Guard was interested.

The information search identified entities actively involved in surveying ocean surface debris and beach litter. Fifty organizations were identified and each was contacted by TSC representatives. Ongoing research efforts were separated and further disaggregated into one-time or continuing efforts. Twelve of the fifty organizations contacted actually had conducted ocean or beach surveys. Five of the twelve that had conducted surveys were doing so on a continuing basis. These five activities appeared to be good sources for data concerning the effects of the new regulations on the environment (U.S. Department of Transportation 1988b).

Information was also needed on the amounts of garbage being generated and prereregulation vessel garbage disposal practices. Eastern Research Group had developed much of this information for the Coast Guard during the regulatory development process as part of the regulatory evaluation (U.S. Department of Transportation 1988c). The estimated garbage generation rates for various vessel types are included in Table 1.

Table 1.--Estimated annual quantities of domestic garbage generated by vessels operating in U.S. waters.

Source	Tons	Percentage of total
Merchant marine	33,574	1.74
Commercial passenger vessels	283,881	14.71
Commercial fishing	256,494	13.29
Recreational boating	1,264,114	65.52
Offshore oil and gas industry	18,381	0.95
Miscellaneous vessels	1,801	0.09
U.S. Navy	63,356	3.28
U.S. Coast Guard	6,782	0.35
U.S. Army	539	0.03
National Oceanic and Atmospheric Administration	349	0.02
Total	1,929,271	100.00

It was apparent, even allowing for errors based on incomplete data, that recreational boating and commercial passenger and fishing vessels create most of the domestic waste (U.S. Department of Transportation 1988a). The garbage disposal practices for different classes of vessels were also estimated and are displayed in Table 2.

It was concluded that recreational boating, commercial fishing, and the U.S. Navy were the leaders in the amount of waste dumped overboard.

METHODOLOGY DEVELOPMENT

Requirements

The most difficult task in planning the report's completion is development of a methodology for measuring compliance with the law and the effects of the new regulations on the coastal and marine environment. Most other portions of the report are straightforward, requiring only data acquisition and formatting. Measuring compliance is another matter. The TSC began with given parameters for developing the methodology. The methodology is to rely on information already being collected by the Coast Guard or other governmental and nongovernmental agencies. The information for measuring compliance should be updated regularly, preferably on an annual basis. In the event that existing data sources and surveys are not adequate to this task, TSC is to recommend ways to acquire the appropriate data. Measurements of the effect of the law on the environment are to use only current data collection and survey efforts.

Table 2.--Estimated amount of vessel-generated domestic garbage disposed of at sea by vessels operating in U.S. waters.

Source	Tons	Percentage of total
Merchant marine	30,493	3.76
Commercial passenger vessels	27,846	3.43
Commercial fishing	256,494	31.63
Recreational boating	421,371	51.96
Offshore oil and gas industry	6,574	0.81
Miscellaneous vessels	1,796	0.22
U.S. Navy	63,356	7.81
U.S. Coast Guard	2,059	0.25
U.S. Army	539	0.07
National Oceanic and Atmospheric Administration	349	0.04
Total	810,877	100.00

Measuring Compliance

Measuring compliance appears to be a fairly simple task. First, find out how many entities are in the regulated population. Then ascertain what portion of the regulated population is obeying the statute. The task is complicated in this instance by the lack of available or collectable data.

The Coast Guard is the primary Federal agency source for information on vessels in the waters of the United States. However, data collection efforts are related directly to legal requirements for maintaining that information. The Coast Guard has a spectrum of information on many types of vessels. Most of the information collected by the Coast Guard is on vessels that are inspected regularly for structural, electrical, safety, and operational requirements. However, Eastern Research Group's estimate is that this population generates less than 20% of the total garbage load entering the waters of the United States. The Coast Guard collects some information on recreational boating, estimated to be the largest garbage contributor. This information is collected through random boardings by Coast Guard regular and reserve personnel and by the Coast Guard Auxiliary. However, less information is collected on recreational vessels than on commercial vessels, and collections are not made on a regular basis. The Coast Guard has virtually no contact with commercial fishing vessels, which are estimated to generate approximately 14% of the total garbage load. Finally, the estimate for the U.S. Navy's contribution to the problem made it the fourth largest contributor. The Coast Guard had no data on or regular contact with the U.S. Navy vessels.

Measurement of the amounts of garbage deposited at ports in the United States before and after implementation of the regulations is likewise hindered by the lack of data. It is apparent that current data collection efforts have to be expanded and other sources of data identified if compliance is to be adequately measured.

Recommended Approach

The study by TSC identified methods by which current data collection shortfalls for measuring compliance with the regulations could be corrected (Department of Transportation 1989). The easiest to address are the data collection efforts associated with commercial vessels currently required to be inspected on a regular basis. Additional data collection would involve requirements associated with the implementation of Annex V. The Coast Guard could also include items associated with Annex V in its examinations of port and terminal facilities.

More information on recreational boating compliance could be obtained by modifying the data collection requirements of the random boarding program to include Annex V requirements. The Coast Guard Auxiliary could also gather information on the level of compliance observed during the courtesy marine examinations that they conduct.

The compliance levels of commercial fishing vessels could be determined by establishing a cooperative effort with the National Marine Fisheries Service (NMFS) for inspecting garbage and fishing net pieces brought in to shore when fishing vessels are off-loading their catches. This would not require an expansion of manpower efforts on NMFS' part.

The compliance level could also be better estimated by conducting in-depth studies of selected ports and vessels on a voluntary basis to observe choices of compliance methods, problems encountered, improvements, and port/vessel interactions.

CURRENT STATUS

The Coast Guard is now processing the recommendations for action. The report is to be developed by a contractor selected through the competitive bidding process. Those entities that are currently conducting studies and surveys are being contacted to ascertain how their information may best be utilized. The Coast Guard is acting internally to effect the changes necessary to collect the additional information needed to measure compliance within the various vessel categories. The NMFS is being approached to ascertain their ability to collect data on fishing vessels returning to off-load their catch. The effort to compile the report has been an interesting exercise in measuring compliance with an environmental regulation that affects a wide range of people. Its success in aiding regulatory efforts will be important, as will the lessons learned in attempting to measure those efforts.

The opinions or assertions contained herein are those of the author and are not to be construed as official or reflecting the opinions of the

Commandant of the Coast Guard, the Chief Counsel of the Coast Guard, or the Coast Guard at large.

REFERENCES

- Public Law 100-220.
1987. Title II. The Marine Plastic Pollution Research and Control Act of 1987.
- U.S. Congress.
1987. House of Representatives, House Report No. 100-360 to accompany H.R. 940, 100th Congress, 1st Session, 39 p.
- U.S. Department of Transportation.
1988a. Development of estimates of garbage disposition in the maritime sectors. U.S. Department of Transportation, U.S. Coast Guard, Transportation Systems Center, submitted by Eastern Research Group, Inc., Cambridge, MA, 139 p.
- 1988b. Information: CG-849 Annex V compliance measurement subtask 3C.1 results: Search for information on surveys of ocean debris and beach litter. U.S. Department of Transportation, Research and Special Programs Administration memorandum, 10 p.
- 1988c. Regulatory evaluation on regulations implementing Annex V to MARPOL. U.S. Department of Transportation, U.S. Coast Guard, submitted by Eastern Research Group, Inc., Wash., D.C., 334 p.
1989. Measurement of compliance with MARPOL Annex V regulations. U.S. Department of Transportation, Transportation System Center, Wash., D.C., 34 p.
- Whitehead, J. R.
1988. Reducing plastic pollution in the marine environment: The U.S. Coast Guard and implementation of Annex V of MARPOL 73/78. Oceans '88 Conference Proceedings 4:1507-1512.

CONTROL REGULATIONS FOR DISCHARGE OF ONBOARD WASTES FROM SHIPS

Junichi Sawaki*
Office of Marine Pollution Control and Waste Management
Water Quality Bureau, Environment Agency
Chiyoda-ku Tokyo, Japan

and

Ichiro Yamaguchi
Environment Division, Transport Policy Bureau
Ministry of Transport
Chiyoda-ku Tokyo, Japan

*Present address: Patrol vessel *Motobu*, 11th Regional Maritime Safety Headquarters, Maritime Safety Agency, 2-11-11 Minatomachi, Naha-city, Okinawa Prefecture, Japan.

ABSTRACT

In Japan, the disposal into the sea of waste from ships is prohibited in principle, except for its disposal in accordance with discharge standards or under emergency conditions as set forth in the Law Relating to Prevention of Marine Pollution and Maritime Disaster. Especially, the disposal of any form of plastics, including but not limited to synthetic ropes, synthetic fishing nets, cups, and bags, is not permitted. These provisions of the law apply to all ships.

INTRODUCTION

Japan introduced regulations on waste discharge from ships by the Law Relating to the Prevention of Marine Pollution and Maritime Disaster (1970). The law went into effect in 1972, when sea areas of discharge and methods of discharge were specified. It was revised in 1988. Annex V of MARPOL 73/78 went into effect on 31 December 1988.

Wastes generated on board ship are classified in two categories: Wastes from the daily life of crews, and wastes arising from the ordinary operation of ships (e.g., fishing and dredging). Regarding wastes from the crews' daily life, the regulations are equivalent to those of Annex V, and the regulations for wastes generated during ordinary operation are in general more stringent than the annex.

OUTLINE OF ANNEX V TO MARPOL 73/78

The garbage subject to regulations under Annex V to MARPOL 73/78 is all kinds of victual, domestic, and operational waste, excluding fresh fish and parts thereof, generated during the normal operation of the ship and liable to be disposed of continuously or periodically. Excluded are those substances which are defined or listed in other annexes to the present convention. As to garbage disposal outside special areas, sea areas of discharge and methods of discharge are outlined for each category of garbage in Annex V. Regulations of discharge are the toughest on plastics, including fishing nets, rope, and nylon bags. Their discharge is prohibited in all areas. Dunnage, lining, and packing materials which will float must not be disposed of into the sea <25 nmi from the nearest land. Food wastes and all other garbage must not be discharged <12 nmi from land. However, they can be discharged within 3 to 12 nmi of land if they are <25 mm in size. Any garbage generated on board must not be discharged <3 nmi from land. All ships are subject to the regulations irrespective of their type, size, or usage, even a one-man pleasure boat.

HISTORY OF DOMESTIC REGULATIONS

Japan has the Law Relating to the Prevention of Marine Pollution and Maritime Disaster (hereinafter referred to as "the marine pollution prevention law") (see Appendix A) to preserve the marine environment and protect the life, health, and assets of the people. The marine pollution prevention law in principle prohibits discharge of garbage into the sea unless it is required for emergency evacuation or force majeure and unless certain discharging standards are met. Substances subject to discharge regulations are oil, noxious liquid substances, and other garbage generated on ships, maritime facilities, and aircraft. Japan established the Law Relating to the Prevention of Maritime Pollution by Oil from Ships in 1967 to make 54 OILPOL, including a 1962 amendment, into domestic law. This law was replaced by the marine pollution prevention law in 1970 to include a 1969 amendment to 54 OILPOL, regulate discharging of garbage from ships and of oil and garbage from marine facilities, and implement measures for preventing maritime pollution by oil and other substances. In 1976, obligations to deploy oil-skimming ships, measures for prevention of maritime disaster, and the establishment of a maritime disaster prevention center were added to the law. At the same time, it was renamed the Law Relating to the Prevention of Marine Pollution and Maritime Disaster in order to specify prevention of maritime disaster as well as pollution.

In response to the London Dumping Convention, Japan revised the marine pollution prevention law in 1980 to toughen regulations by establishing controls on waste discharge from aircraft and incineration at sea, and a waste discharge confirmation system. Japan also modified the marine pollution prevention law in 1983 to subject not only heavy oil but also light oil, noxious liquid substances in bulk, sewage, and other garbage to controls, introduce controls on structures and equipment of ships, and conduct inspection regarding the new controls in order to abide by MARPOL 73/78.

Japan has in this way gradually enhanced regulations for prevention of maritime pollution in response to an international trend and domestic public opinion, and is ready to further toughen and expand such regulations in the future.

CONTROL OF DISCHARGE OF WASTES

Japan's marine pollution prevention law covers both Annex V to MARPOL 73/78 and the London Dumping Convention to regulate discharge of wastes. Annex V regulates garbage generated offshore, while the London Dumping Convention subjects wastes generated on shore to controls. The scope of garbage for Annex V is different from that for the London Dumping Convention, but the marine pollution prevention law unifies regulations under both. It prohibits discharge of wastes from ships in principle, and allows such discharge for emergency evacuation or force majeure and under certain conditions. The marine pollution prevention law defines "wastes" as leavings or rubbish people do not require, excluding oil, noxious liquid substances, and the like. "Discharge" is defined as any action to set afloat or drop matter into the sea. Japan has the same regulations as those under Annex I to MARPOL 73/78 on oil and under Annex II on noxious liquid substances. The marine pollution prevention law thus regulates all wastes other than oil, noxious liquid substances, and the like. It will eventually cover garbage under Annex V to MARPOL 73/78, sewage under Annex IV, and wastes generated on shore under the London Dumping Convention. The law includes waste-discharging standards in accordance with the London Dumping Convention. Hereinafter, we discuss regulations responding to Annex V under the revised marine pollution prevention law put into effect 31 December 1988 and Japan's original controls on discharge of sewage since 1972. Annex IV to MARPOL 73/78, which deals with sewage, has not taken effect, and regulations under this annex have yet to be implemented internationally.

Garbage generated on ships is divided into two categories. One is related to people's daily life and includes trash and sewage. Another covers garbage linked to routine operations of ships. Human life-related trash and sewage are limited in volume and mostly disposable in the sea. Therefore, only on large ships had such garbage been subjected to regulations on discharge. Until Annex IV and V to MARPOL 73/78 take effect, Japan will continue to separate trash from soil in discharging regulations to satisfy the respective annexes. Regulations on human life-related soil have remained unchanged even after Annex V to MARPOL 73/78 went into effect. Japan's regulations on trash were less stringent than those stipulated in the annex, and Japan revised the marine pollution prevention law to introduce regulations conforming to the annex for all ships irrespective of size or type. Under the new regulations, (1) discharge of wastes is prohibited within a distance of <3 nmi from the baseline for territorial waters, (2) discharge of plastics is totally prohibited, but those burned to powder ashes can be discharged at and beyond a 3-nmi distance from the territorial water baseline, and (3) garbage, whose size must be reduced to <25 mm before discharging under MARPOL 73/78, must be burned to ashes or pulverized by machines meeting certain technical standards before discharge. A type certificate system is established for

the pulverizing machines. These machines are required (1) to reduce the size of any garbage put into them to <25 mm, (2) to perform normal functions despite shaking or vibrations, and (3) to be easy to maintain and clean.

"Ship" is defined under MARPOL 73/78 as a vessel of any type whatsoever operating in the marine environment, and includes hydrofoils, air-cushion vehicles, submersibles, floating craft, and fixed or floating platforms. The ship as defined in the convention thus covers both what is defined as ship and what is defined as maritime facility under the Japanese marine pollution prevention law, so that Japan has subjected both ships and maritime facilities to the same discharging regulations. A ship under the Japanese law is defined as any floating craft used for navigation in the sea. A maritime facility is defined as any structure installed in the sea to house people, treat things, transport things, or stockpile things. It excludes structures which are linked to the shore through fixed facilities for free traffic of people and those connected to the shore for discharging oil or other wastes from the shore. As to garbage generated in relation to routine ship operations, including transportation and fishing, discharging regulations under the Japanese marine pollution prevention law were already tough and almost satisfied requirements under Annex V to MARPOL 73/78. The tough regulations already limited garbage for conditional discharge to what must be disposed of in the sea and prohibited discharging of garbage other than animal waste within 50 nmi of the territorial water baseline. Therefore, Japan made only minor changes in respect to garbage generated as a result of routine ship operations. Regulations on plastics were made as tough as those on human life-related plastic trash, and a distance for prohibiting the discharge of garbage other than animal waste was cut to <12 nmi from the territorial water baseline to conform to Annex V to MARPOL 73/78.

As to garbage generated on maritime facilities in relation to their usual operations, a prohibition against discharging had already been established under the law before the latest revision and has been retained. The garbage discharging standards under the Japanese law were different from those under MARPOL 73/78, and the difference has remained even since Annex V to the convention took effect. But the overall Japanese standards sufficiently fulfill all the requirements under the convention. Furthermore, the Japanese marine pollution prevention law includes independent regulations on discharging of garbage generated on aircraft during flights.

Regulations for discharging garbage generated on ships, maritime facilities, and aircraft under the marine pollution prevention law follow.

CONTROL OF DISCHARGE OF WASTE FROM SHIPS

Daily Life-Related Garbage

Daily life-related garbage is garbage generated in relation to daily life of seamen and other persons on board ships.

Waste Plastics

Sea areas of discharge.--Waters at and beyond 3 nmi from the territorial water baseline.

Method of discharge.--The garbage must be transformed into ashes before discharging. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Food Wastes

Sea area of discharge.--Waters between 3 and 12 nmi of the territorial water baseline and waters within 500 m of ships or maritime facilities engaged in mining mineral resources at or under the sea bottom beyond 12 nmi from the baseline.

Method of discharge.--The garbage must be transformed into ashes or processed by pulverizing machines meeting certain technical standards before discharging. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Sea area of discharge.--Waters >12 nmi from the territorial water baseline.

Method of discharge.--Method of discharge is not limited. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Paper, Wood, and Textile Trash and Other Inflammable Garbage

Sea area of discharge.--Waters between 3 and 12 nmi from the territorial water baseline.

Method of discharge.--The garbage must be transformed into ashes or processed by pulverizing machines meeting certain technical standards before discharge. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Sea area of discharge.--Waters >12 nmi from the territorial water baseline.

Method of discharge.--Method of discharge is not limited. Discharge must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in water.

Metal, Glass, and Ceramic Trash and Other Garbage

Sea area of discharge.--Waters between 3 and 12 nmi of the territorial water baseline.

Method of discharge.--Before discharging, the garbage must be processed by pulverizing machines meeting certain technical standards. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Sea area of discharge.--Waters >12 nmi from the territorial water baseline.

Method of discharge.--Method of discharge is not limited. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Sewage

Sea area of discharge.--All waters.

Method of discharge.--Discharging is free.

Soil

Sea area of discharge.--Port waters, waters within 10,000 m of a low-water line on the coast, Ise Bay, and the Seto Inland Sea.

Method of discharge.--The garbage must be pulverized before discharge. Discharging must be done under the sea surface, during navigation, in small amounts, and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Sea area of discharge.--Waters other than the above.

Method of discharge.--Method of discharge is not limited. Discharging must be done in small amounts and as far as possible from the coast, and efforts must be made to promptly diffuse the garbage in the water.

Note: Discharging of soil is allowed in all waters for ships whose maximum accommodation capacity slips under 100 persons. It is also allowed in all waters for ships whose accommodation capacity is 100 persons or more if soil is processed by soil treatment machines meeting certain technical standards.

Sea areas of discharge (except for sewage and soil) do not include waters within 500 m of ships or maritime facilities which are engaged in mining mineral resources at or under the sea bottom. Discharging of

garbage is prohibited in such waters with the exception of food trash discharged in waters beyond 12 nmi from the territorial water baseline.

Operation-Related Garbage

Operation-related garbage is garbage generated in relation to transportation, fishing, and other usual operations of ships.

Garbage

Garbage with an ignition loss of 15% or less (an ash state of waste plastics) and inorganic garbage (e.g., ore powder generated on ore carriers).

Sea area of discharge.--Waters >50 nmi from the territorial water baseline.

Method of discharge.--Specific gravity must be raised to 1.2. Garbage must not be discharged in powder form. Efforts must be made to sink the garbage to the sea bottom as promptly as possible. Discharge should be avoided in waters where it is expected to affect growth of marine animals and plants.

Note: "Ignition loss" indicates how much the mass of a garbage sample (burned garbage) dried at 105°C for 4 h is reduced after the dried sample is heated at 600°C for 2 h. Garbage with an ignition loss of 15% or less has a weight loss of 15% or less. This percentage indicates garbage which has been burned almost completely.

Plant Garbage

Plant garbage consists of wood chips whose sizes have been reduced to 15 cm or less by pulverizing or cutting (e.g., bark generated on timber carriers).

Sea area of discharge.--Waters >50 nmi from the territorial water baseline.

Method of discharge.--Discharging must be done during navigation and in small amounts. Efforts should be made to diffuse the garbage in waters as promptly as possible. Discharging should be avoided in waters where garbage discharge is expected to affect growth of marine animals and plants.

Animal Garbage

Animal garbage (e.g., livestock which died during transport) (excluding fresh fish and parts thereof).

Sea area of discharge.--Waters >12 nmi from the territorial water baseline.

Method of discharge.--Method of discharge is not limited, but discharge should be avoided in waters where it is expected to affect growth of marine animals and plants.

Fish and Other Marine Animals and Their Parts

Sea area of discharge.--Port waters within 10,000 m of a low-water line on the coast, Ise Bay, and the Seto Inland Sea.

Method of discharge.--Method of discharge is not limited, but discharge should be avoided in waters where it is expected to affect growth of marine animals and plants.

Sewage

Sewage--excluding waste water which does not meet the standards in Appendix B (e.g., waste water used for washing cargo holds or the deck). Waste water used for removing garbage on the deck or other places is classified as sewage. Waste water used for removing garbage accumulated on ships is deemed not sewage but the garbage itself.

Sea area of discharge.--All waters.

Method of discharge.--Method of discharge is not limited. Discharge should be avoided in waters where it may affect growth of marine animals and plants.

Notes: Sea areas of discharge for the first three categories of operation-related garbage do not include waters within 500 m of ships or maritime facilities which engage in mining mineral resources at or under the sea bottom. Discharging is prohibited in such waters.

Subject to restrictions set forth in the first three categories of operation-related garbage is garbage generated in relation to ships' usual activities including transportation, fishing, surveying, observation, and rescue operations. If there are two or more categories of garbage meeting different sets of restrictions, the appropriate regulations will be applied.

REGULATIONS ON DISCHARGE FROM MARITIME FACILITIES

Daily Life-Related Garbage

Daily life-related garbage is garbage generated in relation to the daily life of people on marine facilities.

The same regulations as those for control of discharge of waste from ships are applied to maritime facilities, although discharge of sewage and soil is allowed in all waters. Such discharge should be done as gradually as possible. If garbage is discharged in waters within 500 m of facilities

for mining mineral resources beyond a distance of 12 mi from the territorial water baseline, the garbage must be burned to ashes or processed by pulverizing machines meeting certain technical standards. Discharge of food waste is allowed without such processing if discharge is done as gradually as possible.

Operation-Related Garbage

Discharge is prohibited.

CONTROL OF DISCHARGE FROM AIRCRAFT

Daily Life-Related Garbage

Daily life-related garbage is garbage generated in relation to daily life of people on aircraft.

Sewage and Soil

Sea area of discharge--All waters.

Method of discharge--Free discharge is allowed.

Garbage Other Than Above

Discharging is prohibited.

Operation-Related Garbage

Discharging is prohibited.

APPENDIX A

REGULATIONS UNDER THE LAW RELATING TO THE PREVENTION OF MARINE POLLUTION AND MARITIME DISASTER: MEASURES FOR PREVENTION OF MARINE POLLUTION AND MARITIME DISASTER

- I. Preventive measures.
 - A. Regulations on pollution.
 1. Regulations on discharge at sea.
 - a. From ships.
 - b. From maritime facilities and aircraft.
 2. Regulations on incineration at sea.
 3. Regulations on abandonment of ships.
 - B. Improvement of garbage treatment on ships.
 1. Preparation of manual for the prevention of oil pollution and appointment of oil pollution supervisor.
 2. Preparation of manual for the prevention of pollution by noxious liquid substances and appointment of supervisor for prevention of pollution by noxious liquid substances.
 3. Preparation and maintenance of oil record book.
 4. Preparation and maintenance of noxious liquid substance record book.
 - C. Regulations on structures and equipment of ships.
 1. Structure and equipment standards.
 2. Regular checks and receipt of check certificates.
 - D. Improvement of pollutive substance treatment: The establishment of waste oil disposal facilities and their technical standards.
 - E. Surveillance, patrol, and instruction.
 1. Surveillance and patrol using aircraft and patrol boats.
 2. Lectures, instructions.
 3. Measures after pollution and disaster.

- A. Elimination of pollution.
 - 1. Obligations to report on discharge of oil or noxious liquid substances and vessel-contained noxious liquid substances.
 - 2. Emergency measures.
 - 3. Instructions to eliminate oil, noxious liquid substances, and garbage.
 - B. Establishment of pollution elimination setup.
 - 1. Preparation of discharged oil elimination plans.
 - 2. Obligations to set up equipment for eliminating discharged oil.
 - 3. Maritime disaster prevention center.
 - C. Measures for prevention of marine fire.
 - 1. Obligations to report on discharge of dangerous substances and fire.
 - 2. Emergency measures.
 - 3. Allocation of costs.
 - 4. Disposal of assets.
 - 5. Controls on traffic of ships.
 - D. Technical research and survey.
 - 1. Survey on pollution.
 - 2. Research and development of pollution prevention technology.
 - E. Penalty and administrative measures.
-

APPENDIX B

CRITERIA FOR WATER THAT CAN BE DISPOSED OF
AT SEA AS OPERATION-RELATED SEWAGE

Cadmium contents	0.1 mg or less/L
Cyanogen contents	1.0 mg or less/L
Organic phosphorus contents	1.0 mg or less/L
Plumbum contents	1.0 mg or less/L
Hexad chromium contents	0.5 mg or less/L
Arsenic contents	0.5 mg or less/L
Total mercury contents	0.005 mg or less/L
Alkyl mercury contents	Not detected
PCB contents	0.003 mg or less/L
Trichloroethylene contents	0.3 mg or less/L
Perchloroethylene contents	0.1 mg or less/L

MARINE PLASTIC DEBRIS: WHAT WASHINGTON STATE HAS DONE

Robert Rose
Washington Department of Natural Resources QW-21
Olympia, Washington 98504, U.S.A.

ABSTRACT

This paper describes the formation of an interagency task force in Washington State to address the problem of marine plastic debris along its coast and in its waterways, and the resulting action plan.

INTRODUCTION

Early in 1987, Commissioner of Public Lands for the State of Washington, Brian Boyle, focused his interest on how marine plastic debris was affecting Washington State's coastal and inland waters. In his role as president of the Western States Land Commissioners' Association, he was familiar with efforts in Texas to increase public awareness about marine plastic debris in that state. As the elected official charged with administering the proprietary interests in more than 2 million acres (810,000 ha) of Washington's submerged lands, the commissioner saw this issue as one of major environmental importance as well as a way to focus attention on the stewardship of the state's aquatic lands.

Recent passage of Federal legislation to prevent dumping plastic into the nation's coastal waters, and a recognition that these persistent materials ultimately pollute aquatic lands, meant that this was an issue with immediate and long-term implications for the health of the state's aquatic lands. Persistent marine debris is a highly visible warning for a wide range of environmental contaminants.

BACKGROUND

Late in 1987, the Analysis and Planning Section of the Washington State Department of Natural Resources (DNR) researched the issue. At that time, levels of awareness varied in Washington State government about marine plastic debris and the Marine Entanglement Research Program. No comprehensive analysis of the issue had been done in Washington State, particularly in light of the recent signing by the United States of MARPOL Annex V and the passage of the Marine Plastic Pollution Research and Control Act of 1987. The interjurisdictional nature of the problem required an interagency cooperation with significant information provided by affected private and public and volunteer groups. Other state agencies and

organizations expressed interest and a willingness to participate with the DNR in reaching a mutually agreed course of action to develop a framework for state policy.

In January 1988, the commissioner held a briefing for key principals from affected Federal, state and local agencies, the legislature, private industry and organizations, and other interested parties. At that briefing, participants agreed to select representatives to serve on a task force that would develop a state action plan to address marine plastic debris in Washington's waters and on its shores.

THE WASHINGTON STATE MARINE PLASTICS DEBRIS TASK FORCE

The commissioner appointed a chairman (Robert Rose) and directed DNR staff to support the effort. Each of the 30 active task force members represented an agency or organization (Fig. 1). A steering committee of six worked closely with the chairman and support staff.

The marine plastics debris task force met monthly from February through July 1988. A detailed agenda was prepared for each meeting, which lasted approximately 7 h. Meetings were advertised and open to the public, with a chance for comments scheduled at the close of each meeting. In the course of the first two meetings, the task force adopted a 6-month work program, and agreed upon its mission, goals and objectives, and policies. The task force divided itself into three working groups: environment, education, and government and economic impact. Each group was to identify issue areas and possible actions.

In following meetings, the groups examined the diverse programs and regulations administered by agencies dealing with marine debris. Each agency or organization representative developed a "Status of Marine Debris Program" sheet and presented this material to the group (Fig. 2). Each status sheet gave the name of the agency or organization, the contact person, importance of the issue, legal authority to deal with the issue, additional authorities needed, resources currently available, and a description of the 1988 action plan (Appendix A). From this information, the staff developed a matrix of the existing authorities of agencies to deal with marine plastic debris (Fig. 3).

The steering committee assembled specific issue areas identified by the task force and developed a number of draft action recommendations which conformed to the objectives and policies. With subsequent approval by the group as a whole, the chairman and steering committee then developed a brief narrative statement explaining the issue area and identifying the agencies responsible for each issue. The recommendations and preferred-lead agencies to carry out the actions were collated and organized by staff and presented to the committee. Over the course of the next two meetings, bolstered by telephone conferences and mailed-in comments, the chair and staff clarified the committee's intent. At the sixth meeting, the task force gave its final approval to the text of the report. With agreement on issue areas, lead agencies, and participating agencies, a detailed matrix was prepared so that agency or organization directors and staff could

ACKNOWLEDGEMENTS

Marine Plastic Debris Task Force

Members

AGENCY / ORGANIZATION

REPRESENTATIVE

AGENCY / ORGANIZATION

REPRESENTATIVE

FEDERAL

Department of Commerce
National Oceanic and Atmospheric Administration

Alan Bunn, LCDR

Department of Interior
Fish and Wildlife
National Park Service

Don Kase
Chuck Janda

Department of Transportation
13th Coast Guard District

Stephen D. Heath, LCDR

Environmental Protection Agency,
Region X

Michael Rylko

STATE

Department of Community Development

Sandi Benbrook
George Turnbull, Jr.

Department of Ecology

Mikel DeBuse
Brad Everson
Rhonda Hunter
Dennis Johnson
Pam Miller
Leighton Pratt

Department of Fisheries

Sally Hicks
Phil Kaulzloric
Sandra O'Neill

Department of Natural Resources

David G. Bortz
Marge Reed
Bob Rose, Chairman
Glenn Yeary

Department of Wildlife

Steve Jeffries

Office of the Governor

Gerard (Sid) Sidorowicz

Puget Sound Water Quality Authority

Sarah Barten
Richard Tucker

State Parks and Recreation Commission

Nisa Carter

LEGISLATIVE

Rep. Nancy Rust, House Environmental
Affairs Committee

Ya-Yee J. Van

Joint Select Committee Marine and
Ocean Resources

Bob Betts

Senate Environment and
Natural Resources Committee

Laurie Schock

LOCAL

Association of Washington Cities

Andrea Dehl

Northwest Indian Fisheries

R. J. Whitener

Port of Seattle

Linda Cox

Washington Public Ports Association

Eric Johnson
Darrall Russell

PRIVATE

Battelle Marine Research Laboratory

Betsy Brown

Dalco Packaging

Raye Little

Inner Club Boating Association of
Washington

Ray Nelson

Marine Digest

Donna Mooney

Natural Resources Consultants

Jeff June

R. W. Beck and Associates

David E.B. Nightingale

Society of Plastics Engineer

Chal B. Rogers

CITIZEN ORGANIZATIONS

Adopt-A-Beach,
c/o Volunteers for Outdoor Washington

Ken Fritchard

The Mountaineers

Margaret Hansen

Washington Citizens for Recycling

Nancy Pearson

Washington Environmental Council

Lisa Macchio

EDUCATIONAL ORGANIZATIONS

Friday Harbor Laboratories of
University of Washington

Dr. Frederick E. Ellis

Poulsbo Marine Science Center

Laurie Dumdie

Seattle Aquarium

John McMahon
Cherie Williams

University of Washington/Sea Grant

Xan Augerot
Jim Humphreys

Patti Mullin

Other Participants

American President Lines, Ltd.

Board of Clallam County Commissioners

Board of Grays Harbor County Commissioners

Board of Jefferson County Commissioners

Board of Pacific County Commissioners

Board of San Juan County Commissioners

Board of Wahkiakum County Commissioners

Marine Resources Company

Northwest Marine Trade Association

Superintendent of Public Instruction

Washington Association of Counties

Project Staff

Department of Natural Resources

John Bergvall
Marsha Hixson
Jane Grobins
Carol Lind
Camille Blanchette
Gail S. Weir
Karen Wood
Marie Oles
Jacque Encarnacion

Design
Editors

Illustrations
Word Processors

Photos courtesy of:
Center for Environmental Education
Cynthia Feutch
Chuck Fowler
Tom Pani

C.E. O'Clair
Greebel
Bill High

Special thanks to the Department of Ecology Shorelines and Coastal Zone Management Program for supporting the work of this task force by underwriting the mailing of these documents.

Figure 1.--Marine Plastic Debris Task Force members.

understanding of the issue. These will, for the most part, be integrated with existing interpretive programs such as guided walks, evening programs and wayside exhibits. Some activities may be appropriate in conjunction with State and Federal programs and we would be pleased to participate in any way possible.

(c) Projected: There are no projections for future activities at this time. However, the park is prepared to initiate or participate in efforts as opportunities arise and/or funds permit.

(d) Cooperative effort: Aside from our interest in continuing to work with the Task Force and State and Federal programs that will eventually be implemented, the park has only one identifiable cooperative program as a possibility. The National Marine Fisheries Service is interested in Olympic as one of four nationwide sites to be surveyed for marine debris accumulation. As stated in the study proposal, the objective would be: "To develop a program of systematic surveys in each region of the coastal United States to assess the types, quantities and sources of debris arriving on these shores, and to identify trends or changes in these parameters." Final site selection is expected within the next few weeks.



United States Department of the Interior
 NATIONAL PARK SERVICE
 OLYMPIC NATIONAL PARK
 600 East Park Avenue
 Port Angeles, Washington 98126/98

March 17, 1988

MARINE PLASTICS DEBRIS TASK FORCE

1. Agency: National Park Service (Olympic National Park)

2. Why issue is important: Approximately 60 miles of beaches along the open coast of Olympic National Park have become the repository for uncounted tons of plastic debris. The presence of this material is significant in two respects. Aesthetically, the debris creates an ever-present disfigurement of an otherwise all-natural landscape. The bright colors, characteristic shapes and the sheer volume of plastics have visually impacted literally every meter of beach in the park. Of equal importance are the known and unknown, physical and chemical effects this debris has on intertidal organisms, including birds and mammals. The impact of introducing this variety and quantity of plastics into a pristine ecosystem cannot be understated.

3. Authority to deal with the issue: The fact that the National Park Service has exclusive jurisdiction within the park is of minor importance since the debris originates outside park boundaries. One legal recourse that, perhaps, deserves attention is the possibility of invoking the Constitution's property clause. While this is a rarely-used authority within the National Park System, the plastic debris issue, in many key ways, meets the criteria for its implementation, not only at Olympic but also in many National Parks along the Pacific Rim, the Gulf of Mexico and the Atlantic seaboard.

No additional authorities would alleviate the current problem and/or improve our enforcement efforts.

4. Resources available: The park has the authority to allocate operating funds and fee collection revenue to beach cleanup and a wide range of research/inventory/monitoring projects. Consistent with other top priorities and shrinking budgets, FRG's can be assigned to in-park and cooperative projects that deal with the plastics issue. The major workload, however, has been and will likely continue to be borne by volunteers.

5. Description of program effort:

(a) Current: The park will continue to operate an annual beach cleanup program. This year's allocation for the project is \$4,500. Most of the labor will be provided by volunteer groups.

(b) Planned: Although specific plans have yet to be developed, the park will be initiating projects aimed at increasing visitor awareness and

Figure 2. ---Sample agency status sheet.

EXISTING AUTHORITIES AND AGENCIES DEALING WITH MPD

AGENCY	PROGRAM		ENFORCEMENT		STATUTES	FUNDING
	DEVELOPED	POTENTIAL	DEVELOPED	POTENTIAL		
EPA, REGION X	<ul style="list-style-type: none"> ■ National ongoing plastics pollution study ■ 1986-87 Adopt-A-Beach program with Seattle Aquarium; volunteer project hndbk. completed ■ State of the Sound exhibit with Seattle Aquarium 	<ul style="list-style-type: none"> ■ Public Education in cooperation with NOAA and Dept. of Transportation ■ Support establishment of baseline data and monitoring program ■ Additional technical support 			<ul style="list-style-type: none"> ■ Marine Plastics Pollution, Research and Control Act (MPPRCA) (PL 100-220) ■ Clean Water Act, Section 108 	
USFWS	<ul style="list-style-type: none"> ■ Beach cleanup 	<ul style="list-style-type: none"> ■ Cooperative actions ■ Field monitoring ■ Public education materials 			<ul style="list-style-type: none"> ■ Fish and Wildlife Coordination Act ■ Endangered Species Act ■ Migratory Bird Treaty Act ■ Anadromous Fish Conservation Act ■ Rivers and Harbors Act of 1899 ■ Clean Water Act 	
NPS	<ul style="list-style-type: none"> ■ Beach cleanup 	<ul style="list-style-type: none"> ■ Research and monitoring; Olympic Nat'l Park as an accum. monitoring site ■ Cooperative action ■ Visitor information/education 	<ul style="list-style-type: none"> ■ Exclusive jurisdiction in national parks ■ Authority to prosecute 	<ul style="list-style-type: none"> ■ Increased enforcement 	<ul style="list-style-type: none"> ■ Title 16, 18 U.S. Code 	
NMFS	<ul style="list-style-type: none"> ■ Marine entanglement research program 		<ul style="list-style-type: none"> ■ Enforcement personnel available 		<ul style="list-style-type: none"> ■ National Marine Mammal Protection and Control Act 	
COAST GUARD		<ul style="list-style-type: none"> ■ Participation with EPA and NOAA on public education 	<ul style="list-style-type: none"> ■ Fines for vessels ■ Certification of port facilities ■ Developing regulations 	<ul style="list-style-type: none"> ■ Increased enforcement (with increased funds) 	<ul style="list-style-type: none"> ■ MPPRCA 	

Figure 3.--Sample taken from table of existing authorities and agencies.

quickly reference their responsibilities and relationships with other agencies (Fig. 4).

The task force recognized that marine plastic affects fisheries and wildlife resources, endangers boaters and divers, and diminishes aesthetic enjoyment of the state's shorelines. Increasing public awareness and interest about the specific issue of marine plastic debris is a key element of the plan, with a strong emphasis on education. The task force recommendations emphasize recycling and proper disposal of potential marine plastic debris materials. In addition, the coordination of present and future cleanup efforts is critical to use Federal, state, local, and citizen resources efficiently.

The task force wanted its recommendations to be thoroughly examined by the affected agencies and incorporated into future work plans and budgets. Because the issue involves so many agencies and organizations and will involve a commitment of resources over time, the task force recommended designation of an overall coordinating agency.

MARINE PLASTIC DEBRIS: ACTION PLAN

The action plan was completed in October 1988 and presented to the commissioner, the public, and media at a press conference on the Seattle waterfront. The publication of the document was timed to coincide with the last day of Washington State's Coastweeks '88. The Adopt-A-Beach Program, sponsors of Coastweeks (under the auspices of the Washington Department of Ecology), organized the event and collected a representative sampling of debris from the riprapped shore of Elliott Bay.

The 1988 plan is organized into four sections.

Part 1 of the action plan explains how the Marine Plastic Debris Task Force developed a plan to address a major environmental problem: debris in Washington waterways. The floating garbage was not only unsightly but dangerous. How the task force focused on issues and their recommended actions are discussed.

Part 2 of the action plan presents the mission statement for the Marine Plastic Debris Task Force and the goal and objectives for the plan. Policies to guide and coordinate future activities are followed by 20 action recommendations designed to implement the policy statements. These recommendations are divided into three sections: environment, education, and government/economic impact. Part 2 also lists seven required legislative or administrative initiatives. (The mission statement, goal and objectives, and policies of the Marine Plastic Debris Task Force (1988), as documented in the Marine Plastic Debris Action Plan, are found in Appendix A of this paper. A summary of the marine debris action recommendations and the required legislative or administrative initiatives are found in Appendix B of this paper.)

Part 3 of the action plan contains the full text for each of the action recommendations including a short narrative explaining the

ACTION RECOMMENDATIONS AND THE LEAD AGENCIES INVOLVED

ACTION RECOMMENDATIONS	ENVIRONMENT								
	1 Designate coordinating and clearing-house agency	2 Develop environmental baseline and monitoring system	3 Coordinate beach cleanups	4 Conduct research on wildlife and fisheries effects	5 Coordinate ghost net removals	6 When feasible, require and promote recycling and alternative products	7 Recycling and disposal	8 Convene biennial statewide conference	9 Develop and implement a public outreach program
ORGANIZATIONS									
Coordinating agency		L						L	L
NOAA/NMFS	P	P		P	P	P	P		P
USFWS	P	P	P	P					P
NPS	P	P	P						P
CG	P	P					P		P
EPA	P	P		P		P	P		P
DOL	P								P
DCD	P								P
DOE	L	P	L			L	L		P
WDF	P	P	P	L	L	P			P
DNR	L	P	P		L	P			P
WDW	P	P	P	L	P				P
OFM									P
Office of Governor	P								P
DOR									P
PSWQA	P								P
State Bd. of Pilotage	P								P
Parks	L	P				P	L		P
SPI	P								P
IAC	P						L		P
Hse Env. Affrs. Com.	P								P
House Energy & Nat. Res. Coms.									P
Joint Select Com. Mar. & Ocean Res.	P								P
Senate Environ. & Nat'l Res. Com.	P								P
State Legislature	P						L		P
Association of Washington Cities		P	P		P	P	P		P
Wa. Assoc. of Counties		P	P		P	P	P		P
Port of Seattle		P	P				P		P
Wa. Pub. Ports Assoc.		P	P		P		P		P
County Auditors									P
Colleges & Universities	P	P	P						P
Private Companies	P	P	P	P		P	P		P
Aquariums		P		P					P
Marine Labs		P		P					P
Adopt-A-Beach		P	P						P
Indian Tribes		P							P
Dive Groups/Shops					P				P
Environ. Groups									P
Citizen Recycling Org.							P		P

L - Lead agency P - Participating agency

Figure 4.--Sample taken from Table 4 of action recommendations.

background of the issue. For each action recommendation, a lead agency is designated as well as a listing of agencies likely to participate in carrying out the action. A suggested time frame for implementation and an estimate of the duration of the action are proposed for each of the 20 action recommendations.

Part 4 of the action plan consists of three appendixes. Appendix A of the plan is a compilation of reports outlining the status (as of September 1988) of agency and organization activities, authorities, and current and proposed programs. Appendix B of the plan contains an overview and explanation of the Marine Plastic Pollution Research and Control Act of 1987. Appendix C of the plan encapsulates various Washington State programs addressing the problem of marine debris.

Copies of the action plan are available from Photo and Map Sales, Department of Natural Resources, 1055 Capitol Way S, AW-11, Olympia, WA 98504, (206) 753-5338.

IMPLEMENTATION

Since presenting the task force report to Commissioner Boyle, a number of steps have been taken to make the action plan a reality.

Designees of the directors of Parks, Ecology, and Natural Resources met in early 1989 to decide which agency should take the lead for coordinating future state marine plastic debris activity. As a result of that meeting, the Department of Natural Resources, having initiated and staffed the planning effort, was designated lead agency. The relationship among the agencies was formalized in a memorandum of understanding. Based on this agreement, the department prepared legislation for the 1989 Legislature. House Bill 1249 was to authorize the DNR to coordinate implementation of the action plan and to develop rules for cleanup and prevention of pollution in state waters. The department is also authorized to enter into intergovernmental agreements with Federal, state, or private parties, and to hire employees necessary to coordinate the plan.

An important conclusion of the task force deliberations was the recognition that a coordinator was necessary to assure that agencies followed through on the tasks contained in the action plan. This legislation provides the authority and funding source to make the plan a success.

At the same time, the Department of Ecology has funded through the coastal zone management program a baseline inventory survey of selective beaches. This will be carried out by the Adopt-A-Beach program using a newly revised inventory form. Beginning on 25 March, a scientific inventory of selected beaches will occur biweekly for a year, thus providing the first baseline information on marine plastic debris occurrence on Washington's shores.

As a result of widespread publicity about the report, the 1989 and 1990 Department of Transportation's Tide Tables, a popular and necessary document requested by boaters and fishermen along the Washington coast,

contains a special eight-page insert about proper disposal of all marine debris and sewage, with a special emphasis on plastic and other nonbiodegradable materials.

A ghost net task force with representatives from State Departments of Fisheries, Wildlife, and Natural Resources, interested citizens and divers, and the National Marine Fisheries Service, NOAA, is developing an inter-agency memorandum of understanding and protocols for ghost net and crab pot removal.

The Department of Natural Resources will start publishing a marine plastic debris newsletter to bring current activities and opportunities to the attention of boaters, lessees, citizen groups, and others interested in maintaining and improving Washington's aquatic environment.

CONCLUSION

The development and publication of the "Marine Plastic Debris Action Plan for Washington State" was the first and necessary step for addressing the presence and consequences of persistent debris in the state's waters and on its shores. As a result of the plan, institutional arrangements have developed which would not have been possible without the framework proposed by the task force. Positive response by the Washington Legislature portends support for the Department of Natural Resources to develop the necessary educational materials, to create public and private partnerships, and to raise the importance of this issue in the public mind.

REFERENCES

- Marine Plastic Debris Task Force.
1988. Marine plastic debris action plan for Washington State.
Marine Plastic Debris Task Force, Washington State Department of Natural Resources, 2d ed., December 1988, var. pag.

Appendix A.--The mission statement, goal and objectives, and policies of the Marine Plastic Debris Task Force (1988).

MISSION STATEMENT

The purpose of the task force is to increase public awareness and interest about marine plastic debris and to develop a framework for coordinating public and private efforts to ensure an effective response. Plastic debris in the aquatic environment has an adverse impact on wildlife, aesthetics, navigation and overall environmental quality. The task force recognizes that this material is part of a larger problem of waste generation and management. The Washington State plan will link private, local, state and federal efforts in managing the plastic waste stream as it affects the shorelines and aquatic environment.

GOAL AND OBJECTIVES

Goal

To develop a state action plan to address the marine plastic debris issue that affects Washington's shorelines and aquatic environment.

Objectives

1. Identify plastic debris in the aquatic environment as a distinct issue.
2. Develop and support mechanisms to reduce or eliminate marine plastic debris.
3. Focus primarily on marine resources affected by plastic debris, with attention to impacts on other aquatic environments in the state.
4. Provide input to and develop steps for emerging private, local, state and federal policies and actions.
5. Develop and implement mechanisms that will coordinate actions performed by agencies and organizations.
6. Encourage and support private and public policy to increase awareness through education efforts in Washington State.

POLICIES

The Marine Plastic Debris Task Force recommends that the state of Washington, acting through its elected officials, and in cooperation with other appropriate agencies, offices, organizations and the private sector, should seek to:

1. Increase public awareness about effects of marine plastic debris.
 2. Designate a lead agency to act as a clearinghouse and coordinator for marine plastic debris activities in Washington.
 3. Designate appropriate agencies to draft, review and support legislation and/or regulations recommended by the task force.
 4. Empower responsible agencies to implement and participate to the fullest extent in actions recommended by the task force and other actions judged necessary.
 5. Encourage funding measures on all levels to facilitate compliance with MARPOL requirements and to implement task force recommendations.
 6. Coordinate, support and encourage continued volunteer efforts and special events related to the marine plastic debris issue.
 7. Encourage and support efforts by all the state's users of the marine environment to reduce and eliminate marine plastic debris.
 8. Support and encourage the plastics industry and other industries to continue research and development of products which are adaptable to recycling and proper disposal.
 9. Maintain and expand baseline data collection and research on sources, quantities, effects and fates of marine plastic debris.
 10. Encourage ports and local governments to collect and dispose of marine plastic debris in an environmentally sound manner.
 11. Support regional, national and international efforts to reduce and eliminate marine plastic debris.
-

Appendix B.--Summary of the marine debris action recommendations and the required legislative or administrative initiatives (Marine Plastic Debris Task Force (1988)).

SUMMARY OF ACTION RECOMMENDATIONS

- (1) ACTION RECOMMENDATION: Designate an agency or entity to coordinate activities and serve as a clearinghouse for marine plastic debris data reception and information dissemination.

ENVIRONMENT

- (2) ACTION RECOMMENDATION: Develop an environmental baseline and monitoring system for marine plastic debris in Washington.
- (3) ACTION RECOMMENDATION: Coordinate beach cleanup efforts among various agencies.
- (4) ACTION RECOMMENDATION: Conduct additional research about the effects of plastic debris on wildlife and fisheries resources and habitat.
- (5) ACTION RECOMMENDATION: Coordinate information about and removal of ghost nets and other derelict equipment from state waters among DNR, Department of Fisheries, NOAA, and local agencies.
- (6) ACTION RECOMMENDATION: Require and promote recyclable or alternative products when feasible for use in or near the marine environment.
- (7) ACTION RECOMMENDATION: Require recycling and proper disposal of potential marine plastic debris materials for federal, state and local agencies and Indian tribes.

EDUCATION

- (8) ACTION RECOMMENDATION: Convene a biennial statewide conference by the designated marine plastic debris coordinating agency.
- (9) ACTION RECOMMENDATION: Develop and implement a public outreach program using the recommended logo [Illustration 6], public service announcements, publications and special events.
- (10) ACTION RECOMMENDATION: Develop and distribute media materials directed at domestic and foreign maritime communities.

- (11) ACTION RECOMMENDATION: Develop a marine debris curriculum for use at state and private maritime schools training the professional maritime community.
- (12) ACTION RECOMMENDATION: Post permanent information signs about why MPD and other litter is harmful to water quality, wildlife and fish as well as a threat to boating safety at all boat launch ramps, public access sites and public and private marinas.
- (13) ACTION RECOMMENDATION: Require that all state-licensed and registered users of Washington waters receive MPD information materials and display a plaque or decal about proper disposal of MPD and other litter.
- (14) ACTION RECOMMENDATION: Encourage retail and manufacturer cooperation to aid in marine plastic debris public education.
- (15) ACTION RECOMMENDATION: Incorporate the MPD problem into the environmental section of the state's required curriculums for grades K-12 and in other educational material.

GOVERNMENT/ECONOMIC IMPACT

- (16) ACTION RECOMMENDATION: Incorporate an analysis of the true costs of marine debris, including fiscal impacts and nonquantifiable environmental costs into policy decisions.
- (17) ACTION RECOMMENDATION: Conduct an independent cost/benefit analysis of MPD compliance and cleanup.
- (18) ACTION RECOMMENDATION: Integrate MPD disposal with comprehensive solid waste planning.
- (19) ACTION RECOMMENDATION: Increase public awareness about the legal consequences for improper MPD disposal.
- (20) ACTION RECOMMENDATION: Develop a framework of financial incentives to encourage proper disposal of MPD.

REQUIRED LEGISLATIVE OR ADMINISTRATIVE INITIATIVES

1. Marine Debris decal--Departments of Fisheries and Licensing should require that all Washington-licensed boats have a prominently displayed decal regarding proper disposal of marine plastic debris and other wastes.
2. Information for boaters--Require the Department of Licensing include an information packet containing legal requirements for marine plastic debris disposal when issuing boat registrations or renewals.

3. Legislation--Support passage of the State Parks legislation for Boater Recreation Fee Account funds to be used for boater environmental education and to provide capital for sewage pump-out facilities and educational signs.
4. Agreement--Formal agreement among DNR, Department of Fisheries, NOAA (and other appropriate agencies) is necessary to assume prompt location and removal of ghost nets.
5. Funding--Investigate Ecology's Litter Tax program (Chapter 70.93 RCW) for the possibility of directing funding from that program to the marine debris cleanup issue.
6. Clearinghouse--Develop an appropriate memorandum of understanding for the coordinating agency.
7. Staffing--Consider increased state enforcement through WDOE, WDF personnel to carry out these recommendations.

THE PROTECTION OF SPECIFIC SEA AREAS AGAINST MARINE DEBRIS

Gerard Peet
Heemraadssingel 193
NL 3023 CB Rotterdam
The Netherlands

(This paper was presented in April 1989. It has since been updated (e.g., a new title, amended criteria) to include developments occurring at the Marine Environment Protection Committee meeting of March 1990.)

ABSTRACT

The problems associated with debris in the marine environment are being given increased attention. At the international level this attention is reflected in a number of international agreements. Most of the relevant international regulations aim at reduction of debris-associated pollution at the source. In addition to policies aimed at source control, efforts are also made to develop protective measures for specific sea areas. This paper discusses efforts currently under way within the framework of the International Maritime Organization.

In the mid-1980's, the International Maritime Organization decided to develop guidelines for the designation of "special areas" and the identification of "particularly sensitive areas." These guidelines should assist national authorities in developing measures to provide specific areas with additional protection from environmental damage caused by shipping activities. The Baltic Sea became a special area as of 1 October 1989. The United States has announced a proposal to designate the Gulf of Mexico as an Annex V special area as well. The Governments of the North Sea States have formally proposed to do the same for the North Sea. Another major option to protect specific sea areas is the designation "area to be avoided by ships." The Northwestern Hawaiian Islands are an example of such an area.

INTRODUCTION

Several years ago, a photograph of a dead albatross spread out on a beach together with a systematic display of the plastics found in the bird's stomach brought home to me the point that there was more to pollution of the seas than oil or chemicals. Photographs like this one of

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

birds and sea mammals entangled in or killed by plastics and other persistent materials have helped to increase attention for the problems associated with debris in the marine environment worldwide.

My first real encounter with debris-associated pollution was on board a Greek passenger ferry in the Mediterranean. Passengers freely threw plastic bags and other garbage overboard. It emphasized the importance of shipping as a source of this type of pollution.

My second important encounter with debris-associated pollution involved an invitation by an artisanal fisherman in one of the Southeast Asian countries to come and see how plastics rather than fish filled his nets. He took me to the source of these plastics. The waste dump of the town he lived in was located on a waterfront. One of the ironies of this was that waste from this dump would not only fill the fisherman's nets, but would also wash up at the town's beaches. These beaches were cleaned regularly, and the collected waste brought to this waste dump. It reminded me that there were more sources of debris-associated pollution than ships.

Increased attention to the problem has now led to increased attention to measures to control pollution from land-based sources as well as from ships on both national and international levels. This paper concentrates on international measures.

SOURCES OF DEBRIS-ASSOCIATED POLLUTION AND INTERNATIONAL AGREEMENTS FOR POLLUTION CONTROL

Land-Based Pollution

Although the international dimension of debris-associated land-based pollution appears to be limited, there are some important international agreements in this respect. The Convention for the Protection of the Mediterranean Sea Against Pollution and its related protocols (1976) is one of these. "Persistent synthetic materials which may float, sink or remain in suspension and which may interfere with any legitimate use of the sea" are on the Annex I list of the Protocol for the Protection of the Mediterranean Sea Against Pollution from Land-Based Sources (1980); "substances which, though of a non-toxic nature, may become harmful to the marine environment or may interfere with any legitimate use of the sea owing to the quantities in which they are discharged" are on the Annex II list of this protocol. Pollution by Annex I substances should be eliminated (Art. 5 of the protocol); pollution by Annex II substances should be strictly limited (Art. 6 of the protocol). Similar regulations have been included in other international agreements regarding land-based pollution such as the Convention for the Protection of the Marine Environment and Coastal Area of the Southeast Pacific and its Supplementary Agreements (1981, 1983), and the Paris Convention for the Prevention of Marine Pollution from Land-Based Sources (Northwestern Europe 1974).

If effectively implemented, these regulations should provide a basis for sufficient control of debris-associated land-based pollution in the areas where these regulations apply.

Dumping of Wastes at Sea

Another source of debris-associated pollution is the dumping of wastes at sea. On a global as well as on a regional level, international agreements have been developed to regulate the dumping of wastes at sea. The global Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (the so-called London Dumping Convention of 1972) has addressed the dumping of plastics and other types of debris by putting "persistent plastics and other persistent synthetic materials, for example, netting and ropes, which may float or remain in suspension in the sea in such a manner as to interfere materially with fishing, navigation or other legitimate uses of the sea" on its Annex I. The dumping of Annex I substances at sea is prohibited (Art. IV.1.a). "Containers, scrap metals or other bulky wastes liable to sink to the sea bottom which may represent a serious obstacle to fishing or navigation" and "substances which, though of a nontoxic nature, may become harmful due to the quantities in which they are dumped, or which are liable to seriously reduce amenities" were put on Annex II, thus requiring a prior special permit if dumping is to take place (Art. IV.1.b).

Guidelines have been developed for the identification of discharge or dump sites. These guidelines include consideration of such factors as the capacity of the receiving marine environment to receive wastes without undesirable effects and the ecological condition of the area.

Effective implementation of the London Dumping Convention (which, in early 1989, had been ratified by 63 states) could be the basis for controlling this source of pollution.

Operational Pollution From Ships

Research into the origin of plastics and other marine debris suggests that a substantial part of these substances originates from ships at sea (e.g., Dixon and Dixon 1981). A recent report about the German Bight (Schrey 1987) estimates that 95% of the refuse found on beaches of the German Bight can be attributed to shipping.

Any effort to control this source of debris-associated pollution from ships must have an international dimension since most of the world's seas and oceans are international waters. Ships traveling these waters are flying the flags of many nations. International regulations affecting these ships are developed by the International Maritime Organization (IMO), a United Nations agency which as of April 1989 had 133 member states and which consequently is well placed to effectively develop such international regulations.

Operational pollution from ships has been regulated by the IMO in the International Convention for the Prevention of Pollution from Ships (the so-called MARPOL Convention of 1973/78). This convention is now under constant review by IMO's Marine Environment Protection Committee (MEPC). The MARPOL Convention includes regulations to control operational pollution from ships with oil (Annex I), noxious liquid substances in bulk (Annex

II), harmful substances in packaged forms (Annex III), sewage (Annex IV), and garbage (Annex V). Marine debris falls within the scope of Annex V of the MARPOL Convention.

It is important to properly appreciate the MARPOL Convention. This convention was concluded in 1973 and, at that time, already included very strict rules with regard to the disposal of plastics at sea. Yet, its Annex V, the plastic and garbage regulations, did not enter into force until more than 15 years later, on 31 December 1988.

THE EFFECTIVENESS OF EXISTING INTERNATIONAL MEASURES WITH RESPECT TO MARINE DEBRIS

International regulations for control at the source of the introduction of marine debris into the marine environment do exist. The important question, however, is whether they are effective.

As regards land-based pollution, there is only one global international instrument, the so-called Montreal Guidelines. Their effectiveness is limited because they are not binding. Most of the regional international regulations are no more than one or two general articles in a general convention on environmental protection. There are only a few exceptions where special protocols or specific conventions with regard to land-based pollution were developed. The effectiveness of the existing regulations is limited because there is no worldwide coverage of international regulations; many regions do not have such regulations.

There is some doubt about the effectiveness of regional international regulations where these have been developed up to a level of specialized protocols or even specific conventions. One example may illustrate this. With respect to synthetic materials (a blacklisted substance in the Paris Convention), no action at all appears to have been taken by the Commission of the Paris Convention in the first 10 years of its existence (Oslo and Paris Commission 1984; Paris Commission 1987).

As for the dumping of wastes, the London Dumping Convention seems to be an effective instrument. Nevertheless, in at least one instance the convention failed for lack of enforcement.

In April 1988, the car-carrier *Reijin*, with more than 5,000 new cars on board, capsized close to the Portuguese coast. After considering the various salvage options, it was decided to dump two-thirds of the cars from the ship into water 2,000 m deep. The wreckage of the ship together with the remaining cars were then to be sunk in deep water as well (MEPC 1989a). In effect this would mean the dumping at sea of a number of substances from Annex I of the London Dumping Convention for which dumping at sea is prohibited. Plastics and other persistent materials used in the cars were among these Annex I substances. During the 1988 Consultative Meeting of Contracting Parties to the London Dumping Convention, the delegation of Denmark as well as observers from the environmental organizations Greenpeace and Friends of the Earth International raised the issue as being at odds with the regulations and spirit of the London Dumping Convention. The

consultative meeting took no action (pers. observ.). Dumping of cars at sea did start but was stopped after a time.

Finally, with respect to operational discharges by ships, not enough time has elapsed since the entry into force of Annex V of the MARPOL Convention to judge its effectiveness. Doubts have been raised, however, in this regard. Many consider control and enforcement of the provisions of the annex to be extremely difficult. Guidelines for its implementation were completed by MEPC in September 1988 (MEPC 1988f), less than 4 months before its entry into force. It is doubtful whether the necessary reception facilities are available in all ports.

THE PROTECTION OF SENSITIVE SEA AREAS AGAINST DAMAGE BY SHIPPING ACTIVITIES: BACKGROUND

Considering the problems encountered in controlling the discharge of marine debris at the source, one wonders whether a complementary approach of giving special protection to specific sensitive areas might be useful. Within the IMO, efforts are now under way to assess the opportunities such an approach might offer with respect to marine pollution caused by ships.

The IMO has several options for providing additional environmental protection to specific sensitive sea areas. These include the designation of areas as "special areas" under the MARPOL Convention, the designation of areas as "areas to be avoided," or the use of other ship's routing measures such as traffic separation schemes and deep-water routes. With respect to debris-associated pollution, the first two options are especially relevant.

Special areas will normally be larger sea areas. To provide some protection for sea areas which would not qualify as special areas, the International Conference on Tanker Safety and Pollution Prevention in 1978 adopted a resolution which invited the IMO:

"to initiate. . . studies, in collaboration with other relevant international organizations and expert bodies, with a view to making an inventory of sea areas. . . which are in special need of protection against marine pollution from ships and dumping. . . ; assessing. . . the extent of the need of protection, as well as the measures which might be appropriate. . . ; to consider. . . what action will be needed. . . ; to take action. . . within the framework of the relevant conventions. . . ."

In 1985, the IMO started to work on this issue of particularly sensitive sea areas and put it on the agenda of the twenty-third session of the MEPC, which was to take place in 1986. Discussion of the issue at this meeting (MEPC 1986) resulted in the decision to send out a circular letter to IMO member states inviting these states to provide information on the following:

- Criteria which have been used in designating existing marine areas under national jurisdiction which are particularly sensitive with respect to their renewable natural resources

or their importance for scientific purposes, and for which special protection measures are in force.

- National protection measures and restrictions affecting the use of such areas by ships and related maritime activities, and the specific purpose of the restrictions imposed.
- The geographical location of those marine areas which are already protected and of those areas considered for future protection, the seaward limits of which extend beyond the territorial seas established in accordance with international law.

On the basis of the responses to this circular letter, the MEPC developed criteria for the designation of particularly sensitive sea areas and also started work on developing criteria for the designation of special areas. At the twentieth-sixth session of MEPC, a proposal was put before the MEPC on how to proceed and how to make the concept of particularly sensitive sea areas operational (MEPC 1988d).

The proposal did not aim at developing new legal instruments, but at making better use of existing international regulations (such as the designation of special areas) for the protection of specific sea areas against damage caused by shipping activities. The MEPC adopted this proposal (MEPC 1988f) and decided to develop a manual for the designation of particularly sensitive sea areas and special areas. Since the adoption of this proposal some changes have been made to the concept of the manual, including its title, which now is "Guidelines for the designation of special areas and the identification of particularly sensitive areas" (MEPC 1990). The basic concept of providing guidance for better use of existing international regulations is still the same.

GUIDELINES FOR THE DESIGNATION OF SPECIAL AREAS AND THE IDENTIFICATION OF PARTICULARLY SENSITIVE AREAS

The main objective of the guidelines is to provide governments or government departments having limited experience in developing proposals to the IMO with detailed guidance on how to prepare such proposals for environmental protection of specific sea areas. It will also set standards which proposals for the designation of special areas will have to meet if they are to be accepted.

The guidelines will present a range of existing international regulations which could be used better or more frequently for environmental protection purposes. They will be restricted to damage from ships in or in the direct vicinity of an area, and will not address land-based pollution or the dumping of wastes at sea.

They will consist of three parts: (1) a general introduction, (2) criteria and procedures for the designation of special areas as well as some examples of special areas already designated, and (3) criteria for the identification of particularly sensitive areas, criteria and procedures to

provide such areas with additional protection in accordance with IMO regulations, and some examples of areas which have already been given such additional protection.

Guidelines: General Introduction

This first chapter of the guidelines will review their history and background as well as the role the IMO can play in the protection of sensitive sea areas. Attention will also be given to the types of damage ships can cause to sensitive sea areas. One of these will be the discharge of marine debris, including plastics.

The list will, however, not be limited to discharges of the "traditional" substances such as oil or chemicals; it will also include the discharge of ballast water contaminated with "alien" organisms (which has already caused problems near Tasmania, Australia, and in the Great Lakes, Canada, and the United States), the "discharge" of TBT paints from the hull of ships into the marine environment, and even the "discharge" of noise.

Neither will it be limited to damage caused by discharges; physical damage to marine ecosystems (such as the damage to coral reefs caused by the grounding of the *Wellwood* off Key Largo, Florida, United States, in 1984) will also be discussed.

Guidelines: Special Areas

The second chapter of the guidelines will address the designation of special areas. Annex V of the MARPOL Convention defines a special area as "a sea area where for recognized technical reasons in relation to its oceanographic and ecological condition and to the particular character of its traffic the adoption of special mandatory methods for the prevention of sea pollution by garbage is required" (Annex V, Reg. 1.3). The disposal of all plastics, including but not limited to synthetic ropes, synthetic fishing nets, and plastic garbage bags, and of all other garbage, including paper products, rags, glass, metal, bottles, crockery, dunnage, lining, and packing materials is prohibited in special areas. Food wastes can only be disposed of as far as practicable from land but in any case not less than 12 nmi from the nearest land (Annex V, Reg. 5.2).

A word of caution is appropriate with respect to the potential merits of the stricter discharge regime of an Annex V special area: The disposal into the sea of all plastics is prohibited everywhere in the world's seas and oceans. The designation of an area as an Annex V special area will not give any additional protection against plastics pollution beyond that. The merit in this respect would be the increased pressure in such areas to provide the necessary port reception facilities for plastics and other garbage. The disposal into the sea of other types of garbage (e.g., dunnage, lining) would be further limited by a designation as an Annex V special area provided the area is large enough to include areas which are more than, respectively, 25 or 12 nmi from the nearest land. If not, discharge regulations similar to those for a special area will apply anyway within 25 nmi from the nearest land for dunnage, lining, and packing

materials which will float, and within 12 nmi from the nearest land for food wastes and all other garbage.

Five sea areas (the Mediterranean Sea, the Baltic Sea, the Black Sea, the Red Sea, and the "Gulfs Area" (Annex V, Reg. 5.1)) have each been designated as Annex V special areas, but this designation is effective only for the Baltic Sea. For the other sea areas, the special area status will enter into force as soon as there are sufficient reception facilities for garbage in the area (Annex V, Reg. 5.4). In October 1989, the decision was made to designate the North Sea as an Annex V special area. A proposal to designate the Gulf of Mexico as an Annex V special area has been discussed by MEPC but no decision to do so has been made.

Criteria for the designation of special areas were developed during MEPC's twenty-sixth session (MEPC 1988a) and were amended in March 1990 during MEPC's twenty-ninth session (MEPC 1990). They include oceanographic conditions (e.g., particular circulation patterns, long residence times, extreme ice states or adverse ice conditions), ecological conditions (e.g., depleted, endangered, or threatened species; areas of high natural productivity; spawning, breeding, and nursery areas; rare or fragile ecosystems or critical habitats), and vessel traffic characteristics. A special area should also be an area of such a size that, were it not a special area, discharges of garbage could be made in the area in accordance with the discharge criteria of Annex V established for open sea areas.

It was also noted that consideration should be given to the extent to which the condition of a sea area is influenced by nonmaritime sources of pollution. Proposals for the designation of a special area will be strengthened by information on measures that are being or will be taken to prevent, reduce, and control pollution of the marine environment by these other sources of pollution.

The Baltic Sea: The First Annex V Special Area

To date, the Baltic Sea is the only Annex V special area to enter into force. At the twenty-sixth session of the MEPC, the Governments of the Baltic Sea States submitted notification to the IMO that adequate reception facilities had been provided in all ports within the Baltic Sea Area (MEPC 1988e). The MEPC then unanimously decided that the Annex V special area status for the Baltic Sea would take effect on 1 October 1989 (MEPC 1988f).

The North Sea: A New Annex V Special Area

At the same twenty-sixth session of the MEPC, the Governments of the North Sea States submitted a proposal to the MEPC to designate the North Sea as an Annex V special area (MEPC 1988b, 1988c). The proposal was finally adopted in October 1989 at the twenty-eighth session of the MEPC.

The proposal was a result of the second International Conference on the Protection of the North Sea, which was held in London in November 1987. The North Sea States were under considerable pressure from some of their members and environmental organizations to designate the North Sea as a special area for the purposes of Annex I (oil) and Annex II (chemicals in

bulk) of the MARPOL Convention. While no agreement on this could be reached during this conference, they did in the end agree to designate the North Sea as an Annex V special area.

Is this proposal superfluous? No, there is good reason for the designation. Reports on plastics and debris in the North Sea area published about the time of the North Sea Conference (e.g., Schrey 1987) have indicated the seriousness of the situation, a situation underscored during the coffee break that followed the decision on Annex V special area status for the North Sea. As delegates to the twenty-sixth session of the MEPC watched, a rising tide brought an influx of garbage up the Thames. The Annex V special area status will contribute considerably to limiting the input of nonplastic or nonsynthetic garbage into the North Sea.

Guidelines: Particularly Sensitive Areas

Criteria and Options

Criteria for the identification of particularly sensitive areas include ecological criteria (uniqueness, dependency, representativeness, diversity, productivity, naturalness, integrity, vulnerability); social, cultural, and economic criteria (economic benefit, recreation, human dependency); and scientific and educational criteria (research interest, suitable conditions for baseline and monitoring studies, opportunities for educational activities, historic value).

Actions already under way may indicate the need for further protective measures. Consideration should be given to the beneficial effects of such measures, in view of the environmental stress from other sources.

Once an area has been identified by the IMO as a particularly sensitive area, the IMO has several options for providing it with additional protection. These include the introduction of special ships' routing measures to increase safety of navigation in or near the area such as vessel traffic separation schemes, deepwater routes, or even vessel traffic management systems. The most important instrument the IMO can use is the designation "area to be avoided."

Areas to Be Avoided

The guidelines will give substantial information about the designation of a particularly sensitive area as an area to be avoided. It is a ship's routing measure "comprising an area within defined limits in which either navigation is particularly hazardous or it is exceptionally important to avoid casualties and which should be avoided by all ships, or certain classes of ships" (IMO 1984). From a number of existing areas to be avoided, it appears to be accepted practice now that "casualties" are interpreted as environmental damage. Areas which for environmental purposes have been designated as areas to be avoided include waters near Cape Terpeniya (Sakhalin, U.S.S.R.), the waters of Nantucket Shoals (United States), a part of the Great Barrier Reef (Australia), an area near the Bermuda Islands (Great Britain), and the Northwestern Hawaiian Islands (United States) (IMO 1984).

The designation of an area as an area to be avoided is not a measure to limit discharges, yet it is consequential to such a decision that in an area where fewer ships are allowed, ships cannot discharge.

The Northwestern Hawaiian Islands (a U.S. wildlife refuge) was established as an area to be avoided after a shipping accident there pointed up the associated risk of pollution. All vessels of more than 1,000 gross tons (GT) carrying cargoes of oil or hazardous materials should avoid the area, which includes the waters within a circle radius of 50 nmi around Pearl and Hermes Reef, Lisianski Island, Laysan Island, Maro Reef, Gardner Pinnacles, French Frigate Shoals, Necker Island, and Nihoa (IMO 1984). There is a substantial area of ocean space beyond 12 and 25 nmi where certain types of garbage can legally be discharged by ships larger than 1,000 GT if these ships can enter the area.

The prevention of discharge of garbage or other marine debris by ships has, of course, not been an objective of this designation. It would nevertheless be interesting to know whether the status of these waters as an area to be avoided has contributed in any way to limit debris-associated pollution.

CONCLUSION

Reduction of discharges at the source on the basis of globally enforced discharge standards should continue to be the first choice when dealing with pollution by marine debris. However, the "guidelines for the designation of special areas and the identification of particularly sensitive areas" can nevertheless provide a useful tool in the protection of specific sea areas against this type of pollution. The designation of special areas in accordance with MARPOL Annex V should, if effectively implemented, prove to be a valuable instrument against marine debris discharged by ships. Measures such as the designation of areas to be avoided could further add to the opportunities to protect specific sea areas against marine debris.

REFERENCES

- Dixon, T. R., and T. J. Dixon.
1981. Marine litter surveillance. *Mar. Pollut. Bull.* 12:289-295.
- International Maritime Organization.
1984. General provisions on ships' routing. *Ships' routing*.
5th ed, International Maritime Organization, London.
- Marine Environment Protection Committee.
1986. Report of the Marine Environment Protection Committee on its twenty-third session. Document MEPC 23/22. International Maritime Organization.
1988a. Annex 10 of the report of the Marine Environment Protection Committee on its twenty-sixth session. Document MEPC 26/25, 28 September 1988, International Maritime Organization.

- 1988b. Consideration of the North Sea as a special area under Annex V, submitted by the United Kingdom. Document MEPC 26/6, 2 June 1988, International Maritime Organization.
- 1988c. Consideration of the North Sea as a special area under Annex V, submitted by the United Kingdom. Document MEPC 26/6/1, 8 July 1988, International Maritime Organization.
- 1988d. Identification of particularly sensitive sea areas, including development of guidelines for designating special areas under Annexes I, II, and V, submitted by Friends of the Earth International (FOEI). Document MEPC 26/INF.20, 5 August 1988, International Maritime Organization.
- 1988e. Notification to IMO concerning the establishment of the Baltic Sea area as a "special area" for the purposes of Annex V of MARPOL 73/78, submitted by Denmark, Finland, the German Democratic Republic, the Federal Republic of Germany, Poland, Sweden and the USSR. Document MEPC 26/10/1, 17 June 1988, International Maritime Organization.
- 1988f. Report of the Marine Environment Protection Committee on its twenty-sixth session. Document MEPC 26/25, 28 September 1988, International Maritime Organization.
- 1989a. Information from an intervention on this issue by the Portuguese delegation during the twenty-ninth session of the Marine Environment Protection Committee of the International Maritime Organization on 16 March 1989.
1990. Report of the working group on particularly sensitive sea areas. Document MEPC 29/WP.16, 15 March 1990, International Maritime Organization.
- Oslo and Paris Commissions.
1984. The Oslo and Paris Commissions, the first decade. International Co-Operation in Protecting our Marine Environment, London.
- Paris Commission.
1987. The Paris Commission. Ninth Annual Report on the Activities of the Paris Commission, London.
- Schrey, E.
1987. Burdening of the German Bight by ship's refuse. Forschungsbericht 102 04 327 UBA-FB 87-041 by Order of Umweltbundesamt, Berlin, November 1987.
-

USING THE PROTECTIVE PRINCIPLE TO UNILATERALLY
ENFORCE TRANSNATIONAL MARINE POLLUTION STANDARDS

Linda M. Paul
815 Pahumele Place
Kailua, Hawaii 96734, U.S.A.

ABSTRACT

Annex V to the 1973 International Convention for the Prevention of Pollution from Ships (MARPOL) contains a broad prohibition against the marine disposal of all plastics; however, not all polluting states have ratified this treaty. While it is a principle of international law that treaty obligations are binding only upon the ratifying states, it may be possible to hold nonsignatory states equally liable for marine pollution damage prohibited by this convention under principles of customary international law. A case can be made that the general principles of liability for marine pollution damage as codified in this treaty and elsewhere represent, in fact, customary international law. Since customary international law relies on nation states to enforce its principles, liability for marine pollution damage may be enforced by national courts against all states, including those who have not yet signed or refuse to sign the international antipollution conventions.

Furthermore, a case can also be made that neither these conventions nor customary international law limits unilateral enforcement of stricter pollution standards, including enforcement on the high seas. The protective principle of international law permits a state to assert temporary jurisdiction over a person or a commercial entity whose conduct outside a state's territory threatens its national interest. By utilizing this transnational jurisdiction principle in the field of international environmental law enforcement, substance can be added to general admonitions not to pollute the high seas. As a protective principle case study, this paper analyzes the problem of renegade gillnets lost or cast aside on the high seas of the North Pacific.

INTRODUCTION

One of the tragic legacies of the 20th century is that the ocean has become a final resting place of man's waste products: oil, toxic wastes,

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

dredge tailings, sewage, and garbage drift with the currents, wash up upon the beaches, or settle upon once productive fishing grounds. Each night in the North Pacific, fishing fleets from Japan, Korea, and Taiwan set some 48,270 to 64,360 km (30,000 to 40,000 mi) of driftnet for a total of more than 8 million km (5 million mi) a year (Wolfe 1989; Marine Mammal Commission 1990). Additional driftnet fleets set thousands of miles of large mesh driftnets in the South Pacific and driftnet fleets from Taiwan have also been reported in the western Atlantic. In addition to their targeted catch, driftnets annually entrap close to a million seabirds and tens of thousands of marine mammals. It is estimated that 30 to 50% of the fish caught and killed drop out of the nets before being brought on board. In addition, over 965 km (600 mi) of the driftnets are thought to be lost or discarded from these fisheries annually. These monofilament net fragments may continue to entrap animals for years, including populations well within the 200-mi exclusive economic zone (EEZ) and territorial waters of the United States. In an effort to document and control the destructiveness of this fishery, the United States Government has attempted repeatedly, yet for the most part unsuccessfully, to reach agreements on cooperative arrangements for research, monitoring, and enforcement of flag-state restrictions on driftnet vessels fishing the high seas (U.S. Congress 1987a; Wolfe 1989).

With an ever-growing world population, marine pollution is a problem that won't diminish and seemingly defies solution. Numerous treaties have been drafted, signed, and ratified, all of which universally condemn pollution of the marine environment and obligate nation states to honor international pollution standards (London Dumping Convention 1972; Stockholm Declaration 1972; MARPOL 1973). The problem is how to enforce the admonition. Many states have refused to sign these treaties; those that have frequently fail to enforce the treaties' provisions.

This paper begins with a general discussion of existing conventional and customary international laws of liability for transnational pollution damage. Since international law in its current state of development relies on nation states to enforce its principles, this paper focuses on the strategies available to states and their citizens to enforce both international and national marine pollution standards. One such strategy is the use of the protective principle of international law that permits a state to assert temporary jurisdiction over a person or a commercial entity whose conduct outside a state's territory threatens its national interest (Levi 1979). By utilizing this transnational principle in the field of international environmental law enforcement, substance can be added to general admonitions not to pollute the high seas.

This paper also explores one particular pollution problem as a protective principle case study: that of renegade gillnets which have either been lost or cast aside on the high seas of the North Pacific. It examines existing treaties and domestic United States legislation applicable to the gillnet problem and outlines some remedial paths that may be taken.

The paper concludes with a proposal for a liability enforcement program that will encourage both domestic fishing vessels and foreign

vessels fishing in the United States EEZ under a Governing International Fishing Agreement (GIFA) to comply with United States law prohibiting the dumping of fishing gear overboard. The implementation of a liability fund program and its financing by the owners of fishing vessels will help prevent the "loss" of drift gillnets and provide compensation for any resource and property damage resulting from discards, as well as a source of funds for resource conservation activities, including biodegradable gear research.

TREATY AND CUSTOMARY INTERNATIONAL LAW

International law comes from several sources: (1) multilateral and bilateral treaties and conventions; (2) international custom as evidenced by the practice of states; (3) general principles recognized by states and articulated by learned scholars in treatises; and (4) judicial decisions by the International Court of Justice (ICJ), other international tribunals and, to a lesser extent, the national courts of sovereign states (ICJ Statute 1945). Of all these sources, only the first, treaty law, is considered "hard law." The rest form a body collectively regarded as customary international law. Customary law, however, may be equally as binding as treaty law depending on its general acceptance by nation states and evidenced by the extent to which those states honor and enforce it. All international law, treaty, and otherwise, depends on individual sovereign states for enforcement.

Customary international law, like treaty law, is being continuously created and modified. While it used to take many decades, if not centuries, of common practice by states to establish that a certain general principle of international law imparted an obligation on all states regardless of any treaty, that process may now be accomplished in a few years. A good example is the existence of a customary law against airline hijacking. Another is the recognition that coastal states now have a right to exercise jurisdiction over the resources of a 200-mi-wide EEZ once considered part of the high seas. The rapidly increasing exploitation of the world's resources, accompanied by a corresponding degradation of the world's environment, also supports an argument that there is a rapidly growing body of customary international law that governs resource conservation and pollution abatement.

One of the general principles of international law recognized by states is that conventional law may give rights to nonparty states, but may obligate only parties to a convention. To the extent that a treaty codifies customary law, however, it is binding on all states, whether or not they are parties. In the last two decades, treaties dealing with the environment have multiplied. Most of these conventions have incorporated into them certain common principles, including the obligation of "reasonable use" of shared resources (ICJ 1974) and the obligation of a state to ensure that activities within its jurisdiction do not damage its neighbors (Stockholm Declaration 1972).

Although international environmental law may impose general obligations to protect and conserve the world's resources, it is dependent on

sovereign states to enforce the obligations. Furthermore, it is understood that while international and regional cooperative efforts to protect the environment are important and should continue, they tend to be slow and ineffective. Most states recognize that neither they nor the planet can afford to wait for committee consensus. As a result, almost every pollution convention includes provisions whereby the contracting parties not only obligate themselves to carry out the terms of the convention, but also reserve the right to take unilateral measures to protect themselves against the acts of other states that threaten their environment (International Convention for the Prevention of Pollution of the Sea by Oil (OILPOL) 1954; Brussels Liability Convention 1969; Intervention 1969; London Dumping Convention 1972; MARPOL 1973; MARPOL Protocol 1978). This treaty reservation is pervasive, and to the extent that it is backed up by states' practice, it can be considered a rule of customary international environmental law.

The unilateral measures reserved by states may include prescribing stricter environmental standards than those provided by conventional international law. The jurisdiction to enforce these stricter standards, however, can only be exercised where a state has personal or territorial jurisdiction. A state has primary jurisdiction over acts committed within its territory; it may also have jurisdiction over transnational acts that substantially affect its territory. This extraterritorial jurisdiction is based on the protective principle.

THE PROTECTIVE PRINCIPLE IN INTERNATIONAL LAW

Based on the ancient doctrine of self-defense, the protective principle permits a state to assert jurisdiction over a legal person (including corporate individuals) whose conduct outside a state's territory threatens its national interests (Levi 1979). The "Restatement (Third) of Foreign Relations Law of the United States" outlines the scope of this principle in §402:

"A state has jurisdiction to prescribe law with respect to . . . (3) certain conduct outside its territory by persons not its nationals that is directed against the security of the state or against a limited class of other state interests." (American Law Institute 1987)

According to the "Restatement (Third)," §402 comment f, the protective principle is considered to be an application of the territorial effects principle. In the U.S. view, if an extraterritorial act produces an effect within a state's territory, that state can assert jurisdiction on the basis of territoriality. The United Kingdom takes a slightly different and more conservative view. If the commission of at least part of an extraterritorial act occurs inside a state's territory, jurisdiction can be asserted on the basis of an "objective territorial principle." When all acts are concluded outside the territory with only the effects felt within the territory, then it falls within a more limited doctrine of "extraterritorial" jurisdiction (Higgins 1984). Whether there is a real distinction between these two viewpoints depends on how jurisdiction is applied to a

particular set of facts. There apparently is no single rule of international law on the applicability of extraterritorial jurisdiction. Different rules are applied to different situations or legal relationships (Weil 1984).

The "Restatement (Third)," §403(1), however, notes that the exercise of any such jurisdiction must not be unreasonable, and §403(2)(a) further notes that states have an obligation to consider "all relevant factors," including whether the extraterritorial activity has a "substantial, direct, and foreseeable effect upon or in the territory." Additional factors to consider that are particularly relevant to transnational pollution activities are included in §402(2)(c) and (f): the importance of the regulation to the regulating state, the extent to which other states regulate such activities, and the degree to which the desirability of such regulation is generally accepted and consistent with the traditions of the international system.

THE PROTECTIVE PRINCIPLE APPLIED TO CIVIL ACTIONS

The jurisdictional regime described by the "Restatement (Third)," §402 and §403, applies to both civil and criminal actions. The civil application of the territorial effects principle to economic activity has been accepted by most Western European states, as well as Canada and Japan. European Economic Community (EEC) law has been applied to extraterritorial enterprises whose anticompetition activities have affected trade within the EEC. The EEC rules on competition, in particular those contained in articles 85 and 86 of the Treaty of Rome, apply to all agreements and practices of "undertakings" that prevent, restrict, or distort competition within the Common Market, even if those undertakings are situated in third countries (Lasok and Bridge 1987).

The United States applies the territorial effects principle in regulating the resale of technically sensitive U.S. products abroad and in antitrust restraint-of-trade actions. In *United States v. Aluminum Co. of America* (1945) Judge Learned Hand imposed a twofold test for the extraterritorial application of U.S. antitrust law: intent to affect the commerce of the United States and actual effect on that commerce. In *Timberlane Lumber Co. v. Bank of America* (1976), a Sherman Antitrust case, the court held that if commerce had in fact been affected, intent to affect was not required.

The use of the territorial effects principle to obtain relief from damaging acts committed by foreign states, as opposed to private foreign nationals, may be barred by the "act of state" doctrine. Under this doctrine, acts by sovereign states on their own territory are immune from the jurisdiction of foreign courts unless the state has expressly or by implication waived its immunity or the activity falls within the commercial exception (Akehurst 1984). The Foreign Sovereign Immunity Act (FSIA), 28 U.S.C. §1602, states that "[u]nder international law, states are not immune from the jurisdiction of foreign courts insofar as their commercial activities are concerned." Section 1603(d) of the FSIA defines "commercial activity" as "either a regular course of commercial conduct or a particular commercial transaction or act." In *Mannington Mills, Inc. v. Congoleum*

Corp. (1979, p. 1292), another case involving the Sherman Antitrust Act, the court held that the act of state doctrine, which would normally preclude private claims based on the contention that the damaging act of another nation violated either American or international law, does not provide a defense "where the governmental action rises no higher than mere approval." Private foreign nationals, unless they are acting as agent of the state, are not protected by the act of state doctrine and can be sued under principles of private international law.

THE PROTECTIVE PRINCIPLE APPLIED TO CRIMINAL ACTS

The United States Supreme Court first recognized the protective principle in *Strassheim v. Daily* (1911, p. 285). Justice Holmes defined the principle as "acts done outside a jurisdiction, but intended to produce and producing detrimental effects within it." Eleven years later, in *United States v. Bowman* (1922, p. 98), the Supreme Court applied the principle to a case involving a conspiracy to defraud the U.S. Government by American citizens on American ships on the high seas. The lower court had dismissed the case since the controlling criminal statute did not expressly confer jurisdiction on U.S. courts for fraudulent acts committed on the high seas. The Supreme Court reversed, holding that jurisdiction could be inferred since certain criminal statutes are "not logically dependent on their locality for the government's jurisdiction, but are enacted because of the right of the government to defend itself. . . ." In 1968, the Second Circuit Court of Appeals applied the principle in *United States v. Pizzarusso* (1968, p. 10), a case involving false statements made abroad by a foreign national to a U.S. consular officer for purposes of obtaining an immigrant visa. The court held that because the false statements had a "potentially adverse effect" upon U.S. governmental functions, they were sufficient to infer jurisdiction.

The bulk of U.S. case law implementing the criminal application of this principle, however, focuses on violations of customs statutes by drug smugglers and the high seas enforcement of these statutes by the U.S. Coast Guard under "Regular Coast Guard: Functions and Powers," 14 U.S.C. §89(a). Although the State Department routinely requests the consent of the foreign flag state before authorizing the Coast Guard to board a foreign commercial ship on the high seas, Federal courts have ruled that this consent is not essential if the boarding is reasonable under the fourth amendment to the Constitution (*United States v. Conroy* 1979; *United States v. Streifel* 1981; *United States v. Alomia-Riascos* 1987). In *United States v. Verdugo-Urquidez* (1990, p. 1059) the Supreme Court held that the fourth amendment does not apply to search and seizure by U.S. agents of property owned by nonresident aliens and located in a foreign country. Presumably this holding might be applied to searches and seizures of foreign vessels on the high seas under the traditional view that flag vessels are extensions of state territory. The court did not consider whether such warrantless searches and seizures had to be reasonable.

THE PROTECTIVE PRINCIPLE APPLIED TO ACTS ON THE HIGH SEAS

The high seas are a jurisdictional void. International law attempted to fill this void by creating a flag-state regime. Flag states once enjoyed

exclusive jurisdiction over ships flying their flags on the high seas under the old notion that a ship was a floating extension of a state's landed territory (Sohn and Gustafson 1984). This fiction is slowly dissolving under a growing number of exceptions, exceptions based on the protective principle and prompted by abuse of the flag-state prerogative. Many commercial ships fly "flags of convenience." By registering "ownership" in a state such as Panama or Liberia that is not a party to conventions that set international navigation, wage, safety, pollution, and resource conservation standards, shipowners can avoid compliance. This common practice belies the territorial justification for flag-state jurisdiction. In *Cunard Steamship Co. v. Mellon* (1922), the U.S. Supreme Court maintained that flag-state jurisdiction "arises out of the nationality of the ship, as established by her domicile, registry, and use of the flag and partakes more of the characteristics of personal than of territorial sovereignty."

For strictly shipboard matters, the flag-state regime is still an acceptable basis for claiming jurisdiction, but when the effects of a ship's activities spread beyond the confines of the ship, flag-state jurisdiction must be qualified. Ship-generated pollution is only the latest in a long list of activities that have caused nonflag states to assert claims of jurisdiction.

Legal inroads into the tradition of exclusive flag-state jurisdiction can be found in the 1958 United Nations Convention on the High Seas (Law of the Sea (LOS) Convention 1958), a codification of customary international sea law up until that date. Article 22 of the LOS Convention (1958) acknowledges that the exclusive jurisdiction of the flag state over its vessels on the high seas is subject to some exceptions: the boarding of foreign merchant ships by warships is permitted if they are suspected of piracy, slave trading, or if they are in fact the same nationality as the warship. Article 28 of the LOS Convention (1958) also codified the protective principle embodied in the doctrine of "hot pursuit." Hot pursuit of a foreign ship is permitted when authorities of a coastal state have good reason to believe that the ship has violated the laws and regulations of that state. Coastal states are allowed to pursue offenders committing illegal acts in the territorial sea and contiguous zone out into the high seas, subject to the limitation that the pursuit be uninterrupted. Pursuit must be terminated when the ship being pursued enters the territorial sea of another coastal state.

The 1982 United Nations Convention on the LOS codified an expanding application of the protective principle in article 110, adding "unauthorized broadcasting" to the 1958 Convention's list of exceptions (LOS Convention 1982). Hot pursuit was expanded in article 111 to include the right to pursue ships committing illegal acts in the EEZ and over the Continental Shelf. In article 108, the LOS Convention (1958) acknowledges by implication that states are applying the protective principle to high seas drug trafficking and attempts to modify unilateral actions by authorizing flag states to request the cooperation of other states in the suppression of such traffic on the high seas.

THE PROTECTIVE PRINCIPLE AND JURISDICTION OVER HIGH SEAS RESOURCES

The expansion of coastal state jurisdiction over the resources of the high seas is a recent phenomenon prompted in large measure by actions of the United States. In 1945, President Truman issued two proclamations, one claiming jurisdiction over the natural resources of the United States' Continental Shelf, the other over high seas fishery resources adjacent to its territorial sea. Prompted by World War II and a growing dependence on foreign sources of fossil fuels, the President proclaimed:

"[T]he Government of the United States regards the natural resources of the subsoil and sea bed of the continental shelf beneath the high seas but contiguous to the coasts of the United States as appertaining to the United States [and] subject to its jurisdiction and control." (Presidential Proclamation No. 2667, 1945.)

In 1953, this proclamation was enacted into law by Congress in the Outer Continental Shelf Lands Act, 43 U.S.C. §1,331-§1,343.

The second Presidential proclamation announced the policy that conservation zones would be established by treaty in areas of the high seas contiguous to the territorial waters of the United States (Presidential Proclamation No. 2668, 1945). In these zones, all fishing would be subject to U.S. regulation and control. Although nation states had been regulating the exploitation of coastal fishing stocks by treaty for more than a hundred years, overfishing still occurred, prompting the United States to invoke a unilateral policy (Sohn and Gustafson 1984). The fishing proclamation was never enacted into statutes, but the United States did thereafter conclude treaties which extended a regional jurisdictional regime over some high seas fishery resources (North Pacific Convention 1952; Halibut Convention 1953).

The Truman proclamations prompted Chile, Ecuador, and Peru to sign the Declaration of Santiago on the Maritime Zone in 1952, whereby each nation claimed complete sovereignty over a 200-mi-wide zone adjacent to its coast (Sohn and Gustafson 1984). In 1958, the High Seas Fishing Convention attempted to limit unilateral extensions of territorial jurisdiction into the high seas that were based on the protective resource conservation rationale. The convention acknowledges that coastal states have a "special interest" in the high seas resources adjacent to their territorial seas, but grants them limited preferential rights in article 6. Article 7 of the convention permits unilateral conservation measures by coastal states only if resource management negotiations between all interested parties have failed.

The concept of a 200-mi EEZ was developed by the Third United Nations Conference on the Law of the Sea in the 1970's. Simultaneously, other states unilaterally claimed their own exclusive zones. In 1972, Iceland established a 50-mi exclusive fishery zone, which was contested by the United Kingdom and the Federal Republic of Germany before the ICJ (1974). In 1976, the United States unilaterally established a 200-mi fishery conservation zone in which it assumed "exclusive fishery management authority over

all fish, except highly migratory species, and. . . authority beyond such zone over such anadromous species and Continental Shelf fishery resources" (Magnuson Fishery Conservation and Management Act (MFCMA) 1986, 16 U.S.C. §1801-§1882). Article 56 of the LOS Convention (1982) gives coastal states "sovereign rights" to explore, exploit, conserve, and manage the natural resources of a 200-mi-wide EEZ, and article 64 permits coastal state jurisdiction over highly migratory species found within that zone. Furthermore, the right to fish on what is left of the high seas is no longer absolute. The right to fish for anadromous stocks on the high seas is contingent upon negotiated agreement (article 66), and the remaining high seas fishing rights are subject to the interests of coastal states (article 116). The convention imposes on all states the duty to cooperate with other states and to take "such measures for their respective nationals as may be necessary for the conservation of the living resources of the high seas" (article 118).

The protective principle is also applied in the LOS Convention (1982) to the conservation of marine mammals. The convention specifies that "[n]othing in this part restricts the right of a coastal State. . . to prohibit, limit or regulate the exploitation of marine mammals more strictly than provided for in this part" (article 65). It also specifies that these provisions apply to the conservation and management of marine mammals on the high seas (article 120). The LOS Convention (1982) seems to sanction the high seas enforcement of coastal state conservation measures, if the high seas activities of foreign nationals "exploit" protected marine mammals.

POLLUTION ENFORCEMENT AND THE PROTECTIVE PRINCIPLE

The protective principle can also be found in the pollution provisions of the LOS Convention (1982). Exclusive flag-state jurisdiction over vessels beyond the territorial sea had not succeeded in eliminating vessel-source pollution or in protecting the environment of coastal states from it (Boyle 1985). The LOS Convention (1982) authorizes port states to investigate and prosecute foreign vessels voluntarily coming into their jurisdiction who have violated international pollution rules and standards on the high seas (article 218(1)). Presumably these standards are those embodied in the International Convention for the Prevention of Pollution from Ships (MARPOL 1973) and the International Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (London Dumping Convention 1972). This provision represents an expansion of port state jurisdiction beyond that authorized by MARPOL, which limits port states to inspection and the reporting of any violation of the 1973 convention to the flag state.

Coastal states under the LOS Convention (1982) may enforce their own pollution rules and standards against foreign vessels voluntarily coming into their ports, but only with respect to acts committed in their territorial waters and EEZ's (article 220(1)). As for stopping ships in transit passage, a coastal state may enforce its own laws for acts committed in its territorial waters (article 220(2)), but may enforce only international pollution standards for acts committed in its EEZ, provided they result "in a discharge causing major damage or the threat of major damage to the coastline or related interests of the coastal State, or to any resources of its territorial sea or exclusive economic zone" (article 220(6)).

Both port state and coastal state jurisdiction can be preempted by the flag state (article 228(1)), subject to the provision that the flag state continue the proceedings within 6 months and unless there is major damage to the coastal state or the flag state "has repeatedly disregarded its obligations to enforce effectively the applicable international rules and standards in respect of violations committed by its vessels" (article 228(2)).

The 1982 LOS Convention is not yet in force. However, aside from the provisions regarding the deep seabed in part IX, it may be considered to represent a codification of both existing and developing customary international law. Those provisions which are not found in other multilateral treaties or do not represent the practice of a majority of states are, of course, not yet part of international law. Yet most of the provisions cited above represent only an expansion of existing principles already established. For example, LOS Convention (1982) provisions dealing with marine pollution represent an extension of the standards set forth in a plethora of conventions which came into being as a result of maritime oil disasters. The obligations in each of the pollution conventions cited below are applicable only to the states' parties, but taken in aggregate, the conventions establish the right to a pollution-free environment that is shared by all states. Many of the provisions noted in the following conventions represent an emerging regime of customary international pollution law, law that is based on the protective principle.

THE PROTECTIVE PRINCIPLE IN THE MARINE POLLUTION CONVENTIONS

The protective principle is incorporated into the International Convention for the Prevention of Pollution of the Sea by Oil (OILPOL 1954: article 11), the Convention Relating to Intervention on the High Seas in Cases of Oil Pollution Casualties (Intervention 1969, article 1), the International Convention on Civil Liability for Oil Pollution Damage (Brussels Liability Convention 1969), the Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (London Dumping Convention 1972, article 7, 13), the Protocol Relating to Intervention on the High Seas in Cases of Marine Pollution by Substances Other Than Oil (High Seas Protocol 1973, article 1), and the International Convention for the Prevention of Pollution from Ships (MARPOL 1973, article 9). For example, Intervention (1969) specifies:

"Parties to the present convention may take such measures on the high seas as may be necessary to prevent, mitigate or eliminate grave and imminent danger to their coastline or related interests from pollution or threat of pollution of the sea by oil, following upon a maritime casualty or acts related to such a casualty, which may reasonably be expected to result in major harmful consequences" (article 1(1)).

This same provision can also be found in the High Seas Protocol (1973). "Substances other than oil" in this treaty include, inter alia, other substances "which are liable to create hazards to human health, to harm living resources and marine life, to damage amenities or to interfere with other legitimate uses of the sea" (article 1(2)).

Relevant provisions in the 1973 MARPOL Convention, conceived in the early seventies, are considerably more conservative in terms of the protective principle than current states' practice and evolving customary international law as is evidenced by the LOS Convention (1982). The MARPOL Convention, while imposing universal obligations not to pollute, essentially leaves its high seas pollution enforcement up to the flag state, limiting the port state to inspection and reporting activities. Coastal states are given the option of either reporting a violation occurring within their jurisdiction to the flag state or initiating proceedings on their own (MARPOL 1973, article 4(2)). The extent of coastal state jurisdiction, however, was the source of considerable controversy during the drafting of the treaty. The issue was never resolved, and as a result the phrase "within the jurisdiction of any Party" (article 4) must be interpreted according to current interpretations of the term in international law (Timagenis 1980).

The MARPOL negotiators also disagreed over the power of states to take stricter regulatory measures than those expressly provided for in the convention. The compromise text, adopted by a majority of the delegates, but not by the required two-thirds, provided that

"(1) Nothing in the present Convention shall be construed as derogating from the powers of any Contracting state to take more stringent measures where circumstances so warrant, within its jurisdiction, in respect of discharge standards" (article 8, now article 9).

In its final form, MARPOL defers to future decisions by the United Nations Conference on the LOS, stipulating only that nothing in MARPOL "shall prejudice. . .the present or future claims and legal views of any State concerning the law of the sea and the nature and extent of coastal and flag State jurisdiction" (article 9(2)). It leaves interpretation of the term "jurisdiction" up to customary international law (article 9(3)).

With the failure of the compromise text, Australia reserved the right to impose unilateral conditions within its jurisdiction to protect its "adjacent" marine environment from pollution (Timagenis 1980, p. 503). The Canadian delegation stated that since there was no provision restricting the powers of the contracting states to take measures within their jurisdiction, the Canadian Government reserved the right "to take any and all measures within its jurisdiction for the protection of its coasts and the adjacent marine environment from pollution from ships" (Timagenis 1980, p. 505). Ireland and the Philippines made similar statements (Timagenis 1980, p. 506).

THE PROTECTIVE PRINCIPLE AND STATES' PRACTICE--A CANADIAN EXAMPLE

In 1970, Canada unilaterally implemented the protective principle in response to a pollution threat to its Arctic coastline. Fearful of a devastating accident associated with the United States' development of the Alaskan North Slope and the use of the Northwest Passage for transport of crude oil to refineries on the east coast, the Canadian Parliament passed

the Arctic Waters Pollution Prevention Bill, extending Canadian jurisdiction 100 nmi into the Beaufort Sea for purposes of regulating marine pollution. The act held both prospector and ship owner liable for all costs associated with any discharge of wastes into this region (Canadian Arctic Waters Pollution Prevention Bill 1970). In reply to a protest letter from the United States, the Canadian Government responded "that a danger to the environment of a state constitutes a threat to its security. . . . The proposed anti-pollution legislation is based on the overriding right of self-defense of coastal states to protect themselves against grave threats to their environment" (Canadian Reply 1970). This unilateral declaration caused considerable controversy in the world community. The 1982 LOS Convention resolved the controversy with a special provision granting coastal states the right to adopt and enforce laws and regulations to prevent and control pollution in ice-covered areas within the EEZ (article 234).

THE PROTECTIVE PRINCIPLE IN UNITED STATES ENVIRONMENTAL LEGISLATION

In 1961, Congress passed the Oil Pollution Act, 33 U.S.C. §1001-§1015, implementing OILPOL. Over the next decade the convention was amended and more legislation was enacted as marine pollution and public awareness grew. In 1974, the Intervention on the High Seas Act, 33 U.S.C. §1471-§1487, implemented Intervention 1969 and its companion treaty dealing with substances other than oil. This legislation, extending the protective principle to pollution activities on the high seas, empowers the United States to take measures to "prevent, mitigate or eliminate" danger to its coastline caused by oil pollution casualties on the high seas when they "may reasonably be expected to result in major harmful consequences" (§1472).

The 1973 version of MARPOL did not enter into force before it was superseded by the Protocol of 1978, which extensively modified Annex I (Regulation for the Prevention of Pollution by Oil) and Annex II (Regulations for the Control of Pollution by Noxious Liquid Substances in Bulk). The modifications in the pollution prevention standards were the result of an international conference convened at the request of the United States following 16 major tanker accidents in waters around the United States in 1976-77 (U.S. Congress 1980). Although the MARPOL Protocol did not come into force until 1983, many of its provisions were immediately implemented by Congress in the Port and Tanker Safety Act of 1978, 33 U.S.C. §125, §128-§1232 and 46 U.S.C. §391(a), §1221, §1224. The Senate ratified the 1978 MARPOL Protocol in 1980 and implemented the balance of its provisions that year in the Act to Prevent Pollution from Ships, 33 U.S.C. §1901-§1911. The other three original MARPOL annexes are optional. Annex III and Annex IV have not been ratified by the United States and are not in force.

Annex V of the original 1973 MARPOL Convention came into force in December 1988, with U.S. ratification. Not revised by the Protocol of 1978, Annex V prohibits the disposal at sea of garbage and plastic waste; however it excepts the accidental loss of synthetic fishing nets incidental to the making of repairs from its general prohibition. Congress implemented Annex V on 29 December 1987 with an amendment to the Act to Prevent Pollution from Ships, adding simply the word "garbage" to its list of prohibited discharges, 33 U.S.C. §1907(d)(1) and (e)(1). Although the

Annex V amendments continue to defer to the flag-state enforcement scheme of the MARPOL Protocol as enacted into U.S. law in 1980 (§1907(c)), they also include a preemption requirement mandating that "[e]xcept as specifically provided. . . , nothing in this title shall be interpreted or construed to supersede or preempt any other provision of Federal or state law. . . [or] any State from imposing any additional requirements" (33 U.S.C. §1901(a)).

The MFCMA regulations regarding fishing gear disposal were also amended in 1987. Previously these regulations had exempted all accidental gear loss from its general prohibitions. Now any gear discards, including accidental discards, are strictly prohibited except in case of emergency or with specific Coast Guard authorization (50 C.F.R. §611.12). This is a stricter discard standard than that of MARPOL Annex V.

In conjunction with implementation of Annex V, Congress also enacted the Driftnet Impact Monitoring, Assessment, and Control Act of 1987, 16 U.S.C. §1822. This act requires the Secretary of Commerce to assess the impact of driftnets on U.S. marine resources and the Secretary of State to initiate

"negotiations with each foreign government that conducts, or authorizes its nationals to conduct driftnet fishing that results in the taking of marine resources of the United States in waters of the North Pacific Ocean outside of the exclusive economic zone and territorial sea of any nation, for the purpose of entering into agreements for effective enforcement of laws, regulations, and agreements applicable to the location, season, and other aspects of the operations of the foreign government's driftnet fishing vessels." (16 U.S.C. §1822)

The act further specifies that if the foreign government fails to enter into and implement such an agreement within 18 months, certification under the Pelly Amendment will result (16 U.S.C. §1822, 22 U.S.C. §1978(a)). The Pelly Amendment provides that when the nationals of a foreign country, directly or indirectly, are (1) conducting fishing operations in a manner or under circumstances which diminish the effectiveness of an international fishery conservation program, or (2) engaging in trade or taking which diminishes the effectiveness of any international program for endangered or threatened species, the Secretary in charge of the finding shall certify the fact to the President. The President in turn may then direct the Secretary of the Treasury to prohibit the bringing or importation in the United States of fish products of the country whose nationals are engaging in such conduct (22 U.S.C. §1978(a)).

Congressional intent on the extent of jurisdiction can be found in the final committee report to the House. The Merchant Marine and Fisheries Committee expressly stated that no extension of U.S. jurisdiction was to be construed by passage of the legislation. It affirmed, however, "the sovereign rights of the United States to conserve and manage marine resources within its exclusive economic zone and anadromous species on the high seas to the extent provided for in United States law" (U.S. Congress 1987a).

**UNITED STATES MARINE RESOURCE CONSERVATION
ENFORCEMENT AND THE PROTECTIVE PRINCIPLE**

In only a few instances has the United States asserted through active enforcement the protective principle codified in its various conservation and pollution laws. Current case law reveals an enforcement focus on violations of the MFCMA in the EEZ by foreign fishing nations signing GIFA's with the United States. A GIFA is an executive agreement between the United States and a foreign state which gives the foreign state the right to fish for specified quantities of designated species in the United States EEZ for a specified period of time. In return, the foreign fishing nation agrees to abide by the terms of the MFCMA (Japanese GIFA 1982; Korean GIFA 1982; Taiwanese GIFA 1982). Consent to be boarded and inspected for suspected violations, included in the agreement, considerably simplifies jurisdictional questions. Federal courts have assumed GIFA jurisdiction for violations of the MFCMA in the EEZ in *United States v. Tsuda Maru*, 470 F.Supp. 1223 (D. Alaska 1979); *United States v. Marunaka Maru No. 88*, 559 F.Supp. 1365 (D. Alaska 1983).

The signatory state also must agree to comply with any administrative measures taken in accordance with the agreement and to pledge not to kill or harass any marine mammal within the EEZ without express authorization. If the state fails to agree to acceptable monitoring and enforcement arrangements, the Secretary of Commerce has the authority to deny fishing permits to any of its vessels. In addition, Federal regulations implementing the MFCMA prohibit a foreign fishing vessel from throwing overboard any article that may damage any marine resource, including marine mammals and birds (50 C.F.R. §611.12(c)). This regulation is now reinforced by legislation implementing Annex V of the MARPOL treaty.

A GIFA is enforced by both overflight and boarding inspections by the Coast Guard and by placing National Marine Fisheries Service (NMFS) observers on board some foreign vessels fishing in the EEZ. A few observers have also been placed on the Japanese salmon mothership fleet operating in the high seas of the North Pacific (Marine Mammal Commission 1990). In addition, three United States observers were permitted to observe commercial squid driftnet operations on board Japanese vessels in 1982 (Cary and Burgner 1983) and 1986 (Tsunoda 1989) and on a Korean vessel in 1988 (Gooder 1989). A few additional observations have been made by observers from the vantage point of Coast Guard cutters (Ignell et al. 1986) and private vessels. In 1989, following a mandate from Congress (Driftnet Impact Monitoring, Assessment, and Control Act 1987) agreements were concluded with Japan, Korea, and Taiwan to place a few observers on board commercial driftnet vessels. Japan agreed to allow 9 United States and 5 Canadian observers on board 14 of its commercial squid driftnet vessels during the 1989 season (Japan-United States Agreement 2 May 1989). Agreements with Korea and Taiwan were concluded too late to implement a foreign observer program for 1989. Korea agreed to deploy "at least 13" United States observers on board its "commercial driftnet vessels for at least 45 days each to observe 45 or more driftnet retrievals on each vessel" (Republic of Korea-United States Agreement Annex II, 13 September 1989). Taiwan agreed only to deploy "observers of the parties. . . aboard

commercial driftnet vessels. . .for at least 30 days to observe 30 or more driftnet retrievals on each vessel" (American Institute of Taiwan-Coordination Council for North American Affairs Agreement article VII, 13 July 1989).

Data from the joint Canadian-Japan-United States observer program were collected between early June and early October 1989, by observers on board commercial Japanese driftnet vessels chosen by lot fishing between long. 170°E and 145°W, and between lat. 36° and 45°N. The results were released on 30 June 1990 (Joint Report Fisheries Agency of Japan, Canada Department of Fisheries and Oceans, NMFS, and Fish and Wildlife Service (FWS) 1990). In summary 1,402 operations were observed involving 1,427,225 tans of net (50-m each). The catch included 3,119,061 neon flying squid, 914 dolphins, 22 marine turtles, 9,173 seabirds, and 1,580,068 nontargeted fish, including 79 salmonids, 1,433,496 pomfrets, 59,060 albacore, 10,495 yellowtail, 7,155 skipjack tuna, and 58,100 blue shark. The data did not indicate what percentage of the huge incidental catch of assorted fishes were actually brought on board and kept.

Estimates of incidental catch rates of nontargeted species vary greatly depending on the year, the vessel, its location, the time of year, who is doing the reporting, and the proximity of the observer. The total number of sets observed and recorded is extremely small compared to the total amount of driftnet fishing actually done. Obtaining statistically significant data may be an impossible goal. Between 1978 and 1983 the amount of fishing effort by the driftnet fleet rose dramatically and exponentially; since then it has increased much more gradually. Thus the maximum incidental catch of long-lived species such as marine mammals and turtles would have occurred in the early eighties at a time when practically no data were being kept. Incidental catches today probably reflect the impact of driftnet fishing on much reduced populations. Fishermen have also been observed shaking undesired species free of the nets before hauling them on board (Gooder 1989; Tsunoda 1989), a factor which would distort vessel owners' reports of total incidental takings. Although there have been few observations of torn netting being discarded at sea, this may have been influenced by the fact that observers were watching.

In 1984, a NMFS observer on board a Japanese vessel fishing in the EEZ near Dutch Harbor, Alaska, saw its crew members on five separate occasions toss several 0.45 x 0.76-m (1.5 x 2.5-ft) fragments of synthetic trawl netting overboard. At a hearing before a National Oceanic and Atmospheric Administration (NOAA) administrative law judge, expert testimony revealed that some 30% of documented fur seal entanglements involve net fragments in the same size range as those discarded and some 40% of the seals caught in such net fragments die (*In the Matter of Kenji Nakata, Ohoura Gyogyo Co. Ltd.*, 4 O.R.W. 814 (NOAA 1987)). In his decision, the judge indicated that every vessel discarding such net fragments must be held responsible for the cumulative effect on the fur seal population and that the civil penalties assessed for such violations must be large enough to serve as a deterrent and not be considered as a mere cost of doing business. The owners of the offending vessel were fined \$15,000 total for two violations. The MFCMA authorizes up to \$25,000 assessment for each violation (16 U.S.C. §1858(a)).

Although it has been asserted that lost sections of driftnet ball up within a week and thus no longer ghost fish, evidence indicates that tangled driftnets continue to ghost fish, both on the surface (DeGange and Newby 1980; Henderson 1984; von Brandt 1984; Gooder 1989) and on the bottom (Way 1977; Carr and Cooper 1987). In 1978, a 3,500-m section of lost driftnet was found floating in the North Pacific (lat. 49°15'N, long. 168°14'E). Entangled within the 1,500 m brought on board were 75 newly snared salmon and at least twice that many rotten ones, plus assorted other fish and some 99 seabirds (DeGange and Newby 1980). Endangered Hawaiian monk seals have also been found entangled in masses of monofilament drift-net (Henderson 1984).

THE NORTH PACIFIC DRIFTNET FISHERY--A CASE FOR APPLICATION OF THE PROTECTIVE PRINCIPLE

The 20th century development of nylon and plastics has greatly benefited human "progress" and greatly burdened the global environment. Except for the hulls of vessels, plastics dominate the fishing industry: polystyrene and polyurethane foams are used for flotation in lifejackets and fishnet floats; polyethylene is used to make cable coverings, buckets, packaging film, and shipping containers; polyamide (nylon), polyester, polyethylene, and polypropylene are used for ropes and nets (Pruter 1987). Gillnets, one of the most efficient fishing methods ever developed, are made of monofilament nylon. Strong, transparent, and durable, nylon is the perfect material for constructing the miles of pelagic drift gillnets used by high seas salmon and squid fishermen. Gillnets are also cheap, difficult to mend, and easily replaced, usually after only one season of use (Eisenbud 1985). Pelagic gillnets are made in panels 6-8 m deep and may be over a 100 m long. These individual panels are strung together by the fishermen on lines with floats on surface and weights on the bottom to create an invisible wall which may stretch for miles. Set at night, the gillnets are either left to drift free until the following morning, or remain attached to the catcher boat at one end.

There are two Japanese salmon driftnet fisheries: a mothership fishery in the Bering Sea and a land-based fishery that operates south of the Aleutian Islands. They are regulated by the International North Pacific Fisheries Commission, established to implement the provisions of the 1952 North Pacific Convention. The North Pacific Convention is a tripartite agreement between the United States, Canada, and Japan to: "1) ensure cooperation in scientific research and data collection on salmon and other fish species in the North Pacific Ocean and Bering Sea; 2) minimize interceptions of North American origin salmon by Japan; and 3) facilitate cooperation in marine mammal research" (Beasley 1984). Each party agreed to enact and enforce the necessary domestic laws and regulations to implement the convention provisions (North Pacific Convention, article 9(2); 16 U.S.C. §1021-§1035).

In 1976, the MFCMA claimed exclusive management authority over anadromous species of U.S. origin throughout their range, unless they are within the jurisdiction of another nation (16 U.S.C. §1812). The MFCMA required that the North Pacific Convention be renegotiated to the extent that it was in conflict with the act (16 U.S.C. §1822(b)). In 1978 a

protocol was signed amending the North Pacific Convention and shifting the eastern boundary of the salmon mothership fishing area from long. 175°W to 175°E, except for a small enclosed area of high seas in the middle of the Bering Sea known as the donut hole (North Pacific Convention Protocol 1978). The Japanese land-based fleet is limited to an area south of lat. 46°N east of long. 170°E, and south of lat. 48°N west of long. 170°E. The Japanese also have a treaty with the Soviet Union which excludes all salmon fishing within the Soviet EEZ and regulates high seas salmon fishing outside of this zone (Japan-U.S.S.R. 1956). In May 1990, the Soviet Union seized a fleet of more than 10 "North Korean" driftnet vessels for harvesting thousands of tons of Soviet salmon. The fleet had previously been spotted by the U.S. Coast Guard fishing south of the Aleutian Islands. North Korea is not a party to any of the North Pacific fishing agreements. Following arrest it was discovered that the fleet and more than 140 fishermen were actually Japanese and were fishing under the North Korean flag to circumvent treaty restrictions.

The high seas driftnet fisheries are condemned by many as an economically inefficient method of fishing. They catch salmon before they reach full size and can bring the maximum market price, as well as nontargeted species, which are discarded; 50% of the fish caught are estimated to drop out of the nets before they can be brought on board; the various salmon stocks mingle on the high seas making it almost impossible to manage individual stocks; and driftnets are lost or discarded at sea where they continue to entrap fish (Sathre 1986). Furthermore, the high seas squid driftnet fisheries of Japan, the Republic of Korea, and Taiwan have high "incidental" catches of salmon which may not be so incidental (the Japanese squid fishery uses driftnets similar to those used by the salmon fishery) (Anderson 1989; Matsen 1989). Begun in 1978, the Japanese squid fishery now extends across much of the North Pacific and by agreement is supposed to stay in waters too warm for salmon (15°-22°C). However, the Coast Guard has sighted numerous Japanese vessels fishing north of the boundaries established by Japanese regulations (Gordon 1985). Furthermore, neither the Republic of Korea nor Taiwan is bound by the 1978 Protocol's salmon fishing boundary restrictions, although Taiwan has agreed to respect a squid fishing boundary similar to Japan (Gordon 1985). In 1988, the Soviets seized three Taiwan driftnet vessels reportedly carrying 3,357,000 kg (9 million lb) of immature salmon (Anderson 1989). As for MARPOL prohibitions against abandoning old driftnets, only Japan is currently a party to Annex V.

Driftnets fish indiscriminately. While mesh size is targeted for a particular species, the nets are capable of catching almost anything that swims or dives: small whales, porpoises and dolphins, seals and sea lions, turtles, and almost every kind of diving seabird have been found trapped in them (DeGange and Newby 1980). Lost and discarded net sections from high seas gillnet fisheries frequently drift into the U.S. EEZ and continue to entrap large numbers of migratory seabirds and marine mammals. This gear may ghost fish for years before settling on the bottom or washing up on a beach. Although the "act" of losing this deadly gear may have occurred on the high seas, the effect constitutes a taking of migratory seabirds and marine mammals within U.S. EEZ and territorial waters that is in direct

contravention of both treaty and U.S. domestic law. This entrapment of birds and mammals by ghost nets drifting into the EEZ is arguably illegal under the "taking" provisions of the Marine Mammal Protection Act (MMPA), 16 U.S.C. §1361-§1407, the Endangered Species Act (ESA), 16 U.S.C. §1531-§1542, the North Pacific Fur Seals Act, 16 U.S.C. §1151-§1175, and the Migratory Bird Treaty Act (MBTA), 16 U.S.C. §703-§712.

The MMPA defines the term "take" as meaning to harass, hunt, capture, or kill, or attempt to harass, hunt, capture, or kill any marine mammal (16 U.S.C. §1362(12)). Pursuant to this act, a permit is required for any incidental take of marine mammals in the course of commercial fishing operations (§1362(14)). The goal, however, is that incidental takes "be reduced to insignificant levels approaching zero" (§1371(a)(2)). The MMPA also authorizes the Secretary of the Treasury to ban the importation of fish products from fish which have been caught using technology that results in killing or seriously injuring marine mammals "in excess of United States standards" (§1371(a)(2)). Amendments to the MMPA passed in 1988, however, permit exemptions from the incidental taking provisions until 1 October 1993, provided that fishing vessels keep detailed annual records of all such takings (Public Law 100-711, 23 November 1988, adding §114; 50 C.F.R. part 229).

The ESA was created to "provide for the conservation, protection and propagation of endangered species of fish and wildlife by Federal action. . ." (U.S. Congress 1973). The ESA makes it "unlawful for any person subject to the jurisdiction of the United States to take any protected species on the high seas" (16 U.S.C. §1538(a)(1)(C)). Any exceptions require permits (16 U.S.C. §1539 and 50 C.F.R. §10.1 et seq.). The ESA further authorizes the Secretaries of Commerce and State to enter into bilateral or multilateral agreements for the protection and conservation of endangered and threatened species and to encourage citizens of foreign states "who directly or indirectly take fish or wildlife or plants in foreign countries or on the high seas for importation into the United States for commercial purposes. . .to carry out. . .conservation practices designed to enhance such fish or wildlife or plants and their habitat" (16 U.S.C. §1537(b)). The enforcement provisions of the ESA permit any U.S. citizen to bring a civil suit to enjoin any person, including the United States Government, who may be in violation of the ESA (§1540(g)). Worldwide service of process may be implied in the language of the ESA, 16 U.S.C. §1540(c), which specifies that "the several district courts of the United States. . .shall have jurisdiction over any action arising under this chapter." According to reporters' notes 7 and 9 of §421 of the Restatement (Third), where nationwide jurisdiction is conferred by statute, jurisdiction of a Federal court no longer depends on the laws of the state where the court sits. Evidence of an intent to apply the ESA extraterritorily has been found by courts (*Defenders of Wildlife v. Hodel* 1989) in the broad definition of "endangered species," §1532(6), which includes many species not native to or present in the United States, along with §1538(a)(1)(C), which prohibits takings on the high seas, coupled with §1532(13), which defines "persons" to include foreign governments.

Japan and the United States are parties to the bilateral 1972 Migratory Bird Convention, which prohibits the taking of migratory birds in

the territories of both countries (Migratory Bird Convention 1972). The annex to the convention lists 189 protected species; virtually all are species listed in regulations implementing the ESA and the MBTA (50 C.F.R. §10.13). In addition to prohibitions, the convention also includes some affirmative duties. The Migratory Bird Convention mandates that each contracting party "shall (a) [s]eek means to prevent damage to such birds and their environment, including, especially, damage resulting from pollution of the seas. . ." (article 6(a)). Article 6(a) also specifies that "[e]ach Contracting Party agrees to take measures necessary to carry out the purposes of this Convention." Of the 800,000 seabirds estimated to die each year in the gillnets of the Japanese mothership and land-based salmon fisheries of the North Pacific, many belong to species that are listed in the Migratory Bird Convention Annex (U.S. Congress 1987b). Data from the 1989 Canadian-Japan-United States squid driftnet observer program included birds from at least five species listed in the annex (331 Laysan albatrosses, 38 northern fulmars, 20 horned puffins, 5 tufted puffins, and 17 Leach's storm petrels). Ghost driftnet data have also reported catches of listed birds. In one mass of tangled driftnet, DeGange and Newby (1980) identified birds from 6 listed species (4 Laysan albatrosses, 15 northern fulmars, 15 tufted puffins, 14 sooty shearwaters, 40 slender-billed (short tailed) shearwaters, and several fork-tailed storm petrels).

The Migratory Bird Convention is implemented in part in the MBTA (16 U.S.C. §703-§712). The MBTA prohibits the taking of listed migratory birds without a permit from the Department of the Interior (16 U.S.C. §703 and 50 C.F.R. §10.13 and §21.11). In recent testimony given by the Department of the Interior on driftnet takings of migratory seabirds by the Japanese salmon fisheries, there was no mention of any permits having been issued (Lambertson 1985), possibly because the FWS does not believe it has jurisdiction. However, under customary international law, as codified in the 1982 LOS Treaty and recognized as such by Presidential Proclamation No. 5030 on 10 March 1983, the United States has jurisdiction over all "living resources" of its EEZ. The term "marine resources" has been defined by Congress in the 1986 amendments to the MFCMA as including "fish, shellfish, marine mammals, seabirds, and other forms of marine life or waterfowl."

Japan is also a party, along with the Soviet Union, Canada, and the United States, to the 1957 Interim Convention on the Conservation of North Pacific Fur Seals prohibiting the taking of fur seals in the North Pacific Ocean (Fur Seal Convention 1957). The Fur Seal Convention empowers any party to board and inspect another party's fishing vessel anywhere except in territorial waters if there is reasonable cause to suspect that the vessel is violating the prohibition against sealing (article 6). Pelagic "sealing" is defined as the killing, taking, or hunting in any manner whatsoever of fur seals at sea" (article 1). If the suspicion is well founded, the vessel may be seized and the persons on board arrested. Judicial proceedings, however, are left to the flag state. This treaty has been enacted into U.S. law in the North Pacific Fur Seals Act, 16 U.S.C. §1151-§1175. The act, however, permits U.S. flag fishing vessels to refuse boarding if they are within the U.S. EEZ (§1152). Fur seals are, of course, also covered under the MMPA. Although the NMFS has issued permits to GIFA holders for the incidental taking of fur seals and sea lions, NMFS was enjoined by an order of the U.S. District Court in Washington, D.C., from

issuing a permit for the 1988 season (*Kokechik Fishermen's Association v. Secretary of Commerce*, 839 F.2d 795 (D.C. Circuit 1988)). The court ruled that the northern fur seal, *Callorhinus ursinus*, incidentally taken by the Japanese mothership salmon driftnet fleet are depleted below optimum levels and that the MMPA prohibits the issuance of a permit to take any such depleted marine mammals.

ACTION TO BAN DRIFTNETS BY THE INTERNATIONAL COMMUNITY

Several foreign states have already taken action to ban driftnet fishing in waters under their jurisdiction, including Japan, Australia, Canada, the Cook Islands, the Federated States of Micronesia, New Zealand, Peru, French Polynesia, American Samoa, and Vanuatu. In July 1989, the South Pacific Forum states signed the Tarawa Declaration, which expressed regret that Japan and Taiwan "failed to respond to the concerns of regional countries" about pelagic driftnet fishing and called for a ban on driftnet fishing in the South Pacific. Korea had already announced its intention to withdraw its driftnet vessels from the South Pacific area. In November 1989, several South Pacific states signed the Convention for the Prohibition of Fishing with Long Driftnets in the South Pacific. The convention bans driftnet fishing within the 200-mi EEZ's of the signatory states and within certain adjacent high seas areas. In December 1989, the United Nations General Assembly unanimously adopted a resolution that condemns the commercial use of driftnets and calls for a ban on driftnet fishing in the South Pacific, beginning 1 July 1991, and a worldwide ban beginning 30 June 1992. [On 26 June 1990, Taiwan announced that it would prohibit its driftnet fleet from fishing in the South Pacific by July 1991.]

REMEDIES FOR VIOLATIONS OF INTERNATIONAL ENVIRONMENTAL STANDARDS

If activities within a state's jurisdiction cause significant damage to its neighbors or the common environment, that state is responsible to all injured states for violation of its obligations. A state may be obligated by either treaty or customary international law. If the obligation is owed to the international community as a whole, any state may bring a claim before the ICJ without showing that it has suffered a particular injury (American Law Institute 1987, §902(1), §602 comment a). Consent to jurisdiction of the ICJ by the defendant state is not necessary under the Statute of the ICJ. In general, the state seeking redress has the burden of proving the existence of an international obligation and its breach; the responding state has the burden of establishing any justification or excuse. The burden of proof may shift to the party that has control of the evidence (American Law Institute 1987, §901 comment a). Thus any state may call on the violating state to terminate activities which are causing significant injury to the general environment.

If the obligation is owed to a particular state or states, the injured states may also file international claims. Treaties generally include a dispute resolution clause in their text. If the method chosen does not produce results, or if there is no such clause, the injured state(s) may take its grievance before an international tribunal with the consent of the responding state or the ICJ without the responding state's consent. The enforcement power of the ICJ, however, is that of world opinion and

voluntary compliance by sovereign states. If a citizen of a state has a cause of action, it must be taken up by his parent government before the ICJ. Article 34 of the Statute of the ICJ gives it jurisdiction only over states. The injured person must also have exhausted local remedies (American Law Institute 1987, §703 comment b).

A state may also sue a fellow sovereign state in any domestic court that will accept jurisdiction, but because of the principle of sovereign immunity, domestic courts will generally accept jurisdiction only if a state waives its sovereign immunity or if the offending acts attributable to the state involve commercial activities (Foreign Sovereign Immunities Act, 28 U.S.C. §1605(a)(2)). Because of the vagueness of customary international law and the domestic court's general unfamiliarity with it, plus the fact that, at least in the United States, foreign policy issues are reserved to the executive branch, adjudication in a domestic forum stands a better chance of success if arguments are based on domestic law.

Private persons, including organizations, may also bring suit when international environmental standards are violated. Injured persons may bring a claim directly against a foreign state in an international forum when that state has consented to the forum's jurisdiction, or in any domestic court that will accept jurisdiction (American Law Institute 1987, §906, §907). In the United States, Federal courts have jurisdiction over cases arising under international law and international agreements of the United States. However, "international agreements, even those directly benefiting private persons, generally do not create private rights or provide for a private cause of action in domestic courts" unless the provisions are "self-executing" (American Law Institute 1987, §907 comment a).

Whether a treaty of its own force makes law depends on two requirements: (1) the treaty-drafters must have intended that the treaty provision be self-executing as ascertained by the applicable international rules of treaty interpretation (Vienna Convention on the Law of Treaties 1969) and (2) the treaty provision in question must have the force of legislation without any further action by a legislative body (Henry 1928). Treaty obligations not to act, or to act only subject to limitations, are generally self-executing. Self-executing obligations may also arise under customary international law. In conjunction with driftnet fishing there are two customary obligations that may be considered self-executing: (1) "fisheries for anadromous stocks shall be conducted only in waters landward of the outer limits of exclusive economic zones" (LOS Convention 1982, article 66(3)(a)) and (2) "the taking of endangered species is prohibited" (implied in a whole host of multilateral and bilateral treaties). Rules of customary international law are part of U.S. law and as such may permit domestic remedies. Again, suits against foreign states may be barred by sovereign immunity unless there is a waiver or the activity falls within the commercial exception.

Private persons can also bring claims directly against private foreign polluters. The chief hurdle is obtaining personal jurisdiction over the polluter. While filing suit in the foreign polluter's home state may cure this problem, it may cause others. The laws and courts of that state may be less than friendly and the costs and inconvenience of doing so are

usually a severe limitation. Procedural and discovery problems are also magnified in transnational cases.

In the United States, citizens and private organizations may also sue their government, its agencies, and its officials for nonenforcement of environmental laws which implement treaty provisions. The Act to Prevent Pollution From Ships, 33 U.S.C. §1910(a), permits any person having an interest adversely affected to bring an action not only against a private violator, but against the Coast Guard for failure to perform a nondiscretionary act and the Secretary of the Treasury for failure to enforce the ship clearance provision (§1908(e)). The ESA allows a citizen to sue (A) to enjoin any person, including the United States or its agencies, if it is in violation of the ESA; (B) to compel the Secretary of the Interior to apply ESA prohibitions with respect to the taking of any threatened species; and (C) the Secretary if he fails to perform a nondiscretionary duty (16 U.S.C. §1540(g)(1)). Other legislation may also permit citizens to sue public officials for acts which are arbitrary, capricious, involve an abuse of discretion or are contrary to law (Administrative Procedure Act, 5 U.S.C. §553, §701 et seq.). Citizens may or may not be able to recover attorney's fees (33 U.S.C. §1910(d), 16 U.S.C. §1540(g)(4)).

Adjudication has many advantages, including the availability of immediate injunctive relief when it is needed and serving as a catalyst for change by goading inactive commissions, legislatures, and the public into action. Adjudication can also prevent further damage. However, there are other, less confrontational methods of mitigating or preventing environmental harm. One of these methods is the liability fund.

A LIABILITY ENFORCEMENT PROGRAM TO DISCOURAGE GEAR LOSS

Liability funds to compensate for damage caused by environmental pollution are not new to the oil and ultrahazardous waste industries, but are a new idea for curing generally pervasive types of environmental pollution such as acid rain and the disposal of garbage at sea. They have the virtue of serving both as a deterrent to future pollution, if set at punitive levels, and as a source of funds for those damaged by the polluting party's activities. They are also in line with the liability provisions contained in the LOS Convention (1982, article 235). The convention includes compulsory insurance and compensation funds as methods that states should employ in order to provide adequate compensation for pollution damage caused by persons under their jurisdiction (article 235(3)). A liability enforcement program such as the one that follows should encourage both domestic fishing vessels and foreign vessels fishing in the U.S. EEZ under a GIFA to comply with U.S. law prohibiting the dumping of fishing gear overboard. The conclusion of bilateral and multilateral liability fund agreements with high seas fishing nations who have not signed GIFA's should also be considered.

Specifically, the owner of a fishing vessel fishing in the EEZ would be entitled to limit his liability for lost driftnets and any damage caused by them if he constitutes a fund for the total sum representing the limit of his liability. The fund can be constituted either by depositing the sum with the administering agency or by producing an acceptable bank guarantee

or proof of insurance. The liability ceiling would be established according to the risk each fishing enterprise poses to the resource. The risk can be calculated using conventional measures of fishing effort: size and number of vessels, amount and type of gear, and number of days fished.

Control methods would utilize an inventory reporting system to account for all gear purchases, gear retirement, and "lost" gear. Penalties for lost gear would be paid out of the fund. Credit and a lowering of total liability would be awarded for expenditures made for conservation measures such as investing in gear with biodegradable panels and knots. Failure to report could result in a tripling of the normal penalties assessed and a doubling of the liability ceiling. Exceptions may be made for natural disasters and intentional acts by third parties.

A liability fund has several advantages. For the fisherman, contributions can be internalized as a cost of doing business; fishing vessels will no longer be subject to seizure for violations and judgment payments; other assets will be immune from attachment; and fishery allocations will not be jeopardized for inadvertent gear loss. For those specifically injured by renegade gear, the fund will provide a source of compensation for resource and property damage. For the government, proceeds from penalties assessed may be used to finance resource conservation activities and the development of traceable and biodegradable fishing gear. The fund also would be easier to enforce than current methods of trying to catch vessels in the act of discarding gear on the high seas.

CONCLUSION

While additional information on the effects of driftnet fishing is useful, it is not necessary in order to legally proceed against persons whose high seas fishing activities damage the coastal resources of the United States. The effective management of marine resources within the EEZ necessitates that the United States regulate foreign fishing activities on the high seas and in fact legislation already exists that implements this application of the protective principle. Although there are as yet few cases where the United States has extended the protective principle codified in its various conservation and pollution laws to the high seas, such an extension is already legal under both domestic and international law. If the protective principle can be used to enforce narcotics laws on the high seas, it can be used to enforce environmental laws.

The Stockholm Declaration on the Human Environment (1972) directs states to "take all possible steps to prevent pollution of the seas by substances that are liable to . . .harm living resources and marine life. . ." (article 7). The doctrine of "reasonable use" of the high seas, addressed in a 1974 decision by the ICJ, establishes that "[t]he former laissez faire treatment of the living resources of the sea in the high seas has been replaced by recognition of a duty to have due regard to the rights of other states and the need of conservation for the benefit of all" (ICJ 1974, p. 175). It is the role of international treaties and tribunals to establish global environmental standards; it is the role of individual sovereign states to enforce those standards. International and regional cooperative efforts are important and should be promoted, but the global community cannot

afford to wait. Under conventional and customary international law, states not only have the right, they have the obligation to protect the global environment using all the tools available to them.

REFERENCES

- Akehurst, M.
1984. A modern introduction to international law. George Allen and Unwin, London, 5th ed., p. 51, 110-11.
- American Institute of Taiwan and the Coordination Council for North American Affairs.
1989. Agreement between the American Institute of Taiwan and the Coordination Council for North American Affairs for cooperative programs regarding the monitoring, enforcement and regulation of the North Pacific driftnet fishing operations. Article VII, 13 July 1989. [In force until 31 December 1990.]
- American Law Institute.
1987. Restatement of the law, third: the foreign relations law of the United States. American Law Institute Publ., St. Paul, Minn., 2 vols.
- Anderson, R.
1989. Foreign "ghost ship" helps prove claims of high-seas salmon thefts. Natl. Fisherm. 70(1):12.
- Beasley, H. R.
1984. International organizations and agreements division update. F/M31 NMFS p. 15.
- Boyle, A.
1985. Marine pollution under the Law of the Sea Convention. Am. J. Int. Law 79:347-372.
- Brussels Liability Convention.
1969. Resolution on international co-operation concerning pollutants other than oil. Attached to the International Convention on Civil Liability for Oil Pollution Damage. United Nations Doc. ST/LEG/SER.B/16 at 447 (1974). [Reprinted in Int. Legal Mat. 9:45.]
- Canadian Arctic Waters Pollution Prevention Bill.
1970. An act to prevent pollution of areas of the arctic waters adjacent to the mainland and islands of the Canadian arctic. [Reprinted in Int. Legal Mat. 9:543.]
- Canadian Reply to the United States Government.
1970. Reprinted in S. Lay, R. Churchill, and M. Nordquist (editors), New directions in the Law of the Sea, Documents, p. 199. Oceana Publ., Dobbs Ferry, N.Y., 1973, 1.
- Carr, H. A., and R. A. Cooper.
1987. Manned submersible and ROV assessment of ghost gill nets in the Gulf of Maine. In Oceans '87, p. 622-624. Proc. Mar. Tech. Soc.

- Cary, F., and R. L. Burgner.
1983. Observations aboard a Japanese squid drift net fishing vessel in September-October 1982. Univ. Wash. Fish. Res. Inst. FRI-UW-8307, Seattle, WA, 23 p. [Final Report to Pacific Seafood Processors Association.]
- Cunnard Steamship Co. v. Mellon.
1922. 262 U.S. 100, 123; 67 L. Ed. 894, 902.
- Defenders of Wildlife v. Hodel.
1989. 707 F.Supp. 1082 (D. Minn. 1989).
- DeGange, A. R., and T. C. Newby.
1980. Mortality of seabirds and fish in a lost salmon drift net. Mar. Pollut. Bull. 11:322-323.
- Eisenbud, R.
1985. Problems and prospects for the pelagic drift net. Environ. Aff. 12:473, 476-478.
- Foreign Sovereign Immunities Act.
1976. 28 U.S.C. §1605(a)(2). See also European Convention on state immunity and additional protocol, May 16, 1972, chapter I, article 7; 1982 U.N. draft articles on jurisdictional immunities of states and their property, article 12, report of the International Law Commission on the work of its 34th session, 37 U.N. GAOR Supp. (No. 10), U.N. Doc. A/37/10.
- Fur Seal Convention.
1957. Interim Convention on Conservation of North Pacific Fur Seals. U.S. Treaties 8:2284, T.I.A.S. No. 3948.
- Gooder, P.
1989. Observations on board the Korean squid driftnet vessel, *Oyang 53*, 9 June-8 August 1988. Northwest Alaska Fish. Cent., Natl. Mar. Mammal Lab., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. Processed Rep. 89-03, 20 p. + tables.
- Gordon, W. G.
1985. Pelagic driftnet fisheries: Hearing before the National Ocean Policy Study, Senate Commerce, Science, and Transportation Committee, 9 October 1985, p. 13.
- Halibut Convention.
1953. Convention for the Preservation of the Halibut Fishery of the Northern Pacific Ocean and Bering Sea. U.S. Treaties 5:5, T.I.A.S. No. 2900.
- Henderson, J. R.
1984. Encounters of Hawaiian monk seals with fishing gear at Lisianski Island, 1982. Mar. Fish. Rev. 46(3):59-61.
- Henry, L.
1928. When is a treaty self-executing? Mich. Law Rev. 27:776-785.

Higgins, R.

1984. The legal bases of jurisdiction. In C. Olmstead (editor),
Extra-territorial application of laws and responses thereto, p. 7
Int. Law Assoc., Oxford.

High Seas Fishing Convention.

1958. Convention on Fishing and Conservation of the Living Resources
of the High Seas. U.S. Treaties 17:139, T.I.A.S. No. 5969, U.N.
Treaty Series 559:285.

High Seas Protocol.

1973. Protocol relating to Intervention on the High Seas in Cases of
Marine Pollution by Substances Other Than Oil. U.N. Doc ST/LEG/SER.
B/18 at 457 (1976), T.I.A.S. 10561.

Ignell, S., J. Bailey, and J. Joyce.

1986. Observations on high-seas squid gill-net fisheries, North
Pacific Ocean, 1985. U.S. Dep. Commer., NOAA, Tech. Memo. NMFS-
F/NWC-105, 52 p.

In the Matter of Kenji Nakata, Ohoura Gyogyo Co. Ltd., 4 O.R.W. 814 (NOAA
1987) (LEXIS Genfed library, Courts file).

International Court of Justice.

1945. Statute of the International Court of Justice. Reprinted in
U.N. Yearbook 1970:1013.

1974. Fisheries jurisdiction cases (merits). Reports. *United
Kingdom v. Iceland*, p. 3; *Fed. Rep. Germany v. Iceland*, p. 175.

Intervention.

1969. International Convention Relating to Intervention on the High
Seas in Cases of Oil Pollution Casualties. U.S. Treaties 26:765,
T.I.A.S. No. 8068.

Japan-U.S.S.R.

1956. International Convention for the High Seas Fisheries of the
Northwest Pacific, Japan-U.S.S.R. U.N. Treaty Ser. 263:3768.

Japan-United States.

1989. Agreement (2 May 1989) concerning the Japanese monitoring and
enforcement programs for 1989.

Japanese Governing International Fishing Agreement.

1982. Agreement between the Government of the United States of
America and the Government of Japan concerning fisheries off the
coasts of the United States of America, United States-Japan. U.S.
Treaties 28:7507, T.I.A.S. No. 10480. [Renewed 29 December 1987.]

[Japan] Fisheries Agency, Canada Department of Fisheries and Oceans, and the U.S. National Marine Fisheries Service and Fish and Wildlife Service.

1990. Final report of squid and bycatch observations in the Japanese driftnet fishery for neon flying squid (*Ommastrephes bartrami*) June-December 1989 observer program. Joint report Fisheries Agency of Japan, Canada Department of Fisheries and Oceans, and the U.S. National Marine Fisheries Service and Fish and Wildlife Service.

Kokechik Fishermen's Association v. Secretary of Commerce.

1988. 839 F.2d 795 (D.C. Cir. 1988); *Federation of Japan Salmon Fisheries v. Baldrige*, 679 F.Supp. 37 (D.D.C. 1987).

Korean Governing International Fishing Agreement.

1982. Agreement between the Government of the United States of America and the Government of the Republic of Korea concerning Fisheries off the Coasts of the United States, United States-Korea. U.S. Treaties 28:1753, T.I.A.S. No. 8526. [Renewed 10 July 1987.]

Lambertson, R. E.

1985. Pelagic driftnet fisheries. Hearing before the National Ocean Policy Study, Senate Commerce, Science, and Transportation Committee, 9 October 1985, p. 15-17.

Lasok, D., and J. W. Bridge.

1987. Law and institutions of the European communities. Butterworth, London, 428 p.

Law of the Sea Convention.

1958. United Nations Convention on the High Seas. U.S. Treaties 13:2312, T.I.A.S. No. 5200.

1982. United Nations Convention on the Law of the Sea. [Reprinted in Int. Legal Mat. 21:1261.]

Levi, W.

1979. Contemporary international law. Westview Press, Boulder, Colorado, 156 p.

London Dumping Convention.

1972. Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter. U.S. Treaties 26:2403, T.I.A.S. No. 8165.

Magnuson Fishery Conservation and Management Act.

1986. Amendments 1986. Public Law 100-220, Title IV, FF 4001-4009, 29 December 1987, 101 Stat. 1477, §4003(4).

Mannington Mills, Inc. v. Congoleum Corp.

1979. 595 F.2d 1287, 1292-93 (3rd Cir. 1979).

Marine Mammal Commission.

1990. Annual report of the Marine Mammal Commission, calendar year 1989: A report to Congress. Wash., D.C.

MARPOL.

1973. International Convention for the Prevention of Pollution from Ships. U.N. Doc. ST/LEG/SER.B/18. [Reprinted in Int. Legal Mat. 12:1319.]

MARPOL Protocol.

1978. Protocol of 1978 relating to the International Convention for the Prevention of Pollution from Ships, 1973. I.M.C.O. Doc. TSPP/CONF/11. [Reprinted in Int. Legal Mat. 17:546.]

Matsen, B.

1989. Brokers arrested in squid-fleet salmon case. Natl. Fisherm. 69:16.

Migratory Bird Convention.

1972. Protection of Migratory Birds and Birds in Danger of Extinction and Their Environment, United States-Japan. U.S. Treaties 25:3329, T.I.A.S. No. 7990.

North Pacific Convention.

1952. International Convention for the High Seas Fisheries of the North Pacific Ocean, United States-Canada-Japan. U.S. Treaties 4:380, T.I.A.S. No. 2786.

North Pacific Convention Protocol.

1978. Protocol amending the International Convention for the High Seas Fisheries of the North Pacific Ocean, United States-Canada-Japan, annex 1. U.S. Treaties 30:1095, T.I.A.S. No. 9242.

OILPOL.

1954. International Convention for the Prevention of Pollution of the Sea by Oil. U.S. Treaties 12:2989, T.I.A.S. No. 4900.

Presidential Proclamation No. 2667.

1945. Fed. Reg. 10:12303.

Presidential Proclamation No. 2668.

1945. Policy of the United States with respect to coastal fisheries in certain areas of the high seas. Fed. Reg. 10:12304. [Reprinted in Statutes at Large 59:885-86.]

Presidential Proclamation No. 5030.

1983. Fed. Reg. 48:10605.

Pruter, A.

1987. Sources, quantities and distribution of persistent plastics in the marine environment. Mar. Pollut. Bull. 18:305-310.


Republic of Korea-United States.

1989. Agreement Annex II, 13 September 1989. Regulatory program and monitoring program on the Korean squid driftnet fishery for the fishing seasons of 1989 and 1990.


- Sathre, C.
1986. Salmon interception on the high seas. *Environ. Law* 16:731-755.
- Sohn, L., and K. Gustafson.
1984. *The Law of the Sea in a nutshell*. West Publ. Co., St. Paul, Minn., 264 p.
- Springer, A.
1983. *The international law of pollution*. Quorum Books, Westport, Conn., 218 p.
- Stockholm Declaration on the Human Environment.
1972. Principle 21., U.N. Doc. A/CONF.48/14. [Reprinted in *Int. Legal Mat.* 11:1416.]
- Strassheim v. Daily*.
1911. 221 U.S. 280, 285, 55 L.Ed. 735, 738.
- Taiwanese Governing International Fishing Agreement.
1982. Agreement between the Government of the United States of America and the Government of the Republic of China concerning fisheries off the coasts of the United States, 15 September 1976, United States-Republic of China. U.S. Treaties 28:1904, T.I.A.S. No. 8529; agreement between the American Institute in Taiwan and the Coordination Council for North American Affairs, 15 June 1982. [Not renewed in 1987.]
- Timagenis, Gr. J.
1980. *International control of marine pollution*. Oceana Publ., N.Y., 473 p.
- Timberlane Lumber Co. v. Bank of America*.
1976. N.T. and S.A, 549 F.2d 597, 615 (9th Cir.)
- Tsunoda, L. M.
1989. Observations on board a Japanese, high-seas, squid gillnet vessel in the North Pacific Ocean, July 1-August 14, 1989. U.S. Dep. Commer., Northwest Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA. Northwest Fish. Cent. Admin. Rep. NMFS-NOAA-F/NWC-89-02, 23 p. + tables.
- U[nited] S[tates] Congress.
1973. Senate Rep. No. 307, p. 1.
1980. House Rep. No. 1224, p. 4.
1987a. Committee Report on H.R. 3674. House Rep. No. 489, p. 23-24.
1987b. Report on the Driftnet Impact Monitoring, Assessment, and Control Act. Senate Rep. No. 261, p. 2-3.
1987c. Report on the Plastic Pollution Control Act. Senate Rep. No. 270, p. 2.

- United States v. Alomia-Riascos.*
1987. 825 F.2d 769 (4th Cir.).
- United States v. Aluminum Co. of America*
1945. 148 F.2d 416 (2d Cir.).
- United States v. Bowman.*
1922. 260 U.S. 94, 98, 67 L. Ed. 149, 151.
- United States v. Conroy*
1979. 589 F.2d 1258 (5th Cir.), cert. denied, 444 U.S. 831.
- United States v. Marunaka Maru No. 88.*
1983. 559 F.Supp. 1365 (D. Alaska 1983).
- United States v. Pizzarusso*
1968. 388 F.2d 8 (2d Cir.)
- United States v. Streifel*
1981. 665 F.2d 414 (2d Cir.).
- United States v. Tsuda Maru.*
1979. 470 F. Supp. 1223 (D. Alaska 1979).
- United States v. Verdugo-Urquidez*
1990. 494 U.S. ___, 108 L. Ed. 2d 222, 230.
- Vienna Convention on the Law of Treaties.
1969. United Nations Doc. A/Conf. 39/27.
- Von Brandt, A.
1984. Fish catching methods of the world. Fish. News Books, U.K.,
418 p.
- Way, E. W.
1977. Lost gill net (ghost net) retrieval project 1976. Environ.
Canada, Fisheries and Marine Service, Industrial Development Branch,
St. John's, New Foundland, 30 p.
- Weil, P.
1984. International limitations on state jurisdiction. In C.
Olmstead (editor), International law limitations on state
jurisdiction, in extraterritorial application of laws and responses
thereto, p. 33. Int'l. Law Assoc., Oxford.
- Wolfe, E. E.
1989. Hearing before the Subcommittee of Fisheries and Wildlife
Conservation and the Environment, House Committee on Merchant Marine
Fisheries, May 2, 1989.

SESSION VII



SOLUTIONS THROUGH EDUCATION





**THE PLASTICS INDUSTRY AND MARINE
DEBRIS: SOLUTIONS THROUGH EDUCATION**

Ronald G. Bruner
The Society of the Plastics Industry, Inc.
1275 K Street, N.W., Suite 400
Washington, D.C. 20005, U.S.A.

ABSTRACT

Over the past several years, intense media attention and congressional debate have focused on the problems caused by plastics discarded or lost in the marine environment. Less well known are the efforts of the plastics industry, through its major trade association, The Society of the Plastics Industry, Inc. (SPI), to help find solutions to these problems. Since 1986, SPI has been working with the Center for Marine Conservation--formerly the Center for Environmental Education--and the U.S. National Oceanic and Atmospheric Administration to implement public service educational campaigns with specific messages targeted to selected audiences.

When the marine debris problem is examined in detail, it is clear that among the various types of plastic debris, all but one end up in the oceans as the result of activity by individuals beyond the "control" of the plastics industry. The one exception involves resin pellets, the raw material for making plastics products. This presentation will examine how the plastics industry has responded to the marine debris problem in general, and specifically to the presence of resin pellets in the aquatic environment.

INTRODUCTION

Over the past several years, intense media attention and congressional debate have focused on problems caused by plastics that have been intentionally discarded or accidentally lost in the marine environment. Less well known are the efforts of the plastics industry, primarily through its major trade association, The Society of the Plastics Industry, Inc. (SPI), to help find solutions to these problems.

Since 1986, SPI has been working with the Center for Marine Conservation (CMC)--formerly the Center for Environmental Education--and the U.S. National Oceanic and Atmospheric Administration (NOAA) to develop and implement a public service educational campaign with very specific messages

targeted to selected audiences. This paper will examine that segment of the public service campaign targeted to the plastics industry, as well as other activities aimed at educating plastics companies about the role of discarded plastics in the marine environment and what should be done about them.

THE SOCIETY OF THE PLASTICS INDUSTRY POLICY STATEMENT

The SPI's educational efforts have been strongly backed at the highest levels of the organization. Their board of directors developed and passed on 18 September 1987 an official policy statement on marine debris that pledged the industry to help solve the problem.

That policy stated, in part:

"The SPI supports the responsible use of its industry's materials and proper disposal of those products when they become waste. Plastics should not be discarded into the ocean or any other body of water.

- "Plastics Resin Pellets--The SPI is dedicated to working with its member companies to eliminate circumstances that result in resin pellets being lost in manufacturing or transportation and possibly rendering harm to animal or marine life that mistake the pellets for food.
- "MARPOL Ratification--The SPI supports U.S. ratification and implementation of the Annex V of the MARPOL Convention, which would prohibit the dumping of plastics waste into the ocean.
- "Degradability--The SPI endorses continued research and development on degradable plastics. However, it believes there are limitations on what products are suitable to be made degradable. Performance and safety requirements should not be compromised in order to make a product degradable. . . .
- "Public Education--The SPI supports public education encouraging the proper disposal of plastics and other materials as the most effective way to reduce harm to the marine environment. The association is willing to work with other organizations sharing this position on projects to further education on the proper disposal of plastics in the marine environment."

In addition, SPI invited CMC President Roger McManus to make a special presentation on the subject to its board of directors' Issues Management Committee on 12 May 1988, and CMC staffer Kathy O'Hara has addressed the SPI Issues Communication Committee and other industry audiences on several occasions. These presentations were particularly helpful in raising the level of sensitivity and responsiveness within the industry.

THE SOCIETY OF THE PLASTICS INDUSTRY SUPPORTS MARPOL ANNEX V

In 1987, when the U.S. Congress was considering ratification of Annex V, many organizations--including the plastics industry--spoke out in support of it. The SPI was emphatic in its view that the time has come to stop discarding trash of all kinds into the marine environment. Testifying before a subcommittee of the U.S. Senate Committee on Environment and Public Works, SPI's Lew Freeman, vice president of government affairs, said:

" . . . To the extent that plastics are part of the marine debris problem the SPI and the plastics industry will continue to work with government, environmental groups, and other industries to develop responsible and effective solutions. But marine debris is more than just a problem of 'plastics pollution,' the term so frequently used to describe it. It is a broader problem of debris from all types of materials being discarded in the oceans.

"There will be a marine debris problem with or without plastics as long as a growing and affluent world population continues the overt--and sometimes covert--practice of using the oceans of the world as a convenient place to put waste. . . . Clearly, plastics waste does not belong in the oceans. However, neither do glass, metal or even paper wastes" (Freeman 1987).

RESIN PELLETS

When the scope of marine debris is examined in detail, it is clear that the problem stems from many sources. In the CMC's 1987 report for the U.S. Environmental Protection Agency, at least nine distinct sources were identified: commercial fishing operations, merchant shipping, naval and research vessels, plastics manufacturing, offshore drilling operations, recreational boaters, docks and marinas, municipal stormwater and sewage systems, and general littering by beachgoers. All but one of these "sources" are the result of activities that are beyond the sphere of direct influence of the plastics industry. That one exception involves resin pellets, the raw material for making plastics products.

Resin pellets, while not particularly an aesthetic problem in the marine environment, have been identified as a hazard to seabirds who ingest them. Although resin pellets are not as abundant as other debris items in the ocean, they seem to be preferred by seabirds. In studies of plastic pieces in the North Pacific (Wilbur 1987), only 0.5% of the pieces of plastic collected from surface waters were pellets. Yet these pellets form about 70% of the plastic found in the stomachs of seabirds.

BRIEFING FOR RESIN PELLET PRODUCERS

One of SPI's first steps in dealing with the question of resin pellets in the marine environment was to alert companies that produce these materials. In September 1986, SPI hosted a briefing in Washington, D.C. which

featured a presentation by Kathy O'Hara from CMC on the marine debris problem and a videotape on resin pellet reclamation produced by Dow Chemical Co., an SPI member company. Virtually all the major resin-producing companies sent representatives. Discussions at that meeting yielded insights that later would be incorporated into a variety of public service materials. The foundation also was laid for an informal survey of the pellet-handling practices of resin producers. While not a definitive study, the results showed that containment procedures for handling resin pellets generally were implemented in the early-to-mid-1970's and that current practices at resin plants seem to preclude significant losses into the environment. Less was known about the thousands of companies that process the resin pellets into final products; thus materials were developed to inform them of the problem as well as of corrective actions to be taken.

PUBLIC SERVICE ADS AND BROCHURES

As noted in the Introduction, the plastics industry was one of three audiences initially targeted by the SPI/CMC/NOAA public service campaign. Materials developed under the joint logo included a full-page, black-and-white magazine ad (Fig. 1) for use in plastics industry publications and an eight-panel collateral brochure. The ad, which shows an enlargement of a resin pellet, carries the headline "A seabird could mistake this resin pellet for a fish egg. And die." The copy reads:

"One little pellet may be insignificant to your plastics processing operation. But to thousands of seabirds, it could lead to a fatal error.


"These pellets, in many shapes and sizes, can be washed down drains as waste or reject material, or spilled in the course of normal handling. But ultimately, they may find their way to bodies of water, where the real trouble begins.

"When eaten in sufficient quantity by a seabird, they can block digestion or sometimes fool the bird into thinking it is not hungry, causing eventual starvation. Fish and sea turtles can suffer the same fate.

"The growing problem of plastic trash in our oceans threatens more than wildlife. This critical issue is destined to invite increasing public and government scrutiny unless we take action to solve it.

"So please: see that resin pellets are reclaimed or disposed of properly. If we ignore the problem we--like the unfortunate seabird--will be making a serious mistake."

Camera-ready reproduction materials (negatives and veloxes) were made available free-of-charge to plastics trade publications, and readers of the ad were invited to write to SPI for additional information. Initially, that consisted of an eight-panel brochure, which offered these recommendations:



A seabird could mistake this resin pellet for a fish egg. And die.

One little pellet may be insignificant to your plastics processing operation. But to thousands of seabirds, it could lead to a fatal error.

These pellets, in many shapes and sizes, can be washed down drains as waste or reject material, or spilled in the course of normal handling. But ultimately, they may find their way to bodies of water, where the real trouble begins.

When eaten in sufficient quantity by a seabird, they can block digestion or sometimes fool the bird into thinking it is not hungry, causing eventual starvation. Fish and sea turtles

can suffer the same fate.

The growing problem of plastic trash in our oceans threatens more than wildlife. This critical issue is destined to invite increasing public and government scrutiny unless we take action to solve it.

So please: see that resin pellets are reclaimed or disposed of properly. If we ignore the problem, we—like the unfortunate seabird—will be making a serious mistake.

To learn how you can help, write: The Society of the Plastics Industry, 1275 K Street, N.W., Suite 400, Washington, D.C. 20005.

A public service message from:
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 1.--Full-page public service ad developed for plastics industry publications has been published more than 25 times. Value of the donated space exceeds \$100,000.

- Conduct an "audit" of manufacturing facilities to seek out and eliminate practices which could allow pellets to escape into the environment.
- Initiate an awareness and information program within the company.
- Install closed-loop pellet containment and collection systems in resin production facilities.
- If resin pellets are spilled during processing, clean them up promptly and either recycle them or dispose of them in ways that prevent their release into the environment.
- When cleaning hopper cars, do not flush residual pellets into the environment.
- Instruct employees to close valves on the unloading shoes of rail cars and hopper trucks after they have been unloaded. (If left open, small quantities of pellets stuck in the corners may vibrate loose and be scattered along railroad tracks and highways, ready to be washed by rain into the nearest stream.)
- Do not store or dispose of pellets in areas subject to flooding.
- Make sure resin pellets are used only as intended--for manufacture of plastics products.

Subsequently, a 20-piece marine debris information kit (described in detail on the following pages) was provided to all who wrote or called for more information. Individual companies also prepared materials. Dow Chemical Company produced and made available a videotape as well as a special flyer on resin pellet containment procedures.

PLASTICS MEDIA BRIEFINGS

Upon completion of the public service ad, brochure, and a special flip-chart presentation, a two-person SPI/CMC team set out to inform editors and publishers of major plastics industry publications on the importance of the marine debris issue. During the fall of 1987, the briefing team personally met with Bob Martino, editor, and Bob Leaversuch, senior editor, at Modern Plastics; Doug Smock, editor, at Plastics World; Matt Naitove, editor, and Bob Burns, associate editor, Plastics Technology; Bob Forger, publisher, and Abe Schoengood, editor, Plastics Engineering; and Peter Sullivan, group vice president, Suzanne Witzler, executive editor, and Mel Friedman, editor, of Edgell Publications (formerly HBJ Publications, which includes Plastics Machinery and Equipment, Advanced Composites, Plastics Compounding, Plastics Design Forum, and Plastics Packaging). The readers of these publications represent a "Who's Who" in plastics, reaching virtually every segment of the industry.

In each case, the overall marine debris problem was discussed with emphasis on the resin pellet aspect, and the publication was asked not only to write about it but also publish the ad on a public service basis.

PLASTICS MEDIA RESPONSE

The response to the visits by the briefing team was overwhelming, and some publications continue to run the ad nearly 18 months later. (In the summer of 1988, two smaller versions of the ad were developed and reproduction materials provided to each of the publications.) Through February 1989, plastics industry publications have devoted more than 1,000 column inches of space either to the ad or to news coverage of the marine debris issue. The ad alone has been published on at least 25 occasions by 10 different publications. Their combined circulations in conjunction with the number of publication times means that more than 1.1 million magazine pages featuring the resin pellet ad are now circulating within the plastics industry. It is estimated that the donated advertising space alone would have cost in excess of \$100,000 had it been purchased at the regular rates.

Not only was the ad published with great frequency, but it also was read! Plastics Design Forum (circulation: 47,500 design engineers) reported that a Readex study of its January 1988 issue showed the resin pellet ad was "the second best read ad in the whole issue" (second only to a multipage, full-color special section). "I think that says a lot of things--both in terms of the quality of the ad, but maybe even more significantly, in terms of the importance of the message," commented Peter Sullivan in reporting the results of the study to SPI.

As gratifying as the advertising support has been by plastics industry publications, even more crucial has been their editorial endorsement of the need for the plastics industry to respond to the problem of plastics in the marine environment. Within months of the visits by the briefing team, four of the largest publications devoted their editorial pages (Fig. 2) to the subject. An example is this excerpt from Modern Plastics (Martino 1987, p. 41):

"Goodbye, George F. Babbitt.--The era of Babbittry has ended--at least in plastics. Sinclair Lewis' fictional businessman might still find a home in the backwaters of industry, but here in the mainstream Babbitt's boosterism and distrust of skeptics no longer has an audience. We see this clearly in environmental issues. There is a new generation of plastics leadership--mindful that plastics surround citizens in daily life, pragmatic enough to listen to these citizens, and responsible enough to want to listen.

"One hopes that the passing of Babbitt will be mirrored by the passing of those who have made it their personal mission to save the world from plastics. The need for cooperation between industry and citizens on environmental issues is so apparent that responsible people should no longer have patience with this sort of axe grinding. . . .

"Babbitt could never have coped. We, however, can draw encouragement from the Society of the Plastics Industry's dealings with one special form of solid waste problem: marine litter. Instead of knee-jerk defense of plastics, SPI's response to reports that plastics litter kills ocean animals was to take the reports seriously enough to realize that they were accurate accounts from sensible people. It then undertook practical, reasonable measures to help attack the problem. The latest, joining with a government agency and an environmentalist group--that's right an environmental group--to warn about the dangers of plastics marine litter will go far toward convincing outsiders that in plastics Babbitt is dead" (Martino 1987).

Closer to home, SPI has published the full-page public service ad several times in its President's Report to the Members, distributed quarterly to 4,000 leading executives in the plastics industry. In addition, stories have been carried in that same publication as well as in Plastics News Briefs, also published by SPI and distributed to nearly 11,000 people in the industry.

PLASTICS MARINE DEBRIS EDUCATION KIT

While publication of the public service ads, editorials, and news stories in the plastics trade press was raising the awareness level of the issue within the industry, SPI was busy developing additional materials for use with its members. Those materials took the form of the Plastics Marine Debris Education Kit (Fig. 3), which was distributed in May 1988 to some 1,500 resin-producing and processor companies on SPI's membership list. The cover letter, distributed with the kit and signed by SPI President Larry Thomas, said in part:

"As you know, the problem of plastics in the marine environment has gained widespread attention over the past year from both the media and government officials. For the most part, the situation is out of our hands--obviously it is difficult for us to keep track of how the end-user disposes of our products, and the vast majority of the plastics in the marine environment were dumped there by others.

"But the problem of resin pellets in the ocean is ours alone.

"It is time to escalate our education efforts to be sure that pellets are not escaping during routine handling, transportation and shipping procedures. . . . I urge you to make this education effort a priority. Employees who manufacture, ship and handle resin pellets need to become more aware of the importance of careful handling procedures. . . ."

The information kit, which carried the resin pellet theme from the public service ad and brochure, was designed to enable any company to carry out its own internal information campaign. Included in the kit were:




Figure 3.--Plastics Marine Debris Education Kit developed by The Society of the Plastics Industry, Inc., was mailed to some 1,500 of its member companies.

- Posters (in two sizes) for placement on bulletin boards or in well-traveled areas of the plant.
- Stickers for placement on trash cans and other appropriate waste disposal containers.
- A feature article about marine debris and resin pellets for use in company magazines or newsletters.
- Brochures with additional information for distribution to employees.
- Order form to secure additional quantities of any of the printed materials.

The SPI has filled many orders for kits and individual parts of the kit, both from SPI members and nonmembers, many of whom were alerted to the availability through the public service ad. In addition, kits were distributed via other means, such as SPI resin producer member companies to their customers in the processing business.

The information kit also has been distributed to the plastics industry on the international scene. In the summer of 1988, SPI's Thomas presented copies of the kit to attendees at a meeting of the International Plastics Associations Directors (IPAD) held in Berlin, West Germany. The IPAD periodically brings together the chief executive officers of the 50 worldwide plastics associations to exchange information.

THE SOCIETY OF THE PLASTICS INDUSTRY--PLASTICS WORLD ROUNDTABLE

Representing the plastics industry on the marine debris roundtable spearheaded by NOAA's James Coe, SPI gained a greater understanding of many aspects of marine debris--and saw that the solution must be multifaceted. The many hours of discussion convinced SPI's director of technical and regulatory affairs, H. Patrick Toner, that there must be a role for the SPI beyond what it already was doing.

The plastics industry is noted for its entrepreneurial, turn-a-problem-into-an-opportunity approach to business. The question was how to spark that creative problem-solving for the benefit of the marine environment. As a trade association, SPI does not actually develop specific products. That didn't mean, however, that it couldn't facilitate the process with information.

Working with Plastics World magazine, SPI sponsored in July 1988 a roundtable discussion on the marine debris problem, bringing together representatives from major industries and organizations with a stake in the situation. Participants included:

James Coe, U.S. National Marine Fisheries Service
 Joe Cox, American Institute of Merchant Shipping
 William Gordon, New Jersey Marine Sciences Consortium
 Carl Kirkland, Plastics World
 Ralph Rayburn, Texas Shrimpers Association
 Thomas S. Scarano, U.S. Navy
 Gary Schmidt, American President Lines
 Mark D. Sickles, American Association of Port Authorities
 Doug Smock, Plastics World
 H. Patrick Toner, The Society of the Plastics Industry, Inc.

The result was a five-page story by Smock in the September 1988 issue of Plastics World entitled "Are shipboard plastics all washed up?" Smock (1988) examined the scope of the problem and what options were under consideration, and noted: "Many of the changes being considered will create commercial opportunities for alert entrepreneurs. There are visions of recycling industries sprouting at harbors. And who will manufacture the on-board compactors, incinerators and waste processing equipment being contemplated for marine plastics?"

Clearly, the message had been delivered to the industry. Now, only time will reveal the degree of success.

THE SOCIETY OF THE PLASTICS INDUSTRY SYMPOSIUM ON DEGRADABLE PLASTICS

Since marine debris first began showing up in headlines and on the evening news, the notion of degradable plastics often has been posed as a solution. In late 1986 and early 1987, increased references to this prompted SPI to plan a symposium on the subject. As the "voice of the plastics industry," SPI was being called upon to respond to the viability of the idea, but there was not a ready compendium of current information.

The Symposium on Degradable Plastics, held 10 June 1987 in Washington, D.C., helped solve that problem. Ten presentations on the technology of biodegradable and photodegradable plastics, plus additional papers, resulted in a proceedings book that contained the best available public information on the subject at that time.

In his opening remarks to the overflow crowd of nearly 400, C.E. O'Connell, then president of SPI, said:

"I hope that by the end of the day we will have enough information to go away thinking about plastics, and particularly plastics packaging, in a new way. We are being asked to think about product design from the cradle to the grave, and as a responsible industry, we must give serious thought to the ultimate disposal of our products after their useful life is finished.

"I doubt that we will come away with many answers today, but as far as I am concerned, that is not really our goal. If we can come away with an appreciation for the difficult questions, and an awareness of the perceived needs, we will be that much closer to determining just where degradability is truly feasible, both today and in the future."

Perhaps Michael Bean of the Environmental Defense Fund, one of the symposium speakers, summed it up best when he told the audience that day:

"The important environmental questions that must be asked are whether the products of degradation are themselves environmentally safe and whether degradation can occur rapidly enough to reduce significantly the hazards of plastics in the environment. The obvious marketing question is whether plastic products made to degrade can continue to serve their intended function in the marketplace."

Some of the questions posed then still deserve consideration today, but much has happened since that symposium in 1987. Many legislative proposals have been made. Some laws have been passed. A number of companies now offer degradable plastics products.

CONCLUSION

Just as the cause(s) of the marine debris problem is largely an accumulation of careless actions over many years, so too will the solution come in the form of years of dedicated efforts on a variety of fronts. The industry is committed to being a meaningful part of that solution--through both cooperative educational efforts and the development of environmentally safe products.

REFERENCES

- Freeman, L. R.
1987. Statement before U.S. Senate Committee on Environment and Public Works. Soc. Plastics Ind., 17 September 1987.
- Martino, R.
1987. Editorial. Modern Plastics. December, p. 41.
- Smock, D.
1988. "Are shipboard plastics all washed up?" Plastics World Mag., January, 5 p.
- Wilbur, R. J.
1987. Plastics in the North Atlantic. Oceanus 30(3), 8 p.

MARPOL 73/78 INFORMATION, EDUCATION, AND
TRAINING: MEETING THE CHALLENGE

Trevor R. Dixon
Marine Litter Research Program, The Tidy Britain Group
The Pier, Wallgate
Wigan, Lancashire WN3 4EX, United Kingdom

ABSTRACT

In response to growing public concern about the widespread distributions of marine litter, The Tidy Britain Group, the United Kingdom's agency responsible for the prevention and control of mishandled waste, established a Marine Litter Research Program in 1973. The program's overall strategy has been to quantify the nature and scope of the problem in the coastal and oceanic waters of Western Europe, and to persuade governments, intergovernmental organizations, and the shipping industry of the need for remedial actions.

To date a series of information surveys, public awareness campaigns, and educational initiatives has been completed, including a contribution toward an International Maritime Organization (IMO) training package for ships' crews.

This paper provides an insight into these activities, focusing primarily upon the data collection methods developed for the surveillance of marine litter by beach surveys, the approaches adopted in the public awareness campaigns and educational initiatives, and the observed impacts upon coastal environments. These activities have provided background expertise and knowledge to support the information, education, and training recommendations contained in the IMO Guidelines for the Implementation of Annex V of MARPOL 73/78.

An immediate objective of the information surveys was the development of standardized field survey techniques and analytical methods for the surveillance of marine litter. This has been accomplished by the use of appropriate sampling techniques and the establishment of technical support networks. Consequently, major trends in the composition, origins, distributions, and lifetimes of litter in the marine environment have been identified in a series of baseline surveys. Other studies have identified the safety hazards arising from packaged dangerous goods, clinical wastes, munitions, and pyrotechnics washed ashore.

Two national shoreline surveys have been completed since 1979, with observations at 4,000 sites around the coastline. Studies of marine litter have been undertaken at all levels within the formal education system, and beach cleanups are frequently undertaken by environmental land amenity groups.

A more recent development is the Blue Flag Campaign in European Economic Community member states. Beaches which receive these awards are expected to be litter free, and ports or marinas are required to provide facilities for the disposal ashore of garbage from ships and pleasure craft.

INTRODUCTION

The Tidy Britain Group is the United Kingdom's national litter abatement organization, with a broad membership including national and local government, industry, commerce, and voluntary organizations. It functions primarily as an advisory body, making available a range of practical programs which it has developed through research, evaluation, and experience over the years.

The group's broad strategy is to deal with the litter problem by tackling the two main root causes: first, the attitudes and behavior of all people towards littering and the environment generally; and secondly, the incorrect handling of rubbish and waste. This has been achieved by a series of ongoing educational and action programs which involve, and target, the entire community.

The Marine Litter Research Program is an integral part of this structure and approach towards litter abatement. It was established in 1973 with an honorary director supported by specialist advisers in educational institutions, industry, commerce, conservation, and amenity groups.

This paper examines the various activities undertaken by the program during the last 16 years, activities that have made a contribution of background knowledge and expertise to the International Maritime Organization's (IMO's) information, education, and training programs for the successful implementation of Annex V of the International Convention for the Prevention of Pollution from Ships, 1973, as modified by the 1978 Protocol (MARPOL 73/78) (IMO 1988).

INFORMATION

Program Strategy

Against a 1973 baseline, the strategy of the Marine Litter Research Program was to provide systematic data showing qualitatively and quantitatively the nature and scope of the problem, thereby providing a framework for subsequent remedial actions. To achieve this strategy, and in the absence of any comparative studies, priority was given to meeting basic information requirements. In particular, the program needed to produce data

of sufficient accuracy, drawn from a large enough geographical area, to alert national authorities and intergovernmental organizations to the consequences of uncontrolled discharges of ships' garbage at sea.

The immediate objectives were to:

- Develop standardized field survey techniques and analytical methods for the surveillance of marine litter, and
- Identify the major trends in the composition, distribution, and origin of litter occurring in the coastal and oceanic waters of Western Europe.

Use of Beach Surveys

The relative merits of the different approaches used in assessments of marine litter have been discussed (Dixon and Dixon 1981). Beach surveys supported by adequate site selection sampling methods were selected in preference to observations of litter distributions at sea or estimates of discharges from ships for two reasons:

1. Temporal and spatial sampling constraints are more easily overcome, hence representative areas of shoreline can be surveyed more effectively and accurately than sections of water masses, under most weather conditions.
2. Litter tends to accumulate in the short term on beaches and therefore statistically representative samples can be collected at any given time, avoiding uncertain extrapolations from small sample sizes.

A common component of West European beach litter is containers of all types including bottles, cartons, drums, and cans fabricated from glass, metal, paperboard, plastics, and wood. To discover their characteristics, including geographical origins, contents, and dates of production, samples of the most frequently occurring types and brands were collected and their details recorded. Technical support networks were established with packaging and product manufacturers, especially plastics bottle makers, worldwide. The required information was made available from manufacturers' records of container dimensions and designs and the interpretation of article number codes, manufacturers' imprints, individual date codes, and other overprinted or embossed markings.

Consequently, a data base has been compiled for several hundred different containers on the basis of the following selection criteria:

- They had been marketed continuously for more than 10 years in disposable or one-journey packaging, with documented histories of changes in designs or markings.
- The selected brands were market leaders, and therefore produced in large quantities each year, often in several different countries, with distinctive packages.

- The specimen types constituted a representative sample of all containers observed on beaches, by fabrication materials and contents.

Information gathered from shoreline transects has included total weights of all litter collections and each of the main fabrication materials, and the presence or absence of noncontainer items including plastics fragments, fishing gear, plastics sheeting, clothing, rope, and strapping.

Assessment of Major Trends in Surface Marine Litter

Analysis of the combined data has enabled an assessment of major trends in the composition, distributions, geographical origins, and persistence of litter in the surface waters of Western Europe. Findings show:

- Widespread distributions in both coastal and oceanic waters.
- The prominence of plastics materials, especially polyethylene bottles, polythene sheeting, and fragments of expanded polystyrene.
- The identification of the most frequently occurring containers, by fabrication materials and original contents, including polyethylene lavatory cleanser and household cleaner bottles, glass wine and spirits bottles, metal cans and drums originally filled with beverages and petroleum derived products, and wooden fish boxes.
- A global distribution of countries in which containers are manufactured or marketed, with up to 30% from non-European Economic Community (EEC) member states, including Australia, Brazil, Canada, China, Japan, New Zealand, South Africa, the United States, U.S.S.R., and Venezuela.
- The presence of dangerous items of litter, including fragments of broken glass, sharp pieces of metal, nails protruding from timber, pharmaceutical drugs, syringes and other clinical wastes, civilian and military pyrotechnics, munitions, and packaged dangerous goods.
- The absence of any field evidence to show an accumulation of plastics containers in the marine environment. Relative dating methods have indicated that less than 15% of samples have been obsolete types, and absolute dating from individual date codes has shown that between 86 and 94% of container samples have been <4 years old (Dixon and Dixon 1983).

On the basis of these findings and those from more recent studies, discharges of garbage from ships and other craft have been shown to be the major source of litter found on West European beaches (Chaussepeied 1985;

Vauk and Schrey 1987). The contribution from land-based sources is believed to be relatively small, in the region of 30% of the total, and the result of discards from beach users during the summer holiday season.

Baseline surveys at 185 sample sites around the coastline of the United Kingdom were completed between 1979 and 1987. These will be repeated at the same sites during the next 5 years, and the data will be analyzed on a "before" and "after" basis to evaluate the effectiveness of the Annex V regulations. Supplementary studies will also be undertaken during the same time period to examine the efficiency of port reception facilities for ships' garbage.

Other Findings

Questionnaire surveys and desk studies in cooperation with local authorities and government departments, including the Ministry of Defense, have generated more detailed information on the environmental impacts of marine litter.

For example, local authority beach cleaning operations have usually been confined to the summer holiday period, but recently they have been undertaken throughout the year in many locations to remove increasing quantities of litter washed ashore. The costs incurred are met by the local community and have not been recovered in accordance with the "polluter pays" principle.

Local authorities are also responsible for removing and safely disposing of dangerous substances or items recovered on beaches, especially packages of dangerous goods lost from ships' deck cargoes at sea. At least 100 adults and children have undergone precautionary medical examinations or hospital treatment following exposure to the contents of chemical packages washed ashore in southern England between 1976 and 1986 (Dixon and Dixon 1986).

The shipping industry has reported the loss of productive time as well as repair and replacement costs following damage to ships and pleasure craft from surface floating or semisubmerged items of litter.

Munitions are also widely distributed throughout the coastal waters of the United Kingdom following accidental losses or deliberate dumping in the past. Substantial quantities have been discovered by the general public on beaches, and occasional injuries or fatalities have resulted from the improper handling of a small proportion in dangerous condition (Dixon and Dixon 1979).

Marine Litter Research Program Contributions

The Marine Litter Research Program results and subsequent recommendations have made a contribution to debates and discussions on the subject by national authorities and intergovernmental organizations, thereby achieving the program strategy identified above.

For example, the United Kingdom's Royal Commission on Environmental Pollution supported The Tidy Britain Group's recommendations concerning the need for the government to ratify Annexes III and V of MARPOL 73/78 and provide adequate means for the disposal of shipboard-generated garbage ashore (Royal Commission on Environmental Pollution 1984, 1985).

Similarly, The Tidy Britain Group's recommendation concerning the need to designate the North Sea as a special area for the purpose of Annex V was one of the proposals considered at the Second International Conference on the Protection of the North Sea in November 1987. On behalf of all North Sea states, the United Kingdom has since submitted a proposal with the necessary background information and justification for declaring the North Sea a special area at the Twenty-Sixth Session of the IMO's Marine Environment Protection Convention (MEPC) in 1988. The MEPC subsequently approved amendments to Annex V of MARPOL 73/78 in this regard, and it is envisaged they will be formally adopted by the MEPC at its twenty-eighth session.

Recommendations concerning improvements in the extent and durability of markings and labels on packages containing dangerous and polluting substances, and the efficiency of reporting losses of packaged goods overboard were submitted by the Advisory MEPC on Pollution of the Sea to the MEPC in 1985 and 1986 (IMO 1985, 1986).

EDUCATION

The importance of public awareness campaigns and general educational programs in the protection of the marine environment have long been recognized. In the United Kingdom, attitude surveys have consistently suggested that beach pollution and litter are major areas of public concern (National Opinion Poll 1987). Consequently, considerable effort has been invested by The Tidy Britain Group in the development of suitable resources and materials to focus public attention on the marine litter problem.

At the planning stage a series of informal meetings were held with interested parties including educationists, environmentalists, and potential sponsors in order to identify their requirements and enlist support for the public awareness and general education programs. Considerable experience was made available by The Tidy Britain Group's Schools Research Project Team, which specializes in finding ways of increasing environmental awareness and responsibility in children, with particular reference to the awareness of litter.

At these meetings, formal educationists stressed the need for environmental education programs to meet requirements of existing curriculums, especially the promotion of field studies incorporating a variety of different study skills and learning activities across the educational spectrum. Nonformal educationists pressed for detailed information and practical action, as did environmental groups who, in addition, preferred activities which could attract the media. Potential sponsors set forth numerous requirements, often conflicting with each other. Basically, however, they were all prepared to support environmentally worthwhile projects which were structured with clearly defined objectives.

COMMUNITY PARTICIPATION

Public Awareness Activities

A variety of public awareness campaigns and activities have been completed. Some have been directed specifically at professional and nonprofessional seafarers, and port and terminal operators. For example, the owners of pleasure craft were asked to "Stash their trash ashore" in a poster featuring Dame Naomi James, the first lone round-the-world yachts-woman. In another approach, a soft drinks manufacturer sponsored a comic beachcomber competition in which children were asked to record different plastic bottle makers' imprints, and were awarded prizes for their efforts. Beach cleaning competitions organized by local authorities have also been successful, together with more conventional beach cleanups by members of conservation and amenity groups. Among the numerous events and supporting materials is a poster based on a photograph by Linda McCartney, entitled "Please leave nothing but footprints in the sand."

National Shoreline Litter Surveys

The Sunday Times sponsored the First National Shoreline Litter and Refuse Survey, which ran from June 1978 to October 1979. It was questionnaire-based, and participants were asked to record details of litter, especially container markings and codes, from transects between the water's edge and the backshore zone of beaches. Details from 20,000 containers and the presence or absence of other types of litter were recorded at 797 sites around the coastline. The survey report with a foreword written by Marcus Fox, a parliamentary under secretary of state for the environment, generated considerable media interest nationally and therefore attracted the attention of other government departments and shipping interests (Dixon and Hawksley 1980). It was later submitted to the IMO's MEPC by the Department of Transport as an information paper. Some sewage disposal-related issues were included in the report after one participant explained how she had inadvertently flushed a toilet roll holder down the lavatory and recovered it again 2 days later during a local beach survey.

A Second National Shoreline Survey was launched in 1980, sponsored by the Sunday Times and Sealink UK, a major ferry operator, with the added support of network television time. It was again questionnaire-based but the procedures were improved, giving more detailed information on how to identify litter. Observations were undertaken at more than 3,000 sites around the coastline. Among the unexpected discoveries was a sequence of reports from different localities of foreign items of litter, which appeared to have monitored the passage of an East European fishing fleet. Snowmobile bottles were found elsewhere, and these were traced to the discards of Canadian Eskimos near the polar ice cap.

Both shoreline surveys achieved their objectives, including the provision of detailed information on container markings, which in turn has supported the more specialized research. Compared with similar promotions undertaken by The Tidy Britain Group, the shoreline surveys generated considerably more media interest locally and nationally, and were also highly cost-effective. In many instances, educational institutions from

primary schools to universities have adopted the surveys, with suggested learning activities, on a long-term basis.

The Blue Flag Campaign

A major public relations operation in favor of the marine environment is the Blue Flag Campaign, launched during the European Year of the Environment by the Foundation for Environment Education in Europe. The campaign has the financial support and patronage of the Commission of the European Communities. The main objective is to raise awareness about the quality of the marine environment and the need to protect it. During 1988, a total of 373 beaches and 102 ports and marinas in 10 EEC member states were awarded blue flags for the high standards of environmental quality, the provision of services, and the promotion of environmental education (Foundation for Environment Education in Europe 1988). Blue Flag beaches are required to be free from litter, and ports or marinas which qualify are expected to provide facilities for the disposal of garbage ashore from ships and pleasure craft. There is also a Blue Flag charter for boat owners, which is based upon a voluntary code of conduct.

In the United Kingdom, beaches that are unable to qualify for blue flags on the basis of inadequate bathing water quality, are encouraged to enter for Clean Beach awards, a scheme funded by the Department of the Environment.

TRAINING

Throughout its history, the IMO has attached the utmost importance to the training of ships' personnel to help achieve its most important objectives, namely, improvements in maritime safety, and the prevention and control of marine pollution from ships.

Given the rapid changes and developments in world maritime trade and shipping operations, there is an even greater emphasis on the part of the IMO to ensure that the various conventions, codes, and other instruments already adopted are effectively enforced and implemented.

Therefore, in the context of the implementation of Annex V of MARPOL 73/78, considerable effort has been directed towards the preparation of comprehensive and practical guidelines with the following objectives:

- To assist governments in developing and enacting domestic laws which give force to and implement Annex V.
- To assist vessel operators in complying with the requirements set forth in Annex V and domestic laws.
- To assist port and terminal operators in assessing the need for, and providing, adequate reception facilities for garbage generated on different types of ships.

Section 2 of the guidelines recognizes the importance of information, education, and training programs in the implementation process. Given the

need for commercial vessels and other craft to change garbage disposal practices which have persisted for many generations, it is generally believed that comprehensive training packages have an important role in assuring the success of Annex V.

Consequently, with funding provided by the United Kingdom Government and the IMO/Swedish International Development Authority Program for the Protection of the Marine Environment, a training package on marine litter has been prepared for worldwide distribution. Data and other expertise held by The Tidy Britain Group were extensively drawn upon in the production of this material.

The principal aims of the training package are first, to change attitudes and create an awareness of the need to prevent damage to the marine environment from the improper disposal of ships' garbage at sea, and secondly, to provide the necessary background knowledge and identify the practical means of doing so.

The target audience includes not only seafarers and port or terminal operators, but trade associations, shipbuilders, packaging manufacturers, waste management industries, coastal communities, educators, and governments.

With background resource materials contributed by interested parties worldwide, including conservation groups, governments, and the private sector, the training package provides a general overview of the marine litter problem focusing upon:

- the sources and types of garbage generated by ships;
- the environmental impacts of marine litter on wildlife and coastal amenities;
- shipboard garbage handling and storage procedures;
- equipment available for port reception facilities; and
- the Annex V regulations and their implications.

Despite the enormity of the task to be achieved in the successful implementation of Annex V, evidence of new initiatives and innovations has already come to light. Manually operated garbage compactors are now included in the design and construction of some yachts and other pleasure craft in Scandinavian countries, and in some coastal communities of the United States unwanted fishing gear is recycled for decorative purposes instead of being discarded at sea.

REFERENCES

Chaussepeied, M.

1985. Pollution and harmful substances in the coastal environment.

Intensive European course, 16-22 September 1985. Institut Francais de Recherche pour l'Exploitation de la Mer, France.

Dixon, T. R., and T. J. Dixon.

1979. Munitions in British coastal waters. Mar. Pollut. Bull. 10:352-357.

1981. Marine litter surveillance. Mar. Pollut. Bull. 12:289-295.

1983. Marine litter surveillance on the North Atlantic Ocean shores of Portugal and the Western Isles of Scotland. Stage 5. Marine Litter Research Programme. The Tidy Britain Group, Wigan, United Kingdom.

1986. Packaged dangerous goods washed on to beaches of England and Wales. Environmentalist 6:209-218.

Dixon, T. R., and C. Hawksley.

1980. Litter on the beaches of the British Isles. Report of the First National Shoreline Litter and Refuse Survey. Stage 3. Marine Litter Research Programme. The Tidy Britain Group, Wigan, United Kingdom.

Foundation for Environment Education in Europe.

1988. The European Blue Flag. Directorate for the Environment and Consumer Protection, Commission of the European Communities, Brussels, Belgium.

International Maritime Organization.

1985. Twenty-Second Session of the Marine Environment Protection Convention. Agenda item 22/10/INF 2 (7 October 1985). Implementation of Annex III of MARPOL 73/78 and amendments to the IMDG Code to cover pollution aspects. Packaged goods - The need for amendment of MARPOL Annex III. International Maritime Organization, London.

1986. Twenty-Third Session of the Marine Environment Protection Convention. Agenda item 23/8/4 (6 June 1986). Implementation of Annex III of MARPOL 73/78 and amendments to the IMDG Code to cover pollution aspects. International Maritime Organization, London.

1988. Guidelines for the implementation of Annex V of MARPOL 73/78. International Maritime Organization, London.

National Opinion Poll.

1987. Public attitudes to the environment. Department of the Environment, London.

Royal Commission on Environmental Pollution.

1984. Tackling pollution experience and prospects. Tenth report. Her Majesty's Stationery Office, London.

1985. Managing waste: The duty of care. Eleventh report. Her Majesty's Stationery Office, London.

Vauk, G. J. M., and E. Schrey.

1987. Litter pollution from ships in the German Bight. Mar. Pollut. Bull. 18:316-319.

**EDUCATION AND AWARENESS: KEYS TO SOLVING
THE MARINE DEBRIS PROBLEM**

Patty Debenham
Center for Marine Conservation
1725 DeSales Street, N.W.
Washington, D.C. 20036, U.S.A.

ABSTRACT

The Center for Marine Conservation (CMC)--formerly known as the Center for Environmental Education--is a nonprofit conservation organization. The CMC currently conducts a national education campaign on the problems caused by plastic debris in the marine environment. The campaign includes the development and distribution of educational materials to the commercial fishing, merchant shipping, and plastics industries as well as to recreational fishermen, pleasure boaters, and the general public. This program is sponsored in part by the U.S. National Oceanic and Atmospheric Administration's (NOAA), Marine Entanglement Research Program and The Society of the Plastics Industry. Also, under contract to NOAA, CMC distributes information through its National Marine Debris Information Office. The CMC also administers the National Marine Debris Data Base. With support from NOAA, the U.S. Coast Guard, and the U.S. Environmental Protection Agency, the CMC distributed 43,000 data cards to volunteers in all 25 coastal states. Information obtained from volunteer data collection efforts will become part of CMC's national analysis of marine debris data.

INTRODUCTION

There is virtually unanimous agreement that education is necessary to motivate groups and individuals to dispose properly of wastes, especially plastic waste. Several international conferences have stressed the need for marine debris education programs, including the 1984 International Workshop on the Fate and Impact of Marine Debris, the North Pacific Rim Fishermen's Conference on Marine Debris, and the Oceans of Plastic Fishermen's Workshop. Federal legislation entitled the Marine Plastic Pollution Research and Control Act of 1987 mandates that the U.S. National Oceanic and Atmospheric Administration (NOAA) and the U.S. Environmental Protection Agency (EPA) conduct a 3-year public education program. The 1989 Interagency Task Force on Marine Debris encourages marine debris education: "Concerned federal agencies should work with each other, state

and local governments, private industry, and environmental groups to develop comprehensive educational materials on problems caused by marine debris and on ways to solve those problems" (Interagency Task Force on Persistent Marine Debris 1988).

The Center for Marine Conservation (CMC)--formerly the Center for Environmental Education (CEE)--knows that education will be the most effective method to alter 4,000 years of ocean disposal behavior. When mariners realize and understand the effects that their age-old habits have on wildlife and human and vessel safety, they are willing to make a change. Since 1985, the CMC has developed an education campaign that encourages all members of industry, the general public, and the maritime community to get involved. It publishes documents, organizes beach cleanups, and responds to requests for information that encourages groups and individuals to take part in the solution to ocean pollution.

GETTING THE MESSAGE OUT

Early Efforts at Education

Prior to 1984 there were only a handful of programs working to teach people about the problem of marine debris. Judie Neilson's "Get the Drift and Bag It" campaign was not only the first marine debris education program, but also a highly successful coastal cleanup in Oregon (Neilson 1985). Also in 1984, the Workshop on the Fate and Impact of Marine Debris produced several recommendations discussing the need for marine debris education programs (Shomura and Yoshida 1985). The concept of marine debris education was so new that in the workshop proceedings these education recommendations were hidden under the general category of "Report of the Working Group on Management Needs." The subcategory entitled "Public information and education" recommended:

"Recognizing that greater benefits are likely to be realized as a result of positive rather than negative incentives, (Working) Group participants urged that significant emphasis be placed upon public information and education and that steps specifically be taken to:

- "a. Work with fisheries organizations and the fishery management councils to develop and carry out comprehensive information and education programs for foreign fishermen, working within the exclusive economic zone, and U.S. fishermen;
- "b. Work with appropriate national and international organizations to undertake cooperative comprehensive information and education programs; and
- "c. Work with relevant industries, such as has been done with elements of the plastics industry, on public education programs."

The subcategory entitled "Debris cleanup" recommended:

". . . immediate steps to remove existing debris from the environment are clearly needed and concentrated efforts should be directed to reducing the rate at which new debris is deposited. The management steps recommended are:

- "a. To undertake cleanup programs to remove existing debris from shore areas and the water column;
- "b. To assign priority to areas where the density of debris is such that it affects endangered, threatened, or commercially valuable species;
- "c. To require that all potentially harmful debris be retained onboard vessels until proper disposal is possible;
- "d. To encourage the removal of debris from the environment and prevent the discarding of additional debris, positive incentives such as financial rewards for the return of discarded netting material should be considered as should possible negative incentives; and
- "e. To take such actions as may be necessary to assure the proper disposal of unwanted materials in a nonharmful manner."

These recommendations identified a need for marine debris education and provided the baseline for all future education programs.

Plastics Debris Problem

In 1986, the EPA commissioned CMC to prepare a report on the plastics debris problem in the marine and Great Lakes waters of the United States. As the first comprehensive review of available information on marine debris, this document showed that plastics debris is a nationwide problem for marine wildlife. The report identified the major ocean- and land-based sources of plastics debris, and indicated that the total amount of debris generated by these sources is unknown. The report noted the absence of appropriate laws to address the plastic debris problem (CEE 1987a). Finally, the study helped to redirect attention from general marine debris to those problems caused specifically by plastic items.

Upon completion of the document, CMC presented the report to members of The Society of the Plastics Industry (SPI), explaining that plastics debris was threatening wildlife and vessel safety in addition to being unsightly. The plastics industry accepted CMC's invitation to become part of the solution. The result has been the development of numerous brochures, books, and posters geared to promoting proper disposal of plastics at sea.

PUBLIC AWARENESS AND RESPONSE

Beach Cleanups

The CMC's most successful efforts have been its citizen beach cleanups in Texas and Florida. From 1986 to 1988, it organized the largest beach cleanups in American history, accounting for one-third of the nation's total participation. Diverse groups worked toward a common goal: industry provided financial and in-kind support; government and environmental organizations acted as regional coordinators; and the general public together with all these groups removed trash from the beaches.

Since 1984 there has been a steady growth in beach cleanup participation (Fig. 1). In 1984, Judie Neilson encouraged 2,100 Oregonians to clean the beaches. Nationwide during Coastweeks '88, more than 47,000 people participated in coastal cleanups. In 1989, we expect the participation to increase to 60,000, and our goal is to work with the states to encourage 100,000 citizens to clean the beaches by Coastweeks '92.

A beach cleanup is not just a 1-day event, but rather an ongoing education campaign. The CMC's volunteer data collection system is one mechanism that ensures continuing education. Beach cleanup data help identify possible sources and quantify the amounts of debris found by volunteers. Each cleanup volunteer receives a data card and a "Guide to Good Data Collection." The data card lists items volunteers will likely collect during a beach cleanup. The guide explains the importance of data collection and describes items that are found on most beaches yet are difficult to identify.

In Texas and Florida, volunteers work in pairs to share the tasks of debris collection and data recording. From volunteer data, the CMC published two reports on the Texas debris problem (CEE 1987b, 1988b). These reports document the sources of debris and include recommendations to reduce the marine debris problem. In some cases it is possible to attribute certain types of beach debris to a specific source. For example, volunteers found 4,170 plastic light sticks on Texas beaches in 1987. Fishermen commonly use light sticks to attract fish to their hooks. Although it is counterproductive to point accusatory fingers, CMC uses indicative data to encourage possible debris contributors to become active contributors to the solution.

Press generated from a cleanup also helps maintain the long-term education effects. Media coverage reaches people who may not donate their Saturday mornings to clean the beach, but may unconsciously discard their boat or beach trash. The CMC uses the media to remind the public that their plastic trash can have disastrous effects on marine wildlife. The 1986 Texas Coastal Cleanup campaign not only increased awareness among the general public, but also helped the Texas General Land Office initiate a statewide "Adopt-A-Beach" program.



Figure 1.--Volunteer participation during beach cleanups from 1984 to 1990.

National Marine Debris Data Base

In 1988, the CMC expanded its Texas data collection efforts to establish a National Marine Debris Data Base. With support from the EPA, NOAA, and the U.S. Coast Guard, CMC distributed 43,000 data cards to cleanup volunteers in all 25 coastal states. In addition, Spanish data cards were sent to cleanups in Puerto Rico, Costa Rica, and the Dominican Republic. The resulting data base is providing essential information for understanding specific debris problems in each part of the country. O'Hara's paper entitled "National Marine Debris Data Base" gives a detailed analysis of the preliminary results from the first nationwide assessment of coastal debris (O'Hara 1990). The CMC's final report will be out in June 1989.

The CMC's data base relies on volunteer cooperation, and CMC realizes that it is not the same as a rigorous scientific survey. Nonetheless, their data give consistent perspectives of the problem and indicate some

common trends. For example, in all states plastics account for between 55 and 65% of all debris types collected. In 1988, the fact that Florida volunteers collected 489.1 km (304 mi) of monofilament fishing line indicates a large (although not exclusive) contribution from the recreational fishing community. The data from Florida were used by Governor Bob Martinez and his staff to prepare an executive order to provide state enforcement of MARPOL Annex V and to research the use of degradable fishing line. Finally, data collected in 1988, prior to implementation of Annex V, will provide a baseline of information to measure loosely the effectiveness of international and national legislation to reduce ocean pollution.

Marine Debris Information Offices

Under contract to NOAA, CMC coordinates two marine debris information offices (MDIO). The first is in Washington, D.C., to serve the Atlantic coast and Gulf of Mexico. The second, in San Francisco, responds to Pacific coast inquiries. The CMC and NOAA created these offices in response to a growing number of requests for information on the marine debris problem. The MDIO functions to disseminate educational materials and other information on marine debris to government agencies, industry groups, educators, the press, and the general public. In most cases, requests for information fall into specific categories. To respond efficiently to these requests, 16 standardized educational packets were developed :

- General public.
- Teachers and educators.
- Elementary (kindergarten to fifth grade),
- Middle school, high school (6th to 12th grade) and college students.
- Beach cleanup information.
- Recreational fishermen and boaters.
- Press and media personnel.
- Plastics recycling and degradable plastics information.
- Cruise ship passengers.
- Fishermen and fish processors.
- Cargo vessel operators and crews.
- Offshore oil and gas (companies).
- Offshore oil and gas (workers).
- Plastics manufacturers and resin pellet producers.
- Port and terminal operators.
- Charter vessel operators.

All packets contain general information about the marine debris problem, with additional information specific to the requester's interest. From the establishment of the MDIO in October 1988 until 1 April 1989, CMC responded to 842 requests for information. Also available from the MDIO are numerous education materials developed by NOAA, SPI, and CMC as part of a national campaign to promote the proper disposal of plastics.

Chronologically, the first element of the joint educational campaign consisted of print public service advertisements developed for each of the



This discarded net is done fishing. But it's not done killing.

When worn fishing nets or other plastic gear is dumped or lost in the water, something else happens: animals die.

Seabirds get caught in nets when diving for food, and drown. Other marine animals become entangled in them and slowly strangle.

Discarded nets and traps even compete with you, by needlessly catching and killing millions of pounds of potentially valuable fish and shellfish.

In addition, plastic wastes can foul propellers and block cooling intakes, causing costly vessel disablement.

Over 100,000 tons of plastic fishing gear are dumped into our oceans every year. This critical issue is destined to attract increasing public and government scrutiny if we fail to take action to solve it.

So please, alert your dock operators that you'll need trash facilities, because you're saving your plastic trash and worn out gear for proper disposal on land. That's not all you'll be saving.

To learn how you can help, write: Center for Environmental Education, 1725 DeSales Street, N.W., Suite 500, Washington, D.C. 20036.

A public service message from
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 2.--Commercial fishing public service advertisement.

following groups: commercial fisheries (Fig. 2), merchant shippers (Fig. 3), the plastics industry (Fig. 4), recreational boaters (Fig. 5), and recreational fishermen (Fig. 6). To date these ads have appeared in 30 magazines and major trade journals in addition to several regional and local publications including National Fisherman, Marine Log, Modern Plastics, Outdoor Life, and Saltwater Sportsman. Each advertisement directs interested persons to the MDIO for more information about marine debris. The MDIO in turn responds to each request by sending the appropriate information packet and relevant materials.

Each public service advertisement has a corresponding eight-panel brochure with more information on how marine debris affects that particular group. For example, commercial fishermen may be more interested in the fact that discarded gillnets will foul their propellers rather than in the effects the nets may have on seals, which in some cases are viewed as competitors. Groups often request large quantities of brochures for their own distribution. To date, the MDIO has distributed over 60,000 brochures to educators, individuals, and the government, including 15,000 National Safe Boating Week press packets, 8,000 for National Fishing Week, and 3,000 to Coast Guard port captains.

A Citizen's Guide to Plastics in the Ocean: More Than a Litter Problem is another product of the cooperative NOAA, SPI, and CMC campaign that is now available through the MDIO (CEE 1988a). The book informs citizens of the growing problem of plastics in the ocean and gives suggestions on how individuals can become involved in solving this problem. The MDIO has distributed 19,000 copies of this guide since September 1988, including 5,000 copies to the U.S. Navy as part of their educational package on marine debris. Due to the popularity of the book, it is now ready for a second printing that will contain more current information on Annex V and new initiatives to stop plastic pollution at sea.

The MDIO also distributes materials produced by other groups. Recht's (1988) Reference Guide for Ports is a valuable source of information on how ports can comply with the requirements of MARPOL Annex V and the U.S. Marine Plastic Pollution Research and Control Act of 1987. Figure 7 shows the most current version of the MDIO order form with the most frequently requested educational materials.

The CMC staff believe that each person requesting information is a potential grassroots organizer able to educate others about the problems of plastic debris. The staff cultivates each request and acts to network people and information. The elementary school information packet uses a "Playa Pen Pal" program to network even the youngest requests ("playa" means beach in Spanish). All children who ask for information receive the names and addresses of the last three children who wrote in, and are encouraged to exchange information about marine debris in their part of the country with their playa pen pal.



When it's done holding your ship's garbage, it could hold death for some marine animals.

This plastic trash bag may not look like a jellyfish to you. But to a hungry sea turtle, it might. And when the turtle swallows an empty bag, the mistake becomes fatal.

The problem is more than bags. Plastic six-pack holders sometimes become lodged around the necks and bills of pelicans and other seabirds, ultimately strangling or starving them. Other plastic refuse, either through ingestion or entanglement, causes the deaths of thousands of seals, whales, dolphins and other marine mammals every year.

Plastic debris also causes

costly and potentially hazardous delays to shipping when it fouls propellers or clogs intake ports.


It's a critical issue, destined to attract public and government scrutiny if we fail to take action to solve it.

So please, stow your trash, and alert your shipping terminals that you will need proper disposal on land. A sea turtle may not know any better. But now, you do!

To learn how you can help, write: Center for Environmental Education, 1725 DeSales Street, N.W., Suite 500, Washington, D.C. 20036.

A public service message from
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 3.--Commercial shipping public service advertisement.



A seabird could mistake this resin pellet for a fish egg. And die.

One little pellet may be insignificant to your plastics processing operation. But to thousands of seabirds, it could lead to a fatal error.

These pellets, in many shapes and sizes, can be washed down drains as waste or reject material, or spilled in the course of normal handling. But ultimately, they may find their way to bodies of water, where the real trouble begins.

When eaten in sufficient quantity by a seabird, they can block digestion or sometimes fool the bird into thinking it is not hungry, causing eventual starvation. Fish and sea turtles

can suffer the same fate.

The growing problem of plastic trash in our oceans threatens more than wildlife. This critical issue is destined to invite increasing public and government scrutiny unless we take action to solve it.

So please: see that resin pellets are reclaimed or disposed of properly. If we ignore the problem, we—like the unfortunate seabird—will be making a serious mistake.

To learn how you can help, write: The Society of the Plastics Industry, 1275 K Street, N.W., Suite 400, Washington, D.C. 20005.

A public service message from:
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 4.--Plastics industry public service advertisement.



This discarded line is done fishing. But it's not done killing.

Carelessly discarded plastic fishing line can keep working long after you're done with it—entangling birds, seals, sea turtles, and other animals.

And because plastic line is strong and durable, it's nearly impossible for these animals to break free. They strangle, drown, or starve. That's not sporting.

Some birds even use old fishing line in their nests, creating death traps for their young.

Other plastic debris can be dangerous, too. Fish, birds, and seals become entangled in six-pack rings. Sea turtles eat plastic bags—which they mistake for jellyfish—and suffer internal

injury, intestinal blockage, or death by starvation. Birds are known to ingest everything from small plastic pieces to plastic cigarette lighters and bottle caps.

Plastic debris also can foul boat propellers and block cooling intakes, causing annoying—sometimes dangerous—delays and causing costly repairs.

So please, save your old fishing line and other plastic trash for proper disposal.

That's not all you'll be saving. *To learn more about how you can help, write: Center for Environmental Education, 1725 DeSales Street, N.W., Suite 500, Washington, D.C. 20036.*

A public service message from
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 5.--Recreational fishing public service advertisement.



Tossing this trash overboard could leave death in your wake.

Throwing a few plastic items off a boat may seem harmless enough. What's one more six-pack ring, plastic bag, or tangled fishing line?

Actually, it's one more way a fish, bird, seal, or other animal could die.

Fish, birds, and seals are known to strangle in carelessly discarded six-pack rings. Sea turtles eat plastic bags – which they mistake for jellyfish – and suffer internal injury, intestinal blockage, or death by starvation.

Other plastic trash can be dangerous, too. Birds are known to ingest everything from small plastic pieces to plastic cigarette lighters

and bottle caps.

Birds, seals, sea turtles, and whales die when they become trapped in old fishing line, rope, and nets.

Plastic debris also can foul boat propellers and block cooling intakes, causing annoying – sometimes dangerous – delays and causing costly repairs.

So please, save your trash for proper disposal on land.

That's not all you'll be saving.

To learn more about how you can help, write: Center for Environmental Education, 1725 DeSales Street, N.W., Suite 500, Washington, D.C. 20036.

A public service message from
The Center for Environmental Education
The National Oceanic and Atmospheric Administration
The Society of the Plastics Industry

Figure 6.--Recreational boating public service advertisement.

NOAA'S MARINE DEBRIS INFORMATION OFFICE
OPERATED BY THE CENTER FOR MARINE CONSERVATION
EDUCATION MATERIALS LIST AND ORDER FORM

Name _____

Organization _____

Address _____

Phone _____

The National Oceanic and Atmospheric Administration's (NOAA) Marine Debris Information Office distributes educational materials about the effects of plastic in the ocean and about MARPOL Annex V. The materials are free to the public unless otherwise stated. Please order only what you will honestly use because we do have limited quantities.

INFORMATIONAL PACKETS - Choose 4 types (limit 2 packets each)
 We have developed a number of informational packets to meet the needs of different interest groups, however the packets do contain the same basic information.

	Qty.
<u>Elementary School Student</u>	_____
<u>Middle, High and College Students</u>	_____
<u>Teachers or Other Educator</u>	_____
<u>General Public</u>	_____
<u>Beach Cleanup</u>	_____
<u>Plastics Recycling and Degradability</u>	_____
<u>Offshore Oil and Gas Industry</u>	_____
<u>Recreational Boating/Fishing</u>	_____
<u>Commercial Shipping</u>	_____
<u>Port and Terminal Operator</u>	_____
<u>Press and Media Personnel</u>	_____
<u>Plastics Manufacturers/Resin Pellet Producer</u>	_____
<u>Cruise Ship Passenger</u>	_____
<u>Commercial Shipping</u>	_____

BROCHURES - Up to 200 free
 These are 8-panel brochures discussing the problems caused by plastic as related to the interest group mentioned in the title.

<u>Recreational Boating - "Tossing this..."</u>	_____
<u>Recreational Fishing - "This discarded..."</u>	_____
<u>Commercial Shipping - "When it's done..."</u>	_____
<u>General Public - "Our Planet is..."</u>	_____

PACIFIC COAST OFFICE: 312 Sutter St., Suite 606, San Francisco, CA 94108 (415) 391-6204 FAX (415) 956-7441
 ATLANTIC GULF COAST OFFICE: 1725 DeSales St., NW, Washington, DC 20036 (202) 429-5609 FAX (202) 872-0619

Figure 7.--Current version of NOAA's Marine Debris Information Office order form.

**ASSESSING THE IMPACT OF THE CENTER FOR MARINE
CONSERVATION'S EDUCATIONAL PROGRAMS**

Although empirical evidence shows us that hands-on programs are the best form of education, the CMC's 1989 Saltonstall-Kennedy (SK) grant will demonstrate the actual effects of its education programs. Prior to conducting any education events, CMC distributed surveys to a random sample of commercial and recreational fishermen in four designated test areas: Hampton, Virginia; Martin County, Florida; Bayou La Batre, Alabama; and Taylor County, Florida (as a control group). Wallace (1990) of Kearney-Centaur Associates discusses the results of this first SK survey in the paper entitled: "How Much Do Commercial and Recreational Fishermen Know About Marine Debris and Entanglement? Phase 1."

From March to July 1989, the CMC will conduct concentrated educational activities within three of the test areas, excluding the control area of Taylor County, Florida. It will adapt educational activities to accommodate regional differences, style, and events. For example, a large percentage of Martin County, Florida's, recreational fishing community will participate in or attend Arthur Smith Kingfish, Dolphin, Wahoo Fishing Tournament. In addition to providing educational materials to tournament participants, CMC will involve spectators by conducting "Stow It, Don't Throw It" raffle contests for such prizes as boat coolers and fishing reels.

The CMC will distribute a revised version of the survey upon the conclusion of these educational programs. The second survey developed by Kearney-Centaur Associates will assess the impacts of CMC's educational activities. It believes the survey results will statistically demonstrate increased public awareness.

The CMC makes one primary assumption in its approach to solving the marine debris problem. It believes that education will motivate people to alter any harmful disposal behavior. Enforcement of international and national legislation will be very difficult. Marine debris research is both expensive and difficult to conduct in the ocean environment. The CMC feels that education in the form of publicity, books, and if possible hands-on educational events such as beach cleanups will encourage people to keep harmful trash out of the water.

REFERENCES

Center for Environmental Education.

1987a. Plastics in the ocean: More than a litter problem. Center for Environmental Education, Wash., D.C., 128 p.

1987b. 1986 Texas coastal cleanup report. Center for Environmental Education, Wash., D.C., 52 p.

1988a. A citizen's guide to plastics in the ocean: More than a litter problem. Center for Environmental Education, Wash., D.C., 131 p.

1988b. 1987 Texas coastal cleanup report. Center for Environmental Education, Wash., D.C., 105 p.

Interagency Task Force on Persistent Marine Debris.

1988. Report of the Interagency Task Force on persistent marine debris. May 1988, Wash., D.C., 168 p.

Neilson, J.

1985. Get the drift and bag it: A nuts and bolts guide to organizing a beach cleanup campaign the easy way, June 1985; Portland, Oregon, 12 p.

O'Hara, K. J.

1990. National marine debris data base: Findings on beach debris report by citizens. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

Recht, F.

1988. Dealing with Annex V. A reference guide for ports. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-F/NWR-23, 132 p.

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

Wallace, B.

1990. How much do commercial and recreational fishermen know about marine debris and entanglement? Phase 1. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

SHIPPING INDUSTRY MARINE DEBRIS EDUCATION PLAN

Barbara Wallace
Kearney/Centaur Division, A. T. Kearney, Inc.
225 Reinekers Lane, Suite 300
Alexandria, Virginia 22314, U.S.A.

ABSTRACT

The Shipping Industry Marine Debris Education Plan is an education and awareness program on MARPOL Annex V and its consequences for foreign and domestic commercial shippers and cruise lines operating in U.S. waters and for port and terminal operators. The plan is based on the premise that education may be a key factor in gaining voluntary compliance with MARPOL Annex V provisions and, therefore, in reducing the problems caused by marine debris and entanglement. The plan identifies five core and three ancillary activities to increase the shipping industry's awareness of MARPOL Annex V and the consequences of marine debris and entanglement. It was developed in late 1988 and early 1989 under contract to the Marine Entanglement Research Program, U.S. National Marine Fisheries Service. The recommended activities will be implemented over a 9-month period beginning in the spring of 1989. The core activities are: 1) development of case studies of MARPOL Annex V compliance activities, 2) development of a model plastics refuse control and minimization plan, 3) preparation of a MARPOL Annex V kit--a one-source document on MARPOL Annex V implementation in the United States including crew awareness training on the consequences of marine debris, 4) liaison activities with international shipping industry trade associations, and 5) seminars for cruise line owners/operators. The ancillary activities are: 1) placement of public service advertisements on marine debris in trade journals, 2) placement of posters on MARPOL Annex V and marine debris in port areas and on board vessels, and 3) presentations on the plan and its activities at seminars and workshops.

INTRODUCTION

On 31 December 1988, in response to at-sea garbage disposal limitations that went into effect worldwide, the way mariners handle disposal of ship-generated garbage changed. On that date, Annex V of the International Convention for the Prevention of Pollution from Ships, 1973 (MARPOL 73/78) went into effect. Formally, MARPOL Annex V is entitled,

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

"Regulations for the Prevention of Pollution by Garbage from Ships." Simply, Annex V prohibits at-sea disposal of plastic materials and specifies the distance from shore that all other materials may be dumped.

The MARPOL Annex V will be difficult to enforce. Among other things, it changes the traditional way of handling ship-generated garbage. Education may be a key factor in gaining compliance with MARPOL Annex V. In the last several years, marine debris education programs have been developed and implemented for several major marine industry groups (for example, offshore oil and gas workers and commercial fishermen). The current project focuses on marine debris education for the shipping industry.

The amount of garbage generated by the shipping industry varies. It is estimated that large commercial vessels generate between 18 and 40 garbage bags of plastic during a typical voyage. This includes both domestic wastes and plastics included in dunnage. Tow and tugboats are estimated to generate one to three bags of plastic per voyage. Cruise ships are estimated to generate over 70 bags of plastic per day (Eastern Research Group 1988). Until MARPOL Annex V, much of this garbage was disposed of at sea.

The Shipping Industry Marine Debris Education Plan was developed under contract to the Marine Entanglement Research Program, U.S. National Marine Fisheries Service (NMFS) to: 1) ensure that foreign and domestic commercial shippers and cruise lines operating in U.S. waters, and port and terminal operators are aware of the provisions of MARPOL Annex V as it is being implemented in the United States; and 2) encourage voluntary compliance with those provisions. It was developed to help avoid disruption of shipping schedules, which could occur if U.S. prosecution of at-sea garbage disposal violators is necessary. The plan was also designed to be part of the solution to the problems of marine debris by furthering awareness of its consequences.

METHODOLOGY

The Shipping Industry Marine Debris Education Plan was developed using four concurrent tasks (Fig. 1) involving data collection and review of existing information, a synthesis task, and an industry advisory panel. The concurrent tasks were completed using secondary sources and key informant contacts. The major points which emerged from the tasks helped to structure the recommended plan activities.

An industry advisory panel for the project was created to involve the industries affected by the plan in its development. By doing so, the intent was to avoid identifying marine debris education activities which would not work. It was also a mechanism to encourage industry's further involvement with the plan, its implementation, and the issue of marine debris and entanglement. Four trade associations, the U.S. Department of Agriculture, the U.S. Coast Guard, and a maritime professional association participated on the panel.




Figure 1.--Plan development methodology.

A preliminary list of marine debris education activities was developed based on the results of the data collection and synthesis tasks. A packet of materials, including a one-page summary of eight marine debris education activities, was sent to each panel member for review. The industry advisory panel attended a 1-day meeting in Alexandria, Virginia, to discuss MARPOL Annex V implementation and the Shipping Industry Marine Debris Education Plan. The marine debris education activities recommended in the plan reflect the input of the industry advisory panel.

RECOMMENDED MARINE DEBRIS EDUCATION ACTIVITIES FOR THE SHIPPING INDUSTRY

Eight marine debris education activities are recommended for the shipping industry. An overview of these activities is presented in Figure 2. Activities 1 through 5 are the core activities in terms of time and budget. Activities 6 through 8 will have a secondary emphasis, but some time and funds will be allocated to them.

Core Activities

Activity 1: Case Studies of MARPOL Annex V Compliance Activities

No one method of compliance with MARPOL Annex V will work for all vessels because of differences in such things as size, routes, cargo, and owners. The case studies will document policies and activities (including crew awareness training) that operators of different types of vessels as well as port and terminal operators are using to comply with MARPOL Annex V. The approaches different companies are using for MARPOL Annex V compliance are expected to reflect these differences. For example, seagoing vessels, which include commercial shippers and cruise lines, must comply with regulations on food waste coming from foreign waters, as set forth by the Department of Agriculture Animal and Plant Health Inspection Service (APHIS), as well as the provisions of MARPOL Annex V. Some of these vessels may also travel to special areas as defined by MARPOL Annex V, where dumping limitations are further restricted. Some coastal waterway vessels travel far enough from shore to be able to dump some types of garbage at sea legally. These vessels do not leave U.S. waters and, therefore, do not need to comply with APHIS regulations or the dumping restrictions in MARPOL Annex V special areas. Vessels in inland waterways and harbor areas have been prohibited from dumping garbage overboard for almost 100 years. Flexibility in the method used is seen by many in the industry as a key to voluntary compliance. The case studies will document how and to what extent flexibility is used for MARPOL Annex V compliance.

Until recently, much effort in marine debris education was spent identifying the sources of debris and convincing those industries or marine user groups that their practices were causing problems. This had a negative tone. The case studies will be examples of positive actions companies have taken to change the way they handle plastics and garbage disposal. They will demonstrate the range of techniques that companies have used to comply with MARPOL Annex V. They will also identify some of the problems encountered and how these were overcome. The case studies will be developed through key contact interviews and documents provided by these contacts.




Figure 2.--Shipping industry marine debris education plan overview.

Activity 2: Model Shipboard Plastics Refuse Control and Minimization Plan

Each company will need to determine how to handle garbage under MARPOL Annex V. To facilitate that process, some documentation is needed on what to consider in developing strategies for plastic garbage disposal. A formal plastics refuse control and minimization plan can be an important component of a company's policy on garbage handling.

The model plan will outline the development of waste management procedures and the content and possible formats of a plastics refuse control and minimization plan. It will include techniques for increasing crew awareness of the consequences of marine debris and entanglement, and will use existing marine debris education materials. The model plan will be based on experiences profiled in the case studies and in existing waste management plans. Key contact interviews also will be used.

It should be noted that the Coast Guard will require certain vessels to have a waste management plan. Coast Guard guidance will be sought during the preparation of the model plan. However, the model plan may not be fully in keeping with Coast Guard requirements because it is likely to be completed before the Coast Guard issues instructions on the requirements for waste management plans.

Activity 3: MARPOL Annex V Kit

The MARPOL Annex V kit will assemble existing information on MARPOL Annex V implementation in the United States. The kit will provide a focal point for discussion with international groups on what is available on MARPOL Annex V implementation in the United States, and will be a product which can be printed and distributed by different groups (for example, trade associations and the NMFS Marine Debris Information Offices).

The kit, probably in looseleaf binder form, will include such things as the Coast Guard compendium of materials on MARPOL Annex V implementation in the United States, examples of how the requirements are being implemented by different types of vessels (from Activity 1), and guidance on how to set up shipboard waste management procedures and crew awareness training (from Activity 2). The kit will also include examples of existing marine debris education products (brochures, posters, stickers) that can be used on vessels for training or reminders about MARPOL Annex V requirements.

Activity 4: Liaison with International Shipping Industry Trade Associations

There are too many foreign flag vessels operating in U.S. waters to approach each owner and operator directly. However, the international shipping industry trade associations can be used as an indirect means of contact with foreign commercial shipping owners and operators. Personal contact, such as has been used with shipping industry trade associations in the United States, is more effective than telephone and mail contact in soliciting cooperation and exchange of information on MARPOL Annex V implementation. Meetings will be requested with the international shipping trade associations located in London. These meetings will be used to: 1)

make the associations aware of the marine debris education and MARPOL Annex V implementation materials available, 2) encourage them to publish announcements of marine debris education products in their trade publications, 3) encourage them to reproduce some of the materials, 4) increase awareness of the marine debris education activities taking place in the United States, and 5) identify specific needs where existing materials and programs may be of help.

Activity 5: Seminars for Cruise Line Owners and Operators

Experience in marine debris education activities has shown that involvement of the targeted group in development of an education program can be effective in making the program a success. This is the first time in a NMFS-sponsored marine debris education program that the cruise industry is the focus of marine debris education activities. Since the number of cruise lines operating in U.S. waters is small, it is possible to approach these companies directly.

Seminars with cruise line owners and operators will be conducted in Miami, San Francisco, and Washington, D.C. Offices of the majority of the cruise lines are located in or near these cities. These meetings will be used to: 1) solicit individual company support for marine debris education activities; 2) identify specific educational activities (e.g., working with the Centers for Disease Control to include marine debris information in procedures for shipboard inspections); 3) identify mechanisms to distribute information (e.g., include information with paychecks or stubs); and 4) develop educational materials or assemble existing materials in cooperation with company and industry representatives.

Other Activities

Activity 6: Public Service Advertisements

Art work and brochures that made up a previously used public service advertising campaign are available. Since there is new interest in marine debris as a result of MARPOL Annex V implementation, there may be new opportunities to place the ads in appropriate trade publications. The art work for the public service advertisements will be offered to editors of shipping, port, and cruise line trade journals.

Activity 7: Poster Project

There are several posters available which could be used on ships or in port areas as reminders of the consequences of plastics in the ocean. Some vessel owners and operators have expressed interest in having posters, and their use would probably increase if more people knew about them. Written announcement about the posters on marine debris will be sent to: 1) Federal agencies with offices in port districts (e.g., Coast Guard, U.S. Customs, U.S. Immigration and Naturalization Service, and Department of Agriculture); 2) port authorities; 3) terminal operators; and 4) shipping companies.

Activity 8: Seminar and Workshop Presentations

Seminar and workshop presentations are a means of reaching different types of audiences and increasing awareness of the problem of marine debris as well as the ongoing activities on MARPOL Annex V implementation in the United States. The Shipping Industry Marine Debris Education Plan and its implementation will be presented at appropriate forums, should the opportunity arise.

SHIPPING INDUSTRY MARINE DEBRIS EDUCATION PLAN IMPLEMENTATION

The Shipping Industry Marine Debris Education Plan will be implemented during a 9-month period beginning in the spring of 1989. The first half of the implementation phase for the core activities (Activities 1-5) will be used to prepare the new marine debris education products and establish a working relationship with the cruise line owners and operators. The second half of the implementation phase will be used to gain acceptance and distribution of the education products. The optional activities (Activities 6-8) will be undertaken throughout the implementation phase as the opportunity arises and time and budget permit.

REFERENCES

Eastern Research Group, Inc.

1988. An economic evaluation and environmental assessment of regulations implementing Annex V to MARPOL 73/78. Prepared for the U.S. Coast Guard in support of the regulations implementing the pollution prevention requirements of Annex V to MARPOL 73/78, October 1988.

CLEANUP PROGRAM IN JAPAN

Nobuyuki Yagi and Yoshie Otsuka
Division of Fishing Ground Environment Conservation
Fisheries Agency, the Government of Japan
Chiyoda-ku, Tokyo, 100 Japan

ABSTRACT

The amount of marine debris accumulated in Japanese coastal areas has showed an increasing trend from the early 1970's, and this debris has become an obstruction to fisheries operating there, causing destruction of habitats of target species and interfering with fishing operations. In fiscal year 1973, the Fisheries Agency started a program to cope with these issues in coastal areas.

This program consists of two activities. One is the education of local residents including fishermen, and the other is the actual cleanup of the seashore and sea bottom. There are three parts to the cleanup activities: cleaning up the seashore using manpower, cleaning up the fishing grounds using trawl nets, and cleaning up the rocky bottoms using divers.

Cleanup activities were carried out in 137 areas in fiscal year 1987, at a cost to the Japanese Government of ¥342,249,000 (US\$2,738K).

INTRODUCTION

From early times, the Japanese have shown keen interest in various types of drifting objects stranded on beaches. In some places, people even developed habits of worshipping strange-shaped pieces of driftwood as deities. In coastal areas where the Kuroshio passed, people picked up coconuts which drifted ashore on rare occasions and dreamed of islands far to the south.

In recent years, however, great changes have been observed in the types and quantity of debris. Debris items were no longer objects of veneration as in past times but were a nuisance, causing damage to people's livelihood.

PROGRAM BACKGROUND

In the early 1970's, complaints were voiced by fishermen in various parts of Japan that wastes stranded ashore or accumulated on the sea bottom were causing serious damage to fishing activities. Such wastes ranged from man-made objects, including bottles, bottle covers, cans, worn-out tires, plastic bags, and other disposed plastics, to natural objects such as grass, wood, and vegetable garbage carried by rivers after torrential rains. Damage assumed to be caused by debris was varied, according to the fishermen. Some extremists contended that debris not only obstructed fishing operations but even caused a decline in the number of fish as a whole. Fishermen presented these problems in all earnestness because they were experiencing direct economic losses.

The damage seeming to have most apparent causal relations with wastes is summarized as follows:

1. Damage to organisms. Plastic bags attached to rocks on the sea floor (the natural habitat for seashells) make it impossible for the seashells to live there. Nondegradable wastes accumulated in shallow water deprive fish of spawning and nursery grounds. Fishing lines, cut or abandoned at sea or on the beach, break seaweed and entangle seabirds.
2. Damage to ships. Plastic bags obstruct engine cooling water intakes. Lost or discarded nets entangle propellers.
3. Damage to fishing activities. Plastic bags caught in trawl net meshes increase water resistance and thereby damage fishing gear. Increased resistance of gear in the water lessens energy efficiency of fishing vessels and increases fuel cost.

Much time is required to sort out fish from debris caught in trawl nets, cutting down on fishing efficiency. Also, fish taken together with debris frequently bring a lower price.

4. Others. The scenery of beaches is affected, giving an unfavorable impression to visitors who come for sea bathing.

Besides reports on damage, proposed solutions came from various parts of the country. These included requesting the public not to dispose of garbage in the sea and actively collecting debris being accumulated on beaches and the sea floor.

However, several problems complicated the solution. First, there was the difficulty of identifying who actually discarded the wastes. The wide range of potential contaminators included ordinary residents, factories, tourists, ships, and fishermen themselves. Therefore, it was impossible to identify the actual polluters and have

them bear the cost of the damages. Second, the effort of people in any particular area is not sufficient to solve the problem. As marine debris comes from various areas including inland areas and areas hundreds of miles away, a number of municipality offices need to cooperate in efforts against this problem.

ESTABLISHMENT OF THE PROGRAM

In fiscal year 1973, the Fisheries Agency launched a program (hereafter referred to as the "Cleanup Program") aimed at preserving the marine environment and recovering deteriorated fishing grounds. The Fisheries Agency formulated and coordinated the entire program, while regional authorities for each area concerned were responsible for the actual implementation.

Contents of the Program

As for specific program items, local authorities were entitled to select from among the following options, giving due consideration to issues peculiar to their own areas:

1. Alert regional residents including fishermen to the need to preserve the marine environment, conducting appropriate educational activities through television and radio broadcasting and newspapers as well as through calls from aircraft, lectures, public ads, posters, leaflets, calendars, and bathing caps and towels.
2. Cleanup debris accumulated on the sea bottom using trawl vessels.
3. Cleanup debris on the rocky bottom using divers.
4. Eliminate wastes drifting on the sea surface using dipnets.
5. Cleanup the beaches using manpower.
6. Use manpower and machines to cleanup rivers and lakes.
7. Establish councils composed of local authorities, fishermen, and academics to formulate specific cleanup programs.

National Budget for the Program

The cost of implementing the Cleanup Program is covered in part (usually half of project costs) by the National Government, with the remaining amount shouldered by local authorities actually enforcing the program. The government budget for this purpose increased annually, the subsidy for fiscal 1973 (April 1973-March 1974) being ¥96,000,000 (US\$768K), rising to ¥401,126,000 (US\$3,209K) in fiscal 1979. After that the budget remained more or less unchanged (Fig. 1).




Figure 1.--Annual changes of subsidy under the government budget for the Cleanup Program.

CLEANUP TODAY

At a cost to the national treasury of ¥342,249,000 (US\$2,738K) in fiscal year 1987, the program was conducted in a total of 132 areas throughout Japan. The number of areas differed in different prefectures. Some prefectures conducted the program in as many as 23 areas, while some did not carry out any program (Fig. 2).

Following are actual situations in four areas during fiscal year 1987 and the cleanup programs in each area (Fig. 3).

Mutsu Bay, Aomori Prefecture

Mutsu Bay measures about 1,660 km², with a total coastline extension of 251 km. Fishing is a major industry there, notably the scallop fishery, with production worth about ¥13 billion (US\$104 million) in fiscal year 1987. In 1975, a large number of scallops died of an unidentifiable cause in the bay, which up to then was known for its relative cleanliness. This incident prompted local fishermen to request measures to preserve the fishing ground environment, and a cleanup program was implemented in the same year.

In 1987, cleanup activities were conducted in 10 towns and villages covering more or less the entire coast of the bay at an overall cost of ¥11,640,000 (US\$93,120). The scale of the program has been about the




Figure 2.--Distribution of prefectures taking part in the program, shown by number of areas in which the program was conducted in 1987.




Figure 3.--Location of the four areas: Mutsu Bay, Oarai, Mikawa Bay, and Hirao.

same ever since. The program consisted of three major parts: cleaning up the beach, cleaning up the sea bottom, and conducting educational activities.

Beach cleaning was carried out for several days in July and August, with a total of 5,872 residents taking part, including fishermen, housewives, and students. Each participant was supplied with a pair of cotton gloves and a garbage bag, and wastes gathered totaled 356.2 metric tons (MT). Major components were wood fragments and seaweed. Combustible wastes were incinerated at the beaches, and incombustible items were transported by truck to garbage storage areas of local authorities to be used as landfill.

Sea bottom cleaning was conducted for several days mainly in July, with a total of 2,089 small-sized beam trawlers participating. Fishermen used their own nets for collection of wastes. Debris collected amounted to 163.8 MT, and consisted mainly of used cans and seaweed. There was a larger percentage of seashells and used cans than other wastes on the beaches. All the debris was transported to local authorities' garbage storage areas and landfills.

Educational activities were mainly targeted at elementary school pupils and junior high school students with a view to achieving long-term effects. Guidance on the importance of preserving the marine environment was extended in the course of ordinary school curriculums, and pupils and students were encouraged to make posters and catchphrases contributing to environmental preservation. A total of 1,146 posters and 1,904 catchphrases on marine environmental preservation were collected from the children at 37 schools from July to September 1987. Excellent works were publicly commended with commemorative awards worth ¥3,000-5,000 (US\$24-25) (including a book of gift coupons and a painting set). These were distributed for public presentation in the towns and villages concerned. This program has been established as an annual event in the area.

Oarai, Ibaraki Prefecture

Oarai is a Pacific coast town with a population of 21,000, and its major industries are tourism and a coastal fishery. Located relatively close to the Tokyo metropolitan area, Oarai has seen rapid urbanization during the past few years. The amount of wastes has been rising, making their disposal increasingly difficult despite the municipal authorities' effort. The town is situated at the mouth of a big river (Nakagawa River), and it has been pointed out that a great amount of debris flows into the sea from the river. Fishermen are worried that sardine fry, caught with trawl nets, can be easily damaged by garbage netted simultaneously. In the summer, about 8 million people visit this town for sea bathing. The municipal authorities have a hard time disposing of used cans and bottles generated by these seagoers.

Sea bottom 15-30 m deep and totaling 7.84 km² was cleaned on 18 January and again on 20 February 1988 using 57 small (5-ton) trawlers. Fuel expenses of ¥5,400 (US\$43) and ship depreciation expenses of ¥22,000

(US\$176) per day were provided to each vessel. Fishermen used their own trawl nets for cleaning activities. The program was conducted during off seasons of the coastal trawl fishery to avoid an unnecessary by-catch of fish. As a result of the cleanup, 15.5 MT of debris, mostly used cans and plastic bags, were collected and removed to a landfill.

Since the Cleanup Program was launched in 1974, fishermen have seen a decrease in the amount of debris caught in their nets, and they have become increasingly active in many areas in this region.

Mikawa Bay, Aichi Prefecture

Mikawa Bay, with an area of 604 km² and an average depth of 9 m, is a closed area with poor tidal interchange. At the back are located industrial and urban areas, and many rivers which run through these areas flow into the bay. Major industries of the bay area are fishing and tourism, with a resort business targeted at sea bathers.

In cleanup activities implemented in fiscal 1987, a total of 464 residents in 8 areas cleaned up the 30.9 km coastline along the bay, collecting 5,810 kg of wastes from 120,000 m² covered. Principal debris was wood fragments, cans, and bottles (Fig. 4). All the debris collected was either taken to a landfill or incinerated. A total subsidy of ¥1,000,000 (US\$8K) was granted, and was used for purchasing commemorative items, gloves, polyethylene bags, and fuel for incinerating waste.

Hirao, Yamaguchi Prefecture

Hirao is an agricultural and fishing town with a population of 15,000, facing the Seto Inland Sea. The Seto Inland Sea is the largest of its kind in the country, with a latitudinal extension of 445 km, a longitudinal extension of 15-18 km, and a total area of 2,200 km². The average depth is relatively shallow at 37.3 m. This is an area of serious concern when it comes to marine environmental pollution, as it is a closed area and has a population of 29,359,000 along the coast.

Impressive educational activities are being carried out in Hirao. With the collaboration of residents, an experiment was conducted to find out how the garbage arrived at the town by way of sea currents and winds, and where the garbage originating in the town goes. This experiment is known among the residents as "Coconut Strategy," with the capsules used in the experiments being thought of as coconuts. In July 1987, 25 plastic capsules 10 cm in diameter and 40 cm long were set adrift from 25 points along the coast within a radius of 50 km from the center of the town. Posters were placed in the town and surrounding areas in order to draw the attention of residents and ensure that reports would be made when the capsules were found on beaches. So far, a total of 15 capsules have been recovered. Around the areas where capsules were released, the results were publicized using posters saying, "The capsule released from your town arrived at such-and-such an area. This means if you discard garbage into the sea, it will possibly drift to that area and cause trouble. The sea does not belong only to you. Please take heed not to contaminate the sea."




Figure 4.--Composition of debris collected from beaches during the cleanup of Mikawa Bay in July 1987.

A total of ¥322,000 (US\$2,576) was used for producing capsules and posters and for other purposes.

The campaign has grown as town residents have begun to realize the need to raise public ethical standards, and an increasing number of people are participating.

DISCUSSION

Effects of the Program

The Cleanup Program was initiated under the leadership of the administrative authorities, but in many cases, programs were taken over willingly by local residents as years passed.

People's interest has grown as the educational programs have continued, and many people hope that the programs will carry on. Some communities have even formulated their own new and voluntary initiatives, in addition to the basic guidelines. One such example is the establishment of "Fishing Ground Preservation Month," during which stepped-up publicity is conducted.

There have been two different views concerning the effects of the Cleanup Program. One recognizes the effectiveness of the program based on

the fact that the amount of garbage has been steadily declining since the program was launched. The other questions the effectiveness of the Cleanup Program, pointing out that the garbage amount has not decreased substantially even after several years. These differences are due to the fact that the pace of waste accumulation differs from area to area, that amounts collected are subject to weather and sea conditions, and that it is difficult to assess the effects of the program quantitatively. However, even people taking the latter position believe that implementation of the Cleanup Program contributes to an increase in public awareness of environmental preservation. It is therefore concluded that the program as a whole has been significant.

Future Themes

Some future tasks have been pointed out that will improve the effectiveness of the program. First is the need to conduct the program in a comprehensive manner covering a larger area than at present. In many cases, cooperation in the present program has been limited to the level of municipalities along a bay area. In order to cope with environmental problems covering the vast ocean, it will be necessary to step up and expand collaboration. Further, it will be no less important to expand the scope of the program to inland areas. A considerable amount of wastes generated by the people in those areas is transported through rivers to the sea. Garbage originating inland can accumulate on the beaches.

Second, it is necessary to hunt for new ways to promote a voluntary environmental preservation campaign. One potential method is to install garbage processing facilities in areas where individuals voluntarily collect wastes. Such support is expected to further increase the local communities' awareness of the need for environmental preservation. Lastly, some technological problems need to be solved. Although the program does not require any special technology, some problems have arisen as a result of its actual implementation. These include the disposal of incombustible objects containing large amounts of sand and seawater (salt water) and the elimination of wastes accumulated around complex bottoms or man-made structures such as artificial and natural reefs.

ACKNOWLEDGMENTS

We express our gratitude to all those who are striving to promote the Cleanup Program. Special appreciation is due to Messrs. Nagatsu of Aomori Prefecture, Chinone of Ibaraki Prefecture, Matsui of Aichi Prefecture, and Asaka of Yamaguchi Prefecture, who collaborated in collecting materials.

U.S. NAVY'S PLASTICS WASTE EDUCATIONAL EFFORTS

Larry Koss
Office of Chief of Naval Operations (OP-452)
Department of the Navy
Washington, D.C. 20350, U.S.A.

Fred Chitty
Supply Operations and Policy
Commander in Chief, Atlantic Fleet
Department of the Navy
Norfolk, Virginia 23511, U.S.A.

and

William A. Bailey
GEO-CENTERS, Inc.
10903 Indian Head Highway
Fort Washington, Maryland 20744, U.S.A.

ABSTRACT

The U.S. Navy is responding to the at-sea plastics discharge prohibition in the Marine Plastic Pollution Research and Control Act with a program that includes new shipboard solid waste management equipment, changes in the supplies taken on board, and educational efforts.

Basic strategy of the Navy's plastics waste education program focused on motivating the entire chain of command, ships' officers, and ships' crews by providing justification for and information about the new requirements.

Before designing an effective shipboard plastics waste management program, it was necessary to learn (1) the nature and quantities of plastics waste on Navy ships, (2) feasible shipboard plastics waste management practices, and (3) sailors' attitudes about plastics pollution control. The Shipboard Plastics Waste Reduction Demonstration Project gathered this information on seven ships. Then, using recommendations from an ad hoc advisory committee on plastics and actual experiences on the demonstration ships, the Navy designed an educational package to send to all Navy ships.

The Navy's plastics education package includes guidance material, videotapes, posters, and general literature. To educate the ships' officers, a ship's guide contains chapters on (1) problems caused by plastics in the oceans, (2) Navy

requirements, (3) essential elements of a successful shipboard program, (4) example approaches used on the demonstration ships, and (5) general information about related issues. Appendixes to the guide include lists of common plastic and substitute nonplastic items, sample ship instructions to implement the program, descriptions of other materials in the package, and Navy points of contact for further information. To educate the crew members, a 10-min videotape explains the problem, the Navy's program, and appropriate shipboard actions. A series of 30-sec videotapes and written announcements highlight specific aspects of the problem and the program.

Preliminary responses to the education package from Navy ships are very positive.

INTRODUCTION

Impetus for Navy's Plastics Program

Recent and unexpected national regulatory initiatives caused the U.S. Navy to quickly develop and implement shipboard plastics waste management programs. The Navy anticipated more stringent international regulations affecting solid waste disposal at sea, but not a prohibition on plastics discharge.

Annex V of the Protocol of 1978 relating to the International Convention for the Prevention of Pollution from Ships (MARPOL), prohibits disposal of all plastics, and restricts discharges of all types of solid wastes, into the sea from ships. Public vessels are exempt from Annex V restrictions but are expected to comply to the extent possible. However, the U.S. Congress, in passing the U.S. Marine Plastic Pollution Research and Control Act of 1987 (P.L. 100-220), made the plastics discharge prohibition apply to the U.S. Navy. The Congress is requiring the Navy to comply within 5 years or demonstrate why it cannot.

For decades the Navy has worked to eliminate the discharge of floating marine debris from its ships, and for the last 5 years, has been developing suitable shipboard equipment for pulping, compacting, and processing all solid waste (Alig et al. 1990). The equipment will convert all shipboard solid wastes to acceptable forms for overboard disposal at prescribed distances from shore. However, plastics waste would have been processed along with other solid wastes, so the new requirement to eliminate plastics waste disposal at sea created an unexpected and difficult challenge.

Ad Hoc Plastics Advisory Committee

The Navy formed an ad hoc committee on plastics to develop recommendations for reducing plastics waste discharge from Navy ships, and in October 1988, the committee presented its report to the Assistant Secretary at a Washington, D.C., news conference. The report contains 42 recommendations covering 4 areas: supply system, technology, ship operations, and education.

The Navy's shipboard pollution abatement program was already addressing most of the committee's technology recommendations as part of the shipboard solid waste management program. The recommendations in the areas of the supply system, ship operations, and education, however, embodied major changes in Navy practices. Nevertheless, the Navy began implementing many of the recommended actions to help reduce plastics discharge at sea.

Following are the six recommendations made in the area of education:

1. Disseminate information on the plastics problem and the Navy's plans to all levels in the Navy.
2. Develop education materials for ships, supply centers, and procurement offices.
3. Use an education package composed of several key elements on every Navy ship.
4. Assess effectiveness of education efforts through a survey.
5. Use the Navy's environmental award to recognize outstanding ships for their efforts to reduce plastics waste.
6. Sponsor shipboard contests on plastics reduction methods.

Navy Actions

The Navy has already taken significant actions to control plastics pollution at sea. As examples, fleet commanders ordered all ships to separate plastics from nonplastics waste on board and return the plastics waste to port for disposal or recycling. Supply centers have replaced plastic items with nonplastic ones on ships wherever possible. The Chief of Naval Operations office sent a plastics education package to all ships. This paper describes the Navy's initial efforts to educate ships' officers and crews about the new requirement to eliminate plastics waste at sea.

GUIDING PRINCIPLES

The key to success for a new Navy program is to motivate the chain of command and the sailors to participate fully. But in the case of the plastics waste program, the entire chain of command also had to support the program, both financially and authoritatively.

Before the top echelons of the Navy's chain of command would impose such drastic measures as storing wastes on Navy warships, they needed evidence showing shipboard plastics waste programs to be feasible, effective, and safe. Therefore, we had to assess the feasibility, effectiveness, and safety of potential shipboard plastics waste management practices before we could propose a program concept that the chain of command would support.

The program needed careful planning and coordination among many diverse Navy organizations (e.g., Office of the Chief of Naval Operations,

Fleet Commands, Naval Sea Systems Command, Naval Supply System Command, and Navy laboratories). Even then, edicts and emotional appeals do not ensure a successful program. For drastic changes in shipboard waste management practices, a variety of motivation techniques would be needed to get full cooperation on ships. Some officers and crew would be sufficiently motivated by direct orders. Others might respond to the military benefits of new procedures. Sound explanations of the reasons for new requirements would suffice for some. Still others would get personally motivated by appealing to their sense of pride or the need to protect wildlife.

Our guiding principles, then, in developing the educational program were to invoke as many response stimuli as we could.

EDUCATION OF NAVY PROGRAM MANAGERS

Successful reduction of plastics discharge from Navy ships requires two educational efforts. One is educating the Navy chain of command about what shipboard practices are feasible and effective. The other is educating ships' officers and crews about the new practices. To achieve the first step, the Navy needed to learn (1) the nature and quantities of plastics waste on Navy ships, (2) feasible shipboard plastics waste management practices, and (3) sailors' attitudes about plastics pollution control. In 1988, the Navy's Shipboard Plastics Waste Reduction Demonstration Project began collecting this information.

During the demonstration project, Navy researchers traveled on seven ships to quantify shipboard plastics waste, and evaluated prototype equipment and plastics waste management procedures. They sorted and inventoried all plastics waste generated on board. On two ships they evaluated Navy-developed prototypes of a trash compactor and a waste pulper. Other researchers surveyed the crews' knowledge about the plastics problem.

The preliminary conclusions from the demonstration project regarding an education effort are the following:

- Sailors are generally receptive to the major new requirement to separate and store plastics waste on board, once the problem with plastics in the oceans is explained to them.
- The mission, size, crew density, and operating characteristics of different ships differ so much that a single set of prescribed plastics waste management procedures is not appropriate for all ships.
- Ships prefer to find their own approaches to meeting the requirements, i.e., be told what to do, not how to do it.

EDUCATION OF SHIPBOARD PERSONNEL

Development of Education Package

Initially, we planned to prepare a prototype educational package and test its effectiveness on several demonstration ships before sending it to

all Navy ships. However, the Commander of the Atlantic Fleet took a personal interest in the plastics problem and pushed for faster implementation of shipboard programs. His staff prepared instructions for ships to retain plastics waste on board to the extent practicable. He wanted an educational package sent to ships before he would issue the new instructions. Consequently, we had to assemble educational materials quickly without knowing exactly what shipboard procedures were feasible and what educational materials would be effective on ships.

Our basic strategy for the educational package targeted two audiences on ships: executive officers and crew members. We prepared a ship's guide and a new videotape, and supplemented these two key items with available literature, posters, brochures, and videotapes.

The ship's guide serves as a compilation of useful information about the new requirements and suggestions for implementing a shipboard program. To allow ships' officers and crews to find their own ways of meeting the new requirements, the material was presented as a guide, not an instruction.

Because closed circuit television is the prime source of news and entertainment on ships, we believed that videotapes would be the best way to influence and communicate with sailors. Therefore, the Navy produced new videotapes that focused specifically on the Navy's role in the plastics problem. As a general overview, we prepared a 12-min tape that covers all aspects of the problem and the Navy's program. To add authority and credibility to the new requirements, the Vice Chief of Naval Operations made a 3-min statement on videotape. To keep the issue visible to the sailors and reiterate the theme of the plastics program, we produced a series of 30-sec tapes that could be shown on board between movies. For variety, we included in the package existing videotapes produced by other organizations.

Contents of Education Package

The Navy's plastics waste educational package that was sent to all ships included the following items.

- "Ship's Guide to Recent Navy Initiatives for Shipboard Solid and Plastics Waste Management" (U.S. Navy).
- Videotapes in Beta and VHS formats with six selections:
 - "Plastics At Sea, More Than a Litter Problem" (12 min, U.S. Navy).
 - Statement by Admiral Edney, Vice Chief of Naval Operations (3 min).
 - Eight 30-sec public service-type announcements (U.S. Navy).

- Navy Broadcast News segment on the plastics problem (7 min).
- ABC News excerpt on ocean pollution (3 min).
- "Trashing the Oceans" (7 min, U.S. National Marine Fisheries Service).
- "A Citizen's Guide to Plastics in the Ocean: More Than a Litter Problem" (Center for Environmental Education, Washington, D.C.).
- Three posters:
 - "Don't Splash Navy Trash--Others Can Pick It Up" (U.S. Navy).
 - "Our Ocean Is Drowning" (U.S. National Marine Fisheries Service).
 - Photograph of large plastic bag of trash with narrative (Center for Environmental Education, Washington, D.C.).
- "Our Water Planet is Becoming Polluted with Plastic Debris" (brochure prepared by U.S. National Marine Fisheries Service).

The Navy's videotape "Plastics At Sea, More Than a Litter Problem" explains the problems caused by plastics in the oceans, the Navy's overall program for shipboard solid and plastics waste management, and what actions are needed by each ship and crew member. The tape is narrated by an actor in an enlisted man's uniform and contains footage of healthy marine life, entangled animals, Navy ships, shipboard scenes, and computer graphics. To appeal to sailors, segments of the Huey Lewis videotape "Perfect World," showing his band singing in a huge garbage dump, appear throughout the tape.

The ship's guide is similar to a citizen's handbook, except that it contains information applicable to Navy ships and the Navy's plastics waste program. The guide is approximately 50 pages long, with 6 chapters and 9 appendixes. Chapter 1 (Introduction) briefly describes the problems caused by plastics at sea and the international and national regulations restricting disposal at sea. Chapter 2 (Navy Requirements) explains the major changes needed in shipboard solid waste management. Chapter 3 (Essential Elements of Successful Shipboard Program) emphasizes that a shipboard program needs comprehensive operational changes, leadership, crew education, constant reinforcement, and clear definitions of responsibilities. Chapter 4 (Example Approaches Used on Demonstration Ships) provides suggested procedures for ships to use in their own shipboard plastics waste program, based on successful and unsuccessful approaches used on the demonstration ships. Chapter 5 (Medical Wastes) describes the new requirements for managing medical wastes on board. Chapter 6 (Status of Related Issues)

addresses three topics related to the plastics program: degradable plastics, shipboard incinerators, and commercial products.

The nine appendixes are a collection of miscellaneous items to help executive officers develop their ship's program. We provided lists of common plastic items on ships and the available nonplastic substitutes. Brief synopses of information about the plastics problem, sources of marine debris, legislation, and the Navy's program are included as sample "Plan of the Day" announcements for executive officers' use in keeping the program visible to crews. Included also are example ship instructions for implementing a shipboard program and an example message report for reporting any necessary plastics discharge at sea. Other appendixes describe the contents of the educational package and give Navy points of contact for additional information.

PRELIMINARY OBSERVATIONS ON EFFECTIVENESS OF PACKAGE

The Navy sent complete educational packages directly to all ships in the U.S. Fleet in early 1989. Preliminary responses to the education package are very positive. The combination of useful information and multimedia motivational techniques in the package seems appropriate for Navy ships. Each item in the package seems useful for one or more purposes. However, the two items specifically prepared by the Navy for the education package (i.e., ship's guide and videotape) are the most valuable.

The Navy will continue its shipboard plastics educational program and extend it to shore facilities. Plans to modify the videotape and distribute it to shore facilities are ongoing. The future direction of the program will depend on the success of the initial effort and the need to modify the program.

CONCLUSIONS

The requirement for U.S. Navy ships to separate and retain plastics waste on board is so new and the educational packages were distributed so recently that any conclusions about the effectiveness of the program would be premature. However, from our experiences so far, we make the following observations:

1. Sailors generally respond positively to the plastics waste program and the educational materials. We believe the general environmental awareness and young age (average age is about 20 years) of Navy sailors are major factors affecting the positive response.
2. High quality videotapes are an effective means for communicating with shipboard audiences. However, useful guidance material about specific procedural details must accompany the videotapes.
3. Attention by the entire chain of command is needed for successful program implementation.

4. Edicts, emotional appeals, and good educational materials do not ensure successful shipboard plastics waste programs. The program must be carefully planned, coordinated, and implemented. Proposed new plastics waste requirements must be first proved feasible, effective, and safe on ships for the program to succeed.

REFERENCES

- Alig, C. S., L. Koss, T. Scarano, and F. Chitty
1990. Control of plastic wastes aboard naval ships at sea. In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. [See this document.]

HOW MUCH DO COMMERCIAL AND RECREATIONAL FISHERMEN KNOW
ABOUT MARINE DEBRIS AND ENTANGLEMENT? PHASE 1

Barbara Wallace
Kearney/Centaur Division, A. T. Kearney, Inc.
225 Reinekers Lane, Suite 300
Alexandria, Virginia 22314, U.S.A.

ABSTRACT

Education is assumed to be a cost-effective method to encourage shoreside disposal of plastics and other garbage. To test this assumption, a demonstration project is under way at four sites to develop, test, and evaluate marine debris education as a technique for changing the waste management practices of commercial fishermen and recreational boaters. The project is structured in three phases: 1) a baseline survey of commercial fishermen and recreational boaters on their garbage disposal practices and perceptions of the problems of marine debris and entanglement; 2) a targeted marine debris education program for the survey groups at three of the four sites; and 3) a survey of the same groups used in phase 1 on their garbage disposal practices and perceptions of the problems of marine debris after the education program. The project sites are: Bayou La Batre, Alabama; Martin County, Florida; Hampton, Virginia; and Taylor County, Florida. Phase 1 of the project was completed in the spring of 1989. Commercial fishermen and recreational boaters at the four sites were asked questions in six areas: 1) current garbage disposal practices, 2) experiences with plastic marine debris, 3) opinion on the problems caused by plastic marine debris, 4) knowledge of laws on at-sea garbage disposal, 5) opinion on ways to encourage shoreside disposal of plastic garbage, and 6) background information. Results of the phase 1 survey revealed the garbage disposal practices of those surveyed as well as their opinions on and experiences with plastic marine debris.

INTRODUCTION

Annex V of the International Convention for the Prevention of Pollution from Ships, 1973 (MARPOL 73/78) went into effect 31 December 1988. The MARPOL Annex V, formally entitled "Regulations for the Prevention of Pollution by Garbage from Ships," prohibits at-sea dumping of plastics and specifies the distance from shore that all other materials may be dumped. This means overboard disposal of most garbage is no longer an option.

Enforcement of MARPOL Annex V will be difficult because of competing priorities and limited resources of the Federal enforcement agencies. The success of MARPOL Annex V will depend in part on voluntary compliance.

Education is expected to be a key factor in gaining voluntary compliance with MARPOL Annex V. Education is also assumed to be a cost-effective way of encouraging shoreside disposal of plastics and other debris. To test this assumption, a demonstration project is under way to develop, test, and evaluate a marine debris education project for commercial fishermen and recreational boaters. The project is sponsored by a Saltonsall-Kennedy grant from the U.S. National Marine Fisheries Service. This paper outlines the overall structure of the project. Results of a survey of commercial fishermen and recreational boaters at four sites taken before a targeted marine debris education program are also presented.

METHODOLOGY

The project is structured in three phases: 1) a baseline survey of commercial fishermen and recreational boaters on their garbage disposal practices and perceptions of the problems of marine debris and entanglement; 2) a targeted marine debris education program for the survey groups at three of the four sites; and 3) a survey of the same groups used in phase 1 on their garbage disposal practices and perceptions of the problems of marine debris after the education program (Fig. 1). Phase 1, the initial survey, was conducted in the spring of 1989. Phase 2, the marine debris education program, will be conducted from the spring through the summer of 1989. The second and final survey, Phase 3 of the project, will be conducted in the fall of 1989.

The project is being conducted using four sites: Bayou La Batre, Alabama; Martin County, Florida; Hampton, Virginia; and Taylor County, Florida. Taylor County will serve as the control site. Commercial fishermen and recreational boaters in Taylor County will receive no marine debris education as part of this project, but will be surveyed twice. This approach was taken because some of the education activities cannot be restricted to the survey sample.

These sites were selected after consultation with state natural resource agencies. Other factors considered in the selection process included distance from other known marine debris education projects, prior involvement of the state with the marine debris issue, the ability of the state to provide comprehensive lists of commercial fishermen and recreational boaters, the number of commercial fishermen and recreational boaters, and the opportunity to coordinate marine debris education activities with local events or projects. In Martin County, for example, marine debris education activities will be tied to a fishing tournament.

The survey groups were selected from mailing lists of recreational boat registrations and commercial fishing licenses maintained by state agencies. The goal was to obtain 100 completed surveys at each site for both commercial fishermen and recreational boaters. For each site, an initial sample of 300 individuals with registered motorboats used for pleasure was selected




Figure 1.--Overview of project methodology.

using a systematic sampling technique. Surveys returned because of incorrect addresses were replaced with new names drawn using the same sampling technique. An additional sample of up to 150 names was drawn after initial survey returns indicated that fewer than 100 completed surveys would be received. All individuals on the commercial fishing license lists at each site were included in the survey. The number of commercial fishing licenses at the four sites ranged from about 200 to 330. Over 400 surveys were completed and returned--161 (16% of the potential respondents) from commercial fishermen and 257 (20% of the potential respondents) from recreational boaters. The number of completed commercial fishermen surveys ranged from 8% from Bayou La Batre to 21% from Martin County. For recreational boaters, the range was from 7% from Bayou La Batre to 27% from Hampton. None of the sites reached the goal of 100 completed survey forms, although 97 recreational boater surveys from Hampton were completed and returned. The description of the survey responses which follows represents the answers of those responding to the survey only and should not be generalized to a larger population. Unless otherwise noted, the following description of the responses combines the responses from all four sites.

DESCRIPTION OF RESPONDENTS

Commercial Fishermen

About one-quarter of the commercial fishermen responding to the survey were between 30 and 39 years of age. Another quarter of the respondents were 60 years or older. The percentage of income from commercial fishing in 1988 ranged from 0 to 100%, with the average being about 41%. The types of gears used in 1988 in order of percentage of use were as follows: rod and reel (42%), gillnet (38%), traps and pots (26%), tongs (24%), trawl (12%), dredge (4%), and longline (4%). Rod and reel were used at all sites, but most often in Martin County and Taylor County. Gillnets were also used at all sites, but most often in Hampton and Taylor Counties. Surf fishermen and spear fishermen were represented among the respondents from Martin County. Most of the commercial fishermen responding to the survey typically make day trips. However, one or more of the respondents at each of the four sites make longer trips. Some of the trips last up to 14 days at a time. About 28% of the respondents are members of a commercial fishermen's association. Only about 13% of them had participated in an organized beach cleanup in the past 3 years.

Recreational Boaters

The age distribution among the recreational boaters responding to the survey was evenly divided among three groups: 40 to 49 years, 50 to 59 years, and 60 and older. About one-quarter of the respondents fell into each group. About 40% of the respondents fish on every boat trip they make. Only 5% of the respondents never fish from their boats. About 8% of the respondents are members of fishing clubs, and about 11% had participated in an organized beach cleanup within the past 3 years.

CURRENT GARBAGE DISPOSAL PRACTICES

Commercial Fishermen

About 95% of the respondents have one or more trash receptacles on their vessels. The most prevalent types are buckets (52%) and plastic garbage bags (36%). Other receptacles used include trash cans, paper bags, and fish boxes. Only two respondents (1.3%) have compactors and one (0.6%) has an incinerator on board. Over half of the respondents said they consider what plastic items are taken on board which will need to be thrown away. Over 80% of the respondents said they have picked up plastic trash from the ocean, bay, or sound and returned it to shore for disposal. Plastic trash is generally disposed of at home (50%), at the dock or marina (27%), or at the fishhouse (20%). Fewer than 3% of the respondents admitted to disposing of plastic trash in the ocean, bay, or sound. Unwanted gear is generally disposed of at home (54%), at the dock or marina (23%), and at the fishhouse (10%). Fewer than 2% of the respondents admitted to disposing of their fishing gear in the ocean, bay, or sound. Unwanted gear is also left at city and county garbage dumps and landfills, or in dumpsters away from a dock or marina.

Recreational Boaters

Like the commercial fishermen, most recreational boaters responding to the survey have one or more trash receptacles on board their boats. The most common trash receptacles among the respondents are plastic garbage bags (51%) and buckets (43%). Among the other types of receptacles used for trash are coolers and dry and live wells. Over 56% of the respondents said they consider what plastic items will need to be thrown away when deciding what to take on board. About 75% of the respondents have picked up plastic trash from the ocean, bay, or sound and returned it to shore for disposal. Plastic trash from day trips is generally disposed of at home (73%) or at the dock or marina (23%). Fewer than 1% of the respondents said they throw plastics into the ocean, bay, or sound. On overnight trips or longer trips, the respondents generally dispose of their plastic trash at the dock or marina (56%) or at home (38%).

EXPERIENCE WITH PLASTIC MARINE DEBRIS

Commercial Fishermen

The respondents were asked first to identify which of six problems caused by plastic marine debris they had seen or experienced. They were then asked to identify the one they considered to be the most important among the six. Figure 2 summarizes the experience of the respondents with plastic marine debris. More than 95% of the commercial fishermen said they have seen plastic trash floating in the ocean, bay, or sound. This was also identified by the commercial fishermen as the most important problem among the six. About 75% of the respondents said they have seen plastic trash floating near the dock. Many of the respondents have had personal experience with plastic marine debris. Over 45% have had their vessel's propeller caught in plastic. Over 30% have had their gear caught or fouled by




Figure 2. -- Experience with plastic marine debris (based on percent of responses to the question).

plastic debris. Almost 40% have had their boat's cooling intake systems clogged by plastic debris.

About 94% of the respondents said they think plastic trash in the ocean, bay, or sound can kill marine animals, create safety hazards for mariners, and wash ashore as beach litter. Personal experience was cited most frequently as the basis for their opinion.

Recreational Boaters

Figure 2 also summarizes the experience of the recreational boaters with plastic marine debris. About 90% of the respondents said they have seen plastic trash floating in the ocean, bay, or sound. This was also cited most frequently as the most important problem. The recreational boaters responding to the survey have also had personal experience with plastic marine debris, but not to the extent of the commercial fishermen. About 28% of the respondents have had their boat's propeller caught in plastic debris. Between 15 and 20% have had their gear caught or fouled by plastic debris. About 21% of the respondents said that their boat's cooling intake systems have been clogged by plastic marine debris.

Almost all of the respondents (97%) said they think that plastic trash in the ocean, bay, or sound can kill marine animals, create safety hazards for mariners, and wash ashore as beach litter. Like the commercial fishermen, personal experience was cited most frequently as the basis of the respondents' opinion.

KNOWLEDGE OF LAWS ON AT-SEA GARBAGE DISPOSAL

Commercial Fishermen

Just over half (51%) of the commercial fishermen said they know there is a Federal law which prohibits disposal of plastics from vessels and restricts the other types of garbage that may be dumped into the ocean, bays, or sounds. The remainder of the respondents were evenly split between those who were unsure and those who did not know about such a law. Word of mouth was cited most frequently as the source of information on this law, followed by magazine articles, television, and newspapers. Among the sources of information on the law specified by the respondents were training by offshore oil and gas companies for their workers and tickets received from the U.S. Coast Guard or marine patrol. Only 5% of the respondents said they have heard of MARPOL Annex V.

Recreational Boaters

Thirty-eight percent of the respondents said they know there is a Federal law which prohibits disposal of plastic trash from boats and limits the dumping of other types of garbage into the ocean, bays, and sounds. About 33% of the respondents were unsure whether there is such a law, and 30% did not know about the law. Like the commercial fishermen, word of mouth was cited most frequently as the source of the respondents' information. This was followed by television, newspapers, and magazine articles.




Figure 3.--Best ways to encourage shoreside disposal (based on percent of responses to the question).

The Coast Guard, marine patrol, and training by offshore oil and gas companies were also cited as sources of information on this law. Only 3% of the respondents said they have heard of MARPOL Annex V.

OPINION ON THE BEST WAYS TO ENCOURAGE SHORESIDE DISPOSAL OF PLASTIC TRASH

Commercial Fishermen

Commercial fishermen were asked which of seven techniques are the best ways to encourage commercial fishermen to return their plastic trash to shore for disposal. A summary of their responses is shown in Figure 3. Dockside disposal facilities and fines were the most frequent responses. There was some support for each of the techniques. Among the suggestions from the respondents were reminders on the marine radio channel, incorporation of information on proper garbage disposal into the licensing procedure, and use of advertisements and stickers.

Recreational Boaters

Figure 3 summarizes the responses of recreational boaters on which of seven techniques are the best ways to encourage recreational boaters to return their plastic trash to shore for disposal. Like the commercial fishermen, there was some support for each of the techniques, but dockside disposal facilities and fines were cited most often. The recreational boaters also suggested the following techniques: television advertisements and programs, educational material sent with license renewals, information broadcast on the marine weather channel, and peer pressure.

RESULTS

The results of this survey provide a baseline of knowledge on the garbage disposal practices of commercial fishermen and recreational boaters at four sites, and on their opinions and experiences with plastic marine debris. The results indicate that the problem of marine debris is not new to those responding to the survey. Further, the results show there are responsible commercial fishermen and recreational boaters. From the many comments written in the margins and on the backs of the survey form, it is also evident that many of the respondents think the source of marine debris is a group other than their own.

THE OREGON EXPERIENCE--FOUR YEARS LATER

Judie Neilson*
Oregon Department of Fish and Wildlife
Portland, Oregon 97207, U.S.A.

*Present name and address: Judie Hansen, 1502 S. County Road 525W,
Danville, IN 46122.

ABSTRACT

The fifth annual Oregon coastwide cleanup, held 10 October 1988, attracted 2,200 volunteers who collected 14.2 tons of debris. Similar fall cleanups were held in 22 coastal states of the United States, Costa Rica, and Puerto Rico. This paper reviews personal observations about the effectiveness of volunteer beach cleanups and discusses the evolution of data gathering, media coverage of the marine debris problem, changes in attitudes, and advancement in plastic recycling.

I am pleased to be here to share my knowledge on my favorite subject-- "floatable trash." I became interested in this subject just 5 years ago. At the time, it was a challenge to find information on the subject at all. Let me tell you how I became involved.

In 1984, the May-June issue of the Alaska Fish and Game Department's magazine was delivered to my office by mistake. Flipping through it, I was drawn to an article by free-lance writer Tom Paul. Entitled "The plague of plastics," it discussed the proliferation of plastic debris in the natural environment and the resulting ingestion and entanglement by fish and wildlife.

At the time, I had worked at the Oregon Department of Fish and Wildlife for 10 years and been an active birdwatcher for 25 years. I knew birds got entangled in monofilament fishing line and six-pack beverage rings, but I didn't know they had an appetite for polystyrene foam and small bits of plastic.

In talking to birdwatchers, scientists, and friends, I realized others were also unaware of the problem. Since 1984 was the "Year of the Ocean" and we Oregonians love our coast, I had the idea to organize a cleanup of plastic debris on our 563 km (350 mi) of coast to see what we could find.

I put together a small working group; we divided our coast into 14 zones and found people willing to serve as "zone captains." The captains were biologists from the Department of Fish and Wildlife and local residents of the coastal communities. Zone captains coordinate refreshments for celebration parties, assign volunteers to specific stretches of beach, arrange for the pickup of the collected debris, and work with the news media.

We spread the word of that first cleanup and invited the public to the coast to see how much marine debris we could collect on the 241 km (150 mi) of accessible beach.

Saturday, 13 October, was a very cold, wet, dark, blustery day. To my amazement, 2,100 volunteers showed up and collected 26.3 tons of debris in just 3 h. They filled out questionnaires documenting the quantity of fishing gear, six-pack yokes, polystyrene foam, plastic bags and bottles, rope, and strapping bands. We recorded the event on video film and produced a 12-min film entitled "Get the Drift."

Word of the cleanup spread quickly, and that November I was invited to report the results at the Workshop on the Fate and Impact of Marine Debris in Honolulu, Hawaii. Those attending were government scientists and concerned citizens from around the world working on marine mammal entanglement problems. I felt like an imposter with my citizen involvement project results, but the scientists welcomed me with open arms. They came up to me afterwards and thanked me for documenting the volume of trash in a large given area. Many had wanted to do beach surveys for years but were unable to spend time and money on that kind of research.

A number of recommendations came out of the workshop, and one was that beach cleanups are a valid way to document the amount and sources of marine debris. As a result, I was asked to organize and report on the findings of cleanups along the west coast and New England states in 1985.

I prepared the "Nuts and Bolts Guide to Organizing a Beach Cleanup the Easy Way." A "Dear Coastal Colleague" letter was mailed to over 200 organizations and government entities listed in the National Wildlife Federation Conservation Directory. They were asked to take an active part in organizing a cleanup in the study area states. Firm commitments were received from eight states.

State coordinators were mailed a copy of the Nuts and Bolts Guide as a "starter kit" but were encouraged to use special creativity to organize the cleanups. Some interesting logos, posters, and mottos resulted: "Lend a hand in the sand," "Don't be a litter boat," "Be a beach buddy," and "Debris-a-thon," to name a few. The main focus of the national cleanup was to determine the amount of derelict fishing gear, both sport and commercial, which makes its way to the coastal beaches, and to help educate the public.

Following the cleanups, I compiled the results in a report to the U.S. National Marine Fisheries Service.

The coastal cleanup program has grown by leaps and bounds. In 1986, we had 14 states participating. In 1987, there were 19 states. During September and October of 1988, all 22 coastal states, 7 inland states, and Costa Rica and Puerto Rico participated. All of the cleanups were held during Coastweeks, produced by a citizens' network of groups, agencies, and individuals who focus attention on that special place where water meets the land. During the period of Coastweeks, agencies and organizations in coastal states have beach walks, bird identification seminars, beach cleanups, and various activities to call attention to coastal issues. Five years ago it was simply Coastweek, but to accommodate the many states participating, with different weather and tide patterns, the annual campaign now stretches from the middle of September to the middle of October.

So after five cleanups, what are my thoughts about this whole business? I feel the number one value of beach cleanups is raising public awareness. Almost to a person the volunteers remark, "I never realized how much stuff was out there until I had to spend time leaning over to pick it up." And it sticks with them when they go back home. One friend told me that after working on a beach cleanup he couldn't enjoy playing golf because he kept seeing all the polystyrene cups in the ponds on the golf course. In areas where beach cleanups have occurred, government agencies responsible for monitoring trash containers indicate an increase in the amount of plastic debris which is disposed of properly.

Each year, our data gathering gets a little more sophisticated. As you might imagine, turning loose thousands of volunteers for 3 h with no "form-filling-out training" doesn't result in precise accounting of specific materials or number of pieces. But it does give an index of the type of debris and the probable source. The first 4 years we had a very general, short questionnaire. It gave us bulk figures, because volunteers would write "some," "many," "lots," "a few," under number of pieces. In 1988, we worked with the Center for Marine Conservation and used the questionnaire and guide which was used by approximately 43,000 volunteers nationally. The new questionnaire is more complex and specific. Prior to the cleanups, there was virtually no documentation on the amount or source of marine debris. So we have come a long way!

Our volunteers show up to work, are given a large collection bag, questionnaire, reminders about beach safety, and turned loose. Three hours later they come back laden with trash, enthusiasm, and stories of the weird things they found. We treat them to a free lunch to give them an opportunity to share their stories.

In years past, all of the debris from Oregon's cleanup went directly into landfills, thanks to the generosity of the Oregon Sanitary Service Institute. This year we introduced a "beach buddy" system. We asked volunteers to work in pairs and separate the plastic from other debris and place it in a special bag. After the cleanup, all the plastic and polystyrene foam was picked up by Environmental Pacific Corporation, taken to Portland, and analyzed to see how much of it could be recycled. We invited the press to watch us rip open the sacks, not knowing for sure what we would find. Much to my relief, all of it was plastic and most was

recyclable. On the minus side, I think we asked too much of the volunteers. It is not possible for one person to carry two sacks, a clipboard, writing tools, and also to pick up debris. So the system still needs refinement.

Many of our volunteers have participated in all five cleanups, and we are beginning to see more groups who charter buses for the trip from inland cities to the coast. Private industries sponsor employee trips, schools use the cleanup as an official school function, and civic groups organize carpooling and their own potluck picnics on the cleanup day.

Going after marine debris as "litter" on the beaches does not have the same public appeal as focusing on the issue of entanglement and ingestion by fish and wildlife. That focus has attracted the media and gotten new people interested and involved. I am sure everyone in this room knows the risk in getting people stirred up and emotionally involved. All of a sudden, the statistics you gather are used in very creative ways to prove a point on all sides of the marine debris and plastic recycling issues.

One of my earlier recommendations was to get a media blitz in the popular press, not just in obscure technical or professional journals. I am pleased to report there is hardly a week goes by that I don't run across a marine debris article in a commercial fishing industry magazine, conservation organization newsletter, or the newspaper. The state natural resource agency magazines and Sea Grant publications have also done an excellent job to further document the problem through feature articles complete with color photographs of injured wildlife.

I hope this trend continues. There should be repeated articles in newspapers, not just in the outdoor section but in business and science. Even the special newspaper supplements available to tourists along the coast should have articles on marine debris. The amount of trash on beaches has an adverse economic impact for coastal communities and states competing for tourism dollars. That fact was brought home to us in 1988 with the hospital waste showing up on New York and New Jersey beaches.

The publications which have not picked up on the severity of marine debris are those targeted for the sport fishing and recreational boating public. Because of the way licenses and permits are issued, recreationists have been missed by traditional Federal agency notices. We have found a larger percentage of bait containers and recreational gear during our cleanup since commercial vessel owners are better informed. For instance, I don't think the recreational fleet knows about the adoption of Annex V to the International Convention for the Prevention of Pollution from Ships (MARPOL).

A major value of the cleanups is networking with people in coastal states working on marine debris. The networking provides a vehicle to communicate findings and the status of state and Federal legislation; compare how the cleanups are organized; and share artwork, slogans and campaign strategies, what works with the media, and how to get donated materials or funding. Having the cleanups clustered during 1 month in the fall gives everyone higher visibility with the public and news media.

Now that we have the new questionnaire, we can get a better handle on how to identify items that cause the majority of entanglement or ingestion problems and focus our efforts on their source.

One of the most exciting things to come along is the pilot project conducted by the Port of Newport here in Oregon. They set up a recycling program for commercial fishermen and other marine users to separate their trash at the dock. They gave the wood to senior citizens and sold the scrap paper, glass, and metal, and only a small portion was left over for the landfill. The net fragments were recycled by tourists wanting decorations for their patios or local citizens needing supports for their vegetables or backstops for their softball fields. That 1-year program gained support and energy from a small commercial fishing community which has radiated enthusiasm and interest to the entire Pacific Rim fishing industry. Its coordinator, Fran Recht, has a new grant to implement similar projects in Alaska, California, Oregon, and Washington this year and continue educational programs with the commercial fishing industry and port officials. As long as ocean users continue to dump trash overboard, beach cleanups have a transitory value in ridding the beaches of debris. The Newport project has directly reduced the amount of debris on Oregon's central coast.

The adoption of Annex V to MARPOL has provided strong incentives for improved port facilities and less dumping at sea. I suspect it will foster accelerated plastic recycling programs.

In Oregon we have good news. On 8 October 1988, we attracted 2,200 volunteers. But they were only able to collect 14.2 tons compared to the 26.3 tons in 1984. Each year we have seen a steady decline in the amount collected, and there are several contributing factors. Our weather has been mild each fall with no major storms depositing new trash before the cleanup. Luck of the currents, no doubt. Also, the Oregon State Parks Department has held a "Company's Coming" cleanup in the spring for the past 3 years, so we didn't have an entire year's accumulation. As a general rule, beach users are carrying out their own trash and debris they see washed on shore, and on Oregon's central coast, adjacent to the Port of Newport pilot project, there was simply less debris available. The increased public awareness through the cleanups has made a big difference in Oregon.

As an extension to the regularly scheduled cleanups, we launched an "adopt-a-beach" program patterned after the State of Texas, inviting Oregonians to choose a section of the Oregon coast they want to "adopt." All we ask is that they clean it three times a year and tell us what they find.

I am really pleased to visit with you and share my enthusiasm about how one person's idea can make a change. Since being involved in cleaning beaches, I am better informed about entanglement and ingestion by wildlife, environmental monitoring using citizen volunteers, how plastic is made and recycled, and best of all how valuable trash can be.

Thank you.

THE TEXAS ADOPT-A-BEACH PROGRAM: A
PUBLIC/PRIVATE APPROACH TO CLEAN BEACHES

Angela D. Farias
Texas General Land Office
1700 N. Congress Avenue
Austin, Texas 78701, U.S.A.

ABSTRACT

The Texas Adopt-A-Beach Program, founded in 1986 by Texas Land Commissioner Garry Mauro, is made up of private citizens who volunteer to clean Texas beaches three times a year, and is managed by the Texas General Land Office. The program was conceived both as a short-term solution to the problem of trash on Texas beaches and as a means of advancing long-term, permanent solutions to the problem of marine debris: the ratification of Annex V of MARPOL, the International Convention for the Prevention of Pollution from Ships, and the implementation of effective enforcement measures.

The Adopt-A-Beach Program organizes two coastwide beach cleanups a year. The success of these cleanups--nearly 25,000 volunteers have removed 1,000 tons of garbage from Texas beaches--has enabled the program to shift its emphasis to public education about the tremendous economic and environmental damage caused by ocean dumping of garbage. Children are the focus of educational projects that include a coloring book and puppet show featuring the program's mascot, Lucky the Dolphin, and the development of a marine debris curriculum for public schools. The program has also produced videotaped public service announcements for statewide broadcast and has sponsored special awareness events in conjunction with cleanups.

Data collected by volunteers during cleanups were instrumental in building support in Congress for Senate ratification of the international MARPOL Annex V and implementation legislation for the annex in the United States. Annex V prohibits the dumping of plastics in the ocean and sets strict limits on dumping of other solid wastes overboard. The volunteer-collected data are also being used in support of making the Gulf of Mexico a special area under the annex. Dumping of any kind is prohibited in special area waters.

INTRODUCTION

Texas Land Commissioner Garry Mauro founded the Texas Adopt-A-Beach Program in 1986 after taking part in a coastwide beach cleanup organized by the Center for Environmental Education (now the Center for Marine Conservation). In the course of this 3-h event, 2,700 volunteers collected 124 tons of trash from 196 km (122 mi) of beach. It was clear that this volume of trash could not be attributed to beach littering alone; the quantity and the nature of the debris indicated that most of it had washed ashore in tides of waterborne refuse.

Astonished at the quantity of trash on Texas beaches, and recognizing the heavy environmental and economic penalties of this pollution, Commissioner Mauro decided to investigate the problem and determine what role the General Land Office could play in solving it. He began by forming an in-house task force to study the sources of beach debris, to research existing legal and institutional remedies, and to map out an action plan.

Both a 1985 Texas Coastal and Marine Council report and the Center for Environmental Education's report on the 1986 beach cleanup concluded that some 75 to 90% of the trash on Texas beaches originates offshore, about two-thirds of it dumped from ships. Refuse discarded in the Gulf of Mexico is not washed out to sea. What fails to sink is held in gulf waters by loop currents until oblique alongshore currents carry it to shore. For this reason, the beach cleaning efforts of cities and counties along the Texas coast have provided only temporary relief, and at great cost: the State's coastal communities spend about \$14 million each year on the endless task of cleaning their beaches.

The Texas Approach

Two needs were immediately apparent: the need to raise public awareness of the magnitude of the beach garbage problem in Texas, and the need for a broad-based, unified approach to its solution, concentrating on the sources of beach debris.

An obvious objective was to work for U.S. ratification of Annex V to MARPOL 73/78 to prohibit ocean dumping of plastic and restrict the discharge of other types of solid waste at sea. The severity of the beach pollution problem in Texas led to a second: to persuade the International Maritime Organization to designate the Wider Caribbean Region (the Gulf of Mexico and the Caribbean Sea) as a special area where virtually all dumping would be banned. But the General Land Office began with state-level action that would yield positive results more quickly.

As manager of the state's surface and mineral interests in about 1.62 million ha (4 million acres) of submerged land on the Texas gulf coast, the General Land Office issues leases, easements, and permits for a variety of activities. The agency's first step was to adopt emergency rules prohibiting the dumping of solid waste from offshore platforms and seismic vessels operating in Texas waters under state permits. These were later followed by no-dumping rules (and parallel contract provisions) for marinas, wharves, piers, fishing cabins, and all other structures on state-owned coastal land.

Next, to draw public attention to the amount of trash and garbage accumulating on Texas beaches, to augment existing beach cleanup efforts, and to involve citizens in the crusade against marine debris, the General Land Office instituted the Texas Adopt-A-Beach Program.

The program was modeled after the Adopt-A-Highway Program organized by the Texas Department of Highways and Public Transportation. The Adopt-A-Highway Program has proven highly successful both as a cleanup program and as a public awareness campaign to discourage roadside littering. It was felt that the adoption format, already familiar to Texans, would work equally well for the state's beaches. The Highway Department's slogan, "Don't Mess with Texas," was expanded to "Don't Mess With Texas Beaches" for the beach cleanup campaign.

Response to the Program

The Adopt-A-Beach Program won immediate enthusiastic support in Texas. Adopters were secured for all 172 easily accessible Texas beach miles within the first year of the program, and even some segments accessible only by boat or four-wheel-drive vehicle found sponsors. Most adopting groups have renewed their adoption agreements annually. Businesses, philanthropic foundations, entertainers, advertisers, and private citizens have made generous contributions of funding, supplies, services, and promotional assistance.

Since the program's first coastwide cleanup in April 1987, nearly 25,000 volunteers have removed more than 1,000 tons of trash from Texas beaches. An analysis of data collected during coastwide cleanups, prepared by the (then) Center for Environmental Education, was presented to the U.S. Congress and to the International Maritime Organization as evidence documenting the need for U.S. ratification of MARPOL Annex V, the passage of national enforcement legislation, and designation of the Wider Caribbean Region as a special area under the annex.

The spirit of the Adopt-A-Beach Program has spread beyond state boundaries and beyond U.S. borders. The Texas program has been emulated by other coastal states, and it is now spreading to Central America. In September 1988, Texas, Louisiana, Mississippi, Alabama, and Florida joined forces in a "Take Pride Gulfwide" beach cleanup sponsored by the U.S. Minerals Management Service. A symbolic beach adoption agreement entered into by the students of Flour Bluff Junior High School near Corpus Christi, Texas, and the children of Costa Rica in the spring of 1988 immediately led to the establishment of a national Costa Rican beach cleanup program.

PROGRAM STRUCTURE AND OPERATION

The Adopt-A-Beach Program takes its direction from State Government, but it is operated at the county level by an all-volunteer work force and is dependent upon private funding and in-kind donations of supplies and services.

An Adopt-A-Beach Task Force appointed by Commissioner Mauro developed guidelines for the program and oversees its operations. This advisory

body, whose members now number about 55, represents a broad range of coastal interests: Federal, state, and local government; oil, gas, and chemical production; tourism; shipping; agriculture; waste disposal and recycling; scientific research; and conservation. The task force meets periodically in full session for program review and planning. Three subcommittees--Finance, Education, and Legislation--hold independent meetings.

The General Land Office, as administrator of the Adopt-A-Beach Program, coordinates all program activities with the assistance of the Adopt-A-Beach task force. It oversees beach adoptions, promotes the program statewide, solicits funding, organizes two annual coastwide cleanups, and develops educational materials and programs. The agency also maintains a toll-free number for in-state inquiries about the program and publishes a quarterly newsletter, the Texas Beach Bulletin,

The Texas Conservation Foundation, a state agency empowered to manage and expend donated funds, is financial trustee for the Adopt-A-Beach Program. It manages a special fund established to receive tax-deductible contributions for the support of program activities. Monies from the fund are used for:

- the purchase and shipment of cleanup supplies, including trash bags, data cards, and pencils;
- the purchase and installation of beach signs marking adopted beach segments and crediting adopting groups;
- printing and mailing of the program's newsletter, certificates, posters, brochures, and other promotional and educational materials;
- operation of the program's toll-free telephone line; and
- promotional events to publicize the program and to recruit sponsors and cleanup volunteers.

A network of volunteer county coordinators provides grassroots leadership for the program. They recruit and register adopting groups, handle local cleanup logistics, and promote the program within their communities. This structure is not only practical, but also capitalizes on community pride--a powerful force in sustaining the program's momentum.

To join the program, groups sign an adoption agreement that commits them to cleaning a designated beach segment (usually 1.6 km (1 mi)) three times within a 1-year period, participating in the program's two coastwide cleanups and conducting a third cleanup independently. The agreement also releases the Adopt-A-Beach Program from liability for any injury incurred during a beach cleanup. Each adopting group receives an adoption certificate, and the group's name is listed on a sign installed at the access road nearest its adopted beach segment.

Groups now enrolled in the program include civic clubs, sporting clubs, chambers of commerce, large corporations, small businesses, conservation organizations, public schools, colleges, Scout troops, state agencies, cities, property owners' associations, and families.

The program's two annual coastwide cleanups are cosponsored by nonprofit organizations. The "Great Texas Beach Trash-Off," held in April of each year, is cosponsored by Keep Texas Beautiful, Inc. The "Texas Coastal Cleanup," held in September during Coastweeks, is cosponsored by the Center for Marine Conservation. Participants in these cleanups include independent volunteers as well as affiliates of the Adopt-A-Beach Program. Local coordinators distribute trash bags, data cards, and pencils to all cleanup volunteers.

The data cards provide spaces for the tabulation of items within seven broad categories (plastic, glass, Styrofoam, metal, paper, wood, and rubber), for the notation of labels that might indicate the sources of items collected, for recording the number of trash bags filled, and for reporting sightings of stranded or entangled animals.

PROGRAM PROMOTION

The Adopt-A-Beach Program employs a variety of means to publicize the adoption program, advertise coastwide cleanups, and recruit volunteers and sponsors. The program carries its promotional efforts statewide to remind noncoastal Texans that they are residents of a coastal state, that they benefit from the coastal area both directly and indirectly, and that they should share the responsibility for protection of coastal waters and beaches.

Recruitment

When the program was first organized, the General Land Office recruited county coordinators through telephone calls, letters, and personal visits to community leaders and known environmentalists along the coast. The same procedures were used by the General Land Office Adopt-A-Beach staff and county coordinators to enlist adopters. Lists of prospective adopters were compiled from target groups such as garden clubs, 4-H clubs, local branches of oil and gas companies, and waste disposal companies.

Early recruitment was greatly facilitated by television and newspaper coverage of press conferences preceding the program's first coastwide beach cleanup in the spring of 1987 and by establishment of the program's toll-free telephone number (1-800-85-BEACH). Now that the program is well known, it has become less necessary to engage in aggressive recruitment of participants. New county coordinators, new adopting groups, and sponsors often initiate contact with the program.

Solicitation of Financial and In-Kind Support

All public information materials produced by the Adopt-A-Beach Program emphasize that the program is dependent upon private donations for its

operation and that such donations are tax-deductible. The program solicits contributions of money, supplies, or services through direct appeals to companies that have a reputation for supporting environmental causes or that are likely to appreciate the benefits of association with an environmental improvement program. It has also sought grants from foundations known to support conservation efforts.

Financial donations to the Adopt-A-Beach Program have included a 3-year grant of \$50,000 from the Moody Foundation of Galveston, a \$5,000 grant from the Fondren Foundation of Houston, and a contribution of \$10,500 from Browning-Ferris Industries, a waste disposal company. Other contributions have come from oil and gas companies, law firms, and numerous private individuals.

In-kind donations to the program have included bumper stickers, trash bags, celebrity promotions, refreshments for cleanup participants, and pickup of filled trash bags after beach cleanups. The Mobil Corporation donated 100,000 garbage bags valued at \$13,000 to the program in 1988, and Maryland Club Foods donated 9,700 bags for the 1989 Great Texas Beach Trash-Off. One of the first in-kind donations to the program was 15,000 "Don't Mess With Texas Beaches" bumper stickers, provided by a beer distributor.

In 1988, 23 outdoor advertising companies donated space along major routes to the coast for 300 billboards displaying the "Don't Mess With Texas Beaches" slogan and the program's toll-free number. Most of the billboards were printed with money from the Adopt-A-Beach fund and posted by companies that donated the advertising space, but some companies painted billboards free-of-charge.

Media Coverage

The Texas Adopt-A-Beach Program has benefited from both local and national media attention. After the April 1987 coastwide cleanup, articles appeared in the New York Times and the Los Angeles Times, and both Time and Newsweek magazines featured articles about the status of beaches in America as front-cover stories. In August 1987, Commissioner Mauro discussed the problem of beach trash in Texas as a guest on the ABC television news program Good Morning, America.

Press coverage of program activities is invaluable in advertising the program, mustering volunteers for coastwide cleanups, and publicizing cleanup results. Press releases are issued to newspapers, radio stations, and television stations in advance of cleanups and as soon as statistics on the number of volunteers participating and the tons of trash collected are available afterward.

The adoption of a mascot, an Atlantic bottlenose dolphin named Lucky, enlivened the Adopt-A-Beach Program's promotional campaign and gave the program a symbol with appeal for children. Lucky, a performer at Sea-Arama Marineworld in Galveston, Texas, is an especially appropriate mascot for the program because he was a victim of marine debris, barely surviving

entanglement in an abandoned fishing net. He was given his fitting name by the veterinarians of the Marine Mammal Stranding Network who nursed him back to health.

Public service announcements are produced for radio and television broadcast before every coastwide beach cleanup. The program's first videotaped public service announcement featured Lucky and Texas rock musician Joe "King" Carrasco. The second video, featuring actor Randy Quaid, was produced to advertise the 1989 Great Texas Beach Trash-Off. A third was produced to publicize the MARPOL Annex V regulations. All public service announcements are distributed statewide.

Printed Materials

The program produces brochures and posters to advertise coastwide cleanups. These are sent to county coordinators, adopting groups, coastal chambers of commerce, public libraries, hotels and motels, and other businesses for distribution. Cleanup brochures list the names and telephone numbers of county coordinators, the locations of designated beach check-in points for volunteers, and the names of hotels and motels offering discounts to cleanup workers.

The quarterly Texas Beach Bulletin, with a current circulation of about 3,300, summarizes Adopt-A-Beach Program activities, announces cleanup dates, reports results of coastwide cleanups, and acknowledges donations to the program. It serves as an educational as well as a promotional tool, containing articles about national and international efforts to combat marine debris. The newsletter is sent to Adopt-A-Beach Program participants and supporters and is included in the information packet mailed to anyone who calls or writes to inquire about the program.

PUBLIC EDUCATION

With the beach cleanup program well established in all coastal counties, the Adopt-A-Beach Program is concentrating on public education projects, including educational materials and programs for children, an awareness drive targeting recreational boaters, and a campaign to promote recycling. An important element of all these efforts is publicizing the requirements and expected results of MARPOL Annex V.

Outreach to Children

Children are the primary target of educational efforts. In the summer of 1988, the Adopt-A-Beach Program introduced an educational program for preschool and primary-grade children. Its components are a slide show, a puppet show, and a "Don't Mess With Texas Beaches" coloring book.

The slide show contrasts clean and littered beaches, shows how trash reaches the beach and how it can harm birds and marine animals, and tells the story of Lucky the Dolphin. In the puppet show, called "Joey Saves the Day," a boy fishing from a boat rescues Lucky and his friend Clipper the Crab from the Trash Monster. In the 11-page coloring book, Lucky points

out the hazards of beach litter and floating trash and urges children to help keep beaches clean. Simple lyrics for a series of beach cleanup songs set to familiar tunes are printed in the back of the book.

The educational program has been presented to an estimated 4,000 children at day-care centers, public libraries, museums, and elementary schools in Austin and coastal cities and has proven very effective in interesting children in the cause of beach protection. Because the puppet show has been so well received by teachers, librarians, and children, it is being videotaped for distribution throughout the state. Coloring books have been given to all children attending the program, and at least 20,000 more have been distributed by mail. Five hundred Spanish-language versions of the coloring book were sent to children in Costa Rica.

The Adopt-A-Beach Program is now working on a formal marine debris curriculum for kindergartens and elementary schools. It will include lessons about the nature and importance of marine and coastal resources; the damage caused by marine debris, particularly plastics; and recycling as a solution to the problem of solid waste in the environment.

The program staff is preparing a curriculum outline in consultation with the Education Subcommittee of the Adopt-A-Beach Task Force. To ensure that the curriculum meets state requirements for public school use, the staff is also coordinating development of the project with the Texas Education Agency. It is anticipated that the actual writing of the curriculum will be contracted to an educational consultant.

Promoting Awareness Among Recreational Boaters

The Adopt-A-Beach Program is working with the Boating Trades Association of Texas and the Marina Association of Texas to make boaters aware of the need for their cooperation in the battle against marine debris. The program's first public service video was directed at recreational fishermen and boaters, urging them not to discard trash overboard. The Boating Trades Association and the Marina Association have distributed "Stow It-- Don't Throw It" bumper stickers that include the Adopt-A-Beach Program and Center for Environmental Education logos.

Recycling Campaign

The Adopt-A-Beach Program is broadening its mission by promoting recycling as a practical means to reduce solid waste in the environment and alleviate the burden on landfills. The focus of the recycling campaign is plastic, which makes up some 60% of the trash collected in coastwide beach cleanups.

General Land Office staff helped draft container-coding legislation for introduction in the 1989 session of the Texas Legislature. Information about the proposed law, which would require coding of plastic containers by resin type to facilitate recycling, has been published in the program's newsletter.

In conjunction with the April 1988 coastwide cleanup, the Society for the Plastics Industry sponsored a plastics recycling demonstration for cleanup volunteers at South Padre Island. In September 1988, Shell Oil Company sponsored a "Trash Bash" for volunteers who brought glass, paper, aluminum, and plastic to Sea-Arama Marineworld in Galveston for pickup by recycling companies.

To further encourage beach cleanup volunteers to become recyclers, and to take advantage of the opportunity to recycle large amounts of beach debris, participants in the 1989 Great Texas Beach Trash-Off were asked to separate the trash they picked up from the beach, putting recyclable materials into orange bags, and nonrecyclable trash into white bags. Keep Texas Beautiful, Inc., cosponsor of the cleanup, made arrangements for collection of the recyclable trash by local companies.

In conjunction with the 1 April 1989 Great Texas Beach Trash-Off, the Texas Arts Council and Business Volunteers for the Arts/Houston joined the Adopt-A-Beach Program in sponsoring a juried beach trash sculpture contest in Galveston. The contest was conceived as a means of publicizing the sources and types of debris found on Texas beaches, drawing attention to the program's cleanup and recycling campaigns, and adding the arts community to interests endorsing the beach cleanup program.

A brochure containing contest guidelines and an entry form was distributed to members of the Texas Arts Council and to artists on a mailing list supplied by the Texas Commission for the Arts. Posters advertising the contest, called "Trash for Art's Sake," were sent to art museums and galleries. Cash prizes were awarded to the top three winners, and other contestants received honorable mention. Ten prize-winning entries will tour art museums throughout the state before being donated to coastal museums for permanent display.

FUTURE OF THE PROGRAM

Special Area Designation

Data collected by Texas volunteers in future beach cleanups will permit evaluation of the effectiveness of MARPOL Annex V regulations in reducing the amount of plastic and other floating debris reaching Texas shores. The data will also support the Texas Adopt-A-Beach Program's ongoing effort to secure special area designation for the Wider Caribbean Region under the annex.

The Adopt-A-Beach Program will continue to encourage the establishment of parallel beach cleanup programs throughout the Caribbean region. The success of the Costa Rican beach cleanup program has inspired both Panama and Honduras to establish similar programs, and it is hoped that other countries will follow suit. The demonstration of widespread, serious concern about marine debris in the Wider Caribbean should benefit the cause of special area designation.

Continuing Educational and Awareness Projects

The program will continue to stress the importance of recycling as a solution to the solid waste problem and will work to promote the expansion of recycling efforts throughout the state. The Adopt-A-Beach staff is cooperating with the Bryan and College Station, Texas, independent school district in a program that will combine classroom education with an active recycling project for public school students next year.

An expanded awareness campaign will be undertaken to educate recreational boaters about the hazards of marine debris, to inform them about MARPOL Annex V regulations, and to encourage all marinas on the Texas coast to provide garbage reception facilities like those the General Land Office now requires for marinas on state-owned land. The Adopt-A-Beach Program will supply boaters with garbage bags and award certificates of appreciation to volunteers who pledge to participate in an aquatic version of the beach cleanup program.

Research and Planning

The Adopt-A-Beach Program is helping Texas ports prepare to contend with the plastic refuse that will be off-loaded by ships in accordance with MARPOL Annex V regulations. The program is helping the ports locate researchers, waste management companies, and recycling companies that can supply needed planning assistance, equipment, and services.

The Adopt-A-Beach Program is also represented on the Marine Debris Technical Subcommittee of the U.S. Environmental Protection Agency's Gulf of Mexico Program, assisting with data collection, data analysis, and exploration of methods to alleviate both marine and beach debris throughout the gulf region.

Addressing Land-Based Sources of Beach Debris

The reduction--and, it is hoped, eventual elimination--of marine debris in the Gulf of Mexico will not entirely solve the problem of solid waste pollution of Texas beaches. Though offshore sources are to be blamed for most of the trash fouling the state's shoreline and nearshore waters, land-based sources make a substantial contribution. For this reason, the utility of the Adopt-A-Beach Program will not soon be diminished.

Land-based sources--onshore dumping, river-transported trash, and persistent littering--will be the target of future Adopt-A-Beach Program awareness efforts. Data collected by beach cleanup volunteers can provide estimates of the amount of beach trash attributable to these sources and can help to identify chief offenders.

CONCLUSION

The rapid success of the Adopt-A-Beach Program in establishing a strong alliance between government and the private sector in the crusade against marine debris, in stirring citizen activism both within and beyond

Texas, and in becoming a catalyst for the statewide expansion of solid waste control efforts exceeded 1986 expectations. The program's influence can be largely attributed to timing: it was inaugurated at a time when Americans had at last begun to realize that pollution of coastal waters and shorelines had reached a critical stage.

But it cannot be assumed that no one among the volunteers who rallied to the cry "Don't Mess With Texas Beaches" would have been willing to participate in a coastal cleanup effort years earlier. In fact, many program volunteers were already participants in, or supporters of, other conservation efforts. Those who were not may just have been waiting for direction.

The Adopt-A-Beach Program provided that direction. It has succeeded because it offers an easy avenue for citizen participation, enabling citizens from all walks of life to make an important contribution to the coastal cleanup effort. Those who cannot provide hands-on assistance at the beach can support the program by donating money, supplies, or services. Those who can offer neither physical nor financial support for program activities can be of equal help by simply spreading the word--by helping to heighten public awareness of the environmental and economic costs of marine debris. The program's design has facilitated development of the broad base of informed support essential to genuine progress against so widespread and complex a problem.

MARINE DEBRIS DEMONSTRATION AND EDUCATION PROJECT
AT SQUALICUM HARBOR, BELLINGHAM, WASHINGTON, U.S.A.

Jim Humphreys* and Patti Mullin**
Washington Sea Grant Program
Bellingham, Washington 98225, U.S.A.

*Present address: Bornstein Seafoods, P.O. Box 188, Bellingham, WA 98227.
**Present address: 4408 Y Road, Bellingham, WA 98226.

ABSTRACT

Washington Sea Grant's North Puget Sound Office in Bellingham, Washington, began a demonstration and education project about marine debris in January 1988. The objectives of this project were:

1. to develop a demonstration project to collect and recycle vessel-generated wastes from commercial and recreational vessels at Squalicum Harbor, Bellingham, Washington; and
2. to develop an educational program to teach commercial fishermen and boaters about marine debris through a variety of extension education techniques.

Squalicum Harbor provides moorage for about 1,750 boats, of which 1,050 are recreational and 700 are commercial. The commercial fishing fleet is composed mostly of gillnetters (7.6-10.8 m long) and purse seiners (15.3-18.4 m long) that fish in Puget Sound and Alaska. The recreational fleet has about equal numbers of sail and powerboats, and 75-100 boats are used as live-aboard homes.

The demonstration project was coordinated with the Port of Bellingham, which owns and manages Squalicum Harbor. The function of Washington Sea Grant staff in this project was to act as technical advisors to the port staff. They surveyed the boaters, analyzed waste collection facilities and alternatives, and offered suggestions on improvements that could be accomplished in a cost-effective and realistic manner. The educational program focused on using traditional extension education techniques. A poster and three publications on marine debris were developed. These materials were important components of the extension education portion of this project. Because of the time it has taken to implement the changes proposed in the demonstration project, no measurements of their effects have yet been made.

INTRODUCTION

For years, people aboard boats and in coastal areas threw their garbage into the water. Items such as tin cans, food waste, cardboard, and cotton fishing gear and lines quickly sank. These materials generally degraded and caused relatively few problems in the marine environment. Also, as a rule marinas had dumpsters or litter barrels near the docks for garbage disposal, but no specialized waste collection systems.

Today, however, much of the material that is thrown overboard or lost in Puget Sound and in the oceans is made of plastic. Plastics are very useful aboard vessels because they are lightweight, strong, and do not degrade when wet. However, these same qualities can cause problems when plastics are disposed of in the marine environment. Studies from many parts of the world have shown that serious problems result when wildlife encounters plastic marine debris (Shomura and Yoshida 1985; Center for Environmental Education 1987; Alaska Sea Grant 1988; Alverson and June 1988).

As more U.S. and worldwide attention focused on plastic marine debris, Annex V of the MARPOL Convention was ratified internationally in 1987, and to implement that convention in the United States, Congress passed the Marine Plastic Pollution Research and Control Act (MPPRCA) in 1987. The MPPRCA, which became effective on 31 December 1988, prohibits the dumping of plastics at sea and regulates the dumping of other wastes at sea (U.S. Department of Transportation 1988). With the implementation of MPPRCA, boaters and fishermen now must return boat wastes to port, and ports and marinas must have facilities to accept those wastes.

Within Puget Sound, the volume and sources of marine plastic debris and the problems it causes are not well known. Because of Washington's extensive recreational and commercial fishing fleets, one would expect to find debris common to those vessels and activities. Some negative impacts of derelict fishing gear within Puget Sound have been observed (High 1985), but the extent of the problem is unknown.

Despite a lack of specific data about the extent of the marine debris problem within the Puget Sound region, it was felt that marine plastic debris was causing problems in the area. Also, as a result of MPPRCA, boaters were prohibited from disposing of their wastes into Puget Sound, and marinas were mandated to have facilities to accept boat garbage. In general, however, few boaters, fishermen, or marina operators were familiar with the MPPRCA and its provisions. The Squalicum Harbor project was developed with two primary goals:

1. to develop a pilot project to collect and recycle vessel-generated wastes from commercial and recreational vessels at Squalicum Harbor, and
2. to develop an educational program to teach commercial fishermen and boaters in the Puget Sound region about marine plastic debris and its proper disposal.

The Port of Newport (PON), Oregon, had developed a pilot marine debris collection and education project that was fairly successful. With grant funding from the U.S. National Marine Fisheries Service, the Newport project was able to develop a successful waste collection system that was used by the commercial fishing fleet for nets, rope, wood, metal, and other materials (Recht 1988).

The Squalicum Harbor project hoped to build on the experiences of the Newport project. However, four major differences between the projects were evident. First, the PON project was organized as a staff project of the PON. The Squalicum Harbor project was being developed by people outside the Port of Bellingham staff. Second, the PON project provided funding for facilities and maintenance staff, whereas the Squalicum Harbor project did not. (For the project to be successful in the long run, it had to work within the operational budget of Squalicum Harbor.) Third, the types of debris at the harbors were different. In each harbor, the debris reflected the boats that use the marinas: Newport has more trawlers, and Squalicum has more purse seiners, gillnetters, and recreational boats. Fourth, the physical layouts of the harbors are different: Newport has separate marinas for the commercial and recreational fleets, whereas Squalicum Harbor has these fleets within the same marina.

METHODS

Squalicum Harbor is located some 144.8 km (90 mi) north of Seattle on Bellingham Bay. It provides moorage for about 1,750 boats, of which 1,050 are recreational and 700 are commercial. The commercial fishing fleet is composed mostly of gillnetters (7.6-10.8 m long) and purse seiners (15.3-18.4 m long) that fish in Puget Sound and Alaska. The recreational fleet has about equal numbers of sail and powerboats, and 75-100 boats are used as live-aboard homes. Squalicum Harbor has three water entrances and nine ramps to the docks.

Existing waste-handling and collection facilities and procedures were analyzed using: 1) a personal informal interview survey of boaters and fishermen who use the harbor; 2) discussions with Squalicum Harbor staff; 3) a visual survey of the waste-handling facilities; and 4) contacts with waste collection companies, recycling companies, and community agencies.

Educational materials were developed and written by Washington Sea Grant (WSG) staff working on the marine debris project. Original plans called for writing one extension education publication for each of four different audiences: Squalicum Harbor boaters and fishermen, commercial fishermen, recreational boaters, and marina operators. Additionally a poster, a slide show, and a display area were to be developed.

DISCUSSION

Squalicum Harbor Analysis and Proposal

Twenty-seven fishermen and boaters from Squalicum Harbor were interviewed during spring 1987. Tabulation of the respondents' answers

showed that 78% thought Bellingham had a marine debris problem, 52% had experienced problems such as fouled propellers or clogged water intakes caused by plastic debris, 74% indicated that the existing waste facilities at Squalicum Harbor were adequate, and 67% expressed a willingness to sort some of their wastes for a recycling program. Respondents also indicated that management of an oil recycling facility maintained for boaters' use needed improvement.

Squalicum Harbor provided one 4.58 m³ (6 yd³) dumpster at the top of each float ramp, additional dumpsters in the area where commercial fishermen work on their gear, and a 15.29 m³ (20 yd³) dumpster near the dock used for provisioning vessels.

The visual survey of dumpster contents and interviews with Squalicum Harbor staff showed that the composition of the garbage varied with the season and the type of harbor use near that dumpster. For example, the percentage, by volume, of cardboard boxes ranged from 5 to 100%, with a mean of 52%.

Squalicum Harbor had a contract with the local garbage disposal company to empty the dumpster at the harbor. Seasonal fluctuations in quantity and composition of garbage were reflected in different pickup schedules for different dumpsters (Table 1). Rates varied with the frequency of pickup, the size of the dumpster, and whether the garbage went to landfill or to incineration (Table 2). The cost of garbage service at Squalicum Harbor rose dramatically from 1983 to 1987 (Table 3). This increase was caused by marina growth, boater population growth, and increases in garbage pickup rates over the time period.

On analysis, Squalicum Harbor's waste-handling facilities were judged to be adequate. Increased volumes of waste materials generated because of heightened awareness of the facilities could be easily accommodated by increasing the frequency of dumpster pickup. Any attendant cost increases could be minimized by developing a collection and recycling system for cardboard and aluminum. As in the PON project, a significant volume of the wastes in dumpsters at Squalicum Harbor was cardboard.

With this analysis completed, a proposal was written and presented to the Squalicum Harbor staff in August 1988. This proposal provided a detailed plan to improve the waste-handling system at Squalicum Harbor. The major elements of the plan were to maintain all existing dumpsters in the harbor; provide collection boxes or cleared space for netting, cardboard, scrap metal, wood, and aluminum; organize free pickup of materials by local recycling companies; and advertise the program through signs, pamphlets, news articles, presentations, displays, and word of mouth.

Figure 1 locates the proposed waste collection facilities at Squalicum Harbor. These facilities would:

- Provide a central location at the harbor for recycling scrap metal, wood, and netting. This would be a cleared space with signs indicating where to stack different materials.

Table 1.--Frequency of dumpster pickup at Squalicum Harbor.

Season	Size of container	Frequency of pickup
Winter (Oct.-May)	4.58 m ³ (6 yd ³)	1 per week
Winter	15.29 m ³ (20 yd ³)	1 per month
Winter	15.29 m ³ (20 yd ³) (trash compactor)	1 per month
Summer	4.58 m ³ (6 yd ³)	3 per week
Summer	15.29 m ³ (20 yd ³)	On-call basis
Summer	15.29 m ³ (20 yd ³) (trash compactor)	On-call basis

Table 2.--Monthly rates for hauling and disposal of trash from Squalicum Harbor.

Dumpster size	Frequency of pickup		
	1 per week	2 per week	3 per week
4.58 m ³ (6 yd ³)	149.27	276.15	403.03
15.29 m ³ (20 yd ³)	^a 130.75	^a 205.11	^a 279.47
15.29 m ³ (20 yd ³) (trash compactor)	^a 107.20	^a 214.40	^a 321.60

^aAdditional charges for hauling and disposal fee.

Table 3.--Total costs of garbage service in Squalicum Harbor.

Year	Cost
1983	\$18,794
1984	\$22,394
1985	\$30,456
1986	\$34,492
1987	\$41,758
1988	\$50,000 (estimated)

- Provide collection boxes for sorting and recycling of cardboard at five of the nine dumpsters at the harbor, dumpsters used primarily by the commercial fishing fleet. Used wooden fish totes were donated by local seafood companies for collection boxes. Removing the cardboard was expected to reduce the rate at which the dumpsters filled up and thus reduce garbage costs.




Figure 1.--Proposed waste collection facilities at Squalicum Harbor.

- Provide three aluminum recycling facilities at the harbor adjacent to the dumpsters used by the recreational boating fleet. Aluminum is a high value recyclable item, and these recycling revenues could help offset the cost of waste disposal.

The proposed facilities could be used by boaters and fishermen with a minimum of sorting. As a result, use of the facilities was expected to be heavy.

In addition, only materials that had ready markets were included in the recycling program. At no charge to the port, companies in the Bellingham area would pick up one or more of the materials being collected. This was expected to help reduce maintenance needs.

Glass was excluded from the recycling program for two reasons: 1) glass has a low market value and local companies would not pick it up at Squalicum Harbor, and 2) because of the weight of glass, specialized equipment would have been needed to handle it.

The Newport project found that blue color-coding of their recycling facilities was very useful, and the Squalicum Harbor project also color-coded the recycling facilities blue. Many fishermen and boaters travel frequently from port to port on the U.S. west coast and Alaska, and WSG suggests that for ease of recognition blue be adopted as the color for recycling facilities in all ports.

Educational Program

The educational portion of the program was multifaceted and involved working with the Washington State Task Force on Marine Plastic Debris. This task force had representatives from some 40 different governmental, environmental, industrial, educational, and community groups who worked together to develop a Washington State Marine Plastic Debris Action Plan (Washington State Department of Natural Resources 1988). As task force participants, WSG staff developed a logo for statewide marine debris cleanup (Fig. 2). This logo and the slogan "Get a Grip on Marine Debris" are being used throughout Washington State.

The educational portion of the Squalicum Harbor project also included developing and printing a marine debris poster; writing pamphlets directed at Squalicum Harbor boaters, commercial fishermen, and recreational boaters; providing presentations to various community and school (K-12) groups on marine debris; developing a liaison with Western Washington University's plastics technology program; and being available to the media on marine debris-related matters. Using these educational materials, WSG reached a total of 585 people at workshops and other meetings, and distributed more than 2,500 posters and pamphlets.

RESULTS AND CONCLUSIONS

The analysis and proposal conducted by WSG staff were provided to the Squalicum Harbor staff much the way a consultant would provide information.




Figure 2.--Washington State marine debris logo.

However, the Squalicum Harbor staff had to make the actual changes. Six months after receiving the recommendations, they were just beginning to implement the physical changes. The first cardboard collection boxes were installed in early March 1989, and they were immediately used by the boaters and fishermen. The installation of the other facilities were expected to occur shortly thereafter.

We can only theorize about why implementing the proposal took so long. First, the Squalicum Harbor staff appeared to be already working at their maximum level. When a staff is already working at or near capacity, a new project is difficult to start. Second, in spite of the analysis and proposal, the staff appeared reluctant to implement the project for fear of generating more maintenance work for themselves. Third, this project may

have been viewed with some resentment because it was promoted by people outside of the Squalicum Harbor staff.

Because of the time it took to have physical changes made at Squalicum Harbor, no measurements have yet been made on the effects of the changes. This project points out the difficulty of setting up a demonstration project as an "outsider," and should caution others to expect to go slowly in similar projects.

REFERENCES

Alaska Sea Grant.

1988. Oceans of plastic. Report on a workshop on fisheries-generated marine debris and derelict fishing gear. Alaska Sea Grant Rep. 88-7, 68 p.

Alverson, D. L., and J. A. June (editors).

1988. Proceedings of the North Pacific Rim Fisherman's Conference on Marine Debris, Kailua-Kona, Hawaii, 13-16 October 1987. Unpublished report by Natural Resources Consultants, 4055 21st Avenue W., Seattle, WA 98199, 460 p.

Center for Environmental Education.

1987. Plastics in the ocean: More than a litter problem. Center for Environmental Education, Wash., D.C., 128 p.

High, W. L.

1985. Some consequences of lost fishing gear. In R. S. Shomura and H. O. Yoshida (editors), Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii, p. 430-437. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54.

Recht, F.

1988. Dealing with Annex V. A reference guide for ports. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWR-23, 132 p.

Shomura, R. S., and H. O. Yoshida (editors).

1985. Proceedings of the Workshop on the Fate and Impact of Marine Debris, 26-29 November 1984, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-54, 580 p.

U.S. Department of Transportation.

1988. Advance notices of proposed rulemaking: Regulations implementing the pollution prevention requirements of Annex V of MARPOL 73/78. Federal Register, June 24, 1988, p. 23884-23895.

Washington State Department of Natural Resources.

1988. Marine plastic debris action plan for Washington State. Wash. State Dep. Natur. Resour., Olympia, Wash., 45 p.

MARINE DEBRIS: NORTH CAROLINA'S SOLUTIONS THROUGH EDUCATION

Sarah Friday
University of North Carolina Sea Grant College Program
Raleigh, North Carolina 27695, U.S.A.

ABSTRACT

North Carolina began its campaign against marine litter in June 1987. The success of the program has been due largely to an emphasis on interagency cooperation and on education of the public. Five cooperating state agencies have been the University of North Carolina Sea Grant Program, and North Carolina Division of Coastal Management, Division of Parks and Recreation, 4-H, and Office of Marine Affairs. Educational activities have included slide programs given by a Sea Grant marine education specialist to power squadrons, fishing clubs, school groups, and service clubs, and ongoing exhibits and talks on marine debris by the state's three coastal aquariums. The program has also stressed youth-oriented activities relating to marine debris.

INTRODUCTION

North Carolina has nearly 560 km (350 mi) of coastline and 931,500 ha (2.3 million acres) of estuaries, bays, and sounds. Without question, the beaches and coastal waters are vital to the aesthetics and the economy of the state.

But litter and plastics could change this.

That is why North Carolina began its marine litter program in June 1987. Now, 21 months later, the state can show evidence of change and the promise of regulations and educational programs that ultimately will help solve litter problems at the coast.

METHOD AND DISCUSSION

On one Saturday in September 1987 and another in September 1988, the state marine debris program held Beach Sweep, a 1-day coastwide cleanup. Over 4,500 volunteers came to the coast and picked up more than 54 metric tons of trash. The volunteer participation, the amount of litter collected, and the extensive media exposure for Beach Sweep were the result of educational efforts by program coordinators and volunteers.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

Interagency Cooperation

North Carolina patterned Beach Sweep after similar cleanups in other states, but organizer Lundie Spence, Sea Grant's marine education specialist, took a different approach. Like the director of a play, Spence pulled in talent from many of the state agencies to play different roles in Beach Sweep and the marine debris program.

The 1987 Beach Sweep organization committee was composed mainly of representatives from four state government agencies. They were Sea Grant, the North Carolina Division of Coastal Management, Division of Parks and Recreation, and Office of Marine Affairs, which oversees the state's three coastal aquariums. In 1988, North Carolina 4-H Clubs joined the committee.

Full cooperation and contributions of time, money, and services from each agency helped spread the word to thousands of North Carolina citizens.

For example, Sea Grant's Spence kept state government leaders informed of activities concerning marine debris. She sent information on marine debris and Beach Sweep through a Sea Grant newsletter to 2,000 North Carolina teachers, and she gave slide programs to power squadrons, fishing clubs, school groups, and service clubs.

Sea Grant's communications staff handled all of the major publicity for Beach Sweep. This included issuing press releases, writing features and newsletter articles, compiling press kits, and scheduling radio and television interviews.

The Division of Coastal Management took on other tasks. As the state's coastal regulatory agency, this division was able to work closely with the governor's office, legislators, the state's Coastal Resources Commission, and the Marine Science Council to garner support.

Parks and Recreation contributed manpower at the coast for the cleanups and helped with fund-raising and contributions of garbage bags and pencils for tallying data.

The three North Carolina aquariums offered staff that served as regional coordinators for Beach Sweep. Each aquarium provided special exhibits and programs on marine debris to tourists and other interested visitors.

As the state's marine debris program grows, an increasing number of state government agencies, private nonprofit groups, corporations, small businesses, and volunteers are participating.

Targeting groups of all kinds with coastal and environmental interests provides not only a rich pool of talent and services, but also an unlimited resource for ideas.

In this case, the more the merrier.

This year, Beach Sweep and the North Carolina marine debris program still operate with no major corporate funding. The shoestring budget has gotten a bit fatter, but the shoestrings are not yet long enough to help accomplish all of the goals. Therefore, interagency cooperation is even more important to help carry out a comprehensive educational program.

Public Education

Emphasis on Youth

A special emphasis in North Carolina has been placed on creating awareness for the state's youth. Within the school system, the state's marine debris program coordinators have worked with science, environmental, and gifted-and-talented classes. In this area, copies of the marine debris slide and talk program from the Center for Marine Conservation have been made available to the schools.

Teachers have found out about other reference materials through a special newsletter from Sea Grant. In response, children from the fourth to twelfth grades have written Sea Grant for information. This exercise gives younger children the experience of writing for and receiving information on their own. Taking this action is one more step toward increasing awareness by personal involvement.

The North Carolina aquariums, the state maritime history museum, and the parks and recreation system have offered special youth programs on marine debris. Typically, leaders give a short talk or slide show on marine pollution and then take the group out on the beach to collect trash. After 30 to 40 min, they stop and discuss their findings and reactions.

This program has been modified to assist Boy Scout troops at the coast in earning a badge relating to environmental awareness. One troop expanded the idea during Beach Sweep and separated its trash for recycling.

Another excellent idea for increasing youth awareness has been a permanent display on marine debris erected at the North Carolina Maritime History Museum. One panel of the display is just for children. On it are photos of volunteers during Beach Sweep, an award-winning 4-H poster concerning plastics, and a pad and pen for comments. Pages and pages of ideas and comments have told museum educator Patricia Hay that children's eyes are open to the problems at the coast.

Probably the most far-reaching involvement originated with the state 4-H Club. This national youth organization has programs in each of North Carolina's 100 counties. Within this structure, Beach Sweep was promoted through newsletters and electronic mail as a good community service project for its members.

In addition, 4-H implemented a statewide marine debris poster contest for youths ages 9 to 11, and 12 and older. As schools already employ many such contests, this gave the young people a different avenue. The 4-H Club raised \$255 for cash prizes in each category.

Most importantly, 4-H and Sea Grant helped devise five activities for youths concerning plastics and litter. They are currently being used by 4-H Clubs and school teachers throughout the state. But since they are new, 4-H education specialists are still conducting tests on the usability and viability of these activities as a curriculum.

The five projects include:

1. Living labels. This icebreaker invites students to become aware of types of plastic items around them. Students either act out the item they have chosen or give 10 words describing it. Other class members try to guess the item.
2. Why do we use plastics? By listing different kinds of plastics we use daily and discussing them, young people gain a better idea of how much plastic is used and why.
3. How strong is a six-pack ring? Youths test the strength of six-pack rings and correlate this with their durability in the environment. The objective is for them to realize how little chance an animal or fish has of freeing itself after entanglement.
4. Can we make plastics disappear? By comparing degradable and nondegradable six-pack rings, youths
 - a. understand the meanings of photodegradable and biodegradable,
 - b. learn that most plastics are not degradable, and
 - c. learn that plasticlike materials can be made degradable.
5. Turning trash around. By simulating paper and plastics recycling, youths gain a better understanding of the process of and need for recycling in our society.

Other

The Beach Sweep effort has helped make many of North Carolina's citizens aware of marine pollution. Each volunteer that attended the cleanup has seen firsthand what litter can do to our beaches and how he or she can help keep them clean. The social, environmental, and economic impact of cleaner beaches is affecting the behavior of tourists. And pollution has become a political issue because people are concerned.

Since our cleanup began, certain public beaches have been kept cleaner year-round. Municipalities added more trash cans to beach access areas, and some of them added to their cleanup crews. Also, North Carolina is currently considering legislation on degradable six-pack rings.

Public education has not stopped with Beach Sweep, however. Each of the involved agencies has initiated programs concerning marine debris.

CONCLUSION

With these projects and our other efforts, North Carolina has begun its fight against marine pollution. By working together and focusing energies on the state's youth, much has been done.

For the past 2 years, North Carolina Sea Grant has been part of a national Sea Grant marine debris network. Each of the nation's 30 coastal states has contributed toward educating commercial fishermen, teachers, boaters, and the public about pollution and its effects.

Until the problem is conquered, however, much remains to be accomplished.

DESIGNING EFFECTIVE EDUCATIONAL PROGRAMS:
THE ATTITUDINAL BASIS OF MARINE LITTERING

Shirley Laska
Department of Sociology
University of New Orleans
Lakefront, New Orleans, Louisiana 70148, U.S.A.

ABSTRACT

The worldwide concern with marine and coastal debris has sparked recommendations for various abatement interventions. Popular among them have been educational programs. Effectiveness of educational interventions directed toward changing environmental attitudes and behavior has, however, been found wanting, according to some recent assessments. It is argued that problems with educational abatement programs may stem from the lack of appreciation and lack of application of social science knowledge about the basis of environmental beliefs, attitudes, values, and human behavior which affect the environment.

Marine debris abatement efforts can be enhanced by basing them on social science knowledge in three relevant areas: 1) paradigms and the nature of environmental attitude formation; 2) the nature and constraints of the desired nonlittering behavior; and 3) research on attitude and behavior change, including recycling and land litter abatement. Each of these topics is reviewed, with recommendations about its application to marine debris abatement.

INTRODUCTION

At the conference on the topic of fisheries-generated marine debris and derelict fishing gear held approximately 1 year ago in Portland, Oregon, educational programs were assessed by one participant as the most popular marine debris abatement approach. He stated that they are "politically attractive, do not cost much and meet other favorable criteria." However, the author also noted that such programs "often have only modest effectiveness and lack permanence" (Alaska Sea Grant Program 1988, p. 7).

The thesis of this paper is that one of the reasons for the limitation of these educational programs is the failure to understand fully the human and social causes of the problem. It is proposed that this shortcoming is due to the limited participation by social scientists in addressing the

problem. Physical scientists have assessed the extent and nature of the impact of marine debris. From this data base has evolved a determination that a "problem" exists. But the definition of the problem has remained too strongly physical because it lacks the additional perspective of environmental issues provided by the social scientist.

An example of this lack of appreciation of the social component of the marine debris problem is seen in the recommendations made by the Inter-agency Task Force on Persistent Marine Debris formed by the White House Domestic Policy Council. The task force recommended that 1) the (marine debris) problem be quantified, 2) the sources be determined, and 3) ways be found to reduce plastic debris from all sources.

The phrasing of the document suggests that the "source" of the problem is merely a physical location or use or particular economic activity seemingly devoid of human input. Review of this and other documents similarly phrased revealed that there was a missing step in these recommendations. Simply put, the missing step is to ask, "Why do litter and debris exist in the marine and coastal environments?"

The goal of adding this question is to refocus the problem-solving to recognize that human behavior is the cause of the litter and debris and not just a "source." If the human nature of the problem is not addressed in the problem-solving efforts, educational interventions cannot be effective but appear as an afterthought because "something must be done." It is contended that answers to the question of why debris exists must be determined and understood before and if "ways [are to] be found to reduce plastic debris from all sources."

SOCIAL SCIENCE FINDINGS RELEVANT TO MARINE DEBRIS ABATEMENT

With the goal of addressing this issue of the human cause of marine debris, three relevant social science topics will be briefly examined. The three topics are:

1. paradigms and the nature of environmental attitude formation,
2. the nature and constraints of the desired nonlittering behavior in the marine environment, and
3. research on attitude and behavior change including recycling and land litter abatement.

Each of these topics is reviewed with recommendations about its application to marine debris abatement.

The Nature of Environmental Attitude Formation

Societal Paradigm

Members of a society share a common world view embodied in beliefs, attitudes, and values. This world view, frequently referred to as a

paradigm, emanates from the experiences of the members of the society and is functional to the society in that it supports the society's efforts to survive. While the paradigm does not necessarily answer important questions, "it tells us where to look for answers" (Babbie 1989). It also becomes the basis for choosing problems that can be assumed to have solutions (Kuhn 1970). When the existing paradigm no longer serves the members of society, it changes as the established agreed-upon paradigm is modified in favor of a new one(s).

Within each society, subgroups share paradigms useful for supporting their experiences and position in society. For example, different scientific disciplines have different paradigms as do other occupational groups. In order to understand the basis of behavior of members of a society, it is important to appreciate both the general world view of the larger society and those views of the subgroups about which you have specific interest. Both the general and the subgroup will be discussed below.

The paradigm common to the American society has been characterized as the "dominant Western world view" (Dunlap and Van Liere 1984), the "technocratic" paradigm (Drengson 1980), or the "human exemptionalist" paradigm (Catton and Dunlap 1980). This paradigm sees the relationship between humans and the environment as one in which humans utilize the environment for their betterment, even at the expense of the environment. Based on the Judeo-Christian heritage, this paradigm assumes a human superiority over other organisms bequeathed to humans by their special relationship with God. Contained in this perspective of superiority is the belief that any problems which befall the environment during its exploitation can be remedied by humans through technology. A society which holds this paradigm believes that it is exempt from conformity to the natural ecological laws because of its ability to overcome any environmental problems.

The American extension of this paradigm divides the environment into parts which are privately owned for the gain of the owner and parts which are publicly, or commonly, owned (Hardin 1968). The "commons" is shared for the betterment of all members of the society. However, as Hardin (1968) notes, such a common betterment for all becomes impossible once the ratio of the population to the environment surpasses its "carrying capacity." At that point the common good suffers as the individual benefits from use of the commons.

Industrial pollution and littering behavior represent the use of the common for the betterment of individuals and their economic interests. Belief in the right of individuals to so use the commons for their interest is an important belief contained in the American paradigm. Throwing something "away" means simply putting it into the commons when it no longer serves the person's needs.

The strength of this tenet within the western paradigm is evident when we consider the "absurdity" of the idea that all pollution, every single bit, should be totally banned from the commons. To paraphrase, what general societal support would there be for the total prohibition of environmental degradation in all commonly shared environments--air, water,

and public lands? The economic interests would argue that such a position would cause mass bankruptcies. Average citizens would also resist such a ban when they came to realize how it would affect the consumer-oriented lifestyle which has also evolved from the paradigm.

Discarding refuse into the ocean becomes an obvious extension of this paradigm. A series of studies done by sociologists Dunlap and Van Liere (1978, 1984) sought to determine to what extent such a degrading orientation toward the environment was linked to subscription to the western paradigm. In other words, does a person's belief in the dominant western paradigm affect his or her attitudes (and behavior) toward the environment? Van Liere and Dunlap (1980) broke the paradigm down into eight dimensions:

- support for laissez-faire government,
- support for the status quo,
- support for private property rights,
- faith in science and technology,
- support for individual rights,
- support for economic growth,
- faith in material abundance, and
- faith in future prosperity.

Three of these dimensions of the dominant western paradigm were associated negatively with the environmental attitudes scales they also developed. The greater the support for the following, the less the support for the environment:

- support for private property,
- support for economic growth, and
- faith in material abundance.

We may, thus, propose from these research findings that individuals who subscribe closely to the dominant western paradigm--particularly support for private property, economic growth, and belief in material abundance--will more likely not hold proenvironmental attitudes. If such proenvironment attitudes are not held, it is more likely that littering behavior, such as that which results in marine debris, will be exhibited.

An alternative paradigm is evolving as the current paradigm becomes less functional. The paradigm which would reduce the stress on the environment is called the "new environmental paradigm" (Catton and Dunlap 1980) or the "person-planetary" paradigm (Drengson 1980). This paradigm accepts the fact that humans are subject to the same ecological laws as other organisms, and when humans degrade the environment, they are not always able to repair it with technology.

A "paradigm shift" (Kuhn 1970) by the majority of the populace may occur more rapidly than might be anticipated given the escalating environmental problems. Recent political deliberations in southern California were directed at reducing air pollution conditions that are no longer acceptable. The proposal included restricting each family to owning one car and requiring that the family members work near where they live. A

recent proposal by the U.S. Environmental Protection Agency directed toward reducing acid rain indicated that the solution could come with a required dropping of interior household temperature during the winter. While the dominant western paradigm still prevails, the fact that "responsible, mainstream" public officials were making these proposals suggests that the paradigm is shifting.

Educational interventions directed toward antilittering in the marine environment can benefit from an awareness of the target population's paradigm. If it is determined that they subscribe to the traditional paradigm, efforts toward modifying their world view should be included in the communication along with the message directed toward the specific change in littering behavior. If on the other hand, they are shifting their paradigm, then the communication can benefit from "tapping" this new orientation in the message.

Subgroup Paradigms

Besides the general societal orientation toward the environment, subgroups of the population have been found to vary in their attitudes toward the environment depending on their relationship to it. Awareness of these subdifferences can also benefit educational interventions directed at target populations.

For example, Louisiana has both the most active offshore oil extraction activities in the United States and some of the most prolific spawning grounds for fish and shellfish in the United States. Thus, coastal users include large groups of fishermen and offshore oil rig employees. Should we expect to find these groups different in their attitudes toward marine littering or the same? Popular opinion might argue that the oil rig employees would be likely to have less concern for the environment than fishermen whose livelihood depends on a healthy environment.

Research findings, however, suggest the contrary. Rural residents have been found to be less proenvironment than urban residents because they work in agricultural and mining activities which approach the environment in a more utilitarian, exploitative fashion (Lowe and Pinhey 1982). It is thus likely that the fishermen and oil workers will be more similar than different in their environmental attitudes. Louisiana fishermen have a reputation for considerable littering, as do oil rig employees. They also have strongly resisted steps taken by the Federal Government to protect the environment, such as the requirement that they use turtle excluder devices (TED's) while shrimping to protect the Kemp's ridley sea turtles.

The exploitative orientation toward the environment of some jobs may not be the only influence which engenders human exemptionist attitudes in those whose livelihood depends on the environment. Companies for which such individuals work may themselves have company "cultures" reflecting similar orientations. It may be more cost-effective to use the commons for refuse, and this belief is learned as employees learn what is expected of them on the job. Personal worker economic motivation to perform well on the job compounds their own inclinations to litter.

Other individual characteristics have been found to affect attitudes and behavior toward the environment. Small town residents appear to also be less proenvironment. Van Liere and Dunlap (1980) explain this observation with the "progrowth" orientation of small towns. Urban residents have, to the contrary, been found to be more proenvironment. Environmental deprivation theory is used to explain this finding: Urban residents experience higher levels of pollution and environmental degradation and can thus make a comparison between the existence and nonexistence of pollution, which results in developing a proenvironment orientation (Dillman and Christenson 1972). In addition, urban residents are more likely to appreciate a "social solution" to environmental problems because they experience control over their built environment (Lowe and Pinhey 1982). Thus, they are willing to accept the existence of a problem because they perceive it as solvable. Take for example the predominance of urbanites in the groups who are involved in the annual beach cleanups.

In addition to geographic location, education, race, sex (Sigelman and Yanarella 1986), social class (Buttel and Flinn 1978), and age (Hamilton 1985) have been found to predict attitudes toward the environment. Income has not been found so strongly associated (Constantini and Hanf 1972; Sigelman and Yanarella 1986). Van Liere and Dunlap (1980, p. 190), in summarizing the findings from numerous studies of social characteristics, indicate that the association between income and environmental concern is "quite ambiguous and fail[s] to support the hypothesized positive association." This ambiguity may be due to changes which are taking place in the way in which the less affluent view environmental problems. While they value the jobs that come from industry, which often pollutes, they are becoming more aware that the pollution from such activity often is discarded closer to their communities than to those of the more affluent (Bullard and Wright 1986).

Similar changes may also be occurring in some of the relationships found in the subgroups reviewed above. While fishermen may not have traditionally been inclined to be concerned with the environment, the depletion of the resource such as the threat to the Gulf of Mexico redfish population may also begin to change their orientation to the environment as well.

It is important that the educational interventions being developed to reduce beach and marine debris are oriented toward the expected attitudes which various target groups might hold toward the environment, and that support be given to maintain current information on the attitudes which coastal users hold so that the intervention is relevant to the orientation.

Characteristics of Nonlittering Behavior

In addition to appreciating the orientation which coastal users have toward the environment, it is important to understand the nature of the behavior which the educational program attempts to change--i.e., marine littering--and the meaning given to that behavior by those who do it. Conversely, it is also important to recognize the characteristics of the desired behavior--nonlittering--which make conformity to it more difficult than other environment-oriented behaviors.

Complexity of Desired Behavior

Nonlittering behavior in the marine and coastal setting contains both a "don't do" and a "do" component, making its successful implementation somewhat complex. It is desired that individuals not discard that which is no longer of value to them into the commons--i.e., into the water or along the beach. Then, it is desired that individuals maintain the item in their personal space--in their pocket, on the boat, on the oil rig, with camping and fishing gear--until they are able to discard it in an appropriate refuse-collecting device or area such as a trash barrel, dumpster, junk yard, or landfill.

Inconvenience

Nonlittering behavior in the marine and coastal setting is an inconvenience to the individual because the appropriate refuse-collecting device is frequently not in the immediate vicinity. It may be at the beach entrance, at a nearby gas station, at home, or, for larger items, at a special location requiring an even longer trip. The inconvenience is defined by the length of time the individual may have to maintain the item which no longer has utility within his or her personal domain--often extended if out at sea--and by the fact that it is occupying part of a very limited space--fishing boat, oil rig, freighter, camper.

Limitations of Social Control

Littering in the marine and coastal setting is frequently done when there are no other people around to observe the behavior. Or, if others are present, they are experiencing the same need--to discard no longer useful items taking up precious space. Or, the observers are strangers or at such a distance that the litterer can maintain his or her anonymity while violating the norms. This means that the behavior can frequently not be controlled by the knowledge that someone else is observing them doing something contrary to the norm, a manner in which much desired behavior is encouraged.

An anecdote demonstrates norm-control dynamics and the way a litterer can attempt to avoid them. Last summer on a Florida beach, I watched a well-coiffured, expensively dressed beachcomber with cigar in one hand and a soft drink can in the other stop on a crowded beach to dig a small hole in the sand with his toes. He deposited the empty soft drink can in the hole, covered it over, and walked on. This behavior suggested to me that he knew littering was against the norm. If, however, he could hide the object, he would be able to avoid the possible scorn of the onlookers or rationalize that he had not littered because the object was no longer visible--one of the most commonly mentioned qualities of litter being that it is an eyesore.

Impact Not Obvious

The effect which the littering behavior is having on the environment is not seen or appreciated by the litterer because the discarded item frequently disappears under water or sand. Even when it does not disappear,

the beach and the ocean are so vast that the ratio of litter to commons is miniscule. It is difficult to appreciate the significance of one Styrofoam cup tossed off an oil rig into a vast ocean. Or even a piece of fishing gear or the waste from an ocean-going vessel.

Implications for Behavior Change

It is argued that awareness of these particular qualities of marine and coastal littering and nonlittering behavior can contribute to the development of more effective educational interventions directed toward curbing such littering behavior.

First, awareness that there is a lack of a clear antilitter norm in the marine environment and a lack of critical observers to enforce whatever norm there is should reduce emphasis on norm conformity in educational programs.

Second, it would be expected that educational campaigns which present graphic evidence of the impact of littering on wildlife--such as the ones recently developed--would improve antilittering behavior. They would help the individual become aware of the impact on the environment of even one small discarded item, albeit seemingly insignificant and invisible when the littering act is committed.

Third, educational campaigns should be implemented in conjunction with strong efforts to provide very convenient locations for disposal of marine refuse.

Examples of the linking of education and convenient refuse disposal are available in the recycling efforts of some communities. One successful pilot community recycling program in Louisiana has stackable containers for curbside pickup clearly marked for glass, aluminum, and paper. The effort to conform is thus quite minimal.

Similarly, studies should be conducted to determine the most convenient refuse disposal configuration at beaches, boat launches, marinas, and harbors. Some disposal services might best be reached from the water so that refuse does not have to be carried onto land by hand. Also, litter bag dispensers could be placed at convenient locations near boat launches and docks to encourage convenient on-boat refuse storage. With such accommodating refuse disposal facilities in place, beach or dockside antilitter signs would be encouraging a more feasible behavior. Once the marine user is practiced in such nonlittering behavior, the behavior will seem "more natural" and such attention to convenient refuse disposal will not be so important. This would be a case of learning to cope with natural hazards (this hazard being to marine life) through participation, as proposed by Sorensen and Mileti (1987).

An example of such facilitating assistance has been tried with success at several Louisiana fishing rodeos. When registering for the rodeo, each entrant was given a trash bag, with a request that it be used and returned to the registration desk at the end of the day. Those entrants who returned their bags filled with the day's refuse qualified for a special drawing for

several significant prizes. The bags were donated, as were the prizes, in return for public recognition that the companies had performed the public service. While the prize component of the activity may not be conducive to continuing the behavior after the fishing rodeo (see below), it might not have a negative effect if the bag contained a recommendation to always take along a trash bag and if trash bags were conveniently dispensed at docks and launch sites on a regular basis.

Research on Attitude and Behavior Change Including Recycling and Land Litter Abatement

The third social science topic of relevance for improving educational programs directed toward marine debris abatement is the research on attitude and behavior change. Research has been conducted on the content of successful persuasion communication in general and on persuasion directed toward specific attitude and behavior changes. These include self-help behaviors related to health, safety, crime and natural hazards protection (see Weinstein 1987 for a useful review), and energy conservation, recycling, and litter abatement on land. A review of theories useful for attitude and behavior change with regard to solid waste demonstrates the utility of this literature.

There is little systematic theory concerning the social psychological variables which influence littering (Reich and Robertson 1979). However, several theories have been found to be useful in changing littering attitudes and behavior. These include reactance theory (Mazis 1975); cognitive dissonance theory (Weigel and Weigel 1978; Shipee et al. 1980; Cook and Berrenberg 1981) or balance theory (Winham 1972); saliency theory (Cook and Berrenberg 1981); and Bem's self-perception theory (Arbuthnot et al. 1976-77; Pedersen 1979; Pardini and Katzev 1984). Each of these theories explains behavior based on an assessment which people make about themselves or those around them.

Reactance theory asserts that "when a person believes himself free to engage in a given behavior and his freedom is eliminated or threatened with elimination, the individual experiences psychological reactance" (Mazis 1975). When this occurs, the planned intervention results in behavior opposite to what is desired. An example is the TED which has been so strongly resisted by shrimpers in the Gulf of Mexico. While preventing all resistance to the TED's would have been impossible, a greater appreciation of the likelihood of reactance might have engendered different approaches by the environmentalists. Likewise, by knowing what coastal users believe they are free to do in the coastal environment, litter abatement interventions can be developed which will be less likely to cause such reactance.

Dissonance theory also has potential utility. It proposes that dissonance may occur for individuals among various values and beliefs which they hold and observations which they make. When this occurs, a person tries to reduce the dissonance. A person might interpret the observation such that it supports values and beliefs already held. Such dissonance may exist for the marine and coastal user with regard to littering. By determining whether it does, educational programs can be developed encouraging certain attitude and behavior change to "assist" coastal users in reducing their dissonance.

Saliency theory applied to conservation behavior "impl[ies] that the salience of proconservation attitudes will be enhanced primarily through or in anticipation of associations with others" who share proconservation attitudes (Cook and Berrenberg 1981, p. 82). When such persuasions are implemented, the presence of those holding nonlittering attitudes would be likely to increase saliency. Their presence can also be felt by presenting their proconservation statements in their absence or by asking residents to make public commitments to proconservation behavior. Beach cleanups are an example of a way to enhance the saliency of nonlittering behavior. Individuals make a public commitment to proenvironment behavior in the presence of other like-minded individuals.

Likewise, Bem's theory of self-perception can be applied to changing marine littering behavior. Bem proposes that behavior change occurs after a person changes his or her self-image to one capable of the new behavior (Arbuthnot et al. 1976-77). This self-image change can be assisted by educational programs that require small behavior change commitments to start the process of self-image change. Arbuthnot et al.'s successful experiment required minor recycling commitments which then led to a willingness to undertake more extensive recycling. Refuse disposal such as recycling at marinas and harbors could be approached in such an incremental way. Based on this theory, educational programs which encourage refuse disposal by giving prizes would not be expected to work over the long run because individuals do not have to change their self-image. The motivation to dispose correctly remains external to them, i.e., a prize given by someone else.

CONCLUSION

It will not be easy to integrate even the few concepts and theories presented in this paper in addressing the marine debris problem. It will be even more difficult to determine and apply the appropriate theories when social science research on relevant topics is more thoroughly reviewed. However, motivation to address the existing research and to respond to it can be found in recognizing the difficulty of success with educational programs in light of resistance from the existing human exemptionalist paradigm. Human behavior is the result of very complex social psychological processes influenced by the structure of the society in which the person lives and his or her position within that society. To have a modicum of lasting success in behavior change, "one needs all the help one can get." It will require a cooperative effort of both physical and social scientists to provide the knowledge base needed by those working directly with the marine debris problem to address its solution in a timely and successful fashion.


REFERENCES

- Alaska Sea Grant Program.
1988. Oceans of plastic. Report on a workshop on fisheries-generated marine debris and derelict fishing gear. Alaska Sea Grant Rep. 88-7, 68 p.


- Arbuthnot, J., R. Tedeschi, M. Wayner, J. Turner, S. Kressel, and R. Rush.
1976-77. The induction of sustained recycling behavior through foot-in-the-door technique. *J. Environ. Sys.* 6:355-368.
- Babbie, E.
1989. *The practice of social research.* Wadsworth, Belmont, CA.
- Bullard, R. D., and B. H. Wright.
1986. The politics of pollution: Implications for the black community. *Phylon* 47:71-78.
- Buttel, F. H., and W. L. Flinn.
1978. The politics of environmental concern. *Environ. Behav.* 10:17-36.
- Catton, W. R., Jr., and R. Dunlap.
1980. A new ecological paradigm for post-exuberant sociology. *Am. Behav. Scientist* 24:15-47.
- Constantini, E., and K. Hanf.
1972. Environmental concern at Lake Tahoe: A study of elite perceptions, backgrounds, and attitudes. *Environ. Behav.* 4:209-241.
- Cook, S. W., and J. L. Berrenberg.
1981. Approaches to encouraging conservation behavior: A review and conceptual framework. *J. Social Issues* 37:73-107.
- Dillman, D. A., and J. A. Christenson.
1972. The public value for pollution control. In W. R. Burch, Jr., N. H. Creek, Jr., and L. Taylor (editors), *Social behavior, natural resources and the environment*, p. 237-256. Harper & Row, N.Y.
- Drengson, A. R.
1980. Shifting paradigms: From the technocratic to the person-planetary. *Environ. Ethics* 2:221-240.
- Dunlap, R. E., and K. D. Van Liere.
1978. The new environmental paradigm. *J. Environ. Ed.* 9:10-19.

1984. Commitment to the dominant social paradigm and concern for environmental quality. *Soc. Sci. Quart.* 65:1013-1028.
- Hamilton, L. C.
1985. Who cares about water pollution? Opinions in a small-town crisis. *Sociological Inquiry* 55:170-181.
- Hardin, G.
1968. The tragedy of the commons. *Science* 162:1243-1248.
- Kuhn, T.
1970. *The structure of scientific revolution.* Univ. Chicago Press, Chicago.

- Lowe, G. D., and T. K. Pinhey.
1982. Rural-urban differences in support for environmental protection. *Rural Sociology* 47:114-128.
- Mazis, M. B.
1975. Antipollution measures and psychological reactance theory: A field experiment. *J. Pers. Soc. Psychol.* 31:654-660.
- Pardini, A. U., and R. D. Katzev.
1984. The effect of strength of commitment on newspaper recycling. *J. Environ. Sys.* 13:245-256.
- Pedersen, D. M.
1979. Changing beliefs concerning the causes of pollution. *J. Soc. Psychol.* 107:295-296.
- Reich, J. W., and J. L. Robertson.
1979. Reactance and norm appeal in anti-littering messages. *J. Appl. Soc. Psychol.* 9:91-101.
- Shipee, G., J. Burroughs, and S. Wakefield.
1980. Dissonance theory revisited: Perception of environmental hazards in residential areas. *Environ. Behav.* 12:33-51.
- Sigelman, L., and E. J. Yanarella.
1986. Public information on public issues: A multivariate analysis. *Soc. Sci. Quart.* 67:401-410.
- Sorensen, J. H., and D. S. Mileti.
1987. Programs that encourage the adoption of precautions against natural hazards: Review and evaluation. In N. D. Weinstein (editor), *Taking care: Understanding and encouraging self-protective behavior*, p. 208-230. Cambridge Univ. Press, Cambridge.
- Van Liere, K. D., and R. E. Dunlap.
1980. The social bases of environmental concern: A review of hypotheses, explanations and empirical evidence. *Public Opinion Quart.* 44:181-197.
- Weigel, R. H., and J. Weigel.
1978. Environmental concern: The development of a measure. *Environ. Behav.* 10:3-17.
- Weinstein, N. D.
1987. Studying self-protective behavior. In N. D. Weinstein (editor), *Taking care: Understanding and encouraging self-protective behavior*, p. 1-10. Cambridge Univ. Press, Cambridge.
- Winham, G.
1972. Attitudes on pollution and growth in Hamilton or "There's an awful lot of talk these days about ecology." *Can. J. Polit. Sci.* 5:389-401.
-



POSTER PRESENTATIONS



**ANIMALS AS LITTER VICTIMS AT THE
GERMAN NORTH SEA COAST**

G. J. Vauk and Eckart Schrey
North German Academy for Nature Conservation
Norddeutsche Naturschutzakademie
Hof Mohr, D-3043 Schneverdingen
Federal Republic of Germany

ABSTRACT

Over a distance of 65 km at 12 beaches of the German North Sea coast, 64 entangled vertebrates belonging to 14 species (mainly gulls, gannets, guillemots) were found dead from August 1983 to April 1988.

On the island of Helgoland, furthermore, 53 living seabirds belonging to 11 species (most gannets) were observed as entangled with remains of ships' litter. In at least 46 cases, plastic material was involved, so that the chance of survival of the animals must be rated as very low.

The numbers quoted must be considered as minimums, since not all animals affected or entangled with litter reached the shore or were found there. Also those animals which died as a result of litter ingestion are not listed.

MARINE DEBRIS AND EPIPELAGIC FISHES


Jo-Ann N. Kushima
Division of Aquatic Resources
State of Hawaii
1151 Punchbowl Street
Honolulu, Hawaii 96813, U.S.A.

and



Raymond P. Clarke
Southwest Fisheries Science Center Honolulu Laboratory
National Marine Fisheries Service, NOAA
2570 Dole Street
Honolulu, Hawaii 96822, U.S.A.

ABSTRACT

Two species of epipelagic fish caught in 1987 and 1988 by commercial fishing vessels operating in the vicinity of the Hawaiian Archipelago display the impacts of marine debris. The first specimen is a male mahimahi, *Coryphaena hippurus*, 77.5 cm total length and weighing 2.06 kg. Captured during albacore trolling operations, the mahimahi had monofilament net fragments attached to its gills and opercular area. The other instance involves a swordfish, *Xiphias gladius*, measuring approximately 140 cm total length and weighing 10 kg caught during tuna longlining operations. A rubber band cut approximately 3 cm deep into its caudal peduncle; otherwise, the swordfish appeared normal. Presented are photographs of both specimens and speculations as to the possible origins of the impacting debris.



VIDEO PRESENTATIONS



PORTRAIT OF A BARRIER ISLAND BEACH

Anthony F. Amos
Marine Science Institute
University of Texas
750 Channelview Drive
Port Aransas, Texas 78373, U.S.A.

ABSTRACT

Spanning a period of 11 years, 1,800 transects have been made along a 12-km stretch of beach on Mustang Island, Texas, to observe the seasonal and long-term changes to this barrier island beach bordering the northwest Gulf of Mexico. Documented in the ongoing study are bird populations, human disturbance, beach morphology, local oceanographic and weather conditions, stranded marine mammals, turtles, birds, oil spills, fish kills, effects of severe weather, and occurrence of marine debris and litter. This talk illustrates visually the tremendous impact, both aesthetic and as a danger to wildlife, of marine debris and increasing human usage on this otherwise beautiful beach. Also illustrated are several environmental and other factors that complicate efforts to understand the seasonal and long-term trends in the distribution of marine debris.


**U.S. OIL INDUSTRY EFFORTS IN ADDRESSING
BEACH DEBRIS PROBLEM**

Offshore Operators Committee
Amoco Production Company
P.O. Box 3092
Houston, Texas 88253, U.S.A.


ABSTRACT

The United States Gulf of Mexico offshore oil and gas industry has, over the past 2 to 3 years, embarked on an industry-wide effort to eliminate its contribution to marine debris. Beach debris surveys on gulf coast beaches previously identified a significant percentage of the debris originating from offshore exploration and production operations. As regulations which prohibited the discharge of trash from these facilities were already in place, it was thought that carelessness and possibly ignorance were involved.

It was evident that education would be the best way to approach this problem. This presentation will focus on a 12-min video developed by the offshore operators' committee, along with some individual efforts undertaken by companies, which we have used in the education process.



WORKING GROUP REPORTS



**REPORT OF THE WORKING GROUP ON METHODS TO ASSESS
THE AMOUNT AND TYPES OF MARINE DEBRIS**

(Christine A. Ribic, Chair)

INFORMATION NEEDS AND METHODOLOGIES

For the determination of the amount and types of marine debris, the working group distinguished between two types of studies: baseline, studies with low sampling frequency made over large geographic areas; and assessment, studies of a more limited area and having more intensive sampling effort over time. Baseline studies describe existing marine debris and seek to identify the magnitude of a problem. Assessments study the level of pollution.

The group considered various methodologies now in place for determining the amount of debris in the ocean (nearshore, open ocean, and bottom) and on beaches. They agreed that the beach survey is appropriate for assessment studies on a large scale. For limited-scale studies, dedicated surveys using visual observations and neuston tows in nearshore areas (e.g., bays, harbors, and estuaries) or limited ocean areas such as offshore dumping areas can be used for assessment.

Table 1 summarizes the current utility of survey techniques. Use of aircraft, while experimental, is feasible for baseline studies. Techniques to study bottom debris are needed; currently bottom debris studies are categorized as baseline.

Figure 1 is a proposed outline of the stages of a marine debris pollution assessment plan.

GEOGRAPHIC AREAS

The vastness of the oceans makes it necessary to select areas of interest for more intensive studies. On an international level, the working group suggested MARPOL special areas as appropriate geographical areas for more intensive study. On a national or a regional level, special areas of local interest must be developed. Examples of national level areas were the Pribilof Islands, because of the impact of debris on northern fur seals (United States), and national marine sanctuaries (United States). A regional area of interest cited was the Caribbean. Freshwater systems, including estuaries, were not discussed by the working group for lack of time.

Table 1.--Summary of survey techniques.

Survey techniques	Type of study	
	Baseline	Assessment
Nearshore/open ocean		
Surface debris:		
Visual observation (strip/line transect)	Platforms of opportunity	Dedicated surveys in well-defined areas of importance
Neuston nets	Oceanographic surveys	Dedicated surveys in well-defined areas of importance
Photography	Aircraft/heli- copter (limited to large debris items)	
Bottom debris:		
Survey of fishermen	Questionnaire (limited to certain types of debris)	
Bottom trawl	Limited to certain types of communities	
Remote operating vehicles	Expensive to use	
Beach surveys	Volunteer efforts (educational/ public relations)	Planned surveys Estimates of amounts on beaches--random sampling Changes over time--same beach or transect
	Low-flying aircraft	




Figure 1.--Components of a marine debris assessment plan (after G. Kullenberg et al. (1986) Mar. Pollut. Bull. 17:341).

Floating Debris at Sea

At sea, counts of floating debris are made using platforms of opportunity and dedicated surveys.

Visual observation from a viewing platform such as the flying bridge of a vessel is used when counting large debris. Most studies employ a strip transect method and count all debris sighted within a certain distance of the ship, using the glare-free side of the ship for observation. The width of the strip depends on the height of the viewing platform as well as on survey conditions (e.g., Beaufort sea state), and may change during the survey. All debris in the strip is assumed to be sighted. No one has done work on the probability of sighting different debris objects, and there are potential size and color biases that need to be evaluated. The length of a single transect varies as does final total transect length. The variable considered is usually a density estimate, number per square

kilometer or number per square nautical mile. The group recommended that two or three observers be employed in the survey. A single observer should use a strip width of 25 m or less. Calibration runs were recommended to estimate strip width, and experiments were recommended to investigate color and size biases and the probabilities of sighting different debris types.

There has been limited line transect work, but no formal analysis has been published. Problems persist with inaccuracies in the data, notably in the accurate determination of the distance of debris perpendicular to the ship. When accurate distance measurements can be made, the working group recommended the use of the line transect.

Neuston tows (necessarily strip transects) appear to be the most extensively used method for the study of particulate plastic and tar balls. The group agreed on the usefulness of such tows when made from dedicated survey vessels, but questioned whether neuston tows could be made successfully using platforms of opportunity. They require certain speeds--some devices can be used at speeds of only 3 kn or less; others at up to 7 kn--and the group questioned a captain's willingness to slow the ship down sufficiently to accommodate the towing device. Important to the success of a neuston tow is the estimate of time actually towing, or sweep efficiency.

The working group noted the possibility of using low-flying aircraft to survey nearshore areas for debris.

Debris on the Sea Floor

Little is known about bottom debris. Bottom trawls may be used from either dedicated survey vessels or platforms of opportunity to sample sea floor debris. The working group discussed bottom trawls for sampling debris, noting that this area has seen little work. They agreed that composition of debris is measurable using bottom trawl gear, but estimates of density are thought to be questionable.

Remote Operating Vehicles (ROV) were mentioned as a possible sampling tool, but it was agreed that this approach is too expensive for widespread use. Bottom drifter studies were also mentioned.

A potential source of information are fishermen whose gear has become entangled in sea floor debris. The working group recommended the development of a survey form to collect bottom debris from fishermen. This could be a starting point for collecting baseline information on bottom debris.

Debris on Beaches

Beach debris surveys can be carried out in designed or in volunteer programs. Standardization of beach surveys has been attempted for Alaskan beaches and English beaches, with the major difference being the sampling unit. In Alaska the sampling unit is the entire beach (at least 1 km in length). The sampling unit for the English beach is one transect per beach. Based on the working group discussion, it appears that the difference in sampling strategy stems from the types of debris found on the beaches. Entire Alaska beaches have to be surveyed in order to count the trawl web

that dominates the debris. In England, where most of the debris is plastic, transects are used because there is too much debris to count.

To avoid as much as possible counting debris that has originated on land, both approaches emphasize surveying beaches that are away from urban areas and have little recreational use. Beaches of sand and small gravel substrates with moderate slope were selected because they tend to collect debris. In England back beach areas were included; in Alaska they were not. Alaskan beaches facing the open ocean were used. Beaches with complicated topographic features such as partly sheltering reefs should be avoided, as should beaches known to be cleaned periodically. Other variables to consider are prevailing winds and accessibility.

If the intent of the study is to estimate the amount of debris on beaches in a given area, then random beach selection is important. In England a two-stage stratified random sampling scheme was used successfully. To detect changes over time, the majority of the group felt that selected beaches should be sampled repeatedly over time. The kind of change expected should be predicted, an appropriate variable defined to measure that change, and data collected to support or refute the prediction.

For baseline and some assessment studies, total amount, weight, and composition of the debris should be measured. Volume measurements were considered, but the working group felt that they would not be possible in all situations. For studies of changes in debris type over time, there was general agreement that the type of debris would determine whether changes in total amount or changes in composition were of more importance. In Alaska it was considered important to detect a change in the amount of trawl web. In England, composition and age structure of plastic containers were of prime interest. The important point in addressing the question of change over time is to define a variable of interest that can be measured.

For assessment studies, more work is needed to understand beach debris dynamics--for example, local currents and sinks for the debris as well as debris sources.

A suggestion was made to utilize low-flying aircraft, as some current surveys in Alaska are doing.

Debris Emanating From Land

The working group disagreed about the ease of distinguishing debris of land origin from debris originating on board ship. The accuracy of identification may vary from area to area. After some time in the water, debris items lose any paper labels and may acquire encrusting biota. Some items obviously originating on land may include plant seeds.

CATEGORIZATION OF DEBRIS

One suggestion was to categorize debris sizes as follows:

Mega - >2-3 cm
Macro - 5 mm to 2-3 cm

Meso - <5 mm (granule size)
 Micro - powdered (generally unseen).

The working group also made a list of some more common or important types to track. Suggested were:

Nets (by type)
 Other fishing gear
 Strapping bands (open/closed) (cut/uncut)
 Granulated plastic (recycled plastic)
 Particulate plastic
 Fragmented plastic
 Plastic bags
 Plastic containers (country of origin, age)
 Styrofoam
 Medical waste
 Rope
 Entanglement remains (e.g., bones)

Due to the time limitation, the group was unable to decide on broad categories for use in comparing data on an international scale, but recommended a review of existing categories in order to develop a common list that could be tailored to fit individual areas of interest.

MONITORING PROGRAM

Also because of a time limitation, the working group did not address the topics of sampling frequency and sample size requirements. A seasonable variation in the amount of floating debris was noted.

PROCEDURES MANUAL

The working group generally agreed that a procedures manual detailing survey techniques should be written. This manual would provide ideas for those interested in initiating marine debris studies.

LIST OF PARTICIPANTS

Anthony F. Amos, Marine Science Institute, University of Texas
 Douglas G. Chapman, University of Washington
 Trevor R. Dixon, The Tidy Britain Group
 Abraham Golik, Israel Oceanographic and Limnological Research, Ltd.
 Murray R. Gregory, Department of Geology, University of Auckland
 Burr Heneman, Marine Mammal Commission, Bolinas, California
 Kazuhiko Hiramatsu, National Research Institute of Far Seas Fisheries,
 Fisheries Agency, the Government of Japan
 David W. Laist, Marine Mammal Commission, Washington, D.C.
 Julio M. Morrell, Department of Marine Science, University of
 Puerto Rico-Mayaguez
 Christine A. Ribic, Center for Quantitative Science, University of
 Washington
 Satoru Taguchi, Department of Oceanography, University of Hawaii
 Nobuyuki Yagi, Division of Fishing Ground Environment Conservation,
 Fisheries Agency, the Government of Japan

REPORT OF THE WORKING GROUP ON ENTANGLEMENT OF MARINE LIFE

(William R. P. Bourne, Chair)

TERMS OF REFERENCE

It was reported at the first conference on marine debris in 1984 that individual seals, turtles, birds, and fish, some belonging to endangered species, become entangled at times in marine debris. The frequency and severity of these interactions were usually unknown, and no conclusive evidence was demonstrated for any effects on populations. This working group was asked to review the problem and identify the information needed to fill the gaps in current knowledge, notably by devising a model in the light of which current information could be assessed. This should include 1) age, 2) sex, 3) population, 4) numbers, 5) distribution, 6) legal status of victims, 7) activities and materials causing problems, 8) information that is needed to complete the picture and monitor its future development, and 9) the priority that should be given to different aspects of the investigations.

THE WORKING GROUP

The findings at the 1984 conference still seemed valid, so in order to avoid repeating preconceived ideas, the working group first considered materials which cause problems and their impact on different animals. The previous working group reports were then reviewed to see what progress has been made. The first conclusion is that, despite the accumulation of circumstantial evidence that marine debris may have an adverse effect on all sorts of marine wildlife (including cetaceans, which did not receive much attention previously), the information is still insufficient to show clearly the magnitude of the problem.

LIMITATIONS OF THE DATA

Care is needed in the interpretation of the available information. It is substantial only for the most common species of two out of the four main groups affected, pinnipeds and birds. These spend long periods on land at breeding sites, where they can be counted and marked. Even here information is deficient for the important period spent at sea. Much less is known about the turtles, which spend most of their time at sea, and the cetaceans, which never come ashore at all. Owing to the way in which the information is collected, it still remains difficult to distinguish between the effects of a variety of interacting factors. These include oceanic fluctuations, disturbance while breeding, the impact of fishing on both the animals and their food supply, disease, and pollution.

THE NATURE OF THE PROBLEM

Human artifacts must have caused hazards for wildlife since man first went to sea. They cause two types of problem: killing animals in unpleasant ways, a problem whose nature is self-evident so it will not be discussed further here, and harming the status of species and ecosystems. Both uncontrolled hunting or fishing and the incidental capture of unintended animals while taking other species (by-catches) potentially cause conservation problems. Although such factors lie outside our terms of reference, it must be remembered that not only may the losses of marine animals from hunting and fishing be only a fraction of the losses from natural causes, but losses due to marine debris may also be only a fraction of those from hunting and fishing. In such cases, entanglement may be important as a separate entity only to man, because the animals that it kills are deducted from his catch. The situation starts to acquire a wider significance only when the mortality of any animal becomes large enough to affect the welfare of the species and the ecosystem as a whole as well.

Nets

In the past, netting was made of natural materials which were not very durable, so that lost nets were noticed to cause a problem only by fouling vessels. Following the introduction of nylon and other synthetic netting after World War II, there have been growing problems with both lost or abandoned nets and the fragments torn from them by obstructions, discarded during repairs, or used to make small traps which subsequently disintegrate. Some reference collections of different types of netting have been made in the hope of identifying their origin--for example, where strange net which presumably comes from the tropical Atlantic washes ashore in the West Indies. It was thought that it might be useful to consult the fishing industry about the preparation of a guide to different types of fishing gear and its likely origin.

It was considered whether nets should be constructed of, or fastened together with, more rapidly degradable material, so that they would break up and sink sooner when lost. However, it was thought that this would lead to the production of more small fragments of net on the sea floor and along the shore, which would add to the problem. Experiments suggest that, if left intact, whole nets tend to bunch up and may cause less problem for wildlife. Thus it may be better to try and keep the net in one piece so that it is more easily removed or immobilized. It has been reported that floats of unequal size drift at different rates with the wind and current, keeping the net open, so the effect of the buoys and weights used on the performance of nets may deserve more attention.

There still appears to be a need for further study of the way in which animals are caught in nets and the fate of lost nets, using marked trial specimens to see how long they continue ghost fishing at sea and whether they catch more or different animals as they come ashore.

Hooks and Line

These are one of the commoner agents ensnaring wildlife and, occasionally, people. Discarded angling gear often hooks or ensnares birds. Fishing has been banned from some North Sea oil platforms because lost gear has caused problems for divers. Here also the situation has been made worse by the adoption of durable nylon line. It seems likely that the longlines used at sea, which may extend for up to 96.5 km (60 mi), with several thousand hooks and few buoys, may present a more serious problem. Little appears to be known about this, and it deserves more study.

Loops, Sheets, and Sharp Objects

Other potentially dangerous objects may appear occasionally at sea. They range from wrecks and heavy machinery and construction material discarded by the oil industry, through nautical gear and packing materials and containers, to clothing and household equipment. It seems easier to consider the nature of the hazard that they present than the identity of the object concerned.

Anything which incorporates a loop or ring is potentially hazardous. This includes knotted rope, uncut packing bands, containers with holes, loose webbing or fabric, plastic sheeting, and, of course, netting. These present lethal threats to any marine animal up to the largest whale, should their head, jaws, or limbs become ensnared while hastily seizing mobile prey, or hunting and playing around drifting material. Such objects should be disposed of ashore or by incineration.

Anything which includes, or can break down to form, a sharp point or cutting edge also presents a hazard, especially if it is concealed among innocuous materials. Such objects include wrecks and dumped heavy equipment which may catch nets on the sea floor, lesser metalwork, woodwork with projecting nails, tins opened to leave sharp edges, and fragile glass containers. In addition to being a threat to wildlife, these are also a threat to people who are diving, hauling nets which have collected debris, or walking along the shore. All such objects should also be disposed of carefully.

It seems desirable to redesign some objects which regularly cause serious problems. These include perforated plastic six-pack yokes, cans which normally have sharp edges when opened, and openers which leave cans with sharp edges.

VULNERABLE ANIMALS

There is accumulating anecdotal evidence that virtually all marine animals are occasionally entangled in debris, but quantitative data are available for few of them. The main cases where it appears important follow.

Cetaceans

Entanglement appears to be unusual and to be reported most often among the smaller species which are found near the shore. Its impact might be most serious with the North Atlantic right whales, *Eubalaena glacialis*. This is a large whale population reduced to a remnant of a few hundred by commercial fishing which has failed to recover despite half a century of protection, and some are known to be entangled occasionally in nets (Report of the International Whaling Commission, Special Issue 10, p. 116-119, 1986). Nearly a third are also said to be scarred from unknown causes, which might include other whales (notably killer whales), rocks, collisions with ships, and fishing gear, since they frequent areas with concentrations of plankton and these areas are often important fisheries. It seems time that these whales received more study.

Phocid Seals

While these seals are occasionally entangled in netting, the incidence is not usually high. It appears to be worst in the endangered warm water monk seal, *Monachus* sp. The only surviving species in the Mediterranean and North Pacific are both reduced to hundreds. A number of Hawaiian monk seals, *M. schauinslandi*, are known to have been killed by net fragments along the shore, and nearly a quarter of the mortality reported in Greece was also found to be due to fishing gear (J. Jacobs and A. Panou, Conservation of the Mediterranean monk seal, *M. monachus*, in Kefalonia, Ithaca and Lefkada Islands, Ionian Sea, Greece, Institute of Zoology, University of Munich, Seidlstrasse 25, D-8000 Munchen 2, Federal Republic of Germany, 221 p., 1988, per D. E. Sergeant). These species clearly deserve more attention.

Otariid Seals

While many populations of these seals were reduced by exploitation for skins and oil in the past, most now appear to be recovering. Possibly owing to their large numbers, their tendency to feed in areas with important fisheries, and their active behavior and slender physiques, they are also among the marine mammals most prone to entanglement. Young animals which play around nets seem particularly vulnerable. While most species are maintaining their numbers, the northern fur seal, *Callorhinus ursinus*, has been declining in some areas in recent years. A number of other factors which are difficult to study, such as overfishing and climatic change, may also be involved. It seems desirable to continue monitoring the breeding populations, investigate the animals' movements and relationship to nets at sea, and compare the results with those for increasing populations.

Chelonians

Most of the turtles now appear to be endangered, but while they do become entangled occasionally, there appears to be no evidence that this is having any impact on their numbers comparable to such factors as over-exploitation for shell, meat, and eggs, disturbance of the breeding

habitat, losses in fishing nets, and ingestion of plastic material. Pelagic ridley turtles, *Lepidochelys* sp., may be the most vulnerable. The depleted Kemp's ridley turtle, *L. kemp*i, is confined to one breeding beach in the Caribbean; the olive ridley turtle, *L. olivacea*, has been found entangled in the Pacific.

Birds

Birds become entangled in nets, hooks and line, and other debris occasionally, and the reported incidence in British Trust for Ornithology (BTO) banding recoveries of the common guillemot or murre, *Uria aalge*, has increased from 5% before 1970 to 37% since 1987 (C. Mead, BTO News 163, 1989). There is no evidence yet, however, that entanglement is having an important impact on bird numbers when compared with disturbance by man and introduced predators at the breeding places, or losses due to active fishing gear or oil pollution. Some species, especially the Pelecaniformes, are also vulnerable when, to make their nests, they collect floating material which may ensnare either the old birds or their young. The species for which there is the most evidence of damage from pollution of all kinds, the northern gannet, *Sula bassana*, is nonetheless increasing explosively in most areas, even at a small colony on Flamborough Head, England, where more than half the nest material is composed to nylon netting, and where many birds also become entangled at sea.

Fish and Shellfish

While other marine animals may become entangled in debris occasionally, there is no evidence that the resulting mortality amounts to more than a small fraction of that due to fishing. Debris-related mortality therefore seems most important as a loss from fisheries, as discussed by the working group on ghost fishing.

PRIORITIES

The group was asked to arrange its recommendations in order of their importance. Of highest priority is the collection of more information, arranged to cover as many areas and aspects of the subject as possible. There are still many important gaps in the available information, including several inadequately studied major groups of animals where the losses may be important, such as the cetaceans, sirenians, and chelonians. There are also several inadequately studied potential problems, such as long-lines and sharp objects. A large part of the world is still inadequately covered--this conference has lacked any direct representation from not only the Communist and developing nations, but also South America and Australia.

Organization

It was thought that a more permanent organization is required to obtain information from more places. This should be composed of a limited number of representatives who are active in research on different animals, on different aspects of the problem, and in different areas. Its purpose

should be to expand the sphere of activity and maintain more continuity and consistency in recording methods. This would require the identification and enlistment of suitable people, whether in government organizations with their own resources, or voluntary bodies. Providing limited help with administrative expenses and the cost of attending meetings would be useful.

Information

Even in areas where there is already an interest in problems caused by marine debris, there is still a need for more means of circulating information and advice. This could include such matters as the identification of the materials and species encountered, the examination of stranded animals, and the best ways to record comparable observations. It would be useful to have a simple field guide to introduce more people to the subject, supplemented with a newsletter to report further progress and results. (The Marine Pollution Bulletin would be happy to assist.)

Research and Conservation

For purposes of economy it seems desirable to devise proposals that will cover several objectives simultaneously. These should include as far as possible the most vulnerable species in each of the main groups of animals, the most sensitive areas, the most critical threats, and mitigating measures. Five projects which between them might cover most aspects of the subject are monk seals, fur seals, right whales, sea turtles, and man.

Monk Seals

The marine animal for which entanglement appears to pose the worst threat is the Hawaiian monk seal in the Northwestern Hawaiian Islands (NWHI), where several are known to have died as a result of entanglement in stray fragments of net along the shore and where there appears the best chance of practical action to alleviate the threat of entanglement. This area is also important in several other respects. It is one of the first nature reserves of international importance. In addition to studying the impact of entanglement on this most vulnerable phocid seal, investigations there could also cover the impact of debris on a variety of other wildlife, including entanglement of sea turtles and ingestion by albatrosses and other seabirds, in a remote situation in the tropical Pacific. Measures should include the regular collection, evaluation, and destruction of debris on both the beaches and outlying reefs, the liberation of all live entangled animals, and studies of dead ones. The situation of the even rarer Mediterranean monk seal also needs further study, which could also be integrated with a study of related issues in a much more heavily developed area.

Fur Seals

The other marine animal where there is already evidence of serious mortality from entanglement is the northern fur seal. It also inhabits

established nature reserves of international importance. The recovery of its original vast numbers was previously a cause for general satisfaction, and any continuation of its recent decline would cause wide concern. It differs from the preceding species in its more migratory behavior, and may be encountering its most serious problems at sea away from the breeding places. At present it still has a much larger population than the endangered phocids, representing the other main group of otariid seals, so it can be studied more actively with less risk of serious disturbance and might provide interesting comparative results. It is important to continue monitoring the breeding populations, and investigations should be extended out to sea on a larger scale, both tracking animals on their feeding movements and migrations (notably by satellites), and studying their reaction to nets at sea. Such investigations should also yield useful general information about the welfare of other wildlife and the impact of marine debris further north in the Pacific.

Right Whales

The cetaceans for which there is possibly most cause for concern are the North Atlantic right whales (though most right whale stocks are depleted). These provide an instructive contrast to the previous species, since they are much larger, yet feed on plankton in an area with active fisheries in another ocean. Although they were originally very numerous, they were the first species seriously reduced by modern whalers and have failed to recover after half a century of protection. A certain amount is already known about them, such as the location of a small population with many scarred individuals, some of which are occasionally killed in nets. Humpback whales are regularly caught (and sometimes killed) in coastal nets in the same area, so the right whales may be encountering similar problems out at sea. It therefore seems desirable to learn more about the extent to which entanglement is a problem for this species. This might also reveal useful general information about the impact of marine debris on the most important fishing grounds in the northern Atlantic.

Sea Turtles

It is doubtful that entanglement is as important a cause of mortality for turtles as ingestion of debris, and it might be better to investigate the two problems together. One approach might be to study the behavior of turtles in captivity when confronted experimentally with debris. It might also be useful to try to trace individuals from satellites. The most vulnerable species, which might merit attention at an early stage, appear to be the ridley turtles. Any investigation of Kemp's ridley turtle might also yield useful general information about debris-related problems in the Caribbean. It might also be possible to integrate any investigation of the green turtle with research on the Hawaiian monk seal to obtain a better picture of events at sea in actively exploited tropical waters.

Man

It is surprising that there seems to have been little attention paid to the animal whose welfare is of the widest general interest. A certain

amount of harm must be caused to people by marine debris such as netting, hooks, lines, and sharp objects. It seems time for an assessment of the risks posed by various categories of debris on beaches, in shallow waters, and brought up by trawls. It might prove instructive to carry out a trial survey among medical personnel, sailors, fishermen, and divers to discover whether they can supply any information about the incidence, nature, and cost of human injuries due to marine debris. It is possible that marine debris may also cause occasional human fatalities, either directly or by disabling boats, and if so it seems desirable to assess their frequency.

RECOMMENDATIONS

Specific recommendations included:

- Continued monitoring, removal, and destruction of lost or discarded nets and other debris presenting a hazard to monk seals, green turtles, and other wildlife in the NWHI, extending the work to the outlying reefs.
 - Continued monitoring of the numbers, survival, breeding success, and incidence of entanglement of northern fur seals, extending the observations out to sea.
 - Investigation of the impact of entanglement and other possible hazards on right whales in the northwest Atlantic and Kemp's ridley turtles in the Caribbean.
 - A review of the long-term evidence for entanglement provided by bird banding and beach surveys.
 - A survey of the injuries caused to man by marine debris.
 - Observations of the movements and behavior of seals and turtles at sea using satellites.
 - Collection of more information about net use and losses, and means of identifying the origin of different types of net.
 - Studies of the movements and fate of marked debris, including nets, with further observations of the way in which animals react to debris at sea.
 - Reviews of experience with voluntary beach cleaning, artificial reefs, and material left on the sea floor by the oil industry.
 - The use of models to determine the population dynamics of different animals, the way in which processes affecting them are likely to operate, and the best data to collect to elucidate them.
-

- The formulation of standard recording techniques for different types of debris and victims of entanglement in order to facilitate the more systematic collection of records of entangled animals and fouled vessels.
- The preparation of a guide to types of lost or discarded nets and other debris, and the best ways to examine and treat entangled animals and record observations.
- The dissemination of warning against the particular hazards posed by rings and loops, especially uncut packing bands.
- The redesign of six-pack can yokes, so that they are broken up in use, and methods of opening cans, so that they do not leave sharp edges.
- It was concluded that, in view of the number of problems that require investigation and the wide area that needs to be covered, there is a growing need for the establishment of a representative international organization to coordinate the systematic collection and circulation of information about the occurrence and impact of artificial marine debris and possible conservation measures to mitigate its ill effects.

LIST OF PARTICIPANTS

Norihisa Baba, National Research Institute of Far Seas Fisheries,
Fisheries Agency, the Government of Japan

Nelio B. Barros, Rosenstiel School of Marine and Atmospheric Science,
University of Miami

William R. P. Bourne, Department of Zoology, Aberdeen University

Paul A. Breen, New Zealand Ministry of Agriculture and Fisheries

Douglas G. Chapman, University of Washington

John Davis, Canada Department of Fisheries and Oceans

Robert L. DeLong, National Marine Mammal Laboratory, NMFS, NOAA

Charles W. Fowler, National Marine Mammal Laboratory, NMFS, NOAA

Deborah P. French, Applied Science Associates, Inc.

John R. Henderson, Southwest Fisheries Science Center Honolulu
Laboratory, NMFS, NOAA

Minoru Morimoto, Ministry of Agriculture, Forestry, and Fisheries,
Fisheries Agency, the Government of Japan

Samuel S. Sadove, Okeanos Ocean Research Foundation, Inc.

Brent S. Stewart, Sea World Research Institute, Hubbs Marine Research Center

Gordon L. Swartzman, Center for Quantitative Science, University of Washington

Yasuji Tamaki, Fisheries Agency, the Government of Japan

REPORT OF THE WORKING GROUP ON GHOST FISHING

(Paul A. Breen, Chair)

OVERVIEW

Ghost fishing is a potentially serious problem because of the very large volumes of fishing gear now in use. Only a small percentage of this gear lost annually would amount to a very large loss. The increasing use of nondegradable materials such as plastic, vinyl-coated wire, and fiberglass means that lost fishing gear may persist in the marine environment for a long time.

Of the many gear types in use worldwide, the working group considered traps and gillnets to be of primary interest. Ghost fishing is well documented in coastal gillnets and in a few studied trap fisheries; it is much less well documented in pelagic gillnet fisheries. For most trap fisheries, no directed work has been done and whether ghost fishing takes place is not known. Much more work is required to study ghost fishing in specific trap fisheries and in pelagic gillnet fisheries.

Trawls and longline gear types probably cause smaller ghost fishing problems than do traps and gillnets. For other gear types, no evidence exists that ghost fishing takes place.

For American lobsters, the estimated economic waste is several million dollars annually. In other trap fisheries where it has been measured, the loss to ghost fishing appears to be a significant percentage of the reported catch. It seems certain that if other trap fisheries were examined further, serious ghost fishing situations would be discovered.

Mitigation of ghost fishing is technologically simple for traps, but requires situation-specific materials research, legislation, and industry education. Mitigation of ghost fishing by nets is more difficult. Both timed-failure devices and degradable meshes should be developed and tested for nets.

The lower priority problems of ghost fishing by trawls and longlines are poorly documented.

GEAR TYPES

The working group reviewed the various gear types in use with respect to their potential for ghost fishing. Traps, tangle nets, and pelagic gillnets were considered to have the highest potential for ghost fishing because of their passive mode of fishing and the very large quantities

presently in use. Bottom trawls and coastal gillnets were considered to have the next priority. Longlines, both benthic and demersal, and lures were considered at a third level of importance.

With the many other gear types in use, the working group has no reason to suspect significant ghost fishing problems.

FRAMEWORK

The working group devised the following framework for summarizing and evaluating the existing information on ghost fishing impacts. The following questions apply to each species, gear type, and location problem. The working group recognized dangers in generalizing or extrapolating from one species or trap type to another.

- Does ghost fishing take place in a particular situation; i.e., for a particular species/gear type/location combination?
- At what rate does lost gear catch and kill target and nontarget species?
- How is gear lost?
- At what rate is gear lost, or alternatively, how much lost gear is there?
- At what rate does lost gear cease to fish?
- What actions have been taken to reduce gear loss?
- What actions have been taken to reduce the ghost fishing life span of fishing gear?
- What actions have been taken to enhance the recovery of lost gear?

ANALYSIS

The working group reviewed the existing data and analyses by gear type in the context of the framework devised. The analysis is presented in the format outline above. Further information on traps and gillnets can be found in the review by Breen (1990 [this document]).

Traps

Does Ghost Fishing Take Place?

Ghost fishing has been documented in traps in the fisheries for American lobster and Dungeness crab through simulated lost trap studies. For the Western Australian snapper fishery, ghost fishing is suggested by a preliminary lost trap study. Ghost fishing seems likely from observations in fisheries for king crabs, snow crabs, and Pacific sablefish.

Observations and short-term escapements have been used by some workers to suggest that ghost fishing does not occur, for instance in Western Australian snapper. However, the working group considers the published evidence inadequate to reject the hypothesis that lost traps ghost fish for any species.

For most of the world's trap fisheries, the question of ghost fishing is not addressed by published reports. From studies of the few species so far examined, it seems likely that many more trap fisheries suffer from ghost fishing.

At What Rate Does Lost Gear Catch and Kill Target and Nontarget Species?

The rate of capture by lost gear has been measured in American lobsters at 10% of the catch rate by the commercial fishery. For Dungeness crabs in a sheltered British Columbia bay, lost traps killed 10 crabs per trap year. For Atlantic snow crabs, a lost trap fishes only for the life of the bait, then stops fishing. No capture rate estimates are available for other species.

How Is Gear Lost?

Trap gear is lost in a myriad of ways. Vessel traffic and tow boating sever buoylines or drag traps into deep water. Buoylines chafe and break. Buoys may become detached, or can be attacked by marine birds or mammals.

Storms and strong currents may "drown" traps. Traps may be snagged on rocky bottom. Traps are carried away by trawlers or gillnetters. Buoylines are cut by vandals or in fishing disputes.

At What Rate Is Gear Lost, or How Much Lost Gear Is There?

Estimates of annual trap loss rates vary from 5 to 30% based on estimates made from surveys of fishermen (see Breen 1990). For Dungeness crabs annual estimates vary from 11 to 18%. These rates are probably typical of most trap fisheries. In two surveys of Dungeness crab fishermen, it was estimated that about 50% of lost traps continue to ghost fish.

In the Alaska king crab fishery, it is estimated that 30,000 lost traps remain on the fishing grounds. In the U.S. portion of the American lobster fishery, it was conservatively estimated in 1978 that 187,000 traps could be ghost fishing.

No estimates are available for other trap fisheries.

At What Rate Does Lost Gear Cease to Fish?

Traps without timed-failure devices might ghost fish for years. Treated wooden lobster traps may last 2 years; metal king crab traps may last 10-15 years. No experimental results are available.

What Actions Have Been Taken to Reduce Gear Loss?

To reduce trap loss rates, some jurisdictions have regulations that require traps to be buoyed with marked buoys. In some areas, seasonal and area closures create temporal or spatial separation of trap and other fisheries. In Washington State, buoys must be foam-filled and buoylines weighted to prevent losses from buoys sinking or vessels running over buoylines. In Washington State, trap fishermen are notified of potential gear conflicts. High technology navigation systems allow trap gear to be relocated with more precision. Educational programs have reduced gear loss.

All these actions are taken from American lobster and Pacific west coast jurisdictions. Apart from buoyage and marking requirements, little is known about actions to reduce trap loss in fisheries outside North America.

What Actions Have Been Taken to Reduce the Ghost Fishing Life Span of Fishing Gear?

Devices to reduce the ghost fishing life span of a trap are required in all traps in Alaska, Washington, Oregon, and California; in Pacific sablefish traps in Canada; and in American lobster traps in Connecticut.

Actions taken to prevent ghost fishing outside North America have not been published.

What Actions Have Been Taken to Enhance the Recovery of Lost Gear?

The working group is aware of no programs to recover lost traps, except that a small-scale commercial operation once operated in Canada to recover Dungeness crab traps. In Alaska, king crab traps caught by domestic trawlers are slashed before being discarded. Trap recovery is probably opportunistic.

Pelagic Gillnets

Does Ghost Fishing Take Place?

Ghost fishing in a pelagic salmon gillnet was reported in a lost net recovery, and ghost fishing was documented in a simulated lost net study.

At What Rate Does Lost Gear Catch and Kill Target and Nontarget Species?

In one experimental simulation of lost pelagic gillnets, two fish were caught by 1,500 m of net in the first 3 days.

How Is Gear Lost?

Pelagic gillnets may be lost when cut by vessel traffic, broken by storms or large marine mammals, or when the fishing vessel fails to relocate her gear.

At What Rate Is Gear Lost?

The rate of gillnet loss has been estimated in one study at 0.05% per set. This estimate, from the Japanese salmon mothership fishery, appears to be the only available estimate.

Density of lost gillnet fragments observed from passing vessels has been estimated in several studies presented at this conference.

At What Rate Does Lost Gear Cease to Fish?

Two studies suggest that nets less than 2 km long fish for only a short time after loss and then rapidly aggregate into a solid mass. The net tends to remain open longer when attached to a large buoy.

What Actions Have Been Taken to Reduce Gear Loss?

Japan requires gillnets to carry radar reflectors at each end of the unit to prevent cutting by vessel traffic. Radio communication is used to direct vessel traffic around nets. To prevent loss in bad weather, shorter sets are made. Japan requires nets to be marked with radio transmitters, and old gillnets to be recycled. No information is available from other countries.

What Actions Have Been Taken to Reduce the Ghost Fishing Life Span of Fishing Gear?

No actions have been developed to reduce the ghost fishing life span of a pelagic gillnet.

What Actions Have Been Taken to Enhance the Recovery of Lost Gear?

Japan requires nets to carry radio buoys and radar reflectors, and requires old nets to be recycled. Japanese research vessels pick up fishing debris.

Coastal Gillnets**Does Ghost Fishing Take Place?**

Ghost fishing by Pacific salmon gillnets has been documented, and observations of ghost fishing by Pacific herring gillnets have been reported. Experimental results confirm ghost fishing in demersal gillnets in Newfoundland, New England, and New Zealand.

At What Rate Does Lost Gear Catch and Kill Target and Nontarget Species?

Catch rates of lost gillnets have not been estimated in any published study.

How Is Gear Lost?

Coastal gillnets are lost when nets become fouled on the bottom or on snags; broken by storms, marine mammals, or large fishes; cut by vessel traffic; or carried away by trawlers.

At What Rate Is Gear Lost?

Rates of gear loss for coastal gillnets are not immediately available. A submersible survey in a known area in New England found a density of 0.23 nets/ha. A Newfoundland study reports numbers of gillnets retrieved in direct retrieval operations.

At What Rate Does Lost Gear Cease to Fish?

Lost gillnets may become tangled (leadline over corkline) or balled up (tangled in the horizontal plane). Fouling increases visibility and reduces catches. No precise estimates of the rates of these processes are available. Ghost fishing has been observed in Pacific herring gillnets 7 years after net loss.

What Actions Have Been Taken to Reduce Gear Loss?

Most jurisdictions require proper marking and lighting of gillnets to prevent cutting by vessel traffic. Radar reflectors are required on gillnets in New England.

What Actions Have Been Taken to Reduce the Ghost Fishing Life Span of Fishing Gear?

Recent New England experiments have examined degradable corklines and the effect of degradable panels along the net.

What Actions Have Been Taken to Enhance the Recovery of Lost Gear?

A Newfoundland program was conducted in 1975-76 to recover lost gillnets with specially designed recovery gear. In the British Columbia herring fishery, efforts are made to ensure that all gear has been recovered at the end of an open fishing period.

Bottom Trawls

Does Ghost Fishing Take Place?

Ghost fishing has been reported where trawl netting was stretched across bottom features or snags.

At What Rate Does Lost Gear Catch and Kill Target and Nontarget Species?

No estimates of catch rates by lost trawls are available.

How Is Gear Lost?

Trawls are lost when the net or doors become snagged on bottom obstructions. Snagging incidents may result in partial loss of the net. Trawls have been lost when fouled by submarines.

At What Rate Is Gear Lost?

No rates of trawl loss are immediately available. Some logbook programs may contain this information.

At What Rate Does Lost Gear Cease to Fish?

No information.

What Actions Have Been Taken to Reduce Gear Loss?

Snag charts reduce the incidence of net loss on wrecks or other obstructions. High technology navigation systems allow trawls to be set more accurately in known areas.

What Actions Have Been Taken to Reduce the Ghost Fishing Life Span of Fishing Gear?

There appear to be no actions taken with respect to trawls.

What Actions Have Been Taken to Enhance the Recovery of Lost Gear?

The working group uncovered no information on this question.

Longlines**Does Ghost Fishing Take Place?**

Pacific halibut are reported to strike and be caught on bare hooks. Lost halibut longlines may thus ghost fish.

At What Rate Does Lost Gear Catch and Kill Target and Nontarget Species?

No estimate of the rate of ghost fishing by longlines is available.

How Is Gear Lost?

Longlines are lost when snagged on bottom features.

At What Rate Is Gear Lost?

No estimates of loss rate or density of lost gear are immediately available. Some logbook programs may contain information.

At What Rate Does Lost Gear Cease to Fish?

There appears to be no information with respect to longlines.

What Actions Have Been Taken to Reduce Gear Loss?

There appears to be no information with respect to longlines.

What Actions Have Been Taken to Reduce the Ghost Fishing Life Span of Fishing Gear?

There appears to be no information with respect to longlines.

What Actions Have Been Taken to Enhance the Recovery of Lost Gear?

There appears to be no information with respect to longlines.

RECOMMENDATIONS

The working group made recommendations at three levels of priority: high, medium, and low. Within each level no attempt was made to assign priorities.

High Priority Recommendations

1. Fishery agencies responsible for trap and tangle net fisheries should conduct lost gear simulations to determine whether ghost fishing occurs and, if it does, the rate at which target and nontarget species are killed. If a ghost fishing problem is discovered, the rates of gear loss should be estimated through logbook programs or questionnaire surveys. In some situations, useful information might be obtained from surveys of fishing gear manufacturers.
2. Where ghost fishing has been demonstrated or is suspected in a trap fishery, the fishery agency should decide what timed-failure mechanism would be most appropriate to reduce the life span of traps and how soon to cause timed failure to happen. Research under actual fishing conditions should then be conducted to determine the most appropriate regulation for timed-failure devices. Industry should be consulted and involved in this research.
3. Further studies with simulated lost pelagic gillnets should be conducted. In order to simulate the loss of an entire net, studies should use nets approximating the length of commercial nets. More studies on smaller sections are also required. These studies should examine whether ghost fishing takes place and, if it does, then the rate of ghost fishing and the rate at which the nets form a tangled mass or otherwise cease to fish.

4. Direct observations should be made of lost pelagic gillnets to determine their shape and to determine the apparent rate at which ghost fishing for fish, birds, sea turtles, and marine mammals is taking place. These observations should be collated, made available, and distributed by a central agency such as the Food and Agriculture Organization of the United Nations.

Medium Priority Recommendations

1. Research should be continued and new programs developed to examine potential timed-failure mechanisms in gillnets and tangle nets. Both degradable net components and degradable mesh should be considered. In the former case, possible ghost fishing by sunken gillnets must be evaluated with appropriate experiments. In the latter case, the consequences of generating many small fragments must be examined. This research should also address the costs of timed-failure mechanisms for gillnets and tangle nets.
2. Fishery agencies should examine existing data or undertake new programs to estimate the rate of gear loss in fisheries using pelagic or coastal gillnets, trawls, or longlines.
3. Where ghost fishing has been demonstrated or is suspected with any gear type, the responsible fishery agencies should conduct research into the fishing life span of gear after loss.
4. In those fisheries for which an estimate of the impact of ghost fishing is available, ghost fishing should be examined as a mortality source in stock assessments and incorporated in fishery management plans.
5. In those jurisdictions where timed-failure devices are already required in traps, the rate of compliance with such regulations and attitudes of industry should be examined.

Low Priority Recommendations

1. Studies should examine whether ghost fishing takes place by longline gear, especially for Pacific halibut and tunas but also for other species as appropriate. If ghost fishing does occur, then studies should be conducted to measure the rate at which hooks of various kinds cease to catch fish.
 2. Where it has not been done, charting of snags should be carried out to help vessels prevent net loss.
 3. The possibility of encouraging or requiring vessels to retain recovered lost fishing gear for disposal on land should be explored.
-

4. Research should be initiated on possible positive effects of lost gear, especially lost traps acting as habitat for American lobster and floating masses of pelagic gillnet acting as fish aggregating devices.

LIST OF PARTICIPANTS

Norihisa Baba, National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan
Paul A. Breen, New Zealand Ministry of Agriculture and Fisheries
Douglas G. Chapman, University of Washington
John Davis, Canada Department of Fisheries and Oceans
Satsuki Matsumura, National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan
Tracie McKenzie, Northeast Region, NMFS, NOAA
Minoru Morimoto, Ministry of Agriculture, Forestry, and Fisheries, Fisheries Agency, the Government of Japan
Joji Morishita, International Affairs Division, Fisheries Agency, the Government of Japan
David E. Ortman, Friends of the Earth
Yasuji Tamaki, Fisheries Agency, the Government of Japan
Yoh Watanabe, National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan

REPORT OF THE WORKING GROUP ON INGESTION

(Louis Sileo, Chair)

This report includes a summary of the information about ingested plastic presented during the technical sessions and a summary of the working group's discussions. Both are organized by taxa and deal with the prevalence and effects of ingested plastic.

SUMMARY OF TECHNICAL SESSIONS

Sixteen papers concerning ingestion of plastic were presented during the technical sessions. Five of these dealt with birds, two with fish, five with marine turtles, and four with marine mammals. There was one review paper for birds and one partial review each for fish and turtles. The majority (13) of the reports were of a descriptive or anecdotal nature. The latter are very useful for gathering baseline information for hypothesis generating and for defining and attracting attention to an emerging problem. Such anecdotal reports often show associations between observations, such as emaciation and the presence of plastics in stranded marine animal carcasses. However, with anecdotal data it is not possible to determine if such an association is coincidental or cause and effect. Only 4 of the 16 papers reported work with controlled experiments designed to test a hypothesis. There is need for more such studies designed to test hypotheses about the possible cause-effect nature of associations revealed by the anecdotal studies.

Most (14) of the papers presented information about the prevalence of ingested plastics: 9 papers introduced new data about the effects, usually harmful, of ingested plastic on individual animals. There were no data about those effects on the population dynamics of any species, nor about absorption of toxins from ingested plastics.

The nature and extent of the data presented in the technical sessions were summarized by taxa (Appendixes A to D); these summaries provided a basis for the working group's discussions. Crucial knowledge deficiencies were defined by the group and then given priority (Table 1). All priorities were reached by consensus. The areas of expertise of the scientists attending the working group sessions provided an equitable representation of the various taxa.

Future studies should have statistically adequate sampling schemes designed to test hypotheses that the prevalence is increasing or decreasing in given areas or taxa. Future studies of the effects of ingested plastics should also include statistically adequate experimental designs for testing

Table 1.--Knowledge deficiencies and informational needs arranged in order of importance.

Priority	Information needed
First	Effects on marine turtles
Second	Effects on seabirds
Third	Prevalence in marine turtles
Third	Prevalence and effects on manatees
Fourth	Effects on large fish
Fourth	Prevalence in marine mammals
Low	Effects on larval fish
Low	Prevalence in fish
Low	Prevalence in seabirds
Low	Effects on marine mammals

hypotheses. It is possible that estimates of variance from studies already completed will provide the basis for determining sample sizes required for statistical significance in future studies.

Regardless of the taxon, the same three general pathophysiological effects were proposed: (1) mechanical blockages, (2) pseudosatiety, and (3) absorption of toxins from the plastic.

RESEARCH NEEDS

First Priority

Effects of Marine Turtles

Experimental feeding studies are needed to determine (1) diagnostic criteria for interpreting the lethality or other pathologic significance of loads of ingested plastic, and (2) the entire gamut of the pathophysiology of ingested plastic in turtles.

Justification

There are relatively few data available on the prevalence or effects of ingested plastic in turtles, but those data which do exist suggest that the prevalence is high and that ingested plastic causes significant lesions and mortality. The endangered status of marine turtles justifies a prompt look at the role of plastics in mortality. Finally, it seems that a favorable cost/benefit ratio might result from dollars invested in turtle research. So little is known that a relatively small sum may produce considerable new information.

Second Priority

Effects on Seabirds

Controlled experimental work is required to determine if (1) pseudo-satiety does occur, (2) the duration of retention and erosion rates of ingested plastics, and (3) the toxicity of ingested plastics. The results of such studies will establish the need for long-term population studies of things like the postfledgling effect of plastic loading of chicks.

Justification

The available data establish that frequency of ingestion is very high in some species of seabirds and that some individuals contain very large amounts of plastic. There are few data about the effects on individual birds or populations. The few data available show no cause for alarm, but if these preliminary data are misleading, the potential deleterious effects on seabird populations could be severe. Because of the ubiquity of ingested plastic in seabirds and the as-yet-unmeasured potential for harm, it is prudent to identify the effect. Also, this group includes several threatened or endangered surface-feeding seabirds including the short-tailed albatross, *Diomedea albatrus*, which may be at risk.

Third Priority

Prevalence in Marine Turtles

The working group recommends continued monitoring of the prevalence of ingested plastic and its association with lesions. The monitoring efforts should be improved to better determine how often it actually causes harm. The working group recommends that review of the Marine Animal Stranding network be conducted to determine if the network's activities could be enhanced by standardizing necropsy protocols and by including collection of data about ingested plastics. Adequate diagnostic pathology services should be provided for the biologists in the Network. Even though the anecdotal data generated by monitoring programs cannot prove cause-effect relationships, they do provide useful information data bases.

Justification

This is the same as for first priority. Also, the Marine Animal Stranding Network is already in place; it would seem cost efficient to strengthen the program and orient it to collect and analyze data on ingested plastics.

Prevalence and Effects in Manatees

The data presented in the technical session suggested that plastic ingestion is common and was considered the cause of death of one manatee. It is recommended that carcasses found through the Marine Animal Stranding

Network be examined to obtain as much information as possible from each animal recovered.

Justification

There are no data about the impact of ingested plastic on the manatee population. Since this is a remnant population near extinction, any avoidable source of death is unacceptable.

Fourth Priority

Effects on Fishes

The working group recommends that laboratory work be done first with large fish to determine under what conditions they ingest plastics and to determine further the effect of the plastics. For example, will ingested plastic be retained for long periods and cause gastrointestinal tract blockages? Will it induce pseudosatiety, or release toxic chemicals?

Justification

Potential losses to the commercial and recreational fisheries may occur if ingested plastics impair the health of large fish. The working group assigned fourth priority to this issue because field and laboratory evidence available to date are equivocal, and as yet there is no evidence of a significant problem.

Prevalence in Marine Mammals

The working group recommends that monitoring of ingested plastics in stranded marine mammals be continued and improved as much as possible, taking advantage of the Marine Animal Stranding Network.

Justification

The working group generally agreed that available data suggest ingested plastics are a lesser problem in marine mammals and that there are no apparent reasons to elevate this issue to a higher priority at this time. It was also stated that laboratory work might better elucidate the consequences of ingested plastics, but that laboratory work with marine mammals is impractical because of logistics and legal complications.

Low Priority

Low priority issues are not unimportant, but they are less pressing than those above.

Effects on Larval Fish

The working group recommends that additional laboratory feeding experiments be done with larval fish to determine if ingested microparticles reduce growth rates.

Prevalence in Fish

The working group recommends that a specific study be designed to look for plastics in the gastrointestinal tract of large, free-ranging fish and for indications that it causes harm. This might be accomplished by alerting and educating fisheries biologists about the issue. This work could be done in conjunction with other on-going studies.

Prevalence in Seabirds

The working group recommends continued monitoring for benefits accrued (public awareness, time-order trends), but suspects that monitoring will continue without specific, directed guidance.

Effects on Marine Mammals

Laboratory studies of the effects of ingested plastic would provide useful data, but the group generally agreed that such studies are impractical for logistical and legal reasons.

LIST OF PARTICIPANTS

Nelio B. Barros, Rosenstiel School of Marine and Atmospheric Science,
University of Miami
Donald E. Hoss, Southeast Fisheries Science Center, Beaufort
Laboratory, NMFS, NOAA
Peter L. Lutz, Rosenstiel School of Marine and Atmospheric Science,
University of Miami
Haruo Ogi, Japan Research Institute of the North Pacific Fisheries
Pamela T. Plotkin, Marine Science Institute, University of Texas
Richard H. Podolsky, Research and Academics, Island Institute
Peter G. Ryan, Percy FitzPatrick Institute of African Ornithology,
University of Cape Town
Samuel S. Sadove, Okeanos Ocean Research Foundation, Inc.
Paul R. Sievert, Department of Biology, University of Pennsylvania
Louis Sileo, National Wildlife Health Research Center
Raymond J. Tarpley, The Texas Veterinary Medical Center, Texas A&M
University
Yoh Watanabe, Natural Research Institute of Far Seas Fisheries, Fisheries
Agency, the Government of Japan

APPENDIX A

Summary of information from five reports on the prevalence and effect of ingested plastics in seabirds.

Prevalence

Many previously unpublished data were presented at the technical sessions. Ingested plastics are present in many species and the prevalence is high in some species. Prevalence between species is influenced by feeding behavior, feeding location, season, year, resident or migrant status, and whether or not chicks are fed by regurgitation.

Effect of Ingested Plastic on Individual Seabirds

Some die from the lesions caused by impactions. Fledging weights of one species were reduced in chicks having high volumes of ingested plastic.

Effect of Ingested Plastic on Seabird Populations

There are no data available.

APPENDIX B

Summary of information from two reports on the prevalence and effect of ingested plastics in fish.

Prevalence

Ingestion does occur, but the information available suggests that this is uncommon. In one study, 20 of 3,000 larval fish contained ingested plastics. Nothing is known of the prevalence in large fish. In one study, most lancetfish contained one or more pieces of plastic.

Effect on Individual Fish

There is no clear evidence of an effect. In one study, larval fish ate 500 μ spherules, but there was no detectable short-term effect. No data are available about the effect on large fish.

Effect on Fish Populations

No data were presented.

APPENDIX C

Summary of information from five reports on the prevalence and effect of ingested plastics in marine turtles.

Prevalence

The few data available suggest the prevalence is high. In one study, 8 of 15 young pelagic turtles carcasses had intestinal compactions containing hundreds of pieces of debris. There were 3,000 pieces of plastic in 1 compaction. These compactions were the suspected cause of death. In another study, 60 of 111 beach-washed turtle carcasses contained intestinal debris, and 4 died from the effects. Yet another study reported debris in 12 of 168 stranded turtles; 5 of which had blocked pyloruses. Plastic bags or sheeting seemed to be the offenders.

Effect on Individual Turtles

Impactions can kill turtles. The few available data suggest that this is potentially a serious problem. Laboratory studies suggest that low doses of plastic have no effect.

Effect on Turtle Populations

There is no information available.

APPENDIX D

Summary of information from four reports on the prevalence and effect on ingested plastics in marine mammals.

Prevalence

The data are not completely clear. One report suggested the prevalence is low, others that it is high. Plastic was found in 15% of 63 dolphins and in 6 of 82 whales in one study. Another study reported that plastic debris was present in 67% of stranded whales. This study also reported debris in the stomachs of 23 of 86 Baird's beaked whales examined; 30% of this ingested debris was plastic.

Effects on Individual Marine Mammals

Data from both wild and aquarium specimens show that ingested debris can kill cetaceans.

Effects on Marine Mammal Populations

There are no data available.

REPORT OF THE WORKING GROUP ON ECONOMIC ASPECTS OF MARINE DEBRIS

(Kenneth E. McConnell, Chair)

The problem of marine debris is a classic example of markets failing to allocate resources efficiently. When firms and individuals use materials which escape into the marine environment, they impose costs on others. These costs--external costs, as they are known to economists--may be nonmarket, as in the aesthetic degradation of beaches or the killing of noncommercial species of birds and mammals. The external costs may also be incurred by market forces; for example, fishing vessels may have their propellers entangled in abandoned gear. Regardless of who suffers the external cost, its presence indicates a problem which requires some form of public policy to solve.

The marine debris problem is dominated by plastics and other nondegradable materials. Plastics have advantages in production and consumption processes that other materials lack. The replacement of plastics will therefore impose direct or inconvenience costs on consumers and producers. There are three basic ways to reduce marine debris: (1) reduce the loss and disposal of materials that may end up in the marine environment, (2) reduce the production of plastics and other nondegradable materials by using substitutes, and (3) engage in cleanup efforts.

Currently, knowledge of the economic aspects of marine debris is quite limited. This document outlines the basic economic issues and suggests research projects which would help in the process of reducing marine debris. The research projects can also serve as terms of reference for the economic issues of marine debris.

ECONOMIC COSTS

The economic costs of marine debris are the lost economic values that occur when the debris directly influences people and their behavior. These economic costs may be categorized as aesthetic, fouling of gear and vessels, and impact on fish stocks. Knowledge of these costs can help motivate government action. This requires information not only on the physical and biological effects of marine debris but also on the economic costs of these effects. The U.S. Office of Management and Budget, for example, is especially influenced by benefits and costs, not by physical effects. These economic effects require careful research because they do not show up in market transactions.

Aesthetics

Debris makes beaches less attractive. It traps fish and wildlife. Each of these entails an aesthetic loss to some individuals. Currently we

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

know practically nothing of the economic costs of either. We recommend two types of studies to help understand the magnitude of the economic costs of marine debris.

**A Study of the Economic Costs of
Debris on a Specific Set of Beaches**

This study would pick a set of beaches and investigate the economic value of lost services that result from different levels of debris on the beach. The techniques for undertaking this research are well known to economists, but they have not been applied to marine debris.

**A Study of the Economic Costs Incurred When Some
Individuals of a Noncommercial Species (e.g.,
Birds, Mammals) Are Entangled in Marine Debris**

Economists have some experience in estimating the loss in economic value from the threat of extinction of a noncommercial species but little or no experience when only a small number of the species are lost. This study would begin the investigation of this problem.

Fouling of Vessels and Fishing Gear

When vessels and their gear are impaired by contact with marine debris, there are two kinds of costs. The obvious cost is the repair or replacement cost for the damaged gear. Less obvious is the opportunity cost of the vessel and gear when it is not in productive service. Very little is known of this cost in the United States. There is evidence from Japan that the cost to fishing vessels is quite substantial. We suggest the following research.

**Investigate the Incidence of Impairment
and the Magnitude of Costs for One of
the Following Industry Groups: Commercial
Fishing, Shipping, or Recreational Boating**

This research project would involve a small survey among owners or operators in one of these industry groups for a well-defined region.

Impact on Fish Stocks

The biggest impact of marine debris on fish stocks is ghost fishing. But there is also the possibility that consumers' perception of contamination of fish stocks by marine debris can influence the demand and price of selected fish products.

Ghost fishing

The nature of physical or biological effects of ghost fishing is well known, but the quantitative effects are hard to estimate. Ghost fishing has an economic cost in terms of the lost resource. We suggest gauging this cost with a joint project involving economists and biologists.

The Impact of Perceived Contamination on the Price of and Demand for Fish

A project which collects and describes incidence of market effects from perceived contamination would provide especially convincing evidence on the economic costs of marine debris. We suggest a survey of literature and of knowledgeable people to gather these incidents in the form of a research report. This evidence comes in the form of market price changes induced by perceived contamination.

POLICIES TO REDUCE MARINE DEBRIS

Economics, the study of choices and behavior, is sometimes useful, sometimes essential, in thinking about and designing policies to reduce marine debris. Research on the costs of reducing marine debris needs to investigate the direct costs of adopting different techniques, the demand for the use of nondegradable materials, and the rate of compliance with various regulations among different sectors of the public and industry.

Policies to reduce marine debris require people to change their behavior. Behavior can be changed through a variety of means: education, moral suasion, lobbying, incentives, and direct regulation.

Incentives

In 1987, 12 U.S. senators wrote to the President expressing their concern over the marine debris problem. They urged the study of various methods of reducing debris, including instituting deposits "and other incentives for retention and retrieval of debris." Incentive schemes may be especially cost-effective in controlling debris when education and moral suasion fail. The following projects investigate the use of fees and incentives as part of the solution. These projects are not listed in the order of priority.

Deposits on the Return of Nondegradable Products

The efficiency of deposits on beverage containers as a means of controlling land debris is well documented. This research project would investigate the potential for deposits for the return of plastic marine debris. It should focus on coastal states which have experience with deposit systems.

Fees on the Use of Nondegradable Materials

Business firms and households are good at allocating scarce resources when they pay for them. Fees on plastics would induce substitution of other materials. This project would investigate the feasibility of fees on potential debris in the marine environment.

Incentives at the Production Level

Debris in the marine environment is part of the larger social problem of solid-waste management. The disposal of nondegradables is a crucial component of this problem. Fees on the use of nondegradable raw materials, including plastic pellets, in the production process would guide producers to substitute other materials. Such fees would raise the relative costs of nondegradable materials and make it economically more attractive to develop degradable substitutes and to sponsor research in developing substitutes. This research should investigate the demand for raw materials at the production level and the potential for fees and incentives to spur the development of substitutes including recycled plastic materials.

Compliance

The typical approach to solving environmental problems is: to pass a law to prohibit behavior causing the problem, to devote a small amount of money to enforcing the law, and to engage in education and public awareness campaigns to persuade people to comply. This approach frequently fails, raising demands for more enforcement funds, harsher penalties, and so on. It may be fruitful to investigate alternatives to this traditional approach to compliance.

Investigate Alternatives to Traditional Methods of Compliance

Policies combining punishment and reward which partly subsidize the adoption of techniques to help people comply and impose clear penalties for the absence of compliance are used elsewhere. For example, sewage treatment has been enhanced by Federal subsidies to construct waste treatment plants linked to the requirement that all households hook up. Methods of linking compliance to rules and regulations for handling marine debris can be used to access to other beneficial programs. For example, the registration of boats might be linked to evidence that boats have systems for handling solid wastes. This research program would study compliance programs which include education, incentives, and penalties for a specific portion of the industry. The recreational boating industry is an especially good candidate for study because boating is so widespread and boaters so heterogeneous in attitudes.

Evaluate the Effectiveness of Moral Suasion Programs

Public campaigns to reduce pollution by moral suasion have been attempted for other forms of pollution. A study of these campaigns would help understand their failures, which have been many, and their successes, which have been few. This study should cover different countries and different times.

Onshore Disposal

The new laws require that vessels bring their nondegradable wastes to port. Ports are required to handle the solid waste. Within particular

regions, it may be economically very costly for all ports to handle all of the vessel-borne waste. We suggest the following research project.

**Investigate the Economic Gains That Can
Accrue to a Particular Region as a Consequence
of Consolidating Waste Handling Facilities**

Some ports are unable to handle wastes. Other ports may have excess capacity. An economic study of the costs of onshore waste handling would prepare ports for the resource demands and for setting port fees. When the costs differ among ports, there may be incentives to use different ports. Further, there are incentives to dump the trash if fees are based on the trash that is brought ashore.

LIST OF PARTICIPANTS

Norihisa Baba, National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan
 Paul A. Breen, New Zealand Ministry of Agriculture and Fisheries
 Richard A. Buxton, Nova Scotia, Canada
 Jeff Cantin, Eastern Research Group, Inc.
 David Clark, Battelle Human Affairs Research Centers
 Salvatore Comitini, Department of Economics, University of Hawaii
 Thomas A. Grigalunas, Department of Resource Economics, University of Rhode Island
 DeWayne Hollin, Sea Grant College Program, Texas A&M University
 Setsuki Matsumura, National Research Institute of Far Seas Fisheries, Fisheries Agency, the Government of Japan
 Kenneth E. (Ted) McConnell, Department of Agricultural and Resource Economics, University of Maryland
 Norman Meade, Office of Oceanography and Marine Assessment, National Ocean Service, NOAA
 Hal Schlomann, Exhibits Management, Inc.
 John Sutinen, Department of Resource Economics, University of Rhode Island
 Yasuji Tamaki, Fisheries Agency, the Government of Japan

REPORT OF THE WORKING GROUP ON TECHNOLOGY

(William G. Gordon, Chair)

INTRODUCTION

The Working Group on Technology, while recognizing that further work is required to quantify the types and volumes of ship-generated debris, strongly believes that technology/methodology currently exists to address management of the majority of the wastes generated at sea. The group recognizes that a considerable portion of debris originates from such terrestrial sources as careless transportation of garbage, combined storm/sewage outfalls, storm and street drains, industrial activities, beach-goers, and at-sea disposal, and that management of these sources requires application of somewhat different technologies and methodologies. However, a large measure of the ship-generated debris ultimately will be transported ashore. Thus, satisfactory resolution of much of the marine debris issue will require rational resolution of many of the terrestrial waste management issues and problems.

SHIP-GENERATED DEBRIS

Ship-generated debris and sources are categorized in Table 1, Sources and types of ship-generated debris.

SHIP WASTE MANAGEMENT

Technology and/or methodologies for dealing with ship-generated wastes, and their potential application, are shown in the Table 2, available technology and methodology for handling ship-generated waste.

As displayed above, there are currently available a number of techniques for addressing ship-generated wastes. However, all ultimately require some degree of transportation of the waste ashore for disposition. On-land facilities for handling such wastes may not exist, and thus at-sea disposition will continue.

The group stressed that there is no single methodology or technology which will resolve the issue. Regardless of the size of the vessel or craft, a variety of practices will undoubtedly be employed. Brief descriptions follow.

Source Reduction

Source reduction is the use of materials on board the vessels which will reduce both quantity and volume of waste. Such practices will vary with segments of the industry and should be encouraged for all.

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

Table 1.--Sources and types of ship-generated debris.

Sources	Clean plastic	Contaminated plastic	Fishing gear	Paper	Metal and glass	Oils	Food wastes
Merchant marine	Large	Small	--	Medium	Small	Large	Small
Naval vessels	Large	Large	--	Large	Large	Medium	Large
Commercial fishing	Small	Medium	Large	Small	Small	Medium	Small
Recreational craft	Medium	Large	Small	Small	Medium	Small	Medium
Cruise ships	Large	Large	--	Large	Large	Medium	Large
Oil and gas operations	Large	Large	--	Large	Large	Large	Large

Alternate Materials

The use of alternative materials is a simple but effective strategy for plastic waste management at sea. This may involve the replacement of a given plastic product with either a different plastic product or a nonplastic product which is more disposable. This simple approach can have a significant impact on solid waste management options for shipboard use.

Effectiveness and feasibility of incineration of waste on board a vessel can be increased by the selection of plastics which can be safely burned for use at sea. Exclusion of chlorine-containing plastics and, perhaps, the use of compounds with high fractions of fillers (to reduce BTU's per pound) are possible examples.

Recycling of plastics carried to shore is viewed as a plausible means of disposal. Plastics to be used aboard vessels might then be selected to ensure optimum blend compatibility of the disposed plastic waste stream. Alternatively, the plastics can be coded to allow easy separation into the different chemical types. Replacement of nonrecyclable material such as glass and metal (usually disposed of at sea) with recyclable material such as paper and plastic is also feasible.

Reducing the plastic content of a given product can lead to substantial improvements in disposability with minimal loss of performance. Plastic films, plates, and cups, for instance, can be replaced with plastic-coated or composite materials that are more disposable. At the extreme is the replacement of plastics and glass with paper, which can be more easily

Table 2.--Available technology and methodology of disposition of debris generated at sea.

Type of debris	Type of disposal							Source reduction
	Incineration	Disposal at sea	Separation	Compaction	Recycle	Degradation	Alternate	
Plastic	X	--	X	X	X	X	X	X
Foamed plastic	X	--	X	X	--	--	X	X
Fishing gear	X	--	X	--	X	--	--	X
Paper	X	X	X	X	X	--	--	X
Metal and glass	--	X	X	X	X	--	X	--
Oils	X	X	X	--	X	--	--	--
Food wastes	X	X	X	X	--	--	--	--

disposed of. In doing so, of course, the economic factors must be taken into account to make sure that the proposed substitution of materials is realistic. Substitution of materials that are easily compacted and stored on board is stressed by the working group.

Degradation

Deep-sea disposal of many materials remains acceptable. Waste food biodegrades, paper (unless plastic-coated) degrades, metal cans (if punctured) sink, and glass bottles (if broken or if caps are removed) sink. However, plastic materials regardless of form cannot be dealt with in this way as their rate of degradation is unacceptable for at-sea disposal.

Some common plastic and fibers can be made to photooxidize and biodegrade in a controlled fashion. In both film and fiber forms they are already in use in agriculture and some packaging applications. Olefins are among those polymers which can be engineered to degrade. Nylon and polyesters presently available do not degrade in an acceptable time. Nylon can be made to photodegrade, but photodegradable nylon is not yet commercially available. The products of degradation of most polymers are environmentally innocuous.

Photodegradable plastics are compatible with other technologies. Degradable plastics can be burned normally or can be recycled to secondary products. Degradable products can cause environmental problems in landfills, as many cease to degrade under anaerobic conditions.

Degradable plastics have potential disadvantages. For example, few of the existing degradable materials have been cleared by the Food and Drug Administration for contact with food. Some degradable plastics perform differently on land than in the marine environment, and require further evolution. Thus, industry is not yet prepared to supply economically feasible plastics which can be counted on for degradability at sea.

Recycling

Recycling of plastics is a process by which used plastic objects are collected, identified, separated (if necessary), and melt-processed into useful items. Recycling of plastic soda bottles, fishing nets, and general plastics waste is currently being done on a commercial basis. At present, probably less than 10% by weight of the annual production of plastics is being recycled. The impetus brought about by factors such as the increasing costs of dumping at landfills, the influence of environmentally concerned groups, the inherent value of the plastics themselves, which continue to increase in cost, and the profit motive for value-added products made of recycled plastics could well increase the use of recycling to as much as 50% of yearly production.

Except perhaps aboard very large vessels, recycling at sea does not appear to be a viable solution for dealing with plastics waste. A more practical approach is the establishment of recycling centers at ports receiving large volumes of plastic debris. Unless information on how much, what type, and when will it be landed is made available, industry is

unlikely to move. Accordingly there is a need to acquire much more information on the issue in order to encourage the private sector toward a positive response.

Another aspect of recycling is the reuse of plastic items for different purposes. An example of this is the use of fishing nets for decorative or recreational applications.

Recycling of various plastics together (commingling) will result in some difficulties. Commingling will reduce the value of the resultant product as it results in characteristics that are less desirable to most users. This problem must be addressed, particularly in view of the move toward laminated products of various plastics.

Compaction

Compaction technology as wide application for resolution of waste issues within the maritime industry. For many, compaction may be the most appropriate, as all solid waste may be reduced through compaction and stored readily aboard those craft where trips are of short duration. This would include virtually all recreational boats, coastal commercial fishery operations, and coastal maritime shippers. Obviously, on-land disposal must be economical and efficient in order to encourage the practice.

For recreational and small fishing vessels, the initial problem created by the need to retain all wastes aboard ship is that none of these vessels are designed with waste retention compartments. However, they are designed with water and fuel tanks, sleeping compartments, cargo and provision compartments, and most recently with sewage holding tanks. These vessels are designed with these various compartments because either laws require them or they are necessary for the operation of the vessel. Therefore, in a way similar to sewage holding tanks, waste retention compartments could become a necessary requirement of all vessels.

One suggestion for a type of shipboard waste retention compartment might be a "shipboard waste compactor." One could be designed to run off the ship electrical system or be hand-operated. If owner-installation were possible, then some of the burden of cost to the vessel owner would be lessened. Such a device could provide for improved sanitation as well as make transfer of wastes easier. For example, the ports could store compacted wastes more neatly and securely than "untreated wastes," which are often unsightly, cause odors, and are cumbersome to store.

Separation

Separation of all wastes by type will become part of all successful waste management practices within the maritime industries. This methodology must be fully integrated into customary practice in combination with others. All practices must be at reasonable cost and effective.

Combustion

Combustion includes low technology burning and incineration.

Low Technology Burning (e.g., Use of Burners)

Low technology burning has important practicabilities and benefits:

1. It can be an attractive option for compliance with Annex V.
2. It is low in cost.
3. It needs relatively little deck space.

It also has some potential pitfalls:

1. Some separation of hazardous materials is necessary.
2. Products of incomplete combustion (e.g., dioxin, furans) of a wide range of materials (including salt from the marine environment) are environmentally hazardous. Potential hazards include toxic air emissions.
3. There are current problems concerning disposal of toxic ash (which may be resolved in the near future as a result of manufacturing modifications by the plastics industry).
4. The regulatory climate concerning air pollution emissions could change in the future.

Incineration (e.g., Insulated Combustion Chambers With Mechanical Air Control)

Incineration also has important practical aspects and benefits:

1. It is an attractive option for compliance with Annex V for ships such as cruise ships, merchant vessels, and tankers.
2. It provides operational flexibility, given the variable availability of port reception facilities.

It, too, has potential pitfalls:

1. Trained skilled personnel are necessary for proper operation.
 2. It is more costly than low technology burning or overboard dumping.
 3. Combustion of plastics, salt, and other materials produces air pollution emissions.
 4. There are current problems concerning disposal of hazardous ash (which may be resolved in the future as a result of manufacturing changes).
-

5. Regulatory climate concerning air pollution emissions at sea could worsen in the future.

LAND DISPOSAL

The working group recognizes that ultimately ship-generated debris, particularly plastics, will be transported ashore for disposition. Land facilities, therefore, must be expanded or developed to accommodate the increased volume of such materials. Shoreside receivers must be cost effective and convenient in order to encourage maximum use. Shore-based industries should be encouraged to recycle such materials; they will require considerable information on the types, quantities, and location of the materials. Governments should implement policies and incentives permitting and encouraging such initiatives, but this, too, requires a much greater level of knowledge than is currently available.

The group stressed that marine debris is clearly linked to the land and that all efforts to encourage rational waste management must be extended globally.

RECOMMENDATIONS

Data

More information should be obtained about types, quantities, and distribution of the plastic materials which, under MARPOL regulations, will be brought ashore for disposal. Such information should be disseminated throughout the plastics industry to encourage reuse of such material.

Technology

Research and development of new technologies should be encouraged.

Low Technology Burning

1. Research on environmental impacts of air emissions.
2. Research and development of guidelines concerning materials separation and operations.
3. Research concerning environmental implications of ash and methods of disposal.

Incineration

1. Research on environmental impacts of air emissions.
 2. Research concerning environmental implications of ash and methods of disposal.
 3. Research concerning hydrogen chloride corrosion of incinerator units and accompanying potential effects on durability. Measures to address problems, if necessary.
-

Ship Design

1. New design should accommodate waste management strategies.
2. New construction should include facilities and space accommodations for waste management.

Plastics

1. Development or identification of the more desirable plastic mix streams for commingled plastics.
2. Development of performance standards for alternative materials used for specific product applications.
3. Development of trade-off studies on performance versus disposability.

Policy

No single methodology or technology will ensure compliance with waste management regulations. Accordingly, no policy should be established which prohibits technologies which have potential--keep all options open.

Governments should work together to create incentives for on-land disposal wherever feasible. Recycling, for example, will be feasible only where economic conditions are ideal for such practices. Governments should take the lead to assure that new and complementary technologies are created.

Education

Manufacturers of items such as packaging, fiber, and netting, are not aware of the capabilities and potential of programmable degradable plastics. Efforts should be undertaken to achieve broad dissemination of such information. The working group suggests that the plastics industry undertake such efforts.

There is need for a global network to disseminate information on the impact of marine debris widely and to uniformly encourage development of modern technologies or methodologies everywhere.

There is need for education throughout the private sector, aboard all classes of ships as well as shoreside, to encourage good waste management practices.

General Views

The working group strongly endorses the need to conduct global workshops on the issue of marine debris periodically. Sessions should include technical aspects of present and emerging technology for shipboard application. Invitees should include those potentially interested in land-based use of materials.

LIST OF PARTICIPANTS

Terrill J. Chang, SCS Engineers

Paul G. Kelleher, Polymer Processing Institute, Stevens Institute of
Technology

Lissa A. Martinez, Takoma Park, MD

Cathy L. P. Palmer, Mote Marine Laboratory

Bruce Perlson, Quantum Chemical Corporation

Alan Reichman, Greenpeace USA

Akinori Satou, N.S. Environmental Science Consultant

REPORT OF THE WORKING GROUP ON LAW AND POLICY

(David Cottingham, Chair)

The working group reviewed existing legal and institutional arrangements to curtail the disposal and loss of solid wastes into the marine environment. Many nations have signed various international agreements and passed domestic laws that prohibit or limit disposal of plastic and other refuse into the sea, including disposal of wastes from ships. An undetermined, though probably significant, amount of marine debris originates on land. The group, therefore, concluded that solutions to the problem of marine debris should be developed and implemented in concert with efforts to address broader solid waste management issues. The most pressing needs identified include:

1. participating in the relevant international agreements;
2. assuring that adequate reception facilities are available at all ports and harbors to receive ship-generated garbage returned to shore; and
3. adopting national policies and programs, such as recycling and innovative packaging, to reduce the quantities of solid waste generated.

CONTROLLING AT-SEA SOURCES OF MARINE DEBRIS

The international agreement of greatest importance for controlling the discharge of plastics and other solid wastes into marine waters are:

- International Convention for the Prevention of Pollution from Ships, 1973/1978, Annex V (MARPOL Convention); and
- International Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (London Dumping Convention (LDC)).

The MARPOL Annex V regulates the disposal of ship-generated garbage. The LDC restricts transporting land-generated solid wastes to sea for the purpose of dumping.

At least 10 regional conventions control various forms of marine pollution, including the disposal of plastics and other solid wastes, from both sea- and land-based sources. They include:

In R. S. Shomura and M. L. Godfrey (editors), Proceedings of the Second International Conference on Marine Debris, 2-7 April 1989, Honolulu, Hawaii. U.S. Dep. Commer., NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFSC-154. 1990.

- Convention on the Conservation of Antarctic Marine Living Resources, 1980.
- Convention on the Protection of the Marine Environment of the Baltic Sea Area, 1974 (the Helsinki Convention).
- Convention for the Prevention of Marine Pollution from Land-Based Sources, 1970.
- International Convention for the High Seas Fisheries of the North Pacific Ocean, 1952.
- Convention for the Protection of Marine Pollution by Dumping from Ships and Aircraft, 1972 (the Oslo Convention).
- Convention for the Protection and Development of the Wider Caribbean Region, 1983 (the Cartagena Convention).
- Convention for the Protection of the Mediterranean Sea Against Pollution, 1976 (the Barcelona Convention).
- Kuwait Regional Convention for Cooperation on the Protection of the Marine Environment from Pollution, 1978.
- Convention for Cooperation in the Protection and Development of the Marine Environment of the West and Central African Region, 1981 (the Abidjan Convention).
- Regional Convention for the Conservation of the Red Sea and Gulf of Aden Environment, 1982 (the Jeddah Regional Convention).

The latter five are part of the United Nations Environment Program Regional Seas Program.

The working group concluded that these international conventions jointly prohibit the disposal of plastics and other solid materials in the sea, thereby establishing this prohibition as "customary international law."

Of 131 nations that are members of the International Maritime Organization (IMO), 36 have ratified optional Annex V of MARPOL as of April 1989. These countries account for approximately 56% of the world's total gross commercial shipping tonnage. Many signatory nations are still in the process of developing programs to implement Annex V. In particular, many ports do not have adequate or convenient facilities for accepting ship-generated garbage.

Annex V identifies five "special areas"--the Baltic Sea, Red Sea, Mediterranean Sea, Black Sea, and "Gulfs Area." Within these special areas, ships are prohibited from disposing of all solid wastes (except comminuted food wastes beyond 12 nmi from shore). Before these provisions

can become effective, however, 1) each adjoining nation must certify that it agrees to designation of the area as a special area, and 2) that its ports have adequate reception facilities for handling ship-generated garbage. All nations surrounding the Baltic Sea have notified IMO. As of 1 October 1989, special area provisions in the Baltic Sea became effective. The Marine Environment Protection Committee (MEPC) recently added the North Sea as a special area. The United States is proposing to add the Gulf of Mexico as well.

CONTROLLING LAND-BASED SOURCES OF MARINE DEBRIS

Many items which become marine debris wash or blow into oceans and estuaries from landfills, municipal sewer systems, recreational beaches, industrial outfalls, illegal shoreline dumping, and other sources. Controlling land-based sources of persistent marine debris raises difficult problems which all nations must address domestically. International agreements are not well suited for controlling wastes from sites located on land.

Working group participants discussed the problems associated with effectively handling municipal and industrial solid waste throughout the world. Developed nations produce the largest per capita amounts of solid waste and generally have systems in place to dispose of it. Developing nations, on the other hand, frequently lack effective solid waste collection and disposal systems. Problems caused by marine debris in developing nations must be viewed within the context of their overall ability to handle all solid wastes. In such countries, marine debris may be a relatively small component of problems associated with handling solid wastes.

In the United States, responsibility for controlling nonhazardous wastes rests with the state and local governments. State and local agencies throughout the country have antilittering and dumping laws. However, enforcing these laws is difficult and penalties for violating them are not severe. Most local agencies, like the municipally operated Keep America Beautiful and Don't Mess With Texas programs, concentrate efforts on informing the public through antilitter campaigns.

Working group participants discussed the importance of reducing the volume of ship-generated solid waste by modifying ship stores purchasing and having rigorous onboard waste management. Current packaging systems often include throwaway containers, many of which could be recycled if systems and markets existed for the initial products and the recycled material. Japan, Denmark, parts of the United States and Canada, and other countries require people to separate newspapers, aluminum cans, clear and colored glass, and some plastic bottles for recycling. Seattle officials estimate that its recycling program, with curbside residential pickup of separated refuse, has reduced the volume of solid waste by 40%. Recycling has proven to be an effective way to remove aluminum and glass containers from the solid waste stream.

IMPROVED TECHNOLOGY

Although the subject of advanced technology was addressed by another working group, the Law and Policy Working Group briefly discussed ways to encourage research on degradability of single-use items such as plastic cups, plastic eating utensils, plastic bags, and tampon applicators. Degradable products may be a partial solution to the problem of marine debris. However, application of this technology requires further consideration of what happens to them as they break down. For example, wildlife may be just as likely to ingest the smaller fragments produced as plastic items degrade.

WORKING GROUP RECOMMENDATIONS

The Working Group on Law and Policy recommends that the following actions be taken by the IMO, national governments, and private industry:

Loss and Disposal of Garbage From Ships

1. Nations that have not yet ratified MARPOL Annex V and any regional conventions applicable to them which restrict disposal of solid wastes into marine waters should do so as soon as possible.
2. Nations which have ratified MARPOL Annex V should accelerate efforts towards full implementation of required provisions for port reception facilities.
3. The MEPC of IMO should review its guidelines for port reception facilities for garbage to facilitate effective implementation pursuant to Annex V of MARPOL. The MEPC should give particular attention to: 1) recovering or defraying the costs of operating port reception facilities and handling wastes (e.g., through recycling and refuse separation); 2) methods for handling various types of wastes; and 3) simplifying the steps and procedures that vessel owners and operators must follow to use port reception facilities.
4. The MEPC nations should consider measures which could ensure that vessels do not leave ports with garbage on board, for example, consistent fee systems.
5. National governments should provide information and, where possible, economic incentives to help ports comply with port reception facility requirements of Annex V. National governments also should assist local port communities where significant increases in the volume of solid waste result from the installation of new port reception facilities. For example, in the United States, Federal and state officials should expedite reviews of applications and permits for landfills and incinerators in port communities in developing

recycling programs, and expedite review of applications and permits for disposal facilities made necessary by increased wastes from ships.

6. Governments and port authorities should develop incentive systems, such as requiring or including off-loading fees as part of docking fees, rather than penalties to encourage compliance with Annex V.
7. The IMO should consider expanding Annex V guidelines or developing other forms of providing advice on the development and use of vessel logs for tracking the handling and disposal of ship-generated garbage.
8. Nations adjacent to special areas of Annex V should accelerate the development of port reception facilities to enable special area provisions to become effective at the earliest possible dates. In particular, nations bordering the Gulf of Mexico should take steps to designate the gulf as a special area under Annex V.
9. Governments should consider sharing collection and refuse transportation costs.
10. Governments should consider developing uniform signage and coloration standards for refuse and recycling facilities.

Shoreline Sources of Marine Debris

1. Nations should examine ways to prevent garbage and litter from escaping from landfills, industrial outfalls, sewage outfalls, and harbors or washing from beaches and shorelines into coastal waters.
 2. Regional seas programs should provide technical assistance to member nations on siting disposal facilities and handling solid wastes in coastal areas.
 3. Industries manufacturing or transporting plastics and plastic products should ensure that plastic resin pellets are not lost into the marine environment during handling.
 4. All levels of governments should encourage recycling programs to reduce the volume of material which becomes solid waste. For instance, in Japan the plastics industry, in cooperation with fishing villages, has found ways to collect and recycle gillnets and trawl nets.
 5. Governments should require that ships and barges transporting solid wastes be fully covered to prevent debris from dropping or blowing into waterways. Transfer facilities should be required to have booms and skimmers in
-

place to remove refuse which enters waterways, and load limits and maximum heights for barges should be prescribed.

6. The Environmental Protection Agency and coastal municipalities should consider developing and requiring installation of equipment to prevent items in combined sewer overflows from entering waterways.
7. National governments should require that governmental entities preferentially purchase recycled goods.

Compliance and Enforcement

1. Governments party to MARPOL Annex V should develop incentives to encourage vessel owners and operators to comply with garbage disposal provisions.
2. Vessel owners and operators should be encouraged to report to national authorities and IMO those ports and harbors in Annex V signatory nations that do not have adequate port reception facilities. They should also be encouraged to report ports in nations which are not party to Annex V. For example, in the United States, the U.S. Coast Guard should publicize citizen reporting networks and encourage Coast Guard Auxiliary members to report violations of marine pollution regulations, including the absence of required port reception facilities.
3. The IMO should encourage member nations to develop innovative enforcement policies, such as requiring ships to off-load plastic refuse before they sail and providing inexpensive refuse removal.

Technology Improvements

1. National and local governments and private industry should develop institutional arrangements for recycling fishing nets and other large items that may potentially become marine debris.
2. Fisheries agencies should require time-release devices on crab, lobster, and fish traps and pots to avoid long-term ghost fishing by lost gear. Use of degradable material for selected components of drift net gillnets and other types of fishing gear should be required.
3. Private industries should conduct research on enhanced-degradable single-use plastic items such as cups, utensils, packing materials, and tampon applicators. The research also should examine by-products of degradable plastic materials and potential impacts on marine wildlife.

LIST OF PARTICIPANTS

Xan Augerot, Washington Sea Grant
Barbara H. Britten, American Cetacean Society
Richard Buxton, Nova Scotia, Canada
David Cottingham, Ecology and Environmental Conservation Office, NOAA
Mary L. Gessner, National Marine Fisheries Service, NOAA
Timothy Keeney, Office of General Counsel, NOAA
David W. Laist, Marine Mammal Commission, Washington, D.C.
Albert M. Manville II, Defenders of Wildlife
Rod J. Paterson, Canada Department of Fisheries and Oceans
Gerard Peet, Netherlands S.E.A. Foundation
Edward Rymarz, International Maritime Organization

REPORT OF THE WORKING GROUP ON MARINE DEBRIS EDUCATION

(Kathryn J. O'Hara, Chair)

Recognizing the difficulty of enforcing marine debris laws, especially at sea, education has been identified as an important way to help reduce the marine debris problem. Under the assumption that an informed public will be much less inclined to generate marine debris during both commercial and recreational activities, compliance with laws and regulations should be much higher. Education is particularly important in this issue because land-based sources of debris are primarily nonpoint, having diverse contributors that would be difficult to control under regulatory authorities alone. Moreover, longstanding, customary international law has permitted garbage discharge for ships in transit. Therefore, ethics and behavior patterns for individuals both on land and at sea must be changed, and education is the best known means for effecting such changes.

Charged with assembling a comprehensive list of the types of educational materials currently in use, the Working Group on Marine Debris Education identified more than 100 different types of educational materials. This included 21 brochures, 19 reports, factsheets and special documents, 11 posters, 10 videos, 9 curriculums and guides for educators, 6 newsletters, and more than 30 other types of educational materials ranging from public service advertisements to bumper stickers and coloring books. A complete listing of educational materials currently available may be obtained by contacting the National Oceanic and Atmospheric Administration's Marine Debris Information Office operated by the Center for Marine Conservation in Washington, D.C.

The working group was also charged with making recommendations for: (1) production of new educational materials and priority audiences for marine debris education, including the best means for delivery; (2) appropriate methods for the effective dissemination and utilization of these materials; (3) appropriate means for evaluating the success of educational programs; and (4) evaluation of lessons from the development of past marine debris educational programs and materials that may be of value in formulating environmental education programs in the future.

TYPES OF EDUCATIONAL PROGRAMS AND TARGET AUDIENCES

Marine debris education encompasses two key elements: The implementation of educational programs and the development of educational materials. With regard to the former, the group recommended that marine debris education should be incorporated into three primary types of programs:

1. formal education in a structured academic setting;

2. informal education outside a formal academic setting but within structured educational events such as adult education classes and organized youth groups; and
3. general public awareness.

Marine debris education has been or is presently being conducted for many groups, including: plastics manufacturers and processors, offshore oil and gas workers, commercial fishermen and processors, military personnel, politicians, solid waste managers, port and terminal operators, commercial shippers, teachers and educators, elementary, middle, and high school children, college students, recreational fishermen, recreational boaters, charter vessel operators, operators of cruise ships, cruise ship passengers, and the general public.

Several new groups would benefit from education and should be included in future efforts. These are the packaging industry; municipal sewage treatment operators; government officials; government enforcement agencies; coastal tourist industries; tackle manufacturers; operators of small ports, docks, marinas, and yacht clubs; suppliers of stores for vessels; boat manufacturers; employees of retail stores (including fast-food and convenience stores, and fishing and boating stores); environmental and conservation organizations; the media; employees of shipyards; longshoremen; and coastal hunters. Specialized efforts should be directed toward native and rural people.

Among all the groups identified above as target audiences for marine debris education, the working group concluded that five major groups are priority audiences:

1. all marine user groups;
2. the media;
3. teachers and educators;
4. school children; and the
5. general public.

A public awareness campaign is of utmost importance at the present time. Specific elements that should be addressed in developing this campaign are an initial assessment of human behavior and public perception of the marine debris problem. Using this information, a mass media public awareness campaign should then be developed. The working group felt that paramount to the success of this campaign is the development of a comprehensive strategy to use the media effectively as a tool to disseminate educational information. One suggestion was to solicit pro bono support from the National Advertising Council or a similar group. The working group also recognized that substantial funds are necessary to create this campaign.

EFFECTIVE DISSEMINATION AND UTILIZATION OF MATERIALS

After reviewing the list of marine debris educational materials, the working group concluded that there is a wealth of materials currently available but there is a need to facilitate the dissemination of these materials to appropriate groups. In 1988, the National Oceanic and Atmospheric Administration's Marine Entanglement Research Program established two Marine Debris Information Offices, which respond to requests for information on marine debris. The working group suggested that the dissemination of marine debris educational materials would be facilitated if the function of these offices were enhanced by increasing their visibility as an international resource center and providing them with sufficient quantities of educational materials to meet the demand. It was also suggested that an informational vehicle be established to provide updated information on the development of new educational materials and programs. There was an opinion, however, that educational materials should be disseminated in a more decentralized manner.

The dissemination of educational materials could be facilitated with assistance from established education organizations such as the National Marine Educators Association.

Existing government distribution mechanisms such as licensing and registration procedures for fishing and boating should also be used to disseminate materials.

The working group recognized the difficulty of disseminating educational materials on an international level due to the diversity of cultures and languages. However, it was suggested that specific international agencies such as the United Nations Environmental Programme, Food and Agriculture Organization, and the International Maritime Organization should be encouraged to take part in information exchange.

Efforts should be made to include the marine debris issue on the agendas of international conferences and meetings that address the issues of marine pollution and education.

EVALUATING THE SUCCESS OF EDUCATIONAL PROGRAMS

Evaluations could be conducted through long-term monitoring of beach debris and monitoring the usage of shoreside refuse reception facilities.

Formal surveys should be conducted, where possible, to assess changes in attitude and behavior.

LESSONS LEARNED FROM THE DEVELOPMENT OF PAST MARINE DEBRIS EDUCATIONAL PROGRAMS

Specific recommendations made with regard to development of educational programs and materials included:

- Involve members of the target audience in the development of educational materials and distribution.
-

- Identify specific discrete tasks for the involvement of individuals.
- Set realistic goals.
- Make educational experiences positive and enjoyable.
- Be familiar with the audience.

Other experiences shared by group members who have been involved in educational efforts pertained to the content of educational materials. This included the need to:

- Use good photographic materials that show the impact of debris.
- Personalize the message to specific target audiences.
- Emphasize the benefits to a group for their involvement in efforts to reduce the marine debris problem.
- Emphasize the importance of individual efforts.
- Emphasize economic impacts where appropriate.
- Keep the information as locally relevant as possible.
- Keep the message short.
- Highlight positive steps taken by groups or individuals to reduce the marine debris problem.
- Use facts that are updated and verified.

The working group suggested that researchers who are working on the issue of marine debris should be encouraged to make photographic materials available for educational efforts.


FORMULATING FUTURE ENVIRONMENTAL EDUCATION PROGRAMS

The working group also discussed the need to expand educational efforts to include the way in which debris affects estuaries and inland waters. There was considerable discussion with differences of opinion regarding the emphasis of marine debris educational materials and programs. The majority of participants agreed that the primary focus of marine debris educational materials should be to increase awareness of the problems caused by improper disposal of man-made wastes in marine areas. Others suggested that marine debris educational efforts should also emphasize the need for source reduction and the broader issues of wasteful consumer habits. It was agreed by all working group participants that the marine debris issue is part of the larger solid waste problem and, therefore, we should incorporate lessons learned from dealing with solid waste into marine debris education materials and programs.




The working group recognized that the marine debris issue has elicited an unprecedented emotional response and enthusiasm to take action. Therefore, the group sees great potential for using the marine debris issue as a stepping stone to encourage citizen involvement in other environmental issues.

LIST OF PARTICIPANTS

Nancy Azzam, Windstar Foundation
Ronald G. Bruner, The Society of the Plastics Industry
Alan Bunn, Alaska Fisheries Science Center, NMFS, NOAA
James M. Coe, Alaska Fisheries Science Center, NMFS, NOAA
Patty Debenham, Center for Marine Conservation
Trevor R. Dixon, The Tidy Britain Group
Angela Farias, Texas Adopt-A-Beach Program
Sarah Friday, North Carolina Sea Grant
Bernard Griswold, National Sea Grant College
Michael R. Hagler, Greenpeace, New Zealand
Jeffrey June, Natural Resources Consultants
Jill Kauffman, Center for Marine Conservation
Shirley Laska, Department of Sociology, University of New Orleans
Sidney Lasseigne, Pacific Marine Fisheries Commission
Patti Mullin, Washington Sea Grant
Judie Neilson, Oregon Department of Fish and Wildlife
Kathryn J. O'Hara, Center for Marine Conservation
Fran Recht, Pacific States Marine Fisheries Commission
Barbara Schroeder, Florida Department of Natural Resources
Rupert Teves, Pearl City, HI
Barbara Wallace, Kearney/Centaur Division, A. T. Kearney, Inc.
Dottie Wendt, Hawaii Sea Grant
Nobuyuki Yagi, Division of Fishing Grounds Environmental Conservation,
Fisheries Agency, the Government of Japan



APPENDIXES





APPENDIX A

STEERING GROUP

Dayton L. Alverson, Natural Resources Consultants
George W. Boehlert, National Marine Fisheries Service
Douglas G. Chapman, University of Washington
James M. Coe, National Marine Fisheries Service
Jack R. Davidson, Sea Grant College Programs, University of Hawaii
Richard S. Shomura, University of Hawaii

APPENDIX B

AGENDA

1. Opening of the conference
 2. Presentations
 - a. Overview papers
 - b. Session I - Amounts, types, distribution, and sources of marine debris
 - c. Session II - Entanglement of marine life and ghost fishing
 - d. Session III - Ingestion by marine life
 - e. Session IV - Economic impacts on vessels and shorelines
 - f. Session V - Solutions through technology
 - g. Session VI - Solutions through law and policy
 - h. Session VII - Solutions through education
 - i. Poster presentations
 - j. Video presentations
 3. Working group meetings
 4. Plenary session
 5. Conference summary and closing
-

APPENDIX C

LIST OF PARTICIPANTS

Mr. Craig S. Alig
David Taylor Research Center
Department of the Navy
Bethesda, MD 21402

Dr. Dayton L. Alverson
Natural Resources Consultants
4055 - 21st Avenue West
Seattle, WA 98199

Dr. Anthony F. Amos
Marine Science Institute
University of Texas, Austin
750 Channelview Drive
Port Aransas, TX 78373

Dr. Anthony L. Andrady
Research Triangle Institute
Chemistry and Life Sciences
P.O. Box 12194
Research Triangle Park, NC 27709

Ms. Xan Augerot
Washington Sea Grant
3716 Brooklyn Avenue NE
Seattle, WA 98105

Ms. Nancy Azzam
Windstar Foundation
2300 Noble Avenue, N
Golden Valley, MN 55422

Dr. Norihisa Baba
National Research Institute
of Far Seas Fisheries
Fisheries Agency, the
Government of Japan
5-7-1 Orido, Shimizu, 424
Shizuoka, Japan

Dr. William A. Bailey
GEO-CENTERS, Inc.
10903 Indian Head Highway
Fort Washington, MD 20744

Mr. George H. Balazs
SWFSC Honolulu Laboratory
NMFS, NOAA
2570 Dole Street
Honolulu, HI 96822-2396

Ms. Jane Ball
Sea Grant College Program
1000 Pope Road, MSB 219
Honolulu, HI 96822

Dr. John Barlow
Maine Maritime Academy
Castine, ME 04420

Ms. Peigin Barrett, Director
California Marine Mammal Center
Marine Headlands
Sausalito, CA 94965

Mr. Nelio B. Barros
Rosenstiel School of M&AS
University of Miami, DBLR
4600 Rickenbacker Causeway
Miami, FL 33149-1098

Mr. Michael J. Bean
Environmental Defense Fund
1616 P Street, NW
Washington, D.C. 20036

Dr. Mary Bergen
1807 - 13th Street
Sacramento, CA 95814

Dr. George W. Boehlert
SWFSC Honolulu Laboratory
NMFS, NOAA
2570 Dole Street
Honolulu, HI 96822-2396

Dr. W. R. P. Bourne
3 Contlaw Place
Milltimber
Aberdeen AB1 ODS
Scotland, United Kingdom

Mr. Brian Boyle
Washington Dept. Natural Res.
201 John A. Cherby Bldg.
Mail Stop QW-21
Olympia, WA 98504

Dr. Paul A. Breen
Minist. Agric. & Fisheries
Fisheries Research Centre
Greta Point, P.O. Box 297
Wellington, New Zealand

Ms. Barbara H. Britten
American Cetacean Society, #614
1300 S. Arlington Ridge Road
Arlington, VA 22202

Mr. Ronald G. Bruner
Society of The Plastics
Industry, Inc.
1275 K Street NW, Suite 400
Washington, D.C. 20005

Mr. Alan R. Bunn
Alaska Fisheries Science Center
NMFS, NOAA
7600 Sand Point Way NE, BIN C15700
Seattle, WA 98115-0070

Mr. Richard Buxton
1741 Vernon Street
Halifax, Nova Scotia
B3H 3N3 Canada

Mr. Jeff Cantin
Eastern Research Group, Inc.
6 Whittenmore Street
Arlington, MA 02174

Dr. Bong Hwan Chang, Chairman
Korean Distant Water Squid
Gillnet Fisheries Association
1018-1 Bang Bae-Dong, Rm. 302
Suo Cha-Ku, Seoul, Korea

Mr. Terrill Chang
c/o Camp Dresser & McKee
One Woodward Avenue, Ste. 1500
Detroit, MI 48226

Dr. Douglas G. Chapman
Professor & Dean Emeritus
7106 San Bartolo
Carlsbad, CA 92009

Capt. A. A. Clark
Sause Brothers, Pier 20
46-416 Hulupala
Kaneohe, HI 96744

Mr. David Clark
Battelle Human Affairs
Research Centers
4000 NE - 41st Street
Seattle, WA 98115

Mr. Raymond P. Clarke
SWFSC Honolulu Laboratory
NMFS, NOAA
2570 Dole Street
Honolulu, HI 96822-2396

Mr. James M. Coe - F/AKC
Marine Entanglement Program
Alaska Fisheries Science Center
NMFS, NOAA
7600 Sand Point Way NE, BIN C15700
Seattle, WA 98115-0070

Dr. Salvatore Comitini
Department of Economics and
Agricultural Economics
University of Hawaii at Manoa
Honolulu, HI 96822

Mr. David Cottingham
Interagency Task Force PP2
14th & Constitution Avenue NW
Room 6811
Washington, D.C. 20230

LCDR Carl A. Crampton
U.S. Coast Guard (G-MPS-3)
2100 - Second Street, SW
Washington, D.C. 20593-0001

Dr. John Craven
William S. Richardson
School of Law #208
2515 Dole Street
Honolulu, HI 96822

Dr. Jack R. Davidson, Dir.
Sea Grant College Program
University of Hawaii
1000 Pope Road, MSB 220
Honolulu, HI 96822

Dr. John Davis
Canada Department of
Fisheries and Oceans
P.O. Box 6000
Sidney, V8L 4B2
B.C., Canada

Ms. Patty Debenham
Center for Marine
Conservation
1725 DeSales Street, NW
Washington, D.C. 20036

Dr. Robert DeLong F/AKC
National Marine Mammal Laboratory
Alaska Fisheries Science Center
NMFS, NOAA
7600 Sand Point Way, NE, BIN C15700
Seattle, WA 98115-0070

Dr. Trevor R. Dixon
The Tidy Britain Group
The Pier, Wallgate
Wigan, Lancashire WN3 4EX
United Kingdom

Mr. Zoltan J. Dorko
Plastics Division
Polysar, Inc.
P.O. Box 90
Leominster, MA 01453

Ms. Uvetta B. Dozier
103 G Street, SW, Suite B-617
Washington, D.C. 20024

Ms. Angela D. Farias
1700 N. Congress
Austin, TX 78701

Mr. Stewart I. Fefer
Hawaiian/Pacific Islands
NWR Complex
P.O. Box 50167
Honolulu, HI 96850

Mr. R. Barry Fisher
Yankee Fisheries
1626 Coast Highway
Newport, OR 97365

Dr. Charles W. Fowler F/AKC
National Marine Mammal Laboratory
Alaska Fisheries Science Center
NMFS, NOAA
7600 Sand Point Way, NE, BIN C15700
Seattle, WA 98115-0070

Dr. Deborah P. French
Applied Science Assoc., Inc.
70 Dean Knauss Drive
Narragansett, RI 02882-1143

Ms. Sarah Friday
UNC Sea Grant Communications
North Carolina State University
P.O. Box 8605
Raleigh, NC 27695-8605

Dr. Tim Gerrodette
Southwest Fisheries Science Center
NMFS, NOAA
P.O. Box 271
La Jolla, CA 92038

Ms. Mary L. Gessner
National Marine Fisheries
Service, NOAA
1335 East-West Highway
Silver Spring, MD 20910

Dr. Abraham Golik
Israel O & LR, Ltd.
and National Institute
of Oceanography
Tel Shikmona, P.O. Box 8030
Haifa 31080, Israel

Mr. William Gordon
New Jersey Marine Sciences
Consortium, Bldg. 22
Sandy Hook Field Station
Fort Hancock, NJ 07732

Dr. Murray R. Gregory
Department of Geology
The University of Auckland
Private Bag
Auckland, New Zealand

Dr. Thomas A. Grigalunas
Department of Resource Economics
University of Rhode Island
Kingston, RI 02881

Ms. Mary Griggs
1807 - 13th Street
Sacramento, CA 95814

Dr. Bernard Griswold
 Program Director, MAS
 National Sea Grant College
 1001 N. Rochester
 Arlington, VA 22205

Mr. Michael R. Hagler
 Greenpeace, New Zealand
 Private Bag
 Wellesley Street
 Auckland, New Zealand

Ms. Kelly Hamilton
 2284 Redlands Drive
 Newport Beach, CA 92660

Ms. Judie A. Hansen (Ms. Neilson)
 1502 S. County Road 525W
 Sanville, IN 46122

Mr. Robert Harman
 Western Pacific Regional
 Fishery Management Council
 1164 Bishop Street, Ste. 1405
 Honolulu, HI 96813

Mr. John R. Henderson
 SWFSC Honolulu Laboratory
 NMFS, NOAA
 2570 Dole Street
 Honolulu, HI 96822-2396

Mr. Burr Heneman
 Marine Mammal Commission
 35 Horseshoe Hill Road
 Star Route
 Bolinas, CA 94924

Rep. Adelheid Herrmann
 Box 63
 Naknek, AK 99633

Mr. Glenn Higashi
 Division of Aquatic Resources
 Department of Land and
 Natural Resources
 1151 Punchbowl Street
 Honolulu, HI 96813

Dr. Kazuhiko Hiramatsu
 National Research Institute
 of Far Seas Fisheries
 Fisheries Agency, the
 Government of Japan
 5-7-1 Orido, Shimizu-shi, 424
 Shizuoka, Japan

Mr. Dewayne Hollin
 Sea Grant College Program
 Texas A&M University
 College Station, TX 77843-4115

Mr. Randy Honebrink
 Education Coordinator
 Division of Aquatic Resources
 1151 Punchbowl Street, Rm. 330
 Honolulu, HI 96813

Dr. Donald E. Hoss
 Beaufort Laboratory
 Southeast Fisheries Science Center
 NMFS, NOAA
 Beaufort, NC 28516-9722

Mr. Paul Ideker &
 Associates and Company
 1134 La Mirada
 Laguna Beach, CA 92651

Ms. Carole Jaworski
 University of Rhode Island
 Bay Campus
 R.I. Sea Grant Program
 Narragansett, RI 02882-1197

Dr. Scott W. Johnson
 Alaska Fisheries Science Center
 Auke Bay Laboratory
 NMFS, NOAA
 P.O. Box 210155
 Auke Bay, AK 99821

Ms. Colleen Johiro
 Makai Employee - Reporter
 Sea Grant Program, MSB
 University of Hawaii
 Honolulu, HI 96822

Mr. Jeffrey June
Natural Resources Consultants, Inc.
4055 - 21st Avenue West
Seattle, WA 98199

Mr. Manu Kahaialii
Trustee, Office of
Hawaiian Affairs, Maui
217 Akeke Place
Lahaina, HI 96761

Mr. Kenneth S. Kamlet
A. T. Kearney, Inc.
P.O. Box 1405
Alexandria, VA 22313

Mr. Brian Kanenaka
Division of Aquatic Resources
Department of Land and
Natural Resources
1151 Punchbowl Street
Honolulu, HI 96813

Ms. Jill Kauffman
312 Sutter Street, Rm. 316
San Francisco, CA 94108

LCDR Kenneth W. Keane
Chief, Port Operations
433 Ala Moana Boulevard
Honolulu, HI 96813

Mr. Timothy Keeney
Office of General Counsel, Ste. 5814
Department of Commerce, NOAA
15th and Constitution Ave., NW
Washington, D.C. 10130

Mr. Paul G. Kelleher
New Jersey Polymer
Extension Center
Stevens Institute - PPI
Hoboken, NJ 07030

Ms. K. Kendrick-Vanderpool
350 Ward Avenue, Ste. 106
Honolulu, HI 96814

Ms. Leslie Keyt
173 McAulay Place
Laguna Beach, CA 92651

Mr. Han-Mo Kim
Alaska Office Division
Korean Deep Sea
Fishery Association
13439 Karen Street
Anchorage, AK 99515

CDR B. K. Klimek
14th Coast Guard District
Federal Building, Rm. 9149
300 Ala Moana Boulevard
Honolulu, HI 96850

Mr. Larry Koss
Office of Chief of Naval
Operations
Department of the Navy
Washington, D.C. 20350

Mr. August Kotlewski
673 Beach Road, Ste. 3
Sarasota FL 34242

Ms. JoAnn N. Kushima
Division of Aquatic Resources
Department of Land and
Natural Resources
1151 Punchbowl Street
Honolulu, HI 96813

Mr. David Laist
Marine Mammal Commission
1625 Eye Street, NW
Washington, D.C. 20006

Dr. Shirley Laska
Department of Sociology
University of New Orleans
Lakefront, New Orleans, LA 70148

Mrs. Sidney Lasseigne
Pacific States Marine
Fisheries Commission
P.O. Box 221
Depoe Bay, OR 97341

Ms. Marilyn Lee
Sea Life Park
Makapuu Point
Waimanalo, HI 96795

Ms. Elaine F. Leslie
 Marine Mammal Specialist
 Walt Disney World Co.
 P.O. Box 10,000, Lake
 Buena Vista, FL 32830-1000

Ms. Nancy B. Linsley
 46-152 Nahiku Place
 Kaneohe, HI 96744

Dr. Peter Lutz
 Rosenstiel School of M&AS
 University of Miami
 4600 Rickenbacker Causeway
 Miami, FL 33149-1098

Dr. Albert M. Manville II
 Defenders of Wildlife
 1244 - 19th Street, NW
 Washington, D.C. 20036

CDR Martin R. Marks, H.M.S.
 British Navy Staff
 3100 Massachusetts Ave., NW
 Washington, D.C. 20008

Ms. Lissa A. Martinez
 7107 Cedar Avenue
 Takoma Park, MD 20912

CDR M. W. Mastenbrook
 14th Coast Guard District
 Federal Building
 300 Ala Moana Boulevard
 Honolulu, HI 96850-4982

Dr. Satsuki Matsumura
 National Research Institute
 of Far Seas Fisheries
 Fisheries Agency, the
 Government of Japan
 5-7-1 Orido, Shimizu-shi
 Shizuoka, 424 Japan

Dr. Kenneth E. McConnell
 Department of Agriculture
 and Resources
 University of Maryland
 College Park, MD 20742

Ms. Tracie McKenzie
 NOAA Fisheries
 Northeast Region
 Narragansett, RI 02882

Mr. Norman Meade, N/OMA31
 Ocean Assessment Division
 Office of Oceanography
 and Marine Association
 National Ocean Service, NOAA
 Rockville, MD 20852

Ms. Deborah Mercy
 University of Alaska
 Marine Advisory Program
 2221 E. Northern Lights, Ste. 220
 Anchorage, AK 99517

Ms. Suzanne Montgomery
 Washington Communications
 150 N. Muhlenberg Street
 Woodstock, VA 22664

Dr. Julio M. Morell
 Department of Marine Science
 University of Puerto Rico-Mayaguez
 Mayaguez, Puerto Rico 00709-5000

Mr. Minoru Morimoto
 Fisheries Agency, Government of Japan
 Ministry of Agriculture Forest
 and Fisheries
 2-1, 1-chome, Kasumigaseki
 Chiyoda-ku, Tokyo, 100 Japan

Mr. Joji Morishita
 International Affairs Division
 Fisheries Agency, the
 Government of Japan
 2-1, 1-chome, Kasumigaseki
 Chiyoda-ku, Tokyo, 100 Japan

Mr. Clyde Morita
 State Litter Control
 205 Koula Sreet
 Honolulu, HI 96813

Mr. Kenneth S. Mulholland
 7 Ashleigh Crescent, Miramar
 Wellington 6003, New Zealand

Ms. Patti Mullin
 Washington Sea Grant Program
 North Sound Advisory Office
 19 Harbor Mall
 Bellingham, WA 98225

Ms. Glynnis L. Nakai
 1561 Wilhelmina Rise
 Honolulu, HI 96816

Dr. Masayuki Nakajima
Izu-Mito Sea Paradise
3-1 Nagahama
Uchiura, Numazu
Shizuoka, 410-02 Japan

Mr. John J. Naughton
Pacific Area Office
Southwest Region, NMFS, NOAA
2570 Dole Street
Honolulu, HI 96822-2396

Mr. John Nightingale
Maui Ocean Center
101 N. Kihei Road
Kihei, Maui, HI 96753

Dr. Haruo Ogi
Research Institute of the
North Pacific Fisheries
Hakodate, Hokkaido, 141 Japan

Ms. Kathryn J. O'Hara
Marine Debris and
Entanglement Program
Center for Marine Conservation
1725 DeSales Street, NW
Washington, D.C. 20036

Mr. David E. Ortman
Friends of the Earth
Northwest Office
4512 University Way, NE
Seattle, WA 98105

Ms. Cathy L. P. Palmer
SW Florida Coastal Research Center
Mote Marine Laboratory
1600 City Island Park
Sarasota, FL 34236

Dr. Rod J. Paterson, Director
Oceanographic Sciences
Department of Fisheries and Oceans
200 Kent Street
Ottawa, Ontario K1A 0E6, Canada

Mr. William W. Paty, Chairman
Board of Land and Natural
Resources
1151 Punchbowl Street
Honolulu, HI 96817

Ms. Linda Paul
815 Pahumele Place
Kailua, HI 96734

Dr. Gerard Peet
Heemraadssingel 193
023 CB Rotterdam
The Netherlands

Mr. Hank Pennington
University of Alaska
Sea Grant
202 Center Street
Kodiak, AK 99615

Mr. Bruce Perlson (TSV)
11500 Northlake Drive
Cincinnati, OH 45249

Dr. Rose T. Pfund
1077 Kalikimaka Street
Honolulu, HI 96817

Ms. Pamela Plotkin
Department of Biology
Texas A&M University
College Station, TX 77843-3258

Dr. Richard H. Podolsky
Director, Research and
Academics Island Institute
60 Ocean Street
Rockland, ME 04841

Ms. Fran Recht
Pacific States Marine
Fisheries Commission
P.O. Box 221
Depoe Bay, OR 97341

Dr. David P. Redford
3660 Bay Drive
Edgewater, MD 21037

Mr. Alan Reichman
Greenpeace USA
1204 N. 42nd Street
Seattle, WA 98103-7504

Dr. Christine A. Ribic
Environmental Research Laboratory
U.S. Environmental Protection Agency
200 SW 35th Street
Corvallis, OR 97333

Mr. Bob Rose
 Special Assistant to
 Commissioner of Public Lands
 Washington Department of
 Natural Resources, QW-21
 201 John A. Cherberg Bldg.
 Olympia, WA 98504

Mr. C. W. Ross
 Rathrippen Farm
 Castle Frederick Road
 RR #2, Falmouth
 Nova Scotia BOP 1L0, Canada

Mr. Douglas Rost
 344 Slater Street
 Ontario K1A 0N7, Canada

Mr. Peter Ryan
 Percy FitzPatrick Institute
 of African Ornithology
 University of Cape Town
 Rondebosch 7700, South Africa

Capt. Edward Rymarz
 Marine Environment Division
 International Maritime Organization
 4 Albert Embankment
 London SE1 7SR, United Kingdom

Dr. Samuel S. Sadove
 Okeanos Ocean Research
 Foundation, Inc.
 P.O. Box 776
 Hampton Bays, NY 11946

Mr. Ralph S. Saito
 Division of Forestry
 and Wildlife
 Department of Land and
 Natural Resources
 1151 Punchbowl Street
 Honolulu, HI 96813

Dr. Akinori Satou
 Oceanography Office
 N.S. Environmental Science
 Consultant
 2-9-14, Toranomom, Minato-ku
 Tokyo, 105 Japan

Mr. Hal Schlomann
 P.O. Box 129
 Clinton, WA 98236

Mr. Douglas G. Schneider
 Alaska Sea Grant College Program
 University of Alaska
 138 Irving II
 Fairbanks, AK 99775-5040

Ms. Barbara Schroeder
 Florida Department of
 Natural Resources
 P.O. Box 1319
 Stuart, FL 34995

Dr. Steven Schwartz
 Center for Environmental
 Education
 1725 DeSales St., NW
 Washington, D.C. 20036

Dr. Gerald Scott
 Greenridge
 Newby-in-Cleveland
 near Stokesley
 North Yorkshire TS8 0AH
 United Kingdom

LCDR Raymond Seebald
 Commandant (G-MPS-1)
 U.S. Coast Guard Headquarters
 2100 Second Street, SW
 Washington, D.C. 20593-0001

Mr. Richard S. Shomura
 School of Ocean and Earth
 Science and Technology
 University of Hawaii, HIG 131
 Honolulu, HI 96822

Dr. Paul R. Sievert
 Department of Biology
 University of Pennsylvania
 Philadelphia, PA 19104-6018

Dr. Louis Sileo
 National Wildlife Health
 Research Center
 6006 Schroeder Road
 Madison, WI 53711

Ms. Kitty Simonds
 Western Pacific Regional
 Fishery Management Council
 1164 Bishop St., Ste. 1405
 Honolulu, HI 96813

Mr. Richard Sixberry
 Division of Aquatic Resources
 Department of Land and
 Natural Resources
 1151 Punchbowl Street
 Honolulu, HI 96813

Dr. Margarete Steinhauer
 Battelle Ocean Sciences
 397 Washington Street
 Duxbury, MA 02332

Dr. Brent S. Stewart
 Sea World Research Institute
 Hubbs Marine Research Center
 1700 South Shores Road
 San Diego, CA 92109

Mr. Robert Sulnick
 Executive Director
 American Ocean Campaign
 1427 Seventh Street
 Santa Monica, CA 90401

Dr. John Sutinen
 University of Rhode Island
 Department of Resource Economics
 Kingston, RI 02881

Dr. R. Lawrence Swanson
 Waste Management Institute
 Marine Sciences Research Center
 State University of New York
 Stony Brook, NY 11794-5000

Dr. Gordon L. Swartzman
 University of Washington
 7603 - 56th Place, NE
 Seattle, WA 98115 or 98195

Dr. Satoru Taguchi
 Department of Oceanography
 University of Hawaii, MSG 307
 1000 Pope Road
 Honolulu, HI 96822

Mr. Yasuji Tamaki
 Fisheries Agency, the
 Government of Japan
 2-1, 1-chome, Kasumigaseki
 Chiyoda-ku, Tokyo, 100 Japan

Capt. A. E. Tanos
 14th Coast Guard District
 Federal Building, Rm. 9149
 300 Ala Moana Boulevard
 Honolulu, HI 96850-4982

Dr. Raymond Tarpley
 The Texas Veterinary Medical Center
 College of Veterinary Medicine
 Texas A&M University
 College Station, TX 77843-4458

Mr. Rupert A. Teves
 1982 Hoomalolo Street
 Pearl City, HI 96782

Dr. Michael F. Tillman
 National Marine Fisheries
 Service, NOAA
 1335 East-West Highway
 Silver Spring, MD 20910

Mr. Larry Tyler
 948 NW Coast Street
 Newport, OR 97365

Dr. Kathryn D. Wagner
 Department of Political Science
 Swarthmore College
 500 College Avenue
 Swarthmore, PA 19081-1397

Ms. Barbara Wallace
 Kearney/Centaur Division
 A. T. Kearney, Inc.
 225 Reinekers Lane, Ste. 300
 Alexandria, VA 22314

Dr. Yoh Watanabe
 National Research Institute
 of Far Seas Fisheries
 Fisheries Agency, the
 Government of Japan
 5-7-1 Orido, Shimizu-shi
 Shizuoka, 424 Japan

Ms. Dottie Wendt
 389 Kailua Road, #110
 Kailua, HI 96734

Ms. Christine E. Williams
Wellington City Council
Work Department
P.O. Box 2199
Wellington, New Zealand

Ms. Christine Woolaway
Sea Grant College Program
University of Hawaii
1000 Pope Road, MSB 220
Honolulu, HI 96822

Mr. Nobuyuki Yagi
Division of Fishing Ground
Environment Conservation
Fisheries Agency, the
Government of Japan
Chiyoda-ku, Tokyo, 100 Japan

Mr. John Yamauchi
State Litter Control Office
205 Koula Street
Honolulu, HI 96813

Ms. Annette Young
Division of Aquatic Resources
Department of Land
and Natural Resources
1151 Punchbowl Street
Honolulu, HI 96813

Mr. Kendall Zekahi
98-1219 Kaamilo Street
Aiea, HI 96701

RECENT TECHNICAL MEMORANDUMS

Copies of this and other NOAA Technical Memorandums are available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22167. Paper copies vary in price. Microfiche copies cost \$4.50. Recent issues of NOAA Technical Memorandums from the NMFS Southwest Fisheries Center are listed below:

- NOAA-TM-NMFS-SWFSC-144 Atlas of eastern tropical Pacific oceanographic variability and cetacean sightings, 1986-1989.
P.C. FIELDER, L.J. LIERHEIMER, S.B. REILLY, S.N. SEXTON, R.S. HOLT
and D.P. DEMASTER
(July 1990)
- 145 Trends in landings, species composition, length-frequency distributions, and sex ratios of 11 rockfish species (Genus *Sebastes*) from central and northern California ports (1978-88).
D.E. PEARSON and S. RALSTON
(July 1990)
- 146 Field manual for phocid necropsies (specifically *Monachus schauinslandi*).
J. M. WINCHELL
(July 1990)
- 147 Survey of the abundance and distribution of pelagic young-of-the-year rockfishes, *Sebastes*, off central California.
T. W. ECHEVERRIA, W. H. LENARZ and C. A. REILLY
(September 1990)
- 148 United States Agency for International Development and National Marine Fisheries Service workshop on tropical fish stock assessment, 5-26 July 1989, Honolulu, Hawaii.
J. J. POLOVINA and R. S. SHOMURA
(September 1990)
- 149 Summary of the 1988 U.S. tuna/porpoise observer data.
A. R. JACKSON
(September 1990)
- 150 Population monitoring of the Hawaiian monk seal, *Monachus schauinslandi*, and captive maintenance project at Kure Atoll, 1988.
J. R. HENDERSON and M. R. FINNEGAN
(September 1990)
- 151 The Hawaiian monk seal on Laysan Island, 1988.
T. C. JOHANOS, B. L. BECKER, M. A. BROWN,
B. K. CHOY, L. M. HURUKI, R. E. BRAINARD and
R. L. WESTLAKE
(September 1990)
- 152 A personal computer based system for analog-to-digital and serial communication data acquisition.
R. C. HOLLAND
(November 1990)
- 153 The nearshore physical oceanography off the central California coast during May-June, 1989: A summary of CTD from juvenile rockfish surveys.
F. B. SCHWING, S. RALSTON, D. M. HUSBY
and W. H. LENARZ
(December 1990)

